

**$\chi_{b2}(2P)$** 
 $I^G(J^{PC}) = 0^+(2^{++})$   
*J needs confirmation.*

Observed in radiative decay of the  $\Upsilon(3S)$ , therefore  $C = +$ . Branching ratio requires E1 transition, M1 is strongly disfavored, therefore  $P = +$ .

 **$\chi_{b2}(2P)$  MASS**VALUE (MeV)DOCUMENT ID
**102686.5±0.22±0.50 OUR EVALUATION** From  $\gamma$  energy below, using  $\Upsilon(3S)$  mass = 10355.2  $\pm$  0.5 MeV

$m_{\chi_{b2}(2P)} - m_{\chi_{b1}(2P)}$

VALUE (MeV)DOCUMENT IDTECNCOMMENT**13.5±0.4±0.5**

1 HEINTZ

92

CSB2

 $e^+ e^- \rightarrow \gamma X, \ell^+ \ell^- \gamma \gamma$ 
<sup>1</sup> From the average photon energy for inclusive and exclusive events. Supersedes NARAIN 91.
 **$\gamma$  ENERGY IN  $\Upsilon(3S)$  DECAY**VALUE (MeV)EVTSDOCUMENT IDTECNCOMMENT**86.19±0.22 OUR EVALUATION**

Treating systematic errors as correlated

**86.40±0.18 OUR AVERAGE**86.04  $\pm$  0.06  $\pm$  0.27

ARTUSO

05

CLEO

 $\Upsilon(3S) \rightarrow \gamma X$ 86  $\pm$  1

101

CRAWFORD

92B

CLE2

 $e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$ 86.7  $\pm$  0.4

10319

2 HEINTZ

92

CSB2

 $e^+ e^- \rightarrow \gamma X$ 86.9  $\pm$  0.4

157

3 HEINTZ

92

CSB2

 $e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$ 86.4  $\pm$  0.1  $\pm$  0.4

30741

MORRISON

91

CLE2

 $e^+ e^- \rightarrow \gamma X$ 
<sup>2</sup> A systematic uncertainty on the energy scale of 0.9% not included. Supersedes NARAIN 91.

<sup>3</sup> A systematic uncertainty on the energy scale of 0.9% not included. Supersedes HEINTZ 91.
 **$\chi_{b2}(2P)$  DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level
$\Gamma_1 \omega \Upsilon(1S)$	( 1.10 $^{+0.34}_{-0.30}$ ) %	
$\Gamma_2 \gamma \Upsilon(2S)$	( 10.6 $\pm$ 2.6 ) %	S=2.0
$\Gamma_3 \gamma \Upsilon(1S)$	( 7.0 $\pm$ 0.7 ) %	
$\Gamma_4 \pi \pi \chi_{b2}(1P)$	( 5.1 $\pm$ 0.9 ) $\times 10^{-3}$	
$\Gamma_5 D^0 X$	< 2.4 %	CL=90%
$\Gamma_6 \pi^+ \pi^- K^+ K^- \pi^0$	< 1.1 $\times 10^{-4}$	CL=90%
$\Gamma_7 2\pi^+ \pi^- K^- K_S^0$	< 9 $\times 10^{-5}$	CL=90%
$\Gamma_8 2\pi^+ \pi^- K^- K_S^0 2\pi^0$	< 7 $\times 10^{-4}$	CL=90%

$\Gamma_9$	$2\pi^+ 2\pi^- 2\pi^0$	$(3.9 \pm 1.6) \times 10^{-4}$		
$\Gamma_{10}$	$2\pi^+ 2\pi^- K^+ K^-$	$(9 \pm 4) \times 10^{-5}$		
$\Gamma_{11}$	$2\pi^+ 2\pi^- K^+ K^- \pi^0$	$(2.4 \pm 1.1) \times 10^{-4}$		
$\Gamma_{12}$	$2\pi^+ 2\pi^- K^+ K^- 2\pi^0$	$(4.7 \pm 2.3) \times 10^{-4}$		
$\Gamma_{13}$	$3\pi^+ 2\pi^- K^- K_S^0 \pi^0$	$< 4 \times 10^{-4}$	CL=90%	
$\Gamma_{14}$	$3\pi^+ 3\pi^-$	$(9 \pm 4) \times 10^{-5}$		
$\Gamma_{15}$	$3\pi^+ 3\pi^- 2\pi^0$	$(1.2 \pm 0.4) \times 10^{-3}$		
$\Gamma_{16}$	$3\pi^+ 3\pi^- K^+ K^-$	$(1.4 \pm 0.7) \times 10^{-4}$		
$\Gamma_{17}$	$3\pi^+ 3\pi^- K^+ K^- \pi^0$	$(4.2 \pm 1.7) \times 10^{-4}$		
$\Gamma_{18}$	$4\pi^+ 4\pi^-$	$(9 \pm 5) \times 10^{-5}$		
$\Gamma_{19}$	$4\pi^+ 4\pi^- 2\pi^0$	$(1.3 \pm 0.5) \times 10^{-3}$		

## $\chi_{b2}(2P)$ BRANCHING RATIOS

$\Gamma(\omega \Upsilon(1S))/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma$
$\text{VALUE (units } 10^{-2}\text{)}$	$\text{EVTS}$ $\text{DOCUMENT ID}$ $\text{TECN}$ $\text{COMMENT}$

**1.10**<sup>+0.32+0.11</sup><sub>-0.28-0.10</sub>    20.1<sup>+5.8</sup><sub>-5.1</sub> <sup>4</sup> CRONIN-HEN..04    CLE3     $\Upsilon(3S) \rightarrow \gamma \omega \Upsilon(1S)$

<sup>4</sup> Using  $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (11.4 \pm 0.8)\%$  and  $B(\Upsilon(1S) \rightarrow \ell^+ \ell^-) = 2 B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = 2 (2.48 \pm 0.06)\%$ .


$\Gamma(\gamma \Upsilon(2S))/\Gamma_{\text{total}}$	$\Gamma_2/\Gamma$
$\text{VALUE}$	$\text{EVTS}$ $\text{DOCUMENT ID}$ $\text{TECN}$ $\text{COMMENT}$

**0.106±0.026 OUR AVERAGE**    Error includes scale factor of 2.0. See the ideogram below.  
 0.084±0.011±0.010    2.5k    <sup>5</sup> LEES    11J    BABR     $\Upsilon(3S) \rightarrow X \gamma$   
 0.135±0.025±0.035          <sup>6</sup> CRAWFORD    92B    CLE2     $e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$   
 0.173±0.021±0.019          <sup>7</sup> HEINTZ    92    CSB2     $e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$

<sup>5</sup> LEES 11J reports  $[\Gamma(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] = (1.1 \pm 0.1 \pm 0.1) \times 10^{-2}$  which we divide by our best value  $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>6</sup> Using  $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.37 \pm 0.26)\%$ ,  $B(\Upsilon(3S) \rightarrow \gamma \gamma \Upsilon(2S)) \times 2 B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (4.98 \pm 0.94 \pm 0.62) \times 10^{-4}$ , and  $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = 0.135 \pm 0.003 \pm 0.017$ .

<sup>7</sup> Using  $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.44 \pm 0.10)\%$ ,  $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (11.1 \pm 0.5 \pm 0.4)\%$  and assuming  $e\mu$  universality. Supersedes HEINTZ 91.



### $\Gamma(\gamma \gamma(1S))/\Gamma_{\text{total}}$

### $\Gamma_3/\Gamma$

VALUE	EVTS
<b>0.070±0.007 OUR AVERAGE</b>	
0.070±0.004±0.008	11k
0.072±0.014±0.013	
0.070±0.010±0.006	

DOCUMENT ID	TECN	COMMENT
8 LEES	11J BABR	$\gamma(3S) \rightarrow X\gamma$
9 CRAWFORD	92B CLE2	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma\gamma$
10 HEINTZ	92 CSB2	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma\gamma$

<sup>8</sup> LEES 11J reports  $[\Gamma(\chi_{b2}(2P) \rightarrow \gamma \gamma(1S))/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma \chi_{b2}(2P))] = (9.2 \pm 0.3 \pm 0.4) \times 10^{-3}$  which we divide by our best value  $B(\gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>9</sup> Using  $B(\gamma(1S) \rightarrow \mu^+ \mu^-) = (2.57 \pm 0.07)\%$ ,  $B(\gamma(3S) \rightarrow \gamma\gamma \gamma(2S)) \times 2 B(\gamma(1S) \rightarrow \mu^+ \mu^-) = (5.03 \pm 0.94 \pm 0.63) \times 10^{-4}$ , and  $B(\gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = 0.135 \pm 0.003 \pm 0.017$ .

<sup>10</sup> Using  $B(\gamma(1S) \rightarrow \mu^+ \mu^-) = (2.57 \pm 0.07)\%$ ,  $B(\gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = (11.1 \pm 0.5 \pm 0.4)\%$  and assuming  $e\mu$  universality. Supersedes HEINTZ 91.

### $\Gamma(\pi\pi\chi_{b2}(1P))/\Gamma_{\text{total}}$

### $\Gamma_4/\Gamma$

VALUE (units $10^{-3}$ )	EVTS
<b>5.1±0.9 OUR AVERAGE</b>	
4.9±0.7±0.6	17k
6.0±1.6±1.4	

DOCUMENT ID	TECN	COMMENT
11 LEES	11C BABR	$e^+ e^- \rightarrow \pi^+ \pi^- X$
12 CAWLFIELD	06 CLE3	$\gamma(3S) \rightarrow 2(\gamma\pi\ell)$

<sup>11</sup>  $(0.64 \pm 0.05 \pm 0.08) \times 10^{-3}$ . We derive the value assuming  $B(\gamma(3S) \rightarrow \chi_{b2}(2P)X) = B(\gamma(3S) \rightarrow \chi_{b2}(2P)\gamma) = (13.1 \pm 1.6) \times 10^{-2}$ .

<sup>12</sup> CAWLFIELD 06 quote  $\Gamma(\chi_b(2P) \rightarrow \pi\pi\chi_b(1P)) = 0.83 \pm 0.22 \pm 0.08 \pm 0.19$  keV assuming I-spin conservation, no D-wave contribution,  $\Gamma(\chi_{b1}(2P)) = 96 \pm 16$  keV, and  $\Gamma(\chi_{b2}(2P)) = 138 \pm 19$  keV.

$\Gamma(D^0 X)/\Gamma_{\text{total}}$	$\Gamma_5/\Gamma$			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.4 \times 10^{-2}$	90	13,14 BRIERE	08	CLEO $\gamma(3S) \rightarrow \gamma D^0 X$

<sup>13</sup> For  $p_{D^0} > 2.5$  GeV/c.<sup>14</sup> The authors also present their result as  $(0.2 \pm 1.4 \pm 0.1) \times 10^{-2}$ .

$\Gamma(\pi^+ \pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$	$\Gamma_6/\Gamma$			
VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<1.1$	90	15 ASNER	08A	CLEO $\gamma(3S) \rightarrow \gamma \pi^+ \pi^- K^+ K^- \pi^0$

<sup>15</sup> ASNER 08A reports  $[\Gamma(\chi_{b2}(2P) \rightarrow \pi^+ \pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma \chi_{b2}(2P))] < 14 \times 10^{-6}$  which we divide by our best value  $B(\gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = 13.1 \times 10^{-2}$ .

$\Gamma(2\pi^+ \pi^- K^- K_S^0)/\Gamma_{\text{total}}$	$\Gamma_7/\Gamma$			
VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<0.9$	90	16 ASNER	08A	CLEO $\gamma(3S) \rightarrow \gamma 2\pi^+ \pi^- K^- K_S^0$

<sup>16</sup> ASNER 08A reports  $[\Gamma(\chi_{b2}(2P) \rightarrow 2\pi^+ \pi^- K^- K_S^0)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma \chi_{b2}(2P))] < 12 \times 10^{-6}$  which we divide by our best value  $B(\gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = 13.1 \times 10^{-2}$ .

$\Gamma(2\pi^+ \pi^- K^- K_S^0 2\pi^0)/\Gamma_{\text{total}}$	$\Gamma_8/\Gamma$			
VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<7$	90	17 ASNER	08A	CLEO $\gamma(3S) \rightarrow \gamma 2\pi^+ \pi^- K^- 2\pi^0$

<sup>17</sup> ASNER 08A reports  $[\Gamma(\chi_{b2}(2P) \rightarrow 2\pi^+ \pi^- K^- K_S^0 2\pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma \chi_{b2}(2P))] < 87 \times 10^{-6}$  which we divide by our best value  $B(\gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = 13.1 \times 10^{-2}$ .

$\Gamma(2\pi^+ 2\pi^- 2\pi^0)/\Gamma_{\text{total}}$	$\Gamma_9/\Gamma$			
VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$3.9 \pm 1.6 \pm 0.5$	23	18 ASNER	08A	CLEO $\gamma(3S) \rightarrow \gamma 2\pi^+ 2\pi^- 2\pi^0$

<sup>18</sup> ASNER 08A reports  $[\Gamma(\chi_{b2}(2P) \rightarrow 2\pi^+ 2\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma \chi_{b2}(2P))] = (51 \pm 16 \pm 13) \times 10^{-6}$  which we divide by our best value  $B(\gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}$	$\Gamma_{10}/\Gamma$			
VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$0.9 \pm 0.4 \pm 0.1$	11	19 ASNER	08A	CLEO $\gamma(3S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^-$

<sup>19</sup> ASNER 08A reports  $[\Gamma(\chi_{b2}(2P) \rightarrow 2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma \chi_{b2}(2P))] = (12 \pm 4 \pm 3) \times 10^{-6}$  which we divide by our best value  $B(\gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2\pi^+ 2\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{11}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.4±1.0±0.3</b>	16	20 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^- \pi^0$
20 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 2\pi^+ 2\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P))] = (32 \pm 11 \pm 8) \times 10^{-6}$ which we divide by our best value $B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

$\Gamma(2\pi^+ 2\pi^- K^+ K^- 2\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{12}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.7±2.2±0.6</b>	14	21 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^- 2\pi^0$
21 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 2\pi^+ 2\pi^- K^+ K^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P))] = (62 \pm 23 \pm 17) \times 10^{-6}$ which we divide by our best value $B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

$\Gamma(3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{13}/\Gamma$

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;4</b>	90	22 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 3\pi^+ 2\pi^- K^- K_S^0 \pi^0$
22 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P))] < 58 \times 10^{-6}$ which we divide by our best value $B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = 13.1 \times 10^{-2}$ .				

$\Gamma(3\pi^+ 3\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{14}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.9±0.4±0.1</b>	14	23 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 3\pi^+ 3\pi^-$
23 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 3\pi^+ 3\pi^-)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P))] = (12 \pm 4 \pm 3) \times 10^{-6}$ which we divide by our best value $B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

$\Gamma(3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{15}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>12±4±1</b>	45	24 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 3\pi^+ 3\pi^- 2\pi^0$
24 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P))] = (159 \pm 33 \pm 43) \times 10^{-6}$ which we divide by our best value $B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

$\Gamma(3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}$   $\Gamma_{16}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.4±0.7±0.2</b>	12	25 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^-$
25 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P))] = (19 \pm 7 \pm 5) \times 10^{-6}$ which we divide by our best value $B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

**$\Gamma(3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$**   **$\Gamma_{17}/\Gamma$**

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>4.2 \pm 1.7 \pm 0.5</math></b>	16	26 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^- \pi^0$

26 ASNER 08A reports  $[\Gamma(\chi_{b2}(2P) \rightarrow 3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P))] = (55 \pm 16 \pm 15) \times 10^{-6}$  which we divide by our best value  $B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

**$\Gamma(4\pi^+ 4\pi^-)/\Gamma_{\text{total}}$**   **$\Gamma_{18}/\Gamma$**

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.9 \pm 0.4 \pm 0.1</math></b>	9	27 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 4\pi^+ 4\pi^-$

27 ASNER 08A reports  $[\Gamma(\chi_{b2}(2P) \rightarrow 4\pi^+ 4\pi^-)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P))] = (12 \pm 5 \pm 3) \times 10^{-6}$  which we divide by our best value  $B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

**$\Gamma(4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}$**   **$\Gamma_{19}/\Gamma$**

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>13 \pm 5 \pm 2</math></b>	27	28 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 4\pi^+ 4\pi^- 2\pi^0$

28 ASNER 08A reports  $[\Gamma(\chi_{b2}(2P) \rightarrow 4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P))] = (165 \pm 46 \pm 50) \times 10^{-6}$  which we divide by our best value  $B(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

**$\chi_{b2}(2P)$  Cross-Particle Branching Ratios**

$$\Gamma(\chi_{b2}(2P) \rightarrow \gamma \Gamma(1S))/\Gamma_{\text{total}} \times \Gamma(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P))/\Gamma_{\text{total}} \\ \Gamma_3/\Gamma \times \Gamma_{19}^{\Gamma(3S)}/\Gamma^{\Gamma(3S)}$$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>9.2 \pm 0.3 \pm 0.4</math></b>	11k	LEES	11J BABR	$\Gamma(3S) \rightarrow X\gamma$

$$\Gamma(\chi_{b2}(2P) \rightarrow \gamma \Gamma(2S))/\Gamma_{\text{total}} \times \Gamma(\Gamma(3S) \rightarrow \gamma \chi_{b2}(2P))/\Gamma_{\text{total}} \\ \Gamma_2/\Gamma \times \Gamma_{19}^{\Gamma(3S)}/\Gamma^{\Gamma(3S)}$$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>1.1 \pm 0.1 \pm 0.1</math></b>	2.5k	LEES	11J BABR	$\Gamma(3S) \rightarrow X\gamma$

$$B(\chi_{b2}(2P) \rightarrow \chi_{b2}(1P)\pi^+\pi^-) \times B(\Gamma(3S) \rightarrow \chi_{b2}(2P)X)$$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.64 \pm 0.05 \pm 0.08</math></b>	17k	LEES	11C BABR	$e^+ e^- \rightarrow \pi^+ \pi^- X$

## $\chi_{b2}(2P)$ REFERENCES

LEES	11C	PR D84 011104	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ASNER	08A	PR D78 091103	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BRIERE	08	PR D78 092007	R.A. Briere <i>et al.</i>	(CLEO Collab.)
CAWLFIELD	06	PR D73 012003	C. Cawlfield <i>et al.</i>	(CLEO Collab.)
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)
CRONIN-HEN... ... HENNESSY	04	PRL 92 222002	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)
CRAWFORD	92B	PL B294 139	G. Crawford, R. Fulton	(CLEO Collab.)
HEINTZ	92	PR D46 1928	U. Heintz <i>et al.</i>	(CUSB II Collab.)
HEINTZ	91	PRL 66 1563	U. Heintz <i>et al.</i>	(CUSB Collab.)
MORRISON	91	PRL 67 1696	R.J. Morrison <i>et al.</i>	(CLEO Collab.)
NARAIN	91	PRL 66 3113	M. Narain <i>et al.</i>	(CUSB Collab.)