

# $\eta_c(2S)$

$$I^G(J^{PC}) = 0^+(0^{-+})$$

Quantum numbers are quark model predictions.

## $\eta_c(2S)$ MASS

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>3639.4 ± 1.3 OUR AVERAGE</b> Error includes scale factor of 1.2.				
3646.9 ± 1.6 ± 3.6	57 ± 17	ABLIKIM	13K BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$
3637.6 ± 2.9 ± 1.6	127 ± 18	<sup>1</sup> ABLIKIM	12G BES3	$\psi(2S) \rightarrow \gamma K^0 K \pi,$ $K K \pi^0$
3638.5 ± 1.5 ± 0.8	624	<sup>2</sup> DEL-AMO-SA..11M BABR		$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$
3640.5 ± 3.2 ± 2.5	1201	<sup>2</sup> DEL-AMO-SA..11M BABR		$\gamma\gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
3636.1 <sup>+3.9+0.7</sup> <sub>-4.2-2.0</sub>	128	<sup>3</sup> VINOKUROVA 11	BELL	$B^\pm \rightarrow K^\pm (K_S^0 K^\pm \pi^\mp)$
3626 ± 5 ± 6	311	<sup>4</sup> ABE	07 BELL	$e^+ e^- \rightarrow J/\psi (c\bar{c})$
3645.0 ± 5.5 <sup>+4.9</sup> <sub>-7.8</sub>	121 ± 27	AUBERT	05C BABR	$e^+ e^- \rightarrow J/\psi c\bar{c}$
3642.9 ± 3.1 ± 1.5	61	ASNER	04 CLEO	$\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0 K^\pm \pi^\mp$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
3639 ± 7	98 ± 52	<sup>5</sup> AUBERT	06E BABR	$B^\pm \rightarrow K^\pm X_{c\bar{c}}$
3630.8 ± 3.4 ± 1.0	112 ± 24	<sup>6</sup> AUBERT	04D BABR	$\gamma\gamma \rightarrow \eta_c(2S) \rightarrow K \bar{K} \pi$
3654 ± 6 ± 8	39 ± 11	<sup>7</sup> CHOI	02 BELL	$B \rightarrow K K_S K^- \pi^+$
3594 ± 5		<sup>8</sup> EDWARDS	82C CBAL	$e^+ e^- \rightarrow \gamma X$

<sup>1</sup> From a simultaneous fit to  $K_S^0 K^\pm \pi^\mp$  and  $K^+ K^- \pi^0$  decay modes.  
<sup>2</sup> Ignoring possible interference with continuum.  
<sup>3</sup> Accounts for interference with non-resonant continuum.  
<sup>4</sup> From a fit of the  $J/\psi$  recoil mass spectrum. Supersedes ABE,K 02 and ABE 04G.  
<sup>5</sup> From the fit of the kaon momentum spectrum. Systematic errors not evaluated.  
<sup>6</sup> Superseded by DEL-AMO-SANCHEZ 11M.  
<sup>7</sup> Superseded by VINOKUROVA 11.  
<sup>8</sup> Assuming mass of  $\psi(2S) = 3686$  MeV.

## $\eta_c(2S)$ WIDTH

<u>VALUE (MeV)</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>11.3<sup>+</sup><sub>-</sub> 3.2<sup>+</sup><sub>2.9</sub> OUR AVERAGE</b>					
9.9 ± 4.8 ± 2.9		57 ± 17	ABLIKIM	13K BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$
16.9 ± 6.4 ± 4.8		127 ± 18	<sup>9</sup> ABLIKIM	12G BES3	$\psi(2S) \rightarrow \gamma K^0 K \pi,$ $K K \pi^0$
13.4 ± 4.6 ± 3.2		624	<sup>10</sup> DEL-AMO-SA..11M BABR		$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$
6.6 <sup>+8.4+2.6</sup> <sub>-5.1-0.9</sub>		128	<sup>11</sup> VINOKUROVA 11	BELL	$B^\pm \rightarrow K^\pm (K_S^0 K^\pm \pi^\mp)$
6.3 ± 12.4 ± 4.0		61	ASNER	04 CLEO	$\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0 K^\pm \pi^\mp$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 23	90	98 ± 52	<sup>12</sup> AUBERT	06E BABR	$B^\pm \rightarrow K^\pm X_c \bar{c}$
22 ± 14		121 ± 27	AUBERT	05C BABR	$e^+ e^- \rightarrow J/\psi c \bar{c}$
17.0 ± 8.3 ± 2.5		112 ± 24	<sup>13</sup> AUBERT	04D BABR	$\gamma\gamma \rightarrow \eta_c(2S) \rightarrow$ $K \bar{K} \pi$
< 55	90	39 ± 11	<sup>14</sup> CHOI	02 BELL	$B \rightarrow K K_S K^- \pi^+$
< 8.0	95		<sup>15</sup> EDWARDS	82C CBAL	$e^+ e^- \rightarrow \gamma X$

<sup>9</sup> From a simultaneous fit to  $K_S^0 K^\pm \pi^\mp$  and  $K^+ K^- \pi^0$  decay modes.

<sup>10</sup> Ignoring possible interference with continuum.

<sup>11</sup> Accounts for interference with non-resonant continuum.

<sup>12</sup> From the fit of the kaon momentum spectrum. Systematic errors not evaluated.

<sup>13</sup> Superseded by DEL-AMO-SANCHEZ 11M.

<sup>14</sup> For a mass value of  $3654 \pm 6$  MeV. Superseded by VINOKUROVA 11.

<sup>15</sup> For a mass value of  $3594 \pm 5$  MeV

### $\eta_c(2S)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1$ hadrons	not seen	
$\Gamma_2$ $K \bar{K} \pi$	( 1.9 ± 1.2 ) %	
$\Gamma_3$ $2\pi^+ 2\pi^-$	not seen	
$\Gamma_4$ $\rho^0 \rho^0$	not seen	
$\Gamma_5$ $3\pi^+ 3\pi^-$	not seen	
$\Gamma_6$ $K^+ K^- \pi^+ \pi^-$	not seen	
$\Gamma_7$ $K^{*0} \bar{K}^{*0}$	not seen	
$\Gamma_8$ $K^+ K^- \pi^+ \pi^- \pi^0$	( 1.4 ± 1.0 ) %	
$\Gamma_9$ $K^+ K^- 2\pi^+ 2\pi^-$	not seen	
$\Gamma_{10}$ $K_S^0 K^- 2\pi^+ \pi^- + c.c.$	seen	
$\Gamma_{11}$ $2K^+ 2K^-$	not seen	
$\Gamma_{12}$ $\phi\phi$	not seen	
$\Gamma_{13}$ $p\bar{p}$	< 2.9 × 10 <sup>-4</sup>	90%
$\Gamma_{14}$ $\gamma\gamma$	( 1.9 ± 1.3 ) × 10 <sup>-4</sup>	
$\Gamma_{15}$ $\pi^+ \pi^- \eta$	not seen	
$\Gamma_{16}$ $\pi^+ \pi^- \eta'$	not seen	
$\Gamma_{17}$ $K^+ K^- \eta$	not seen	
$\Gamma_{18}$ $\pi^+ \pi^- \eta_c(1S)$	< 25 %	90%

### $\eta_c(2S)$ PARTIAL WIDTHS

$\Gamma(\gamma\gamma)$	VALUE (keV)	DOCUMENT ID	TECN	COMMENT	$\Gamma_{14}$
------------------------	-------------	-------------	------	---------	---------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.3 ± 0.6	<sup>16</sup> ASNER	04	CLEO	$\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0 K^\pm \pi^\mp$
-----------	---------------------	----	------	---

<sup>16</sup> They measure  $\Gamma(\eta_c(2S)\gamma\gamma) B(\eta_c(2S) \rightarrow K\bar{K}\pi) = (0.18 \pm 0.05 \pm 0.02) \Gamma(\eta_c(1S)\gamma\gamma) B(\eta_c(1S) \rightarrow K\bar{K}\pi)$ . The value for  $\Gamma(\eta_c(2S) \rightarrow \gamma\gamma)$  is derived assuming that the branching fractions for  $\eta_c(2S)$  and  $\eta_c(1S)$  decays to  $K_S K\pi$  are equal and using  $\Gamma(\eta_c(1S) \rightarrow \gamma\gamma) = 7.4 \pm 0.4 \pm 2.3$  keV.

### $\eta_c(2S) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(2\pi^+2\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_3\Gamma_{14}/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<6.5	90	UEHARA	08	BELL	$\gamma\gamma \rightarrow \eta_c(2S) \rightarrow 2(\pi^+\pi^-)$

$\Gamma(K\bar{K}\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_2\Gamma_{14}/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>41±4±6</b>	624	<sup>17</sup> DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$	

<sup>17</sup> Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.

$\Gamma(K^+K^-\pi^+\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_6\Gamma_{14}/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<5.0	90	UEHARA	08	BELL	$\gamma\gamma \rightarrow \eta_c(2S) \rightarrow K^+K^-\pi^+\pi^-$

$\Gamma(K^+K^-\pi^+\pi^-\pi^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_8\Gamma_{14}/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>30±6±5</b>	1201	<sup>18</sup> DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$	

<sup>18</sup> Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.

$\Gamma(2K^+2K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_{11}\Gamma_{14}/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<2.9	90	UEHARA	08	BELL	$\gamma\gamma \rightarrow \eta_c(2S) \rightarrow 2(K^+K^-)$

$\Gamma(\pi^+\pi^-\eta_c(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_{18}\Gamma_{14}/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<133	90	LEES	12AE	BABR	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\eta_c$

### $\eta_c(2S) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma^2(\text{total})$

$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_{13}/\Gamma \times \Gamma_{14}/\Gamma$
VALUE (units $10^{-8}$ )	CL%	DOCUMENT ID	TECN	COMMENT	
< 5.6	90 <sup>19,20,21</sup>	AMBROGIANI 01	E835	$\bar{p}p \rightarrow \gamma\gamma$	
• • •				We do not use the following data for averages, fits, limits, etc. • • •	
< 8.0	90 <sup>19,20,22</sup>	AMBROGIANI 01	E835	$\bar{p}p \rightarrow \gamma\gamma$	
<12.0	90	<sup>20,22</sup> AMBROGIANI 01	E835	$\bar{p}p \rightarrow \gamma\gamma$	

<sup>19</sup> Including the measurements of of ARMSTRONG 95F in the AMBROGIANI 01 analysis.

<sup>20</sup> For a total width  $\Gamma=5$  MeV.

<sup>21</sup> For the resonance mass region 3589–3599 MeV/ $c^2$ .

<sup>22</sup> For the resonance mass region 3575–3660 MeV/ $c^2$ .

## $\eta_c(2S)$ BRANCHING RATIOS

### $\Gamma(\text{hadrons})/\Gamma_{\text{total}}$

$\Gamma_1/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>not seen</b>	ABREU	980	DLPH $e^+e^- \rightarrow e^+e^- + \text{hadrons}$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
seen	<sup>23</sup> EDWARDS	82c	CBAL $e^+e^- \rightarrow \gamma X$

<sup>23</sup> For a mass value of  $3594 \pm 5$  MeV

### $\Gamma(K\bar{K}\pi)/\Gamma_{\text{total}}$

$\Gamma_2/\Gamma$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.9±0.4±1.1</b>	$59 \pm 12$	<sup>24</sup> AUBERT	08AB	BABR $B \rightarrow \eta_c(2S)K \rightarrow K\bar{K}\pi K$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	$127 \pm 18$	ABLIKIM	13K	BES3 $\psi(2S) \rightarrow \gamma K\bar{K}\pi$
seen	$39 \pm 11$	<sup>25</sup> CHOI	02	BELL $B \rightarrow K K_S K^- \pi^+$

<sup>24</sup> Derived from a measurement of  $[B(B^+ \rightarrow \eta_c(2S)K^+) \times B(\eta_c(2S) \rightarrow K\bar{K}\pi)] / [B(B^+ \rightarrow \eta_c K^+) \times B(\eta_c \rightarrow K\bar{K}\pi)] = (9.6^{+2.0}_{-1.9} \pm 2.5)\%$  and using  $B(B^+ \rightarrow \eta_c(2S)K^+) = (3.4 \pm 1.8) \times 10^{-4}$ , and  $[B(B^+ \rightarrow \eta_c K^+) \times B(\eta_c \rightarrow K\bar{K}\pi)] = (6.88 \pm 0.77^{+0.55}_{-0.66}) \times 10^{-5}$ .

<sup>25</sup> For a mass value of  $3654 \pm 6$  MeV

### $\Gamma(2\pi^+2\pi^-)/\Gamma_{\text{total}}$

$\Gamma_3/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>not seen</b>	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S)$

### $\Gamma(\rho^0\rho^0)/\Gamma_{\text{total}}$

$\Gamma_4/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>not seen</b>	ABLIKIM	11H	BES3 $\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$

### $\Gamma(K^+K^-\pi^+\pi^-)/\Gamma_{\text{total}}$

$\Gamma_6/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>not seen</b>	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S)$

### $\Gamma(K^+K^-\pi^+\pi^-\pi^0)/\Gamma(K\bar{K}\pi)$

$\Gamma_8/\Gamma_2$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.73±0.17±0.17</b>	1201	<sup>26</sup> DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$

<sup>26</sup> We have multiplied the value of  $\Gamma(K^+K^-\pi^+\pi^-\pi^0)/\Gamma(K_S^0 K^\pm \pi^\mp)$  reported in DEL-AMO-SANCHEZ 11M by a factor 1/3 to obtain  $\Gamma(K^+K^-\pi^+\pi^-\pi^0)/\Gamma(K\bar{K}\pi)$ . Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.

### $\Gamma(K^{*0}\bar{K}^{*0})/\Gamma_{\text{total}}$

$\Gamma_7/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>not seen</b>	ABLIKIM	11H	BES3 $\psi(2S) \rightarrow \gamma K^+K^-\pi^+\pi^-$

### $\Gamma(K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$

$\Gamma_{10}/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	$57 \pm 17$	ABLIKIM	13K	BES3 $\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$

$\Gamma(2K^+2K^-)/\Gamma_{\text{total}}$				$\Gamma_{11}/\Gamma$
VALUE	DOCUMENT ID	TECN	COMMENT	
not seen	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S)$	

$\Gamma(\phi\phi)/\Gamma_{\text{total}}$				$\Gamma_{12}/\Gamma$
VALUE	DOCUMENT ID	TECN	COMMENT	
not seen	ABLIKIM	11H	BES3 $\psi(2S) \rightarrow \gamma K^+ K^- K^+ K^-$	

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$				$\Gamma_{14}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<5 \times 10^{-4}$	90	<sup>27</sup> WICHT	08	BELL	$B^\pm \rightarrow K^\pm \gamma\gamma$
not seen		AMBROGIANI	01	E835	$\bar{p}p \rightarrow \gamma\gamma$
$<0.01$	90	LEE	85	CBAL	$\psi' \rightarrow \text{photons}$

<sup>27</sup> WICHT 08 reports  $[\Gamma(\eta_c(2S) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \eta_c(2S) K^+)] < 0.18 \times 10^{-6}$  which we divide by our best value  $B(B^+ \rightarrow \eta_c(2S) K^+) = 3.4 \times 10^{-4}$ .

$\Gamma(\pi^+ \pi^- \eta_c(1S))/\Gamma(K\bar{K}\pi)$				$\Gamma_{18}/\Gamma_2$	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<b><math>&lt;3.33</math></b>	90	<sup>28</sup> LEES	12AE	BABR	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \eta_c$

<sup>28</sup> We divided the reported limit by 3 to take into account isospin relations.

### $\eta_c(2S)$ CROSS-PARTICLE BRANCHING RATIOS

$\Gamma(\eta_c(2S) \rightarrow 2\pi^+ 2\pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$				$\Gamma_3/\Gamma \times \Gamma_{122}^{\psi(2S)}/\Gamma\psi(2S)$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT

<b><math>&lt;14.6 \times 10^{-6}</math></b>	90	<sup>29</sup> CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$
---	----	------------------------------	------	---

<sup>29</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$\Gamma(\eta_c(2S) \rightarrow \rho^0 \rho^0)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$				$\Gamma_4/\Gamma \times \Gamma_{122}^{\psi(2S)}/\Gamma\psi(2S)$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT

<b><math>&lt;12.7 \times 10^{-7}</math></b>	90	ABLIKIM	11H	BES3 $\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$
---	----	---------	-----	--

$\Gamma(\eta_c(2S) \rightarrow 3\pi^+ 3\pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$				$\Gamma_5/\Gamma \times \Gamma_{122}^{\psi(2S)}/\Gamma\psi(2S)$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT

<b><math>&lt;13.2 \times 10^{-6}</math></b>	90	<sup>30</sup> CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma 3\pi^+ 3\pi^-$
---	----	------------------------------	------	---

<sup>30</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow K^+ K^- \pi^+ \pi^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}} \times \Gamma_6 / \Gamma \times \Gamma_{122}^{\psi(2S)} / \Gamma\psi(2S)$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<9.6 \times 10^{-6}$	90	31 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$

<sup>31</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow K^{*0} \bar{K}^{*0}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}} \times \Gamma_7 / \Gamma \times \Gamma_{122}^{\psi(2S)} / \Gamma\psi(2S)$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<19.6 \times 10^{-7}$	90	ABLIKIM	11H BES3	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$

$$\Gamma(\eta_c(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}} \times \Gamma_8 / \Gamma \times \Gamma_{122}^{\psi(2S)} / \Gamma\psi(2S)$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<43.0 \times 10^{-6}$	90	32 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^- \pi^0$

<sup>32</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow K^+ K^- 2\pi^+ 2\pi^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}} \times \Gamma_9 / \Gamma \times \Gamma_{122}^{\psi(2S)} / \Gamma\psi(2S)$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<9.7 \times 10^{-6}$	90	33 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- 2\pi^+ 2\pi^-$

<sup>33</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}} \times \Gamma_{10} / \Gamma \times \Gamma_{122}^{\psi(2S)} / \Gamma\psi(2S)$$

VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$7.03 \pm 2.10 \pm 0.7$	60		ABLIKIM	13K BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 15.2$	90	34 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}$
----------	----	-------------------	------	--

<sup>34</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow \phi\phi) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}} \times \Gamma_{12} / \Gamma \times \Gamma_{122}^{\psi(2S)} / \Gamma\psi(2S)$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.8 \times 10^{-7}$	90	ABLIKIM	11H BES3	$\psi(2S) \rightarrow \gamma K^+ K^- K^+ K^-$

$$\Gamma(\eta_c(2S) \rightarrow \pi^+ \pi^- \eta) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}} \times \Gamma_{15} / \Gamma \times \Gamma_{122}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.3 \times 10^{-6}$	90	35 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- \eta$

<sup>35</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow \pi^+ \pi^- \eta') / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}} \times \Gamma_{16} / \Gamma \times \Gamma_{122}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<14.2 \times 10^{-6}$	90	36 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- \eta'$

<sup>36</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow K^+ K^- \eta) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}} \times \Gamma_{17} / \Gamma \times \Gamma_{122}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.9 \times 10^{-6}$	90	37 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- \eta$

<sup>37</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow \pi^+ \pi^- \eta_c(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}} \times \Gamma_{18} / \Gamma \times \Gamma_{122}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.7 \times 10^{-4}$	90	38 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- \eta_c(1S)$

<sup>38</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

## $\eta_c(2S)$ REFERENCES

ABLIKIM	13K	PR D87 052005	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	12G	PRL 109 042003	M. Ablikim <i>et al.</i>	(BES III Collab.)
LEES	12AE	PR D86 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	11H	PR D84 091102	M. Ablikim <i>et al.</i>	(BES III Collab.)
DEL-AMO-SA...	11M	PR D84 012004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
VINOKUROVA	11	PL B706 139	A. Vinokurova <i>et al.</i>	(BELLE Collab.)
CRONIN-HEN...	10	PR D81 052002	D. Cronin-Hennessey <i>et al.</i>	(CLEO Collab.)
AUBERT	08AB	PR D78 012006	B. Aubert <i>et al.</i>	(BABAR Collab.)
UEHARA	08	EPJ C53 1	S. Uehara <i>et al.</i>	(BELLE Collab.)
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)
ABE	07	PRL 98 082001	K. Abe <i>et al.</i>	(BELLE Collab.)
AUBERT	06E	PRL 96 052002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	05C	PR D72 031101	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABE	04G	PR D70 071102	K. Abe <i>et al.</i>	(BELLE Collab.)
ASNER	04	PRL 92 142001	D.M. Asner <i>et al.</i>	(CLEO Collab.)
AUBERT	04D	PRL 92 142002	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABE,K	02	PRL 89 142001	K. Abe <i>et al.</i>	(BELLE Collab.)
CHOI	02	PRL 89 102001	S.-K. Choi <i>et al.</i>	(BELLE Collab.)
AMBROGIANI	01	PR D64 052003	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)
ABREU	98O	PL B441 479	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ARMSTRONG	95F	PR D52 4839	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
LEE	85	SLAC 282	R.A. Lee	(SLAC)
EDWARDS	82C	PRL 48 70	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)