

$f_2(1810)$

$I^G(J^{PC}) = 0^+(2^{++})$

OMITTED FROM SUMMARY TABLE

Needs confirmation.

$f_2(1810)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1815 ± 12 OUR AVERAGE		Error includes scale factor of 1.4. See the ideogram below.		
$1737 \pm 9^{+198}_{-65}$		1 UEHARA	10A BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \eta\eta$
1800 ± 30	40	ALDE	88D GAM4	$300 \pi^- p \rightarrow \pi^- p 4\pi^0$
1806 ± 10	1600	ALDE	87 GAM4	$100 \pi^- p \rightarrow 4\pi^0 n$
1870 ± 40		2 ALDE	86D GAM4	$100 \pi^- p \rightarrow \eta\eta n$
1857^{+35}_{-24}		3 COSTA...	80 OMEG	$10 \pi^- p \rightarrow K^+ K^- n$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1858^{+18}_{-71}		4 LONGACRE	86 RVUE	Compilation
1799 ± 15		5 CASON	82 STRC	$8 \pi^+ p \rightarrow \Delta^{++} \pi^0 \pi^0$


¹ Breit-Wigner mass.

² Seen in only one solution.

³ Error increased by spread of two solutions. Included in LONGACRE 86 global analysis.

⁴ From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.

⁵ From an amplitude analysis of the reaction $\pi^+ \pi^- \rightarrow 2\pi^0$. The resonance in the $2\pi^0$ final state is not confirmed by PROKOSHKIN 97.



$f_2(1810)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
197 ± 22 OUR AVERAGE		Error includes scale factor of 1.5. See the ideogram below.		
228 ± 21	21 + 234	6 UEHARA	10A BELL	10.6 $e^+ e^- \rightarrow e^+ e^- \eta\eta$
20 - 153				
160 ± 30	40	ALDE	88D GAM4	300 $\pi^- p \rightarrow \pi^- p 4\pi^0$
190 ± 20	1600	ALDE	87 GAM4	100 $\pi^- p \rightarrow 4\pi^0 n$
250 ± 30		7 ALDE	86D GAM4	100 $\pi^- p \rightarrow \eta\eta n$
185 ± 102		8 COSTA...	80 OMEG	10 $\pi^- p \rightarrow K^+ K^- n$
- 139				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
388 ± 15	15	9 LONGACRE	86 RVUE	Compilation
- 21				
280 ± 42	42	10 CASON	82 STRC	8 $\pi^+ p \rightarrow \Delta^{++} \pi^0 \pi^0$
- 35				


⁶ Breit-Wigner width.

⁷ Seen in only one solution.

⁸ Error increased by spread of two solutions. Included in LONGACRE 86 global analysis.

⁹ From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.

¹⁰ From an amplitude analysis of the reaction $\pi^+ \pi^- \rightarrow 2\pi^0$. The resonance in the $2\pi^0$ final state is not confirmed by PROKOSHKIN 97.



$f_2(1810)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \pi\pi$	
$\Gamma_2 \eta\eta$	
$\Gamma_3 4\pi^0$	seen
$\Gamma_4 K^+K^-$	
$\Gamma_5 \gamma\gamma$	seen

$f_2(1810) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(\eta\eta) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_2\Gamma_5/\Gamma$
<u>VALUE (eV)</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
5.2$^{+0.9+37.3}_{-0.8-4.5}$	11 UEHARA 10A BELL 10.6 $e^+e^- \rightarrow e^+e^-\eta\eta$

¹¹ Including interference with the $f'_2(1525)$ (parameters fixed to the values from the 2008 edition of this review, PDG 08) and $f_2(1270)$. May also be the $f_0(1500)$.

$f_2(1810)$ BRANCHING RATIOS **$\Gamma(\pi\pi)/\Gamma_{\text{total}}$** **$\Gamma_1/\Gamma$**

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	AMSLER 02	CBAR	0.9 $\bar{p}p \rightarrow \pi^0 \eta \eta, \pi^0 \pi^0 \pi^0$
not seen	PROKOSHKIN 97	GAM2	38 $\pi^- p \rightarrow \pi^0 \pi^0 n$
$0.21^{+0.02}_{-0.03}$	12 LONGACRE 86	RVUE	Compilation
0.44 ± 0.03	13 CASON 82	STRC	$8 \pi^+ p \rightarrow \Delta^{++} \pi^0 \pi^0$
12 From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.			
13 Included in LONGACRE 86 global analysis.			

 $\Gamma(\eta\eta)/\Gamma_{\text{total}}$ **Γ_2/Γ**

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.008^{+0.028}_{-0.003}$	14 LONGACRE 86	RVUE	Compilation
14 From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.			

 $\Gamma(\pi\pi)/\Gamma(4\pi^0)$ **Γ_1/Γ_3**

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
<0.75	ALDE 87	GAM4	$100 \pi^- p \rightarrow 4\pi^0 n$

 $\Gamma(4\pi^0)/\Gamma(\eta\eta)$ **Γ_3/Γ_2**

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.8 ± 0.3	ALDE 87	GAM4	$100 \pi^- p \rightarrow 4\pi^0 n$

 $\Gamma(K^+ K^-)/\Gamma_{\text{total}}$ **Γ_4/Γ**

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.003^{+0.019}_{-0.002}$	15 LONGACRE 86	RVUE	Compilation
seen	COSTA... 80	OMEG	$10 \pi^- p \rightarrow K^+ K^- n$
15 From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.			

 $f_2(1810)$ REFERENCES

UEHARA PDG	10A 08	PR D82 114031 PL B667 1	S. Uehara <i>et al.</i> C. Amsler <i>et al.</i>	(BELLE Collab.) (PDG Collab.)
AMSLER	02	EPJ C23 29	C. Amsler <i>et al.</i>	
PROKOSHKIN	97	SPD 42 117 Translated from DANS 353 323.	Y.D. Prokoshkin <i>et al.</i>	(SERP)
ALDE	88D	SJNP 47 810 Translated from YAF 47 1273.	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+)
ALDE	87	PL B198 286	D.M. Alde <i>et al.</i>	(LANL, BRUX, SERP, LAPP)
ALDE	86D	NP B269 485	D.M. Alde <i>et al.</i>	(BELG, LAPP, SERP, CERN+)
LONGACRE	86	PL B177 223	R.S. Longacre <i>et al.</i>	(BNL, BRAN, CUNY+)
CASON	82	PRL 48 1316	N.M. Cason <i>et al.</i>	(NDAM, ANL)
COSTA...	80	NP B175 402	G. Costa de Beauregard <i>et al.</i>	(BARI, BONN+)