

$f_2(2150)$ $I^G(J^{PC}) = 0^+(2^{++})$

OMMITTED FROM SUMMARY TABLE

This entry was previously called T_0 . **$f_2(2150)$ MASS** **$f_2(2150)$ MASS, COMBINED MODES (MeV)**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

2157±12 OUR AVERAGE Includes data from the 2 datablocks that follow this one.

• • • We do not use the following data for averages, fits, limits, etc. • • •

2170± 6 80k 1 UMAN 06 E835 5.2 $\bar{p}p \rightarrow \eta\eta\pi^0$ ¹ Statistical error only. **$\eta\eta$ MODE**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

The data in this block is included in the average printed for a previous datablock.

2157±12 OUR AVERAGE

2151±16	BARBERIS	00E	450 $p p \rightarrow p_f \eta\eta p_s$
2175±20	PROKOSHKIN	95D GAM4	300 $\pi^- N \rightarrow \pi^- N 2\eta$, 450 $p p \rightarrow p p 2\eta$
2130±35	SINGOVSKI	94 GAM4	450 $p p \rightarrow p p 2\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2140±30 2 ABELE 99B CBAR

2104±20 3 ARMSTRONG 93C E760 $\bar{p}p \rightarrow \pi^0 \eta\eta \rightarrow 6\gamma$ ² Spin not determined.³ No J^{PC} determination. **$\eta\pi\pi$ MODE**

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

• • • We do not use the following data for averages, fits, limits, etc. • • •

2135±20±45 4 ADOMEIT 96 CBAR 0 1.94 $\bar{p}p \rightarrow \eta 3\pi^0$ ⁴ ANISOVICH 00E recommends to withdraw ADOMEIT 96 that assumed a single $J^P = 2^+$ resonance. **$\bar{p}p \rightarrow \pi\pi$**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 2090	5 OAKDEN	94 RVUE	0.36–1.55 $\bar{p}p \rightarrow \pi\pi$
~ 2120	6 OAKDEN	94 RVUE	0.36–1.55 $\bar{p}p \rightarrow \pi\pi$
~ 2170	7 MARTIN	80B RVUE	
~ 2150	7 MARTIN	80C RVUE	
~ 2150	8 DULUDE	78B OSPK	1–2 $\bar{p}p \rightarrow \pi^0 \pi^0$

⁵ OAKDEN 94 makes an amplitude analysis of LEAR data on $\bar{p}p \rightarrow \pi\pi$ using a method based on Barrelet zeros. This is solution A. The amplitude analysis of HASAN 94 includes earlier data as well, and assume that the data can be parametrized in terms of towers of nearly degenerate resonances on the leading Regge trajectory. See also KLOET 96 and MARTIN 97 who make related analyses.

⁶ From solution B of amplitude analysis of data on $\bar{p}p \rightarrow \pi\pi$.

⁷ $I(J^P) = 0(2^+)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^-\pi^+$ and $\pi^0\pi^0$.

⁸ $I^G(J^P) = 0^+(2^+)$ from partial-wave amplitude analysis.

S-CHANNEL $\bar{p}p$, $\bar{N}N$ or $\bar{K}K$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2139 ^{+ 8} _{- 9}	9 EVANGELIS...	97	SPEC	0.6-2.4 $\bar{p}p \rightarrow K_S^0 K_S^0$
~2190	9 CUTTS	78B	CNTR	0.97-3 $\bar{p}p \rightarrow \bar{N}N$
2155 \pm 15	9,10 COUPLAND	77	CNTR 0	0.7-2.4 $\bar{p}p \rightarrow \bar{p}p$
2193 \pm 2	9,11 ALSPECTOR	73	CNTR	$\bar{p}p$ S channel

⁹ Isospins 0 and 1 not separated.

¹⁰ From a fit to the total elastic cross section.

¹¹ Referred to as T or T region by ALSPECTOR 73.


$K\bar{K}$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2200 \pm 13	VLADIMIRSK...06	SPEC	40 $\pi^- p \rightarrow K_S^0 K_S^0 n$
2150 \pm 20	ABLIKIM 04E	BES2	$J/\psi \rightarrow \omega K^+ K^-$
2130 \pm 35	BARBERIS 99	OMEG 450	$p p \rightarrow p_s p_f K^+ K^-$

$f_2(2150)$ WIDTH

$f_2(2150)$ WIDTH, COMBINED MODES (MeV)

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
152\pm30 OUR AVERAGE		Includes data from the 2 datablocks that follow this one. Error includes scale factor of 1.4. See the ideogram below.		
• • • We do not use the following data for averages, fits, limits, etc. • • •				
182 \pm 11	80k	12 UMAN	06 E835	5.2 $\bar{p}p \rightarrow \eta\eta\pi^0$
12 Statistical error only.				



$\eta\eta$ MODE


VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
The data in this block is included in the average printed for a previous datablock.			

152±30 OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below.

280±70	BARBERIS 00E	450 $p p \rightarrow p_f \eta\eta p_s$
150±35	PROKOSHKIN 95D	300 $\pi^- N \rightarrow \pi^- N 2\eta$, 450 $p p \rightarrow p p 2\eta$
130±30	SINGOVSKI 94	GAM4 450 $p p \rightarrow p p 2\eta$
• • • We do not use the following data for averages, fits, limits, etc. • • •		
310±50	13 ABELE 99B	CBAR
203±10	14 ARMSTRONG 93C	E760 $\bar{p} p \rightarrow \pi^0 \eta\eta \rightarrow 6\gamma$

¹³ Spin not determined.

¹⁴ No JPC determination.



$\eta\pi\pi$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
The data in this block is included in the average printed for a previous datablock.				

• • • We do not use the following data for averages, fits, limits, etc. • • •

$250 \pm 25 \pm 45$ ¹⁵ ADOMEIT 96 CBAR 0 $1.94 \bar{p}p \rightarrow \eta 3\pi^0$

¹⁵ ANISOVICH 00E recommends to withdraw ADOMEIT 96 that assumed a single $J^P = 2^+$ resonance.

$\bar{p}p \rightarrow \pi\pi$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
250 OUR ESTIMATE			

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 70 ¹⁶ OAKDEN 94 RVUE $0.36-1.55 \bar{p}p \rightarrow \pi\pi$

~ 250 ¹⁷ MARTIN 80B RVUE

~ 250 ¹⁷ MARTIN 80C RVUE

~ 250 ¹⁸ DULUDE 78B OSPK 1-2 $\bar{p}p \rightarrow \pi^0\pi^0$

¹⁶ See however KLOET 96 who fit $\pi^+\pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.

¹⁷ $I(J^P) = 0(2^+)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^-\pi^+$ and $\pi^0\pi^0$.

¹⁸ $I^G(J^P) = 0^+(2^+)$ from partial-wave amplitude analysis.

S-CHANNEL $\bar{p}p$, $\bar{N}N$ or $\bar{K}K$

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
56 ⁺³¹ ₋₁₆	19 EVANGELIS...	97	SPEC	0.6-2.4 $\bar{p}p \rightarrow K_S^0 K_S^0$
135±75	20,21 COUPLAND	77	CNTR 0	0.7-2.4 $\bar{p}p \rightarrow \bar{p}p$
98± 8	21 ALSPECTOR	73	CNTR	$\bar{p}p$ S channel

19 Isospin 0 and 2 not separated.

20 From a fit to the total elastic cross section.

21 Isospins 0 and 1 not separated.

 $K\bar{K}$ MODE

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
91±62	VLADIMIRSK...06	SPEC	40 $\pi^- p \rightarrow K_S^0 K_S^0 n$
150±30	ABLIKIM 04E	BES2	$J/\psi \rightarrow \omega K^+ K^-$
270±50	BARBERIS 99	OMEG 450	$p p \rightarrow p_s p_f K^+ K^-$

 $f_2(2150)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi\pi$	
Γ_2 $\eta\eta$	seen
Γ_3 $K\bar{K}$	seen
Γ_4 $f_2(1270)\eta$	seen
Γ_5 $a_2(1320)\pi$	seen
Γ_6 $p\bar{p}$	seen

 $f_2(2150)$ BRANCHING RATIOS

<u>$\Gamma(K\bar{K})/\Gamma(\eta\eta)$</u>	<u>Γ_3/Γ_2</u>
1.28±0.23	BARBERIS 00E 450 $p p \rightarrow p_f \eta\eta p_s$
• • • We do not use the following data for averages, fits, limits, etc. • • •	
<0.1	95 22 PROKOSHIN 95D GAM4 300 $\pi^- N \rightarrow \pi^- N 2\eta$, 450 $p p \rightarrow p p 2\eta$

22 Using data from ARMSTRONG 89D.

<u>$\Gamma(\pi\pi)/\Gamma(\eta\eta)$</u>	<u>Γ_1/Γ_2</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •	
<0.33	95 23 PROKOSHIN 95D GAM4 300 $\pi^- N \rightarrow \pi^- N 2\eta$, 450 $p p \rightarrow p p 2\eta$

23 Derived from a $\pi^0 \pi^0/\eta\eta$ limit.

$\Gamma(f_2(1270)\eta)/\Gamma(a_2(1320)\pi)$				Γ_4/Γ_5
VALUE	DOCUMENT ID	TECN	COMMENT	
0.79±0.11	²⁴ ADOMEIT	96	CBAR	$1.94 \bar{p}p \rightarrow \eta 3\pi^0$
²⁴ Using $B(a_2(1320) \rightarrow \eta\pi) = 0.145$				
$\Gamma(p\bar{p})/\Gamma_{\text{total}}$				Γ_6/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	73	ALEXANDER	10	CLEO $\psi(2S) \rightarrow \gamma p\bar{p}$

$f_2(2150)$ REFERENCES

ALEXANDER	10	PR D82 092002	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
UMAN	06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)
VLADIMIRSKY	06	PAN 69 493	V.V. Vladimirska <i>et al.</i>	(ITEP, Moscow)
		Translated from YAF 69 515.		
ABLIKIM	04E	PL B603 138	M. Ablikim <i>et al.</i>	(BES Collab.)
ANISOVICH	00E	PL B477 19	A.V. Anisovich <i>et al.</i>	
BARBERIS	00E	PL B479 59	D. Barberis <i>et al.</i>	(WA 102 Collab.)
ABELE	99B	EPJ C8 67	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
BARBERIS	99	PL B453 305	D. Barberis <i>et al.</i>	(Omega Expt.)
EVANGELISTI	97	PR D56 3803	C. Evangelista <i>et al.</i>	(LEAR Collab.)
MARTIN	97	PR C56 1114	B.R. Martin, G.C. Oades	(LOUC, AARH)
ADOMEIT	96	ZPHY C71 227	J. Adomeit <i>et al.</i>	(Crystal Barrel Collab.)
KLOET	96	PR D53 6120	W.M. Kloet, F. Myhrer	(RUTG, NORD)
PROKOSHKIN	95D	SPD 40 495	Y.D. Prokoshkin	(SERP) IGJPC
		Translated from DANS 344 469.		
HASAN	94	PL B334 215	A. Hasan, D.V. Bugg	(LOQM)
OAKDEN	94	NP A574 731	M.N. Oakden, M.R. Pennington	(DURH)
SINGOVSKI	94	NC 107A 1911	A.V. Singovsky	(SERP)
ARMSTRONG	93C	PL B307 394	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
ARMSTRONG	89D	PL B227 186	T.A. Armstrong, M. Benayoun	(ATHU, BARI, BIRM+)
MARTIN	80B	NP B176 355	B.R. Martin, D. Morgan	(LOUC, RHEL) JP
MARTIN	80C	NP B169 216	A.D. Martin, M.R. Pennington	(DURH) JP
CUTTS	78B	PR D17 16	D. Cutts <i>et al.</i>	(STON, WISC)
DULUDE	78B	PL 79B 335	R.S. Dulude <i>et al.</i>	(BROW, MIT, BARI) JP
COUPLAND	77	PL 71B 460	M. Coupland <i>et al.</i>	(LOQM, RHEL)
ALSPECTOR	73	PRL 30 511	J. Alspector <i>et al.</i>	(RUTG, UPNJ)