

**$\Sigma(1620)$**  $I(J^P) = 1(\frac{1}{2}^-)$  Status: \*

## OMITTED FROM SUMMARY TABLE

The  $S_{11}$  state at 1697 MeV reported by VANHORN 75 is tentatively listed under the  $\Sigma(1750)$ . CARROLL 76 sees two bumps in the isospin-1 total cross section near this mass. GAO 12 sees no evidence for this resonance.

Production experiments are listed separately in the next entry.

 **$\Sigma(1620)$  MASS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>≈ 1620 OUR ESTIMATE</b>			
1600 ± 6	<sup>1</sup> MORRIS 78	DPWA	$K^- n \rightarrow \Lambda\pi^-$
1608 ± 5	<sup>2</sup> CARROLL 76	DPWA	Isospin-1 total $\sigma$
1633 ± 10	<sup>3</sup> CARROLL 76	DPWA	Isospin-1 total $\sigma$
1630 ± 10	LANGBEIN 72	IPWA	$\bar{K}N$ multichannel
1620	KIM 71	DPWA	K-matrix analysis

 **$\Sigma(1620)$  WIDTH**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>87 ± 19</b>			
15	<sup>1</sup> MORRIS 78	DPWA	$K^- n \rightarrow \Lambda\pi^-$
10	<sup>2</sup> CARROLL 76	DPWA	Isospin-1 total $\sigma$
65 ± 20	<sup>3</sup> CARROLL 76	DPWA	Isospin-1 total $\sigma$
40	LANGBEIN 72	IPWA	$\bar{K}N$ multichannel
	KIM 71	DPWA	K-matrix analysis

 **$\Sigma(1620)$  DECAY MODES**

Mode
$\Gamma_1 N\bar{K}$
$\Gamma_2 \Lambda\pi$
$\Gamma_3 \Sigma\pi$

 **$\Sigma(1620)$  BRANCHING RATIOS**

$\Gamma(N\bar{K})/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	$\Gamma_1/\Gamma$
0.22 ± 0.02	LANGBEIN 72	IPWA	$\bar{K}N$ multichannel	
0.05	KIM 71	DPWA	K-matrix analysis	

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1620) \rightarrow \Lambda\pi$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
0.12 ± 0.02	<sup>1</sup> MORRIS 78	DPWA	$K^- n \rightarrow \Lambda\pi^-$	
not seen	BAILLON 75	IPWA	$\bar{K}N \rightarrow \Lambda\pi$	
0.15	KIM 71	DPWA	K-matrix analysis	

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1620) \rightarrow \Sigma\pi$	$(\Gamma_1 \Gamma_3)^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
not seen	HEPP	76B	$K^- N \rightarrow \Sigma\pi$
$0.40 \pm 0.06$	LANGBEIN	72	$\bar{K}N$ multichannel
0.08	KIM	71	DPWA K-matrix analysis

## $\Sigma(1620)$ FOOTNOTES

<sup>1</sup> MORRIS 78 obtains an equally good fit without including this resonance.

<sup>2</sup> Total cross-section bump with  $(J+1/2) \Gamma_{\text{el}} / \Gamma_{\text{total}}$  is 0.06 seen by CARROLL 76.

<sup>3</sup> Total cross-section bump with  $(J+1/2) \Gamma_{\text{el}} / \Gamma_{\text{total}}$  is 0.04 seen by CARROLL 76.

## $\Sigma(1620)$ REFERENCES

GAO	12	PR C86 025201	P. Gao, J. Shi, B.S. Zou	(BHEP, BEIJT)
Also		NP A867 41	P. Gao, B.S. Zou, A. Sibirtsev	(BHEP, BEIJT+)
MORRIS	78	PR D17 55	W.A. Morris <i>et al.</i>	(FSU) IJP
CARROLL	76	PRL 37 806	A.S. Carroll <i>et al.</i>	(BNL) I
HEPP	76B	PL 65B 487	V. Hepp <i>et al.</i>	(CERN, HEIDH, MPIM) IJP
BAILLON	75	NP B94 39	P.H. Baillon, P.J. Litchfield	(CERN, RHEL) IJP
VANHORN	75	NP B87 145	A.J. van Horn	(LBL) IJP
Also		NP B87 157	A.J. van Horn	(LBL) IJP
LANGBEIN	72	NP B47 477	W. Langbein, F. Wagner	(MPIM) IJP
KIM	71	PRL 27 356	J.K. Kim	(HARV) IJP
Also		Duke Conf. 161	J.K. Kim	(HARV) IJP
Hyperon Resonances, 1970				