




$$I(J^P) = \frac{1}{2}(0^-)$$

K⁰ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
497.614 ± 0.024 OUR FIT				Error includes scale factor of 1.6.
497.614 ± 0.022 OUR AVERAGE				Error includes scale factor of 1.5. See the ideogram below.
497.583 ± 0.005 ± 0.020	35k	AMBROSINO	07B	KLOE e ⁺ e ⁻ → K _L ⁰ K _S ⁰
497.625 ± 0.001 ± 0.031	655k	LAI	02	NA48 K _L ⁰ beam
497.661 ± 0.033	3713	BARKOV	87B	CMD e ⁺ e ⁻ → K _L ⁰ K _S ⁰
497.742 ± 0.085	780	BARKOV	85B	CMD e ⁺ e ⁻ → K _L ⁰ K _S ⁰
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
497.44 ± 0.50		FITCH	67	OSPK
498.9 ± 0.5	4500	BALTAY	66	HBC K ⁰ from $\bar{p}p$
497.44 ± 0.33	2223	KIM	65B	HBC K ⁰ from $\bar{p}p$
498.1 ± 0.4		CHRISTENS...	64	OSPK

WEIGHTED AVERAGE
497.614 ± 0.022 (Error scaled by 1.5)



$m_{K^0} - m_{K^\pm}$

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	
3.937±0.028 OUR FIT	Error includes scale factor of 1.8.					
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
3.95 ±0.21	417	HILL	68B	DBC	+	$K^+ d \rightarrow K^0 p p$
3.90 ±0.25	9	BURNSTEIN	65	HBC	-	
3.71 ±0.35	7	KIM	65B	HBC	-	$K^- p \rightarrow n \bar{K}^0$
5.4 ±1.1		CRAWFORD	59	HBC	+	
3.9 ±0.6		ROSENFELD	59	HBC	-	

K^0 MEAN SQUARE CHARGE RADIUS

<u>VALUE (fm²)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
-0.077±0.010 OUR AVERAGE					
-0.077±0.007±0.011	5037	ABOUZAID	06	KTEV	$K_L^0 \rightarrow \pi^+ \pi^- e^+ e^-$
-0.090±0.021		LAI	03C	NA48	$K_L^0 \rightarrow \pi^+ \pi^- e^+ e^-$
-0.054±0.026		MOLZON	78		K_S regen. by electrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
-0.087±0.046		BLATNIK	79		VMD + dispersion relations
-0.050±0.130		FOETH	69B		K_S regen. by electrons

T-VIOLATION PARAMETER IN K^0 - \bar{K}^0 MIXING

The asymmetry $A_T = \frac{\Gamma(\bar{K}^0 \rightarrow K^0) - \Gamma(K^0 \rightarrow \bar{K}^0)}{\Gamma(\bar{K}^0 \rightarrow K^0) + \Gamma(K^0 \rightarrow \bar{K}^0)}$ must vanish if T invariance holds.

ASYMMETRY A_T IN K^0 - \bar{K}^0 MIXING

<u>VALUE (units 10⁻³)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
6.6±1.3±1.0	640k	¹ ANGELOPO... 98E	CPLR

¹ ANGELOPOULOS 98E measures the asymmetry $A_T = [\Gamma(\bar{K}_{t=0}^0 \rightarrow e^+ \pi^- \nu_{t=\tau}) - \Gamma(K_{t=0}^0 \rightarrow e^- \pi^+ \bar{\nu}_{t=\tau})] / [\Gamma(\bar{K}_{t=0}^0 \rightarrow e^+ \pi^- \nu_{t=\tau}) + \Gamma(K_{t=0}^0 \rightarrow e^- \pi^+ \bar{\nu}_{t=\tau})]$ as a function of the neutral-kaon eigentime τ . The initial strangeness of the neutral kaon is tagged by the charge of the accompanying charged kaon in the reactions $p\bar{p} \rightarrow K^- \pi^+ K^0$ and $p\bar{p} \rightarrow K^+ \pi^- \bar{K}^0$. The strangeness at the time of the decay is tagged by the lepton charge. The reported result is the average value of A_T over the interval $1\tau_S < \tau < 20\tau_S$. From this value of A_T ANGELOPOULOS 01B, assuming CPT invariance in the $e\pi\nu$ decay amplitude, determine the T -violating as $\Delta S = \Delta S$ conserving parameter (for its definition, see Review below) $4\text{Re}(\epsilon) = (6.2 \pm 1.4 \pm 1.0) \times 10^{-3}$.

A REVIEW GOES HERE – Check our WWW List of Reviews

CP-VIOLATION PARAMETERS

Re(ϵ)

VALUE (units 10^{-3})	DOCUMENT ID	TECN
1.596 ± 0.013	² AMBROSINO 06H	KLOE

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.664 ± 0.010	³ LAI	05A NA48
---------------	------------------	----------

² AMBROSINO 06H uses Bell-Steinberger relations with the following measurements: $B(K_L^0 \rightarrow \pi^+ \pi^-)$ in AMBROSINO 06F, $B(K_S^0 \rightarrow \pi^0 \pi^0 \pi^0)$ in AMBROSINO 05B, the K_S^0 -semileptonic charge asymmetry in AMBROSINO 06E, and K^0 -semileptonic results in ANGELOPOULOS 98F.

³ LAI 05A values are obtained through unitarity (Bell-Steinberger relations), improving determination of η_{000} and combining other data from PDG 04 and APOSTOLAKIS 99B.

CPT-VIOLATION PARAMETERS

In K^0 - \bar{K}^0 mixing, if CP -violating interactions include a T conserving part then

$$|K_S\rangle = [|K_1\rangle + (\epsilon + \delta) |K_2\rangle] / \sqrt{1 + |\epsilon + \delta|^2}$$

$$|K_L\rangle = [|K_2\rangle + (\epsilon - \delta) |K_1\rangle] / \sqrt{1 + |\epsilon - \delta|^2}$$

where

$$|K_1\rangle = [|K^0\rangle + |\bar{K}^0\rangle] / \sqrt{2}$$

$$|K_2\rangle = [|K^0\rangle - |\bar{K}^0\rangle] / \sqrt{2}$$

and

$$|\bar{K}^0\rangle = CP |K^0\rangle.$$

The parameter δ specifies the CPT -violating part.

Estimates of δ are given below assuming the validity of the $\Delta S = \Delta Q$ rule. See also THOMSON 95 for a test of CPT -symmetry conservation in K^0 decays using the Bell-Steinberger relation.

REAL PART OF δ

A nonzero value violates CPT invariance.

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.51 ± 2.25		⁴ ABOUZAIID 11	KTEV	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.3 ± 2.7		⁵ AMBROSINO 06H	KLOE	
2.4 ± 2.8		⁶ APOSTOLA...	99B RVUE	
2.9 ± 2.6 ± 0.6	1.3M	⁷ ANGELOPO...	98F CPLR	
180 ± 200	6481	⁸ DEMIDOV 95		$K_{\ell 3}$ reanalysis

⁴ ABOUZAIID 11 uses Bell-Steinberger relations.

⁵ AMBROSINO 06H uses Bell-Steinberger relations with the following measurements: $B(K_L^0 \rightarrow \pi^+ \pi^-)$ in AMBROSINO 06F, $B(K_S^0 \rightarrow \pi^0 \pi^0 \pi^0)$ in AMBROSINO 05B, the K_S^0 -semileptonic charge asymmetry in AMBROSINO 06E, and K^0 -semileptonic results in ANGELOPOULOS 98F.

⁶ APOSTOLAKIS 99B assumes only unitarity and combines CPLEAR and other results.

⁷ ANGELOPOULOS 98F use $\Delta S = \Delta Q$. If $\Delta S = \Delta Q$ is not assumed, they find $\text{Re}\delta = (3.0 \pm 3.3 \pm 0.6) \times 10^{-4}$.

⁸ DEMIDOV 95 reanalyzes data from HART 73 and NIEBERGALL 74.

IMAGINARY PART OF δ

A nonzero value violates *CPT* invariance.

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
– 1.5± 1.6		⁹ ABOUZAIID	11	KTEV
••• We do not use the following data for averages, fits, limits, etc. •••				
0.4± 2.1		¹⁰ AMBROSINO	06H	KLOE
– 0.2± 2.0		¹¹ LAI	05A	NA48
2.4± 5.0		¹² APOSTOLA...	99B	RVUE
– 90 ± 290 ± 100	1.3M	¹³ ANGELOPO...	98F	CPLR
2100 ± 3700	6481	¹⁴ DEMIDOV	95	$K_{\ell 3}$ reanalysis

⁹ ABOUZAIID 11 uses Bell-Steinberger relations.

¹⁰ AMBROSINO 06H uses Bell-Steinberger relations with the following measurements: $B(K_L^0 \rightarrow \pi^+ \pi^-)$ in AMBROSINO 06F, $B(K_S^0 \rightarrow \pi^0 \pi^0 \pi^0)$ in AMBROSINO 05B, the K_S^0 -semileptonic charge asymmetry in AMBROSINO 06E, and K^0 -semileptonic results in ANGELOPOULOS 98F.

¹¹ LAI 05A values are obtained through unitarity (Bell-Steinberger relations), improving determination of η_{000} and combining other data from PDG 04 and APOSTOLAKIS 99B.

¹² APOSTOLAKIS 99B assumes only unitarity and combines CPLEAR and other results.

¹³ If $\Delta S = \Delta Q$ is not assumed, ANGELOPOULOS 98F finds $\text{Im}\delta = (-15 \pm 23 \pm 3) \times 10^{-3}$.

¹⁴ DEMIDOV 95 reanalyzes data from HART 73 and NIEBERGALL 74.

Re(y)

A non-zero value would violate *CPT* invariance in $\Delta S = \Delta Q$ amplitude. Re(y) is the following combination of K_{e3} decay amplitudes:

$$\text{Re}(y) = \text{Re} \left(\frac{A(\bar{K}^0 \rightarrow e^- \pi^+ \bar{\nu}_e)^* - A(K^0 \rightarrow e^+ \pi^- \nu_e)}{A(\bar{K}^0 \rightarrow e^- \pi^+ \bar{\nu}_e)^* + A(K^0 \rightarrow e^+ \pi^- \nu_e)} \right)$$

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
0.4±2.5	13k	¹⁵ AMBROSINO	06E KLOE

••• We do not use the following data for averages, fits, limits, etc. •••

0.3±3.1 ¹⁶ APOSTOLA... 99B CPLR

¹⁵ They use the PDG 04 for the K_L^0 semileptonic charge asymmetry and PDG 04 (*CP* review, *CPT* NOT ASSUMED) for Re(ϵ).

¹⁶ Constrained by Bell-Steinberger (or unitarity) relation.

Re(x₋)

A non-zero value would violate *CPT* invariance in decay amplitudes with $\Delta S \neq \Delta Q$. x_- , used here to define Re(x_-), and x_+ , used below in the $\Delta S = \Delta Q$ section are the following combinations of K_{e3} decay amplitudes:

$$x_{\pm} = \frac{1}{2} \left(\frac{A(\bar{K}^0 \rightarrow \pi^- e^+ \nu_e)}{A(K^0 \rightarrow \pi^- e^+ \nu_e)} \pm \frac{A(K^0 \rightarrow \pi^+ e^- \bar{\nu}_e)^*}{A(\bar{K}^0 \rightarrow \pi^+ e^- \bar{\nu}_e)^*} \right)$$

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-2.9± 2.0		¹⁷ AMBROSINO	06H	KLOE

••• We do not use the following data for averages, fits, limits, etc. •••

-0.8± 2.5 13k ¹⁸ AMBROSINO 06E KLOE

-0.5± 3.0 ¹⁹ APOSTOLA... 99B CPLR Strangeness tagged

2 ± 13 ± 3 650k ANGELOPO... 98F CPLR Strangeness tagged

- ¹⁷ AMBROSINO 06H uses Bell-Steinberger relations with the following measurements: $B(K_L^0 \rightarrow \pi^+ \pi^-)$ in AMBROSINO 06F, $B(K_S^0 \rightarrow \pi^0 \pi^0 \pi^0)$ in AMBROSINO 05B, the K_S^0 -semileptonic charge asymmetry in AMBROSINO 06E, and K^0 -semileptonic results in ANGELOPOULOS 98F.
¹⁸ Uses PDG 04 for the K_L^0 semileptonic charge asymmetry and $\text{Re}(\delta)$ from CPLEAR, ANGELOPOULOS 98F.
¹⁹ Constrained by Bell-Steinberger (or unitarity) relation.

$$|m_{K^0} - m_{\bar{K}^0}| / m_{\text{average}}$$

A test of *CPT* invariance. "Our Evaluation" is described in the "Tests of Conservation Laws" section. It assumes *CPT* invariance in the decay and neglects some contributions from decay channels other than $\pi\pi$.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
$<6 \times 10^{-19}$	90	PDG	12

• • • We do not use the following data for averages, fits, limits, etc. • • •

$(-3 \pm 4) \times 10^{-18}$ ²⁰ ANGELOPO... 99B RVUE

²⁰ ANGELOPOULOS 99B assumes only unitarity and combines CPLEAR and other results.

$$(\Gamma_{K^0} - \Gamma_{\bar{K}^0}) / m_{\text{average}}$$

A test of *CPT* invariance.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
$(7.8 \pm 8.4) \times 10^{-18}$	²¹ ANGELOPO... 99B	RVUE

²¹ ANGELOPOULOS 99B assumes only unitarity and combines CPLEAR with other results. Correlated with $(m_{K^0} - m_{\bar{K}^0}) / m_{\text{average}}$ with a correlation coefficient of -0.95 .

TESTS OF $\Delta S = \Delta Q$ RULE

$\text{Re}(x_+)$

A non-zero value would violate the $\Delta S = \Delta Q$ rule in *CPT* conserving transitions. x_+ is defined above in the $\text{Re}(x_-)$ section.

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
-0.9 ± 3.0 OUR AVERAGE			
-2 ± 10		²² BATLEY 07D	NA48
-0.5 ± 3.6	13k	²³ AMBROSINO 06E	KLOE
-1.8 ± 6.1		²⁴ ANGELOPO... 98D	CPLR

²² Result obtained from the measurement $\Gamma(K_S^0 \rightarrow \pi e \nu) / \Gamma(K_L^0 \rightarrow \pi e \nu) = 0.993 \pm 0.34$, neglecting possible *CPT* non-invariance and using PDG 06 values of $B(K_L^0 \rightarrow \pi e \nu) = 0.4053 \pm 0.0015$, $\tau_L = (5.114 \pm 0.021) \times 10^{-8}$ s and $\tau_S = (0.8958 \pm 0.0005) \times 10^{-10}$ s.

²³ $\text{Re}(x_+)$ can be shown to be equal to the following combination of rates:

$$\text{Re}(x_+) = \frac{1}{2} \frac{\Gamma(K_S^0 \rightarrow \pi e \nu) - \Gamma(K_L^0 \rightarrow \pi e \nu)}{\Gamma(K_S^0 \rightarrow \pi e \nu) + \Gamma(K_L^0 \rightarrow \pi e \nu)}$$

which is valid up to first order in terms violating *CPT* and/or the $\Delta S = \Delta Q$ rule.

²⁴ Obtained neglecting *CPT* violating amplitudes.

K^0 REFERENCES

PDG	12	PR D86 010001	J. Beringer <i>et al.</i>	(PDG Collab.)
ABOUZAID	11	PR D83 092001	E. Abouzaid <i>et al.</i>	(FNAL KTeV Collab.)
AMBROSINO	07B	JHEP 0712 073	F. Ambrosino <i>et al.</i>	(KLOE Collab.)
BATLEY	07D	PL B653 145	J.R. Batley <i>et al.</i>	(CERN NA48 Collab.)
ABOUZAID	06	PRL 96 101801	E. Abouzaid <i>et al.</i>	(KTeV Collab.)
AMBROSINO	06E	PL B636 173	F. Ambrosino <i>et al.</i>	(KLOE Collab.)
AMBROSINO	06F	PL B638 140	F. Ambrosino <i>et al.</i>	(KLOE Collab.)
AMBROSINO	06H	JHEP 0612 011	F. Ambrosino <i>et al.</i>	(KLOE Collab.)
PDG	06	JP G33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)
AMBROSINO	05B	PL B619 61	F. Ambrosino <i>et al.</i>	(KLOE Collab.)
LAI	05A	PL B610 165	A. Lai <i>et al.</i>	(CERN NA48 Collab.)
PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)
LAI	03C	EPJ C30 33	A. Lai <i>et al.</i>	(CERN NA48 Collab.)
LAI	02	PL B533 196	A. Lai <i>et al.</i>	(CERN NA48 Collab.)
ANGELOPO... 01B	EPJ C22 55	A. Angelopoulos <i>et al.</i>	(CPLEAR Collab.)	
ANGELOPO... 99B	PL B471 332	A. Angelopoulos <i>et al.</i>	(CPLEAR Collab.)	
APOSTOLA... 99B	PL B456 297	A. Apostolakis <i>et al.</i>	(CPLEAR Collab.)	
ANGELOPO... 98D	PL B444 38	A. Angelopoulos <i>et al.</i>	(CPLEAR Collab.)	
Also	EPJ C22 55	A. Angelopoulos <i>et al.</i>	(CPLEAR Collab.)	
ANGELOPO... 98E	PL B444 43	A. Angelopoulos <i>et al.</i>	(CPLEAR Collab.)	
ANGELOPO... 98F	PL B444 52	A. Angelopoulos <i>et al.</i>	(CPLEAR Collab.)	
Also	EPJ C22 55	A. Angelopoulos <i>et al.</i>	(CPLEAR Collab.)	
DEMIDOV	95	PAN 58 968	V. Demidov, K. Gusev, E. Shabalin	(ITEP)
From YAF	58 1041.			
THOMSON	95	PR D51 1412	G.B. Thomson, Y. Zou	(RUTG)
BARKOV	87B	SJNP 46 630	L.M. Barkov <i>et al.</i>	(NOVO)
BARKOV	85B	JETPL 42 138	L.M. Barkov <i>et al.</i>	(NOVO)
Translated from	YAF 46 1088.	ZETFP 42 113.		
BLATNIK	79	LNC 24 39	S. Blatnik, J. Stahov, C.B. Lang	(TUZL, GRAZ)
MOLZON	78	PRL 41 1213	W.R. Molzon <i>et al.</i>	(EFI+)
NIEBERGALL	74	PL 49B 103	F. Niebergall <i>et al.</i>	(CERN, ORSAY, VIEN)
HART	73	NP B66 317	J.C. Hart <i>et al.</i>	(CAVE, RHEL)
FOETH	69B	PL 30B 276	H. Foeth <i>et al.</i>	(AACH, CERN, TORI)
HILL	68B	PR 168 1534	D.G. Hill <i>et al.</i>	(BNL, CMU)
FITCH	67	PR 164 1711	V.L. Fitch <i>et al.</i>	(PRIN)
BALTAY	66	PR 142 932	C. Baltay <i>et al.</i>	(YALE, BNL)
BURNSTEIN	65	PR 138 B895	R.A. Burnstein, H.A. Rubin	(UMD)
KIM	65B	PR 140B 1334	J.K. Kim, L. Kirsch, D. Miller	(COLU)
CHRISTENS...	64	PRL 13 138	J.H. Christenson <i>et al.</i>	(PRIN)
CRAWFORD	59	PRL 2 112	F.S. Crawford <i>et al.</i>	(LRL)
ROSENFELD	59	PRL 2 110	A.H. Rosenfeld, F.T. Solmitz, R.D. Tripp	(LRL)