

**$f_1(1285)$**  $I^G(J^{PC}) = 0^+(1^{++})$  **$f_1(1285)$  MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>1281.9 ± 0.5 OUR AVERAGE</b>		Error includes scale factor of 1.8. See the ideogram below.			
1281.16 ± 0.39 ± 0.45		1 LEES	12X BABR	$\tau^- \rightarrow \pi^- f_1(1285) \nu_\tau$	
1285.1 ± 1.0 ± 1.6		2 ABLIKIM	11J BES3	$J/\psi \rightarrow \omega(\eta\pi^+\pi^-)$	
1281 ± 2 ± 1		AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow f_1(1285)\pi^+\pi^-\gamma$	
1276.1 ± 8.1 ± 8.0	203	BAI	04J BES2	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$	
1274 ± 6	237	ABDALLAH	03H DLPH	$91.2 e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp + X$	
1280 ± 4		ACCIARRI	01G L3		
1288 ± 4 ± 5	20k	ADAMS	01B B852	$18 \text{ GeV } \pi^- p \rightarrow K^+ K^- \pi^0 n$	
1284 ± 6	1400	ALDE	97B GAM4	$100 \pi^- p \rightarrow \eta\pi^0\pi^0 n$	
1281 ± 1		BARBERIS	97B OMEG	$450 pp \rightarrow pp2(\pi^+\pi^-)$	
1281 ± 1		BARBERIS	97C OMEG	$450 pp \rightarrow ppK_S^0 K^\pm \pi^\mp$	
1280 ± 2		3 ANTINORI	95 OMEG	$300,450 pp \rightarrow pp2(\pi^+\pi^-)$	
1282.2 ± 1.5		LEE	94 MPS2	$18 \pi^- p \rightarrow K^+ \bar{K}^0 2\pi^- p$	
1279 ± 5		FUKUI	91C SPEC	$8.95 \pi^- p \rightarrow \eta\pi^+\pi^- n$	
1278 ± 2	140	ARMSTRONG	89 OMEG	$300 pp \rightarrow K\bar{K}\pi pp$	
1278 ± 2		ARMSTRONG	89G OMEG	$85 \pi^+ p \rightarrow 4\pi\pi p, pp \rightarrow 4\pi pp$	
1280.1 ± 2.1	60	RATH	89 MPS	$21.4 \pi^- p \rightarrow K_S^0 K_S^0 \pi^0 n$	
1285 ± 1	4750	4 BIRMAN	88 MPS	$8 \pi^- p \rightarrow K^+ \bar{K}^0 \pi^- n$	
1280 ± 1	504	BITYUKOV	88 SPEC	$32.5 \pi^- p \rightarrow K^+ K^- \pi^0 n$	
1280 ± 4		ANDO	86 SPEC	$8 \pi^- p \rightarrow \eta\pi^+\pi^- n$	
1277 ± 2	420	REEVES	86 SPEC	$6.6 p\bar{p} \rightarrow K\bar{K}\pi X$	
1285 ± 2		CHUNG	85 SPEC	$8 \pi^- p \rightarrow N\bar{K}\bar{K}\pi$	
1279 ± 2	604	ARMSTRONG	84 OMEG	$85 \pi^+ p \rightarrow K\bar{K}\pi\pi p, pp \rightarrow K\bar{K}\pi pp$	
1286 ± 1		CHAUVAT	84 SPEC	ISR 31.5 pp	
1278 ± 4		EVANGELIS...	81 OMEG	$12 \pi^- p \rightarrow \eta\pi^+\pi^-\pi^- p$	
1283 ± 3	103	DIONISI	80 HBC	$4 \pi^- p \rightarrow K\bar{K}\pi n$	
1282 ± 2	320	NACASCH	78 HBC	$0.7, 0.76 \bar{p}p \rightarrow K\bar{K}3\pi$	
1279 ± 5	210	GRASSLER	77 HBC	$16 \pi^\mp p$	
1286 ± 3	180	DUBOC	72 HBC	$1.2 \bar{p}p \rightarrow 2K4\pi$	
1283 ± 5		DAHL	67 HBC	$1.6-4.2 \pi^- p$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

1281.9 $\pm$ 0.5	<sup>5</sup> SOSA	99	SPEC	$p p \rightarrow p_{\text{slow}} (K_S^0 K^+ \pi^-) p_{\text{fast}}$
1282.8 $\pm$ 0.6	<sup>5</sup> SOSA	99	SPEC	$p p \rightarrow p_{\text{slow}} (K_S^0 K^- \pi^+) p_{\text{fast}}$
1270 $\pm$ 10	AMELIN	95	VES	$37 \pi^- N \rightarrow \pi^- \pi^+ \pi^- \gamma N$
1280 $\pm$ 2	ABATZIS	94	OMEG	$450 p p \rightarrow p p 2(\pi^+ \pi^-)$
1282 $\pm$ 4	ARMSTRONG	93C	E760	$\bar{p} p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$
1270 $\pm$ 6 $\pm$ 10	ARMSTRONG	92C	OMEG	$300 p p \rightarrow p p \pi^+ \pi^- \gamma$
1281 $\pm$ 1	ARMSTRONG	89E	OMEG	$300 p p \rightarrow p p 2(\pi^+ \pi^-)$
1279 $\pm$ 6 $\pm$ 10	BECKER	87	MRK3	$e^+ e^- \rightarrow \phi K \bar{K} \pi$
1286 $\pm$ 9	GIDAL	87	MRK2	$e^+ e^- \rightarrow e^+ e^- \eta \pi^+ \pi^-$
1287 $\pm$ 5 $\sim 1279$	BITYUKOV <sup>6</sup> TORNQVIST	353 82B	SPEC RVUE	$32 \pi^- p \rightarrow K^+ K^- \pi^0 n$
1275 $\pm$ 6	BROMBERG	31	SPEC	$100 \pi^- p \rightarrow K \bar{K} \pi X$
1288 $\pm$ 9	GURTU	200	HBC	$4.2 K^- p \rightarrow n \eta 2\pi$
$\sim 1275.0$	STANTON	46	CNTR	$8.5 \pi^- p \rightarrow n 2\gamma 2\pi$
1271 $\pm$ 10	CORDEN	34	OMEG	$12-15 \pi^- p \rightarrow K^+ K^- \pi n$
1295 $\pm$ 12	CORDEN	85	OMEG	$12-15 \pi^- p \rightarrow n 5\pi$
1292 $\pm$ 10	DEFOIX	150	HBC	$0.7 \bar{p} p \rightarrow 7\pi$
1280 $\pm$ 3	THUN	500	MMS	$13.4 \pi^- p$
1303 $\pm$ 8	BARDADIN-...	71	HBC	$8 \pi^+ p \rightarrow p 6\pi$
1283 $\pm$ 6	BOESEBECK	71	HBC	$16.0 \pi p \rightarrow p 5\pi$
1270 $\pm$ 10	CAMPBELL	69	DBC	$2.7 \pi^+ d$
1285 $\pm$ 7	LORSTAD	69	HBC	$0.7 \bar{p} p$ , 4,5-body
1290 $\pm$ 7	D'ANDLAU	68	HBC	$1.2 \bar{p} p$ , 5-6 body

<sup>1</sup> Using the  $2\pi^+ 2\pi^-$  and  $\pi^+ \pi^- \eta$  modes of  $f_1(1285)$  decay.

<sup>2</sup> The selected process is  $J/\psi \rightarrow \omega a_0(980)\pi$ .

<sup>3</sup> Supersedes ABATZIS 94, ARMSTRONG 89E.


<sup>4</sup> From partial wave analysis of  $K^+ \bar{K}^0 \pi^-$  system.

<sup>5</sup> No systematic error given.

<sup>6</sup> From a unitarized quark-model calculation.

<sup>7</sup> From phase shift analysis of  $\eta \pi^+ \pi^-$  system.

<sup>8</sup> Seen in the missing mass spectrum.



$f_1(1285)$  mass (MeV)

### $f_1(1285)$ WIDTH

Only experiments giving width error less than 20 MeV are kept for averaging.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>24.2± 1.1 OUR AVERAGE</b>				Error includes scale factor of 1.3. See the ideogram below.
$22.0 \pm 3.1 \pm 2.0$	9	ABLIKIM	11J BES3	$J/\psi \rightarrow \omega(\eta\pi^+\pi^-)$
$35 \pm 6 \pm 4$		AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow f_1(1285)\pi^+\pi^-\gamma$
$40.0 \pm 8.6 \pm 9.3$	203	BAI	04J BES2	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
$29 \pm 12$	237	ABDALLAH	03H DLPH	$91.2 e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp + X$
$45 \pm 9 \pm 7$	20k	ADAMS	01B B852	$18 \text{ GeV } \pi^- p \rightarrow K^+ K^- \pi^0 n$

55 $\pm$ 18	1400	ALDE	97B	GAM4	$100 \pi^- p \rightarrow \eta \pi^0 \pi^0 n$
24 $\pm$ 3		BARBERIS	97B	OMEG	$450 pp \rightarrow pp2(\pi^+ \pi^-)$
20 $\pm$ 2		BARBERIS	97C	OMEG	$450 pp \rightarrow ppK_S^0 K^\pm \pi^\mp$
36 $\pm$ 5	10	ANTINORI	95	OMEG	$300,450 pp \rightarrow pp2(\pi^+ \pi^-)$
29.0 $\pm$ 4.1		LEE	94	MPS2	$18 \pi^- p \rightarrow K^+ \bar{K}^0 2\pi^- p$
25 $\pm$ 4	140	ARMSTRONG	89	OMEG	$300 pp \rightarrow K\bar{K}\pi pp$
22 $\pm$ 2	4750	11 BIRMAN	88	MPS	$8 \pi^- p \rightarrow K^+ \bar{K}^0 \pi^- n$
25 $\pm$ 4	504	BITYUKOV	88	SPEC	$32.5 \pi^- p \rightarrow K^+ K^- \pi^0 n$
19 $\pm$ 5		ANDO	86	SPEC	$8 \pi^- p \rightarrow \eta \pi^+ \pi^- n$
32 $\pm$ 8	420	REEVES	86	SPEC	$6.6 p\bar{p} \rightarrow K K \pi X$
22 $\pm$ 2		CHUNG	85	SPEC	$8 \pi^- p \rightarrow N K \bar{K} \pi$
32 $\pm$ 3	604	ARMSTRONG	84	OMEG	$85 \pi^+ p \rightarrow K\bar{K}\pi\pi p, pp \rightarrow K\bar{K}\pi pp$
24 $\pm$ 3		CHAUVAT	84	SPEC	ISR 31.5 $pp$
29 $\pm$ 10	103	DIONISI	80	HBC	$4 \pi^- p \rightarrow K\bar{K}\pi n$
28.3 $\pm$ 6.7	320	NACASCH	78	HBC	$0.7, 0.76 \bar{p}p \rightarrow K\bar{K}3\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

18.2 $\pm$ 1.2	12 SOSA	99	SPEC	$pp \rightarrow p_{\text{slow}} (K_S^0 K^+ \pi^-)$	
19.4 $\pm$ 1.5	12 SOSA	99	SPEC	$pp \rightarrow p_{\text{slow}} (K_S^0 K^- \pi^+)$	
40 $\pm$ 5	ABATZIS	94	OMEG	$450 pp \rightarrow pp2(\pi^+ \pi^-)$	
31 $\pm$ 5	ARMSTRONG	89E	OMEG	$300 pp \rightarrow pp2(\pi^+ \pi^-)$	
41 $\pm$ 12	ARMSTRONG	89G	OMEG	$85 \pi^+ p \rightarrow 4\pi\pi p, pp \rightarrow 4\pi pp$	
17.9 $\pm$ 10.9	60	RATH	89	MPS	$21.4 \pi^- p \rightarrow K_S^0 K_S^0 \pi^0 n$
14 $\pm$ 10	16	BECKER	87	MRK3	$e^+ e^- \rightarrow \phi K\bar{K}\pi$
26 $\pm$ 12		EVANGELIS...	81	OMEG	$12 \pi^- p \rightarrow \eta \pi^+ \pi^- \pi^- p$
25 $\pm$ 15	200	GURTU	79	HBC	$4.2 K^- p \rightarrow n\eta 2\pi$
$\sim 10$	13 STANTON	79	CNTR	$8.5 \pi^- p \rightarrow n2\gamma 2\pi$	
24 $\pm$ 18	210	GRASSLER	77	HBC	$16 \pi^\mp p$
28 $\pm$ 5	150	14 DEFOIX	72	HBC	$0.7 \bar{p}p \rightarrow 7\pi$
46 $\pm$ 9	180	14 DUBOC	72	HBC	$1.2 \bar{p}p \rightarrow 2K4\pi$
37 $\pm$ 5	500	15 THUN	72	MMS	$13.4 \pi^- p$
10 $\pm$ 10		BOESEBECK	71	HBC	$16.0 \pi p \rightarrow p5\pi$
30 $\pm$ 15		CAMPBELL	69	DBC	$2.7 \pi^+ d$
60 $\pm$ 15	14 LORSTAD	69	HBC	$0.7 \bar{p}p, 4,5\text{-body}$	
35 $\pm$ 10	14 DAHL	67	HBC	$1.6\text{--}4.2 \pi^- p$	

<sup>9</sup> The selected process is  $J/\psi \rightarrow \omega a_0(980)\pi$ .

<sup>10</sup> Supersedes ABATZIS 94, ARMSTRONG 89E.


<sup>11</sup> From partial wave analysis of  $K^+ \bar{K}^0 \pi^-$  system.

<sup>12</sup> No systematic error given.

<sup>13</sup> From phase shift analysis of  $\eta \pi^+ \pi^-$  system.

<sup>14</sup> Resolution is not unfolded.

<sup>15</sup> Seen in the missing mass spectrum.



$f_1(1285)$  width (MeV)

### $f_1(1285)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level
$\Gamma_1 \quad 4\pi$	$(33.1 \pm 2.1) \%$	$S=1.3$
$\Gamma_2 \quad \pi^0 \pi^0 \pi^+ \pi^-$	$(22.0 \pm 1.4) \%$	$S=1.3$
$\Gamma_3 \quad 2\pi^+ 2\pi^-$	$(11.0 \pm 0.7) \%$	$S=1.3$
$\Gamma_4 \quad \rho^0 \pi^+ \pi^-$	$(11.0 \pm 0.7) \%$	$S=1.3$
$\Gamma_5 \quad \rho^0 \rho^0$	seen	
$\Gamma_6 \quad 4\pi^0$	$< 7 \times 10^{-4}$	$CL=90\%$
$\Gamma_7 \quad \eta \pi^+ \pi^-$	$(35 \pm 15) \%$	
$\Gamma_8 \quad \eta \pi \pi$	$(52.4 \pm 1.9) \%$	$S=1.2$
$\Gamma_9 \quad a_0(980)\pi$ [ignoring $a_0(980) \rightarrow K\bar{K}$ ]	$(36 \pm 7) \%$	
$\Gamma_{10} \quad \eta \pi \pi$ [excluding $a_0(980)\pi$ ]	$(16 \pm 7) \%$	
$\Gamma_{11} \quad K\bar{K}\pi$	$(9.0 \pm 0.4) \%$	$S=1.1$

$\Gamma_{12}$	$K\bar{K}^*(892)$	not seen
$\Gamma_{13}$	$\pi^+\pi^-\pi^0$	$(3.0 \pm 0.9) \times 10^{-3}$
$\Gamma_{14}$	$\rho^\pm\pi^\mp$	$< 3.1 \times 10^{-3}$ CL=95%
$\Gamma_{15}$	$\gamma\rho^0$	$(5.5 \pm 1.3)\%$ S=2.8
$\Gamma_{16}$	$\phi\gamma$	$(7.4 \pm 2.6) \times 10^{-4}$
$\Gamma_{17}$	$\gamma\gamma^*$	
$\Gamma_{18}$	$\gamma\gamma$	

## CONSTRAINED FIT INFORMATION

An overall fit to 7 branching ratios uses 16 measurements and one constraint to determine 5 parameters. The overall fit has a  $\chi^2 = 24.7$  for 12 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$ , in percent, from the fit to the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

$x_9$	-17			
$x_{10}$	-8	-95		
$x_{11}$	46	-9	-4	
$x_{15}$	-36	-4	-2	-34
	$x_1$	$x_9$	$x_{10}$	$x_{11}$

### $f_1(1285) \Gamma(i) \Gamma(\gamma\gamma) / \Gamma(\text{total})$

$\Gamma(\eta\pi\pi) \times \Gamma(\gamma\gamma) / \Gamma_{\text{total}}$		$\Gamma_8 \Gamma_{18} / \Gamma = (\Gamma_9 + \Gamma_{10}) \Gamma_{18} / \Gamma$		
VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.62	95	GIDAL	87	MRK2 $e^+ e^- \rightarrow e^+ e^- \eta\pi^+\pi^-$

$\Gamma(\eta\pi\pi) \times \Gamma(\gamma\gamma^*) / \Gamma_{\text{total}}$		$\Gamma_8 \Gamma_{17} / \Gamma = (\Gamma_9 + \Gamma_{10}) \Gamma_{17} / \Gamma$		
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.4 ± 0.4 OUR AVERAGE</b>	Error includes scale factor of 1.4.			
1.18 ± 0.25 ± 0.20	26	16,17 AIHARA	88B TPC	$e^+ e^- \rightarrow e^+ e^- \eta\pi^+\pi^-$
2.30 ± 0.61 ± 0.42		16,18 GIDAL	87 MRK2	$e^+ e^- \rightarrow e^+ e^- \eta\pi^+\pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.8 ± 0.3 ± 0.3	420	19 ACHARD	02B L3	$183-209 e^+ e^- \rightarrow e^+ e^- \eta\pi^+\pi^-$

<sup>16</sup> Assuming a  $\rho$ -pole form factor.

<sup>17</sup> Published value multiplied by  $\eta\pi\pi$  branching ratio 0.49.

<sup>18</sup> Published value divided by 2 and multiplied by the  $\eta\pi\pi$  branching ratio 0.49.

<sup>19</sup> Published value multiplied by the  $\eta\pi\pi$  branching ratio 0.52.

**$f_1(1285)$  BRANCHING RATIOS** **$\Gamma(K\bar{K}\pi)/\Gamma(4\pi)$** 

$$\Gamma_{11}/\Gamma_1$$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.271±0.016 OUR FIT</b>	Error includes scale factor of 1.3.		
<b>0.271±0.016 OUR AVERAGE</b>	Error includes scale factor of 1.2.		
0.265±0.014	20 BARBERIS 97C OMEG 450 $p p \rightarrow p p K_S^0 K^\pm \pi^\mp$		
0.28 ± 0.05	21 ARMSTRONG 89E OMEG 300 $p p \rightarrow p p f_1(1285)$		
0.37 ± 0.03 ± 0.05	22 ARMSTRONG 89G OMEG 85 $\pi p \rightarrow 4\pi X$		
20 Using $2(\pi^+ \pi^-)$ data from BARBERIS 97B.			
21 Assuming $\rho\pi\pi$ and $a_0(980)\pi$ intermediate states.			
22 $4\pi$ consistent with being entirely $\rho\pi\pi$ .			

 **$\Gamma(\pi^0\pi^+\pi^-)/\Gamma_{\text{total}}$** 

$$\Gamma_2/\Gamma = \frac{2}{3}\Gamma_1/\Gamma$$

VALUE	DOCUMENT ID
<b>0.220<sup>+0.014</sup><sub>-0.012</sub> OUR FIT</b>	Error includes scale factor of 1.3.

 **$\Gamma(2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$** 

$$\Gamma_3/\Gamma = \frac{1}{3}\Gamma_1/\Gamma$$

VALUE	DOCUMENT ID
<b>0.110<sup>+0.007</sup><sub>-0.006</sub> OUR FIT</b>	Error includes scale factor of 1.3.

 **$\Gamma(\rho^0\pi^+\pi^-)/\Gamma_{\text{total}}$** 

$$\Gamma_4/\Gamma = \frac{1}{3}\Gamma_1/\Gamma$$

VALUE	DOCUMENT ID
<b>0.110<sup>+0.007</sup><sub>-0.006</sub> OUR FIT</b>	Error includes scale factor of 1.3.

 **$\Gamma(\rho^0\pi^+\pi^-)/\Gamma(2\pi^+ 2\pi^-)$** 

$$\Gamma_4/\Gamma_3$$

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.0±0.4	GRASSLER 77 HBC 16 GeV $\pi^\pm p$		

 **$\Gamma(\rho^0\rho^0)/\Gamma_{\text{total}}$** 

$$\Gamma_5/\Gamma$$

VALUE	DOCUMENT ID	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •		
seen	BARBERIS 00C 450 $p p \rightarrow p_f 4\pi p_s$	

 **$\Gamma(4\pi^0)/\Gamma_{\text{total}}$** 

$$\Gamma_6/\Gamma$$

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<7	90	ALDE 87	GAM4	100 $\pi^- p \rightarrow 4\pi^0 n$

 **$\Gamma(\pi^+\pi^-\pi^0)/\Gamma(\eta\pi^+\pi^-)$** 

$$\Gamma_{13}/\Gamma_7$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.86±0.16±0.20</b>	2.3k	23 DOROFEEV 11 VES		$\pi^- N \rightarrow \pi^- f_1(1285) N$

<sup>23</sup> Value obtained selecting the region corresponding to  $f_0(980)$  in the  $\pi^+\pi^-$  mass spectrum.

$\Gamma(\eta\pi\pi)/\Gamma_{\text{total}}$ VALUE**0.524<sup>+0.019</sup><sub>-0.022</sub> OUR FIT** Error includes scale factor of 1.2.DOCUMENT ID $\Gamma_8/\Gamma = (\Gamma_9 + \Gamma_{10})/\Gamma$  $\Gamma(4\pi)/\Gamma(\eta\pi\pi)$ VALUEDOCUMENT IDTECNCOMMENT**0.63<sup>+0.06</sup><sub>-0.06</sub> OUR FIT** Error includes scale factor of 1.2.**0.41<sup>+0.14</sup><sub>-0.14</sub> OUR AVERAGE** $\Gamma_1/\Gamma_8 = \Gamma_1/(\Gamma_9 + \Gamma_{10})$ 0.37 $\pm$ 0.11 $\pm$ 0.11 BOLTON 92 MRK3  $J/\psi \rightarrow \gamma f_1(1285)$ 0.64 $\pm$ 0.40 GURTU 79 HBC 4.2  $K^- p$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.93 $\pm$ 0.30 <sup>24</sup> GRASSLER 77 HBC 16  $\pi^\mp p$ <sup>24</sup> Assuming  $\rho\pi\pi$  and  $a_0(980)\pi$  intermediate states. $\Gamma(2\pi^+ 2\pi^-)/\Gamma(\eta\pi\pi)$ VALUEDOCUMENT IDTECNCOMMENT**0.28<sup>+0.02</sup><sub>-0.02</sub><sup>+0.02</sup><sub>-0.02</sub>** <sup>25</sup> LEES 12X BABR  $\tau^- \rightarrow \pi^- f_1(1285) \nu_\tau$ <sup>25</sup> Assuming  $B(f_1(1285) \rightarrow \pi\pi\eta) = 3/2 B(f_1(1285) \rightarrow \pi^+\pi^-\eta)$ . $\Gamma_3/\Gamma_8$  $\Gamma(a_0(980)\pi [\text{ignoring } a_0(980) \rightarrow K\bar{K}])/\Gamma(\eta\pi\pi)$  $\Gamma_9/\Gamma_8 = \Gamma_9/(\Gamma_9 + \Gamma_{10})$ VALUECL%DOCUMENT IDTECNCOMMENT**0.69<sup>+0.13</sup><sub>-0.12</sub> OUR FIT****0.69<sup>+0.13</sup><sub>-0.12</sub> OUR AVERAGE**0.72 $\pm$ 0.15 GURTU 79 HBC 4.2  $K^- p$ 0.6 $^{+0.3}_{-0.2}$  CORDEN 78 OMEG 12–15  $\pi^- p$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

>0.69 95 318 ACHARD 02B L3  $183\text{--}209 e^+ e^- \rightarrow e^+ e^- \eta\pi^+\pi^-$ 0.28 $\pm$ 0.07 1400 ALDE 97B GAM4 100  $\pi^- p \rightarrow \eta\pi^0\pi^0 n$ 1.0  $\pm$ 0.3 GRASSLER 77 HBC 16  $\pi^\mp p$  $\Gamma(K\bar{K}\pi)/\Gamma(\eta\pi\pi)$  $\Gamma_{11}/\Gamma_8 = \Gamma_{11}/(\Gamma_9 + \Gamma_{10})$ VALUEDOCUMENT IDTECNCOMMENT**0.171<sup>+0.013</sup><sub>-0.013</sub> OUR FIT** Error includes scale factor of 1.1.**0.170<sup>+0.012</sup><sub>-0.012</sub> OUR AVERAGE**0.166 $\pm$ 0.01  $\pm$ 0.008 BARBERIS 98C OMEG 450  $p p \rightarrow p_f f_1(1285) p_s$ 0.42  $\pm$ 0.15 GURTU 79 HBC 4.2  $K^- p$ 0.5  $\pm$ 0.2 <sup>26</sup> CORDEN 78 OMEG 12–15  $\pi^- p$ 0.20  $\pm$ 0.08 <sup>27</sup> DEFOIX 72 HBC 0.7  $\bar{p}p \rightarrow 7\pi$ 0.16  $\pm$ 0.08 CAMPBELL 69 DBC 2.7  $\pi^+ d$ <sup>26</sup> CORDEN 78 assumes low-mass  $\eta\pi\pi$  region is dominantly  $1^{++}$ . See BARBERIS 98C and MANAK 00A for discussion.<sup>27</sup>  $K\bar{K}$  system characterized by the  $I = 1$  threshold enhancement. (See under  $a_0(980)$ ).

$\Gamma(K\bar{K}^*(892))/\Gamma_{\text{total}}$   $\Gamma_{12}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>not seen</b>	NACASCH 78	HBC	$0.7, 0.76 \bar{p}p \rightarrow K\bar{K}3\pi$
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>			
seen	28 ACHARD 07 L3		$183-209 e^+e^- \rightarrow e^+e^- K_S^0 K^\pm \pi^\mp$
28 A clear signal of $19.8 \pm 4.4$ events observed at high $Q^2$ .			

 $\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{13}/\Gamma$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.30±0.055±0.074</b>	2.3k	29 DOROFEEV	11 VES	$\pi^- N \rightarrow \pi^- f_1(1285) N$
29 Value obtained selecting the region corresponding to $f_0(980)$ in the $\pi^+\pi^-$ mass spectrum. The systematic error includes the uncertainty on the partial width $f_1 \rightarrow \eta\pi\pi$ obtained from PDG 10 data.				

 $\Gamma(\rho^\pm\pi^\mp)/\Gamma_{\text{total}}$   $\Gamma_{14}/\Gamma$ 

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.31</b>	95	DOROFEEV	11 VES	$\pi^- N \rightarrow \pi^- f_1(1285) N$

 $\Gamma(\gamma\rho^0)/\Gamma_{\text{total}}$   $\Gamma_{15}/\Gamma$ 

VALUE (units $10^{-2}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>5.5±1.3 OUR FIT</b>		Error includes scale factor of 2.8.		
<b>2.8±0.7±0.6</b>		AMELIN	95 VES	$37 \pi^- N \rightarrow \pi^-\pi^+\pi^-\gamma N$
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>				
<5	95	BITYUKOV	91B SPEC	$32 \pi^- p \rightarrow \pi^+\pi^-\gamma n$

 $\Gamma(\gamma\rho^0)/\Gamma(2\pi^+2\pi^-)$   $\Gamma_{15}/\Gamma_3 = \Gamma_{15}/\frac{1}{3}\Gamma_1$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.50±0.13 OUR FIT</b>	Error includes scale factor of 2.5.		
<b>0.45±0.18</b>	30 COFFMAN	90 MRK3	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
30 Using $B(J/\psi \rightarrow \gamma f_1(1285) \rightarrow \gamma\gamma\rho^0) = 0.25 \times 10^{-4}$ and $B(J/\psi \rightarrow \gamma f_1(1285) \rightarrow \gamma 2\pi^+ 2\pi^-) = 0.55 \times 10^{-4}$ given by MIR 88.			

 $\Gamma(\eta\pi\pi)/\Gamma(\gamma\rho^0)$   $\Gamma_8/\Gamma_{15} = (\Gamma_9 + \Gamma_{10})/\Gamma_{15}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>9.5±2.0 OUR FIT</b>	Error includes scale factor of 2.5.		
<b>7.9±0.9 OUR AVERAGE</b>			

10.0±1.0±2.0	BARBERIS	98C OMEG	$450 \bar{p}p \rightarrow p_f f_1(1285) p_s$
7.5±1.0	31 ARMSTRONG	92C OMEG	$300 \bar{p}p \rightarrow pp\pi^+\pi^-\gamma, pp\eta\pi^+\pi^-$

31 Published value multiplied by 1.5.

 $\Gamma(\gamma\rho^0)/\Gamma(K\bar{K}\pi)$   $\Gamma_{15}/\Gamma_{11}$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>				
>0.035	90	32 COFFMAN	90 MRK3	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
32 Using $B(J/\psi \rightarrow \gamma f_1(1285) \rightarrow \gamma\gamma\rho^0) = 0.25 \times 10^{-4}$ and $B(J/\psi \rightarrow \gamma f_1(1285) \rightarrow \gamma K\bar{K}\pi) < 0.72 \times 10^{-3}$ .				

$\Gamma(\phi\gamma)/\Gamma(K\bar{K}\pi)$  $\Gamma_{16}/\Gamma_{11}$ 

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.82±0.21±0.20</b>		19	BITYUKOV	88	SPEC $32.5 \pi^- p \rightarrow K^+ K^- \pi^0 n$
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>					
<0.50	95		BARBERIS	98C	OMEG $450 pp \rightarrow p_f f_1(1285) p_s$
<0.93	95		AMELIN	95	VES $37 \pi^- N \rightarrow \pi^- \pi^+ \pi^- \gamma N$

**f<sub>1</sub>(1285) REFERENCES**

LEES	12X	PR D86 092010	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	11J	PRL 107 182001	M. Ablikim <i>et al.</i>	(BES III Collab.)
DOROFEEV	11	EPJ A47 68	V. Dorofeev <i>et al.</i>	(SERP, MIPT)
PDG	10	JP G37 075021	K. Nakamura <i>et al.</i>	(PDG Collab.)
ACHARD	07	JHEP 0703 018	P. Achard <i>et al.</i>	(L3 Collab.)
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)
BAI	04J	PL B594 47	J.Z. Bai <i>et al.</i>	(BES Collab.)
ABDALLAH	03H	PL B569 129	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
ACHARD	02B	PL B526 269	P. Achard <i>et al.</i>	(L3 Collab.)
ACCIARRI	01G	PL B501 1	M. Acciarri <i>et al.</i>	(L3 Collab.)
ADAMS	01B	PL B516 264	G.S. Adams <i>et al.</i>	(BNL E852 Collab.)
BARBERIS	00C	PL B471 440	D. Barberis <i>et al.</i>	(WA 102 Collab.)
MANAK	00A	PR D62 012003	J.J. Manak <i>et al.</i>	(BNL E852 Collab.)
SOSA	99	PRL 83 913	M. Sosa <i>et al.</i>	
BARBERIS	98C	PL B440 225	D. Barberis <i>et al.</i>	(WA 102 Collab.)
ALDE	97B	PAN 60 386	D. Alde <i>et al.</i>	(GAMS Collab.)
		Translated from YAF 60	458.	
BARBERIS	97B	PL B413 217	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BARBERIS	97C	PL B413 225	D. Barberis <i>et al.</i>	(WA 102 Collab.)
AMELIN	95	ZPHY C66 71	D.V. Amelin <i>et al.</i>	(VES Collab.)
ANTINORI	95	PL B353 589	F. Antinori <i>et al.</i>	(ATHU, BARI, BIRM+)
ABATZIS	94	PL B324 509	S. Abatzis <i>et al.</i>	(ATHU, BARI, BIRM+)
LEE	94	PL B323 227	J.H. Lee <i>et al.</i>	(BNL, IND, KYUN, MASD+)
ARMSTRONG	93C	PL B307 394	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
ARMSTRONG	92C	ZPHY C54 371	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)
BOLTON	92	PL B278 495	T. Bolton <i>et al.</i>	(Mark III Collab.)
BITYUKOV	91B	SJNP 54 318	S.I. Bityukov <i>et al.</i>	(SERP)
		Translated from YAF 54	529.	
FUKUI	91C	PL B267 293	S. Fukui <i>et al.</i>	(SUGI, NAGO, KEK, KYOT+)
COFFMAN	90	PR D41 1410	D.M. Coffman <i>et al.</i>	(Mark III Collab.)
ARMSTRONG	89	PL B221 216	T.A. Armstrong <i>et al.</i>	(CERN, CDEF, BIRM+) JPC
ARMSTRONG	89E	PL B228 536	T.A. Armstrong, M. Benayoun	(ATHU, BARI, BIRM+)
ARMSTRONG	89G	ZPHY C43 55	T.A. Armstrong <i>et al.</i>	(CERN, BIRM, BARI+)
RATH	89	PR D40 693	M.G. Rath <i>et al.</i>	(NDAM, BRAN, BNL, CUNY+)
AIHARA	88B	PL B209 107	H. Aihara <i>et al.</i>	(TPC-2 $\gamma$ Collab.)
BIRMAN	88	PRL 61 1557	A. Birman <i>et al.</i>	(BNL, FSU, IND, MASD) JP
BITYUKOV	88	PL B203 327	S.I. Bityukov <i>et al.</i>	(SERP)
MIR	88	Photon-Photon 88, 126	R. Mir	(Mark III Collab.)
Conference				
ALDE	87	PL B198 286	D.M. Alde <i>et al.</i>	(LANL, BRUX, SERP, LAPP)
BECKER	87	PRL 59 186	J.J. Becker <i>et al.</i>	(Mark III Collab.)
GIDAL	87	PRL 59 2012	G. Gidal <i>et al.</i>	(LBL, SLAC, HARV)
ANDO	86	PRL 57 1296	A. Ando <i>et al.</i>	(KEK, KYOT, NIR, SAGA+) IJP
REEVES	86	PR D34 1960	D.F. Reeves <i>et al.</i>	(FLOR, BNL, IND+) JP
CHUNG	85	PRL 55 779	S.U. Chung <i>et al.</i>	(BNL, FLOR, IND+) JP
ARMSTRONG	84	PL 146B 273	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+) JP
BITYUKOV	84B	PL 144B 133	S.I. Bityukov <i>et al.</i>	(SERP)
CHAUVAT	84	PL 148B 382	P. Chauvat <i>et al.</i>	(CERN, CLER, UCLA+)
TORNQVIST	82B	NP B203 268	N.A. Tornqvist	(HELS)
EVANGELIS...	81	NP B178 197	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)
BROMBERG	80	PR D22 1513	C.M. Bromberg <i>et al.</i>	(CIT, FNAL, ILLC+)
DIONISI	80	NP B169 1	C. Dionisi <i>et al.</i>	(CERN, MADR, CDEF+)
GURTU	79	NP B151 181	A. Gurtu <i>et al.</i>	(CERN, ZEEM, NIJM, OXF)

STANTON	79	PRL 42 346	N.R. Stanton <i>et al.</i>	(OSU, CARL, MCGI+) JP
CORDEN	78	NP B144 253	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+) JP
NACASCH	78	NP B135 203	R. Nacasch <i>et al.</i>	(PARIS, MADR, CERN)
GRASSLER	77	NP B121 189	H. Grassler <i>et al.</i>	(AACH3, BERL, BONN+)
DEFOIX	72	NP B44 125	C. Defoix <i>et al.</i>	(CDEF, CERN)
DUBOC	72	NP B46 429	J. Duboc <i>et al.</i>	(PARIS, LIVP)
THUN	72	PRL 28 1733	R. Thun <i>et al.</i>	(STON, NEAS)
BARDADIN-...	71	PR D4 2711	M. Bardadin-Otwinowska <i>et al.</i>	(WARS)
BOESEBECK	71	PL 34B 659	K. Boesebeck	(AACH, BERL, BONN, CERN, CRAC+)
CAMPBELL	69	PRL 22 1204	J.H. Campbell <i>et al.</i>	(PURD)
LORSTAD	69	NP B14 63	B. Lorstad <i>et al.</i>	(CDEF, CERN) JP
D'ANDLAU	68	NP B5 693	C. d'Andlau <i>et al.</i>	(CDEF, CERN, IRAD+) IJP
DAHL	67	PR 163 1377	O.I. Dahl <i>et al.</i>	(LRL) IJP