

$\phi(1680)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

$\phi(1680)$ MASS

e^+e^- PRODUCTION

| VALUE (MeV) | EVTs | DOCUMENT ID | TECN | COMMENT |
|---|------|----------------------------|------|--|
| 1680±20 OUR ESTIMATE | | | | |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 1689± 7±10 | 4.8k | ¹ SHEN 09 | BELL | 10.6 $e^+e^- \rightarrow K^+K^- \pi^+ \pi^- \gamma$ |
| 1709±20±43 | | ² AUBERT 08S | BABR | 10.6 $e^+e^- \rightarrow$ hadrons |
| 1623±20 | 948 | ³ AKHMETSHIN 03 | CMD2 | 1.05–1.38 $e^+e^- \rightarrow K_L^0 K_S^0$ |
| ~ 1500 | | ⁴ ACHASOV 98H | RVUE | $e^+e^- \rightarrow \pi^+ \pi^- \pi^0, \omega \pi^+ \pi^-,$ $K^+ K^-$ |
| ~ 1900 | | ⁵ ACHASOV 98H | RVUE | $e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$ |
| 1700±20 | | ⁶ CLEGG 94 | RVUE | $e^+e^- \rightarrow K^+ K^-, K_S^0 K \pi$ |
| 1657±27 | 367 | BISELLO 91C | DM2 | $e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$ |
| 1655±17 | | ⁷ BISELLO 88B | DM2 | $e^+e^- \rightarrow K^+ K^-$ |
| 1680±10 | | ⁸ BUON 82 | DM1 | $e^+e^- \rightarrow$ hadrons |
| 1677±12 | | ⁹ MANE 82 | DM1 | $e^+e^- \rightarrow K_S^0 K \pi$ |

¹ From a fit with two incoherent Breit-Wigners.

² From the simultaneous fit to the $K\bar{K}^*(892) + c.c.$ and $\phi\eta$ data from AUBERT 08S using the results of AUBERT 07AK.

³ From the combined fit of AKHMETSHIN 03 and MANE 81 also including ρ, ω , and ϕ . Neither isospin nor flavor structure known.

⁴ Using data from IVANOV 81, BARKOV 87, BISELLO 88B, DOLINSKY 91, and ANTONELLI 92.

⁵ Using the data from BISELLO 91C.

⁶ Using BISELLO 88B and MANE 82 data.

⁷ From global fit including ρ, ω, ϕ and $\rho(1700)$ assume mass 1570 MeV and width 510 MeV for ρ radial excitation.

⁸ From global fit of ρ, ω, ϕ and their radial excitations to channels $\omega \pi^+ \pi^-, K^+ K^-, K_S^0 K_L^0, K_S^0 K^\pm \pi^\mp$. Assume mass 1570 MeV and width 510 MeV for ρ radial excitations, mass 1570 and width 500 MeV for ω radial excitation.

⁹ Fit to one channel only, neglecting interference with $\omega, \rho(1700)$.

PHOTOPRODUCTION

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|---|------------------------|------|--|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| 1753± 3 | ¹⁰ LINK | 02K | FOCS 20–160 $\gamma p \rightarrow K^+ K^- p$ |
| 1726±22 | ¹⁰ BUSENITZ | 89 | TPS $\gamma p \rightarrow K^+ K^- X$ |
| 1760±20 | ¹⁰ ATKINSON | 85C | OMEG 20–70 $\gamma p \rightarrow K\bar{K}X$ |
| 1690±10 | ¹⁰ ASTON | 81F | OMEG 25–70 $\gamma p \rightarrow K^+ K^- X$ |

¹⁰ We list here a state decaying into $K^+ K^-$ possibly different from $\phi(1680)$.

$\rho\bar{\rho}$ ANNIHILATION

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|---|----------------------|------|---|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 1700±8 | ¹¹ AMSLER | 06 | CBAR 0.9 $\bar{p}p \rightarrow K^+ K^- \pi^0$ |
| ¹¹ Could also be $\rho(1700)$. | | | |

 $\phi(1680)$ WIDTH **e^+e^- PRODUCTION**

| VALUE (MeV) | EVTs | DOCUMENT ID | TECN | COMMENT |
|---|------|--------------------------|------|--|
| 150±50 OUR ESTIMATE | | | | This is only an educated guess; the error given is larger than the error on the average of the published values. |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 211±14± 19 | 4.8k | ¹² SHEN | 09 | BELL 10.6 $e^+e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$ |
| 322±77±160 | | ¹³ AUBERT | 08S | BABR 10.6 $e^+e^- \rightarrow$ hadrons |
| 139±60 | 948 | ¹⁴ AKHMETSHIN | 03 | CMD2 1.05–1.38 $e^+e^- \rightarrow K_L^0 K_S^0$ |
| 300±60 | | ¹⁵ CLEGG | 94 | RVUE $e^+e^- \rightarrow K^+ K^-, K_S^0 K \pi$ |
| 146±55 | 367 | BISELLO | 91C | DM2 $e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$ |
| 207±45 | | ¹⁶ BISELLO | 88B | DM2 $e^+e^- \rightarrow K^+ K^-$ |
| 185±22 | | ¹⁷ BUON | 82 | DM1 $e^+e^- \rightarrow$ hadrons |
| 102±36 | | ¹⁸ MANE | 82 | DM1 $e^+e^- \rightarrow K_S^0 K \pi$ |

¹² From a fit with two incoherent Breit-Wigners.¹³ From the simultaneous fit to the $K\bar{K}^*(892) +$ c.c. and $\phi\eta$ data from AUBERT 08S using the results of AUBERT 07AK.¹⁴ From the combined fit of AKHMETSHIN 03 and MANE 81 also including ρ , ω , and ϕ . Neither isospin nor flavor structure known.¹⁵ Using BISELLO 88B and MANE 82 data.¹⁶ From global fit including ρ , ω , ϕ and $\rho(1700)$ ¹⁷ From global fit of ρ , ω , ϕ and their radial excitations to channels $\omega\pi^+\pi^-$, K^+K^- , $K_S^0 K_L^0$, $K_S^0 K^\pm \pi^\mp$. Assume mass 1570 MeV and width 510 MeV for ρ radial excitations, mass 1570 and width 500 MeV for ω radial excitation.¹⁸ Fit to one channel only, neglecting interference with ω , $\rho(1700)$.**PHOTOPRODUCTION**

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|---|------------------------|------|--|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 122±63 | ¹⁹ LINK | 02K | FOCS 20–160 $\gamma p \rightarrow K^+ K^- p$ |
| 121±47 | ¹⁹ BUSENITZ | 89 | TPS $\gamma p \rightarrow K^+ K^- X$ |
| 80±40 | ¹⁹ ATKINSON | 85C | OMEG 20–70 $\gamma p \rightarrow K\bar{K}X$ |
| 100±40 | ¹⁹ ASTON | 81F | OMEG 25–70 $\gamma p \rightarrow K^+ K^- X$ |
| ¹⁹ We list here a state decaying into $K^+ K^-$ possibly different from $\phi(1680)$. | | | |

 $\rho\bar{\rho}$ ANNIHILATION

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|---|----------------------|------|---|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 143±24 | ²⁰ AMSLER | 06 | CBAR 0.9 $\bar{p}p \rightarrow K^+ K^- \pi^0$ |
| ²⁰ Could also be $\rho(1700)$. | | | |

$\phi(1680)$ DECAY MODES

| Mode | Fraction (Γ_i/Γ) |
|--|--------------------------------|
| Γ_1 $K\bar{K}^*(892) + \text{c.c.}$ | dominant |
| Γ_2 $K_S^0 K\pi$ | seen |
| Γ_3 $K\bar{K}$ | seen |
| Γ_4 $K_L^0 K_S^0$ | |
| Γ_5 $e^+ e^-$ | seen |
| Γ_6 $\omega\pi\pi$ | not seen |
| Γ_7 $\phi\pi\pi$ | |
| Γ_8 $K^+ K^- \pi^+ \pi^-$ | seen |
| Γ_9 $\phi\eta$ | |
| Γ_{10} $K^+ K^- \pi^0$ | |

$\phi(1680) \Gamma(i)\Gamma(e^+ e^-)/\Gamma^2(\text{total})$

This combination of a branching ratio into channel (*i*) and branching ratio into $e^+ e^-$ is directly measured and obtained from the cross section at the peak. We list only data that have not been used to determine the branching ratio into (*i*) or $e^+ e^-$.

$$\Gamma(K_L^0 K_S^0)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \qquad \Gamma_4/\Gamma \times \Gamma_5/\Gamma$$

| VALUE (units 10^{-6}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-----------------------------|------|---|
| 0.131 ± 0.059 | 948 | ²¹ AKHMETSHIN 03 | CMD2 | 1.05–1.38 $e^+ e^- \rightarrow K_L^0 K_S^0$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

²¹ From the combined fit of AKHMETSHIN 03 and MANE 81 also including ρ , ω , and ϕ . Neither isospin nor flavor structure known. Recalculated by us.

$$\Gamma(K\bar{K}^*(892) + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \qquad \Gamma_1/\Gamma \times \Gamma_5/\Gamma$$

| VALUE (units 10^{-6}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|---------------------------|------|--|
| $1.15 \pm 0.16 \pm 0.01$ | | ²² AUBERT 08S | BABR | 10.6 $e^+ e^- \rightarrow K\bar{K}^*(892)\gamma + \text{c.c.}$ |
| 3.29 ± 1.57 | 367 | ²³ BISELLO 91C | DM2 | 1.35–2.40 $e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp$ |

²² From the simultaneous fit to the $K\bar{K}^*(892) + \text{c.c.}$ and $\phi\eta$ data from AUBERT 08S using the results of AUBERT 07AK.

²³ Recalculated by us with the published value of $B(K\bar{K}^*(892) + \text{c.c.}) \times \Gamma(e^+ e^-)$.

$$\Gamma(\phi\pi\pi)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \qquad \Gamma_7/\Gamma \times \Gamma_5/\Gamma$$

| VALUE (units 10^{-7}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-----------------------|------|---|
| $1.86 \pm 0.14 \pm 0.21$ | 4.8k | ²⁴ SHEN 09 | BELL | 10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$ |

²⁴ Multiplied by 3/2 to take into account the $\phi\pi^0\pi^0$ mode. Using $B(\phi \rightarrow K^+ K^-) = (49.2 \pm 0.6)\%$.

$\Gamma(\phi\eta)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_9/\Gamma \times \Gamma_5/\Gamma$

VALUE (units 10^{-6}) DOCUMENT ID TECN COMMENT

••• We do not use the following data for averages, fits, limits, etc. •••

$0.43 \pm 0.10 \pm 0.09$ ²⁵ AUBERT 08S BABR $10.6 e^+e^- \rightarrow \phi\eta\gamma$

²⁵ From the simultaneous fit to the $K\bar{K}^*(892) + \text{c.c.}$ and $\phi\eta$ data from AUBERT 08S using the results of AUBERT 07AK.

$\phi(1680)$ BRANCHING RATIOS

$\Gamma(K\bar{K}^*(892) + \text{c.c.})/\Gamma(K_S^0 K\pi)$ Γ_1/Γ_2

VALUE DOCUMENT ID TECN COMMENT

dominant MANE 82 DM1 $e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$

$\Gamma(K\bar{K})/\Gamma(K\bar{K}^*(892) + \text{c.c.})$ Γ_3/Γ_1

VALUE DOCUMENT ID TECN COMMENT

••• We do not use the following data for averages, fits, limits, etc. •••

0.07 ± 0.01 BUON 82 DM1 e^+e^-

$\Gamma(\omega\pi\pi)/\Gamma(K\bar{K}^*(892) + \text{c.c.})$ Γ_6/Γ_1

VALUE DOCUMENT ID TECN COMMENT

<0.10 BUON 82 DM1 e^+e^-

$\Gamma(\phi\eta)/\Gamma(K\bar{K}^*(892) + \text{c.c.})$ Γ_9/Γ_1

VALUE DOCUMENT ID TECN COMMENT

••• We do not use the following data for averages, fits, limits, etc. •••

≈ 0.37 ²⁶ AUBERT 08S BABR $10.6 e^+e^- \rightarrow \text{hadrons}$

²⁶ From the fit including data from AUBERT 07AK.

$\phi(1680)$ REFERENCES

| | | | | |
|------------|------|-------------------------------|--|---------------------------------|
| SHEN | 09 | PR D80 031101 | C.P. Shen <i>et al.</i> | (BELLE Collab.) |
| AUBERT | 08S | PR D77 092002 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 07AK | PR D76 012008 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AMSLER | 06 | PL B639 165 | C. Amsler <i>et al.</i> | (CBAR Collab.) |
| AKHMETSHIN | 03 | PL B551 27 | R.R. Akhmetshin <i>et al.</i> | (Novosibirsk CMD-2 Collab.) |
| Also | | PAN 65 1222 | E.V. Anashkin, V.M. Aulchenko, R.R. Akhmetshin | |
| | | Translated from YAF 65 1255. | | |
| LINK | 02K | PL B545 50 | J.M. Link <i>et al.</i> | (FNAL FOCUS Collab.) |
| ACHASOV | 98H | PR D57 4334 | N.N. Achasov, A.A. Kozhevnikov | |
| CLEGG | 94 | ZPHY C62 455 | A.B. Clegg, A. Donnachie | (LANC, MCHS) |
| ANTONELLI | 92 | ZPHY C56 15 | A. Antonelli <i>et al.</i> | (DM2 Collab.) |
| BISELLO | 91C | ZPHY C52 227 | D. Bisello <i>et al.</i> | (DM2 Collab.) |
| DOLINSKY | 91 | PRPL 202 99 | S.I. Dolinsky <i>et al.</i> | (NOVO) |
| BUSENITZ | 89 | PR D40 1 | J.K. Busenitz <i>et al.</i> | (ILL, FNAL) |
| BISELLO | 88B | ZPHY C39 13 | D. Bisello <i>et al.</i> | (PADO, CLER, FRAS+) |
| BARKOV | 87 | JETPL 46 164 | L.M. Barkov <i>et al.</i> | (NOVO) |
| | | Translated from ZETFP 46 132. | | |
| ATKINSON | 85C | ZPHY C27 233 | M. Atkinson <i>et al.</i> | (BONN, CERN, GLAS+) |
| BUON | 82 | PL 118B 221 | J. Buon <i>et al.</i> | (LALO, MONP) |
| MANE | 82 | PL 112B 178 | F. Mane <i>et al.</i> | (LALO) |
| ASTON | 81F | PL 104B 231 | D. Aston | (BONN, CERN, EPOL, GLAS, LANC+) |
| IVANOV | 81 | PL 107B 297 | P.M. Ivanov <i>et al.</i> | (NOVO) |
| MANE | 81 | PL 99B 261 | F. Mane <i>et al.</i> | (ORSAY) |