

$\psi(4160)$

$I^G(J^{PC}) = 0^-(1^{--})$

$\psi(4160)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
4191 ± 5 OUR AVERAGE			
4191 + 9 - 8	AAIJ	13BC LHCb	$B^+ \rightarrow K^+ \mu^+ \mu^-$
4191.7 ± 6.5	¹ ABLIKIM	08D BES2	$e^+ e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
4193 ± 7	² MO	10 RVUE	$e^+ e^- \rightarrow$ hadrons
4151 ± 4	³ SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
4155 ± 5	⁴ SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
4159 ± 20	BRANDELIK	78C DASP	$e^+ e^-$

¹ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (293 \pm 57)^\circ$.

² Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects.

³ From a fit to Crystal Ball (OSTERHELD 86) data.

⁴ From a fit to BES (BAI 02C) data.

$\psi(4160)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
70 ±10 OUR AVERAGE			
65 +22 -16	AAIJ	13BC LHCb	$B^+ \rightarrow K^+ \mu^+ \mu^-$
71.8±12.3	⁵ ABLIKIM	08D BES2	$e^+ e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
79 ±14	⁶ MO	10 RVUE	$e^+ e^- \rightarrow$ hadrons
107 ±10	⁷ SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
107 ±16	⁸ SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
78 ±20	BRANDELIK	78C DASP	$e^+ e^-$

⁵ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (293 \pm 57)^\circ$.

⁶ Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects.

⁷ From a fit to Crystal Ball (OSTERHELD 86) data.

⁸ From a fit to BES (BAI 02C) data.

$\psi(4160)$ DECAY MODES

Due to the complexity of the $c\bar{c}$ threshold region, in this listing, “seen” (“not seen”) means that a cross section for the mode in question has been measured at effective \sqrt{s} near this particle’s central mass value, more (less) than 2σ above zero, without regard to any peaking behavior in \sqrt{s} or absence thereof. See mode listing(s) for details and references.

Mode	Fraction (Γ_i/Γ)	Confidence level
$\Gamma_1 e^+ e^-$	$(6.9 \pm 3.3) \times 10^{-6}$	
$\Gamma_2 \mu^+ \mu^-$	seen	
$\Gamma_3 D\bar{D}$	seen	
$\Gamma_4 D^0 \bar{D}^0$	seen	
$\Gamma_5 D^+ D^-$	seen	
$\Gamma_6 D^* \bar{D} + \text{c.c.}$	seen	
$\Gamma_7 D^*(2007)^0 \bar{D}^0 + \text{c.c.}$	seen	
$\Gamma_8 D^*(2010)^+ D^- + \text{c.c.}$	seen	
$\Gamma_9 D^* \bar{D}^*$	seen	
$\Gamma_{10} D^*(2007)^0 \bar{D}^*(2007)^0$	seen	
$\Gamma_{11} D^*(2010)^+ D^*(2010)^-$	seen	
$\Gamma_{12} D^0 D^- \pi^+ + \text{c.c. (excl.)}$ $D^*(2007)^0 \bar{D}^0 + \text{c.c.},$ $D^*(2010)^+ D^- + \text{c.c.})$	not seen	
$\Gamma_{13} D \bar{D}^* \pi + \text{c.c. (excl. } D^* \bar{D}^*)$	seen	
$\Gamma_{14} D^0 D^{*-} \pi^+ + \text{c.c. (excl.)}$ $D^*(2010)^+ D^*(2010)^-$	not seen	
$\Gamma_{15} D_s^+ D_s^-$	not seen	
$\Gamma_{16} D_s^{*+} D_s^- + \text{c.c.}$	seen	
$\Gamma_{17} J/\psi \pi^+ \pi^-$	$< 3 \times 10^{-3}$	90%
$\Gamma_{18} J/\psi \pi^0 \pi^0$	$< 3 \times 10^{-3}$	90%
$\Gamma_{19} J/\psi K^+ K^-$	$< 2 \times 10^{-3}$	90%
$\Gamma_{20} J/\psi \eta$	$< 8 \times 10^{-3}$	90%
$\Gamma_{21} J/\psi \pi^0$	$< 1 \times 10^{-3}$	90%
$\Gamma_{22} J/\psi \eta'$	$< 5 \times 10^{-3}$	90%
$\Gamma_{23} J/\psi \pi^+ \pi^- \pi^0$	$< 1 \times 10^{-3}$	90%
$\Gamma_{24} \psi(2S) \pi^+ \pi^-$	$< 4 \times 10^{-3}$	90%
$\Gamma_{25} \chi_{c1} \gamma$	$< 7 \times 10^{-3}$	90%
$\Gamma_{26} \chi_{c2} \gamma$	$< 1.3 \%$	90%
$\Gamma_{27} \chi_{c1} \pi^+ \pi^- \pi^0$	$< 2 \times 10^{-3}$	90%
$\Gamma_{28} \chi_{c2} \pi^+ \pi^- \pi^0$	$< 8 \times 10^{-3}$	90%
$\Gamma_{29} h_c(1P) \pi^+ \pi^-$	$< 5 \times 10^{-3}$	90%
$\Gamma_{30} h_c(1P) \pi^0 \pi^0$	$< 2 \times 10^{-3}$	90%

Γ_{31}	$h_c(1P)\eta$	< 2	$\times 10^{-3}$	90%
Γ_{32}	$h_c(1P)\pi^0$	< 4	$\times 10^{-4}$	90%
Γ_{33}	$\phi\pi^+\pi^-$	< 2	$\times 10^{-3}$	90%

$\psi(4160)$ PARTIAL WIDTHS

$\Gamma(e^+e^-)$	Γ_1		
VALUE (keV)	DOCUMENT ID	TECN	COMMENT
0.48±0.22	⁹ ABLIKIM	08D BES2	$e^+e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.4 to 1.1	¹⁰ MO	10 RVUE	$e^+e^- \rightarrow$ hadrons
0.83±0.08	¹¹ SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
0.84±0.13	¹² SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
0.77±0.23	BRANDELIK	78C DASP	e^+e^-
⁹ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (293 \pm 57)^\circ$.			
¹⁰ Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects. Four sets of solutions are obtained with the same fit quality, mass and total width, but with different e^+e^- partial widths. We quote only the range of values.			
¹¹ From a fit to Crystal Ball (OSTERHELD 86) data.			
¹² From a fit to BES (BAI 02C) data.			

$\psi(4160)$ $\Gamma(i)\Gamma(e^+e^-)/\Gamma^2(\text{total})$

$\Gamma(J/\psi\eta)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{20}/\Gamma \times \Gamma_1/\Gamma$		
VALUE (units 10^{-8})	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2.8±0.9±0.9	¹³ WANG	13B BELL	$e^+e^- \rightarrow J/\psi\eta\gamma$
12.8±1.7±2.0	¹⁴ WANG	13B BELL	$e^+e^- \rightarrow J/\psi\eta\gamma$
¹³ Solution I of two equivalent solutions in a fit using two interfering resonances. Mass and width fixed at 4153 MeV and 103 MeV, respectively.			
¹⁴ Solution II of two equivalent solutions in a fit using two interfering resonances. Mass and width fixed at 4153 MeV and 103 MeV, respectively.			

$\psi(4160)$ BRANCHING RATIOS

$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$	Γ_2/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹⁵ AAIJ	13BC LHCb	$B^+ \rightarrow K^+\mu^+\mu^-$
¹⁵ AAIJ 13BC report $B(B^+ \rightarrow K^+\psi(4160)) B(\psi(4160) \rightarrow \mu^+\mu^-) = (3.5^{+0.9}_{-0.8}) \times 10^{-9}$.			

$\Gamma(D\bar{D})/\Gamma(D^*\bar{D}^*)$	Γ_3/Γ_9		
VALUE	DOCUMENT ID	TECN	COMMENT
0.02±0.03±0.02	AUBERT	09M BABR	$e^+e^- \rightarrow \gamma D^{(*)}\bar{D}^{(*)}$

$\Gamma(D^0\bar{D}^0)/\Gamma_{\text{total}}$

VALUE

seen

seen

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen

DOCUMENT ID

CRONIN-HEN..09

PAKHLOVA 08

AUBERT

TECN

CLEO

BELL

09M

BABR

COMMENT

$e^+e^- \rightarrow D^0\bar{D}^0$

$e^+e^- \rightarrow D^0\bar{D}^0\gamma$

Γ_4/Γ

$\Gamma(D^+D^-)/\Gamma_{\text{total}}$

VALUE

seen

seen

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen

DOCUMENT ID

CRONIN-HEN..09

PAKHLOVA 08

AUBERT

TECN

CLEO

BELL

09M

BABR

COMMENT

$e^+e^- \rightarrow D^+D^-$

$e^+e^- \rightarrow D^+D^-\gamma$

Γ_5/Γ

$\Gamma(D^*(2007)^0\bar{D}^0 + \text{c.c.})/\Gamma_{\text{total}}$

VALUE

seen

seen

DOCUMENT ID

AUBERT 09M

CRONIN-HEN..09

TECN

BABR

CLEO

COMMENT

$e^+e^- \rightarrow D^{*0}\bar{D}^0\gamma$

$e^+e^- \rightarrow D^{*0}\bar{D}^0$

Γ_7/Γ

$\Gamma(D^*(2010)^+D^- + \text{c.c.})/\Gamma_{\text{total}}$

VALUE

seen

seen

seen

DOCUMENT ID

AUBERT 09M

CRONIN-HEN..09

PAKHLOVA 07

TECN

BABR

CLEO

BELL

COMMENT

$e^+e^- \rightarrow D^{*+}D^-\gamma$

$e^+e^- \rightarrow D^{*+}D^-$

$e^+e^- \rightarrow D^{*+}D^-\gamma$

Γ_8/Γ

$\Gamma(D^*\bar{D} + \text{c.c.})/\Gamma(D^*\bar{D}^*)$

VALUE

0.34 ± 0.14 ± 0.05

DOCUMENT ID

AUBERT 09M

TECN

BABR

COMMENT

$e^+e^- \rightarrow \gamma D^{(*)}\bar{D}^{(*)}$

Γ_6/Γ_9

$\Gamma(D^*(2007)^0\bar{D}^*(2007)^0)/\Gamma_{\text{total}}$

VALUE

seen

seen

DOCUMENT ID

AUBERT 09M

CRONIN-HEN..09

TECN

BABR

CLEO

COMMENT

$e^+e^- \rightarrow D^{*0}\bar{D}^{*0}\gamma$

$e^+e^- \rightarrow D^{*0}\bar{D}^{*0}$

Γ_{10}/Γ

$\Gamma(D^*(2010)^+D^*(2010)^-)/\Gamma_{\text{total}}$

VALUE

seen

seen

seen

DOCUMENT ID

AUBERT 09M

CRONIN-HEN..09

PAKHLOVA 07

TECN

BABR

CLEO

BELL

COMMENT

$e^+e^- \rightarrow D^{*+}D^{*-}\gamma$

$e^+e^- \rightarrow D^{*+}D^{*-}$

$e^+e^- \rightarrow D^{*+}D^{*-}\gamma$

Γ_{11}/Γ

$\Gamma(D^0D^-\pi^+ + \text{c.c.} \text{ (excl. } D^*(2007)^0\bar{D}^0 + \text{c.c., } D^*(2010)^+D^- + \text{c.c.})) / \Gamma_{\text{total}}$

VALUE

not seen

DOCUMENT ID

PAKHLOVA 08A

TECN

BELL

COMMENT

$e^+e^- \rightarrow D^0D^-\pi^+\gamma$

Γ_{12}/Γ

$\Gamma(D\bar{D}^*\pi + \text{c.c.} \text{ (excl. } D^*\bar{D}^*)) / \Gamma_{\text{total}}$

VALUE

seen

DOCUMENT ID

CRONIN-HEN..09

TECN

CLEO

COMMENT

$e^+e^- \rightarrow D\bar{D}^*\pi$

Γ_{13}/Γ

$\Gamma(D^0 D^{*-} \pi^+ + \text{c.c. (excl. } D^*(2010)^+ D^*(2010)^-)) / \Gamma_{\text{total}}$ Γ_{14}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
not seen	PAKHLOVA 09	BELL	$e^+ e^- \rightarrow D^0 D^{*-} \pi^+ \gamma$

 $\Gamma(D_s^+ D_s^-)/\Gamma_{\text{total}}$ Γ_{15}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
not seen	PAKHLOVA 11	BELL	$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$
not seen	DEL-AMO-SA..10N	BABR	$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$
not seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D_s^+ D_s^-$

 $\Gamma(D_s^{*+} D_s^- + \text{c.c.}) / \Gamma_{\text{total}}$ Γ_{16}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
seen	PAKHLOVA 11	BELL	$e^+ e^- \rightarrow D_s^{*+} D_s^- \gamma$
seen	DEL-AMO-SA..10N	BABR	$e^+ e^- \rightarrow D_s^{*+} D_s^- \gamma$
seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D_s^{*+} D_s^-$

 $\Gamma(J/\psi \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{17}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<3	90	COAN	06	CLEO 4.12–4.2 $e^+ e^- \rightarrow$ hadrons

 $\Gamma(J/\psi \pi^0 \pi^0) / \Gamma_{\text{total}}$ Γ_{18}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<3	90	COAN	06	CLEO 4.12–4.2 $e^+ e^- \rightarrow$ hadrons

 $\Gamma(J/\psi K^+ K^-) / \Gamma_{\text{total}}$ Γ_{19}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2	90	COAN	06	CLEO 4.12–4.2 $e^+ e^- \rightarrow$ hadrons

 $\Gamma(J/\psi \eta) / \Gamma_{\text{total}}$ Γ_{20}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<8	90	COAN	06	CLEO 4.12–4.2 $e^+ e^- \rightarrow$ hadrons

 $\Gamma(J/\psi \pi^0) / \Gamma_{\text{total}}$ Γ_{21}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1	90	COAN	06	CLEO 4.12–4.2 $e^+ e^- \rightarrow$ hadrons

 $\Gamma(J/\psi \eta') / \Gamma_{\text{total}}$ Γ_{22}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<5	90	COAN	06	CLEO 4.12–4.2 $e^+ e^- \rightarrow$ hadrons

 $\Gamma(J/\psi \pi^+ \pi^- \pi^0) / \Gamma_{\text{total}}$ Γ_{23}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1	90	COAN	06	CLEO 4.12–4.2 $e^+ e^- \rightarrow$ hadrons

$\Gamma(\psi(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{24}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<4	90	COAN	06	CLEO 4.12–4.2 $e^+e^- \rightarrow$ hadrons

 $\Gamma(\chi_{c1}\gamma)/\Gamma_{\text{total}}$ Γ_{25}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<7	90	COAN	06	CLEO 4.12–4.2 $e^+e^- \rightarrow$ hadrons

 $\Gamma(\chi_{c2}\gamma)/\Gamma_{\text{total}}$ Γ_{26}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<13	90	COAN	06	CLEO 4.12–4.2 $e^+e^- \rightarrow$ hadrons

 $\Gamma(\chi_{c1}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{27}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2	90	COAN	06	CLEO 4.12–4.2 $e^+e^- \rightarrow$ hadrons

 $\Gamma(\chi_{c2}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{28}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<8	90	COAN	06	CLEO 4.12–4.2 $e^+e^- \rightarrow$ hadrons

 $\Gamma(h_c(1P)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{29}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<5	90	16 PEDLAR	11	CLEO $e^+e^- \rightarrow h_c(1P)\pi^+\pi^-$

¹⁶ At $\sqrt{s} = 4170$ MeV, PEDLAR 11 measures $\sigma(e^+e^- \rightarrow h_c(1P)\pi^+\pi^-) = 15.6 \pm 2.3 \pm 1.9 \pm 3.0$ pb, where the errors are statistical, systematic, and due to uncertainty in $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$, respectively.

 $\Gamma(h_c(1P)\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_{30}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2	90	17 PEDLAR	11	CLEO $e^+e^- \rightarrow h_c(1P)\pi^0\pi^0$

¹⁷ At $\sqrt{s} = 4170$ MeV, PEDLAR 11 measures $\sigma(e^+e^- \rightarrow h_c(1P)\pi^0\pi^0) = 3.0 \pm 3.3 \pm 1.1 \pm 0.6$ pb, where the errors are statistical, systematic, and due to uncertainty in $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$, respectively.

 $\Gamma(h_c(1P)\eta)/\Gamma_{\text{total}}$ Γ_{31}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2	90	18 PEDLAR	11	CLEO $e^+e^- \rightarrow h_c(1P)\eta$

¹⁸ At $\sqrt{s} = 4170$ MeV, PEDLAR 11 measures $\sigma(e^+e^- \rightarrow h_c(1P)\eta) = 4.7 \pm 1.7 \pm 1.0 \pm 0.9$ pb, where the errors are statistical, systematic, and due to uncertainty in $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$, respectively.

 $\Gamma(h_c(1P)\pi^0)/\Gamma_{\text{total}}$ Γ_{32}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.4	90	19 PEDLAR	11	CLEO $e^+e^- \rightarrow h_c(1P)\pi^0$

¹⁹ At $\sqrt{s} = 4170$ MeV, PEDLAR 11 measures $\sigma(e^+e^- \rightarrow h_c(1P)\pi^0) = -0.7 \pm 1.8 \pm 0.7 \pm 0.1$ pb, where the errors are statistical, systematic, and due to uncertainty in $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$, respectively.

$\Gamma(\phi\pi^+\pi^-)/\Gamma_{\text{total}}$	Γ_{33}/Γ			
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<2	90	COAN	06	CLEO 4.12–4.2 $e^+e^- \rightarrow$ hadrons

$\psi(4160)$ REFERENCES

AAIJ	13BC	PRL 111 112003	R. Aaij <i>et al.</i>	(LHCb Collab.)
WANG	13B	PR D87 051101	X.L. Wang <i>et al.</i>	(BELLE Collab.)
PAKHLOVA	11	PR D83 011101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PEDLAR	11	PRL 107 041803	T. Pedlar <i>et al.</i>	(CLEO Collab.)
DEL-AMO-SA...	10N	PR D82 052004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
MO	10	PR D82 077501	X.H. Mo, C.Z. Yuan, P. Wang	(BHEP)
AUBERT	09M	PR D79 092001	B. Aubert <i>et al.</i>	(BABAR Collab.)
CRONIN-HEN...	09	PR D80 072001	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)
PAKHLOVA	09	PR D80 091101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
ABLIKIM	08D	PL B660 315	M. Ablikim <i>et al.</i>	(BES Collab.)
PAKHLOVA	08	PR D77 011103	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PAKHLOVA	08A	PRL 100 062001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PAKHLOVA	07	PRL 98 092001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
COAN	06	PRL 96 162003	T.E. Coan <i>et al.</i>	(CLEO Collab.)
SETH	05A	PR D72 017501	K.K. Seth	
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)
OSTERHELD	86	SLAC-PUB-4160	A. Osterheld <i>et al.</i>	(SLAC Crystal Ball Collab.)
BRANDELIK	78C	PL 76B 361	R. Brandelik <i>et al.</i>	(DASP Collab.)