

$D_2^*(2460)^0$

$$I(J^P) = \frac{1}{2}(2^+)$$

$J^P = 2^+$ assignment strongly favored(ALBRECHT 89B, ALBRECHT 89H), natural parity confirmed by the helicity analysis(DEL-AMO-SANCHEZ 10P),

$D_2^*(2460)^0$ MASS

The fit includes D^\pm , D^0 , D_s^\pm , $D^{*\pm}$, D^{*0} , D_s^{*0} , $D_s^{*\pm}$, $D_1(2420)^0$, $D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2462.6±0.6 OUR FIT	Error includes scale factor of 1.2.			
2461.8±0.7 OUR AVERAGE	Error includes scale factor of 1.1.			
2462.5±2.4 ^{+1.3} _{-1.1}	2.3k	¹ ABRAMOWICZ13	ZEUS	$e^\pm p \rightarrow D^{(*)+} \pi^- X$
2462.2±0.1±0.8	243k	DEL-AMO-SA..10P	BABR	$e^+ e^- \rightarrow D^+ \pi^- X$
2460.4±1.2±2.2	3.4k	AUBERT	09AB BABR	$B^- \rightarrow D^+ \pi^- \pi^-$
2461.6±2.1±3.3	² ABE	04D BELL	$B^- \rightarrow D^+ \pi^- \pi^-$	
2464.5±1.1±1.9	5.8k	² LINK	04A FOCS	γA
2465 ± 3 ± 3	486	AVERY	94C CLE2	$e^+ e^- \rightarrow D^+ \pi^- X$
2453 ± 3 ± 2	128	FRABETTI	94B E687	$\gamma Be \rightarrow D^+ \pi^- X$
2461 ± 3 ± 1	440	AVERY	90 CLEO	$e^+ e^- \rightarrow D^{*+} \pi^- X$
2455 ± 3 ± 5	337	ALBRECHT	89B ARG	$e^+ e^- \rightarrow D^+ \pi^- X$
2459 ± 3 ± 2	153	ANJOS	89C TPS	$\gamma N \rightarrow D^+ \pi^- X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2469.1±3.7 ^{+1.2} _{-1.3}	1560 ± 230	³ CHEKANOV	09 ZEUS	$e^\pm p \rightarrow D^{(*)+} \pi^- X$
2463.3±0.6±0.8	20k	ABULENCIA	06A CDF	$1900 p\bar{p} \rightarrow D^+ \pi^- X$
2461 ± 6	126	⁴ ABREU	98M DLPH	$e^+ e^-$
2466 ± 7	1	ASRATYAN	95 BEBC	$53,40 \nu(\bar{\nu}) \rightarrow pX, dX$

¹ From the combined fit of the $M(D^+ \pi^-)$ and $M(D^{*+} \pi^-)$ distributions. and A_{D_2} fixed to the theoretical prediction of -1.

² Fit includes the contribution from $D_0^*(2400)^0$.

³ Calculated using the mass difference $m(D_2^{*0}) - m(D^{*+})_{PDG}$ reported below and $m(D^{*+})_{PDG} = 2010.27 \pm 0.17$ MeV. The 0.17 MeV uncertainty of the PDG mass value should be added to the experimental uncertainty of $^{+1.2}_{-1.3}$ MeV.

⁴ No systematic error given.

$m_{D_2^{*0}} - m_{D^+}$

The fit includes D^\pm , D^0 , D_s^\pm , $D^{*\pm}$, D^{*0} , D_s^{*0} , $D_1(2420)^0$, $D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
593.0±0.6 OUR FIT	Error includes scale factor of 1.2.			
593.9±0.6±0.5	20k	ABULENCIA	06A CDF	$1900 p\bar{p} \rightarrow D^+ \pi^- X$

$m_{D_2^{*0}} - m_{D^{*+}}$

The fit includes D^\pm , D^0 , D_s^\pm , $D^{*\pm}$, D^{*0} , $D_s^{*\pm}$, $D_1(2420)^0$, $D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.


VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
452.3±0.6 OUR FIT		Error includes scale factor of 1.2.		
458.8±3.7^{+1.2}_{-1.3}	1560±230	CHEKANOV 09	ZEUS	$e^\pm p \rightarrow D^{(*)} + \pi^- X$

$D_2^*(2460)^0$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
49.0± 1.3 OUR AVERAGE		Error includes scale factor of 1.5. See the ideogram below.		
46.6± 8.1 ^{+ 5.9} _{- 3.8}	2.3k	⁵ ABRAMOWICZ13	ZEUS	$e^\pm p \rightarrow D^{(*)} + \pi^- X$
50.5± 0.6± 0.7	243k	DEL-AMO-SA..10P	BABR	$e^+ e^- \rightarrow D^+ \pi^- X$
41.8± 2.5± 2.9	3.4k	AUBERT	09AB BABR	$B^- \rightarrow D^+ \pi^- \pi^-$
49.2± 2.3± 1.3	20k	ABULENCIA	06A CDF	1900 $p\bar{p} \rightarrow D^+ \pi^- X$
45.6± 4.4± 6.7		⁶ ABE	04D BELL	$B^- \rightarrow D^+ \pi^- \pi^-$
38.7± 5.3± 2.9	5.8k	⁶ LINK	04A FOCS	γA
28 ^{+ 8} _{- 7} ± 6	486	AVERY	94C CLE2	$e^+ e^- \rightarrow D^+ \pi^- X$
25 ± 10 ± 5	128	FRABETTI	94B E687	$\gamma Be \rightarrow D^+ \pi^- X$
20 ^{+ 9} _{- 12} ^{+ 9} _{- 10}	440	AVERY	90 CLEO	$e^+ e^- \rightarrow D^{*+} \pi^- X$
15 ^{+ 13} _{- 10} ^{+ 5} _{- 10}	337	ALBRECHT	89B ARG	$e^+ e^- \rightarrow D^+ \pi^- X$
20 ± 10 ± 5	153	ANJOS	89C TPS	$\gamma N \rightarrow D^+ \pi^- X$

⁵ From the combined fit of the $M(D^+ \pi^-)$ and $M(D^{*+} \pi^-)$ distributions. and A_{D_2} fixed to the theoretical prediction of -1.

⁶ Fit includes the contribution from $D_0^*(2400)^0$.



$D_2^*(2460)^0$ DECAY MODES

$\overline{D}_2^*(2460)^0$ modes are charge conjugates of modes below.

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 D^+ \pi^-$	seen
$\Gamma_2 D^*(2010)^+ \pi^-$	seen
$\Gamma_3 D^0 \pi^+ \pi^-$	not seen
$\Gamma_4 D^{*0} \pi^+ \pi^-$	not seen

$D_2^*(2460)^0$ BRANCHING RATIOS

$\Gamma(D^+ \pi^-)/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
seen	3.4k	AUBERT	09AB BABR	$B^- \rightarrow D^+ \pi^- \pi^-$	
seen	337	ALBRECHT	89B ARG	$e^+ e^- \rightarrow D^+ \pi^- X$	
seen		ANJOS	89C TPS	$\gamma N \rightarrow D^+ \pi^- X$	

$\Gamma(D^*(2010)^+ \pi^-)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
seen	ACKERSTAFF 97W	OPAL	$e^+ e^- \rightarrow D^{*+} \pi^- X$	
seen	AVERY 90	CLEO	$e^+ e^- \rightarrow D^{*+} \pi^- X$	
seen	ALBRECHT 89H	ARG	$e^+ e^- \rightarrow D^* \pi^- X$	

$\Gamma(D^+\pi^-)/\Gamma(D^*(2010)^+\pi^-)$

Γ_1/Γ_2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.54±0.15 OUR AVERAGE				
1.4 ± 0.3 ± 0.3	2.3k	7 ABRAMOWICZ13	ZEUS	$e^\pm p \rightarrow D^{(*)}+\pi^- X$
1.47±0.03±0.16	379k	DEL-AMO-SA..10P	BABR	$e^+ e^- \rightarrow D^{(*)}+\pi^- X$
2.8 ± 0.8 ± 0.5	1560 ± 230	CHEKANOV 09	ZEUS	$e^\pm p \rightarrow D^{(*)}+\pi^- X$
2.2 ± 0.7 ± 0.6		AVERY 94C	CLE2	$e^+ e^- \rightarrow D^*+\pi^- X$
2.3 ± 0.8		AVERY 90	CLEO	$e^+ e^-$
3.0 ± 1.1 ± 1.5		ALBRECHT 89H	ARG	$e^+ e^- \rightarrow D^*\pi^- X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.9 ± 0.5		ABE 04D	BELL	$B^- \rightarrow D^{(*)}+\pi^-\pi^-$

⁷ From the combined fit of the $M(D^+\pi^-)$ and $M(D^*+\pi^-)$ distributions. and A_{D_2} fixed to the theoretical prediction of -1.

$\Gamma(D^+\pi^-)/[\Gamma(D^+\pi^-)+\Gamma(D^*(2010)^+\pi^-)]$

$\Gamma_1/(\Gamma_1+\Gamma_2)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.62±0.03±0.02	8414	8 AUBERT 09Y	BABR	$B^+ \rightarrow D_2^{*0}\ell^+\nu_\ell$
⁸ Assuming $\Gamma(\gamma(4S) \rightarrow B^+ B^-) / \Gamma(\gamma(4S) \rightarrow B^0 \bar{B}^0) = 1.065 \pm 0.026$ and equal partial widths for charged and neutral D_2^* mesons.				

$D_2^*(2460)^0$ POLARIZATION AMPLITUDE A_{D_2}

A polarization amplitude A_{D_2} is a parameter that depends on the initial polarization of the D_2 . For D_2 decays the helicity angle, θ_H , distribution varies like $1 + A_{D_2} \cos^2(\theta_H)$, where θ_H is the angle in the D^* rest frame between the two pions emitted by the $D_2 \rightarrow D^* \pi$ and $D^* \rightarrow D \pi$.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-1.16±0.35	2.3k	9 ABRAMOWICZ13	ZEUS	$e^\pm p \rightarrow D^{(*)}+\pi^- X$
consistent with -1	243k	DEL-AMO-SA..10P	BABR	$e^+ e^- \rightarrow D^+\pi^- X$
-0.74 ^{+0.49} _{-0.38}		10 AVERY 94C	CLE2	$e^+ e^- \rightarrow D^*+\pi^- X$

⁹ From the combined fit of the $M(D^+\pi^-)$ and $M(D^*+\pi^-)$ distributions.

¹⁰ Systematic uncertainties not estimated.

$D_2^*(2460)^0$ REFERENCES

ABRAMOWICZ 13	NP B866 229	H. Abramowicz <i>et al.</i>	(ZEUS Collab.)
DEL-AMO-SA..10P	PR D82 111101	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
AUBERT 09AB	PR D79 112004	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT 09Y	PRL 103 051803	B. Aubert <i>et al.</i>	(BABAR Collab.)
CHEKANOV 09	EPJ C60 25	S. Chekanov <i>et al.</i>	(ZEUS Collab.)
ABULENCIA 06A	PR D73 051104	A. Abulencia <i>et al.</i>	(CDF Collab.)
ABE 04D	PR D69 112002	K. Abe <i>et al.</i>	(BELLE Collab.)
LINK 04A	PL B586 11	J.M. Link <i>et al.</i>	(FOCUS Collab.)
ABREU 98M	PL B426 231	P. Abreu <i>et al.</i>	(DELPHI Collab.)

ACKERSTAFF	97W	ZPHY C76 425	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
ASRATYAN	95	ZPHY C68 43	A.E. Asratyan <i>et al.</i>	(BIRM, BELG, CERN+)
AVERY	94C	PL B331 236	P. Avery <i>et al.</i>	(CLEO Collab.)
FRAZETTI	94B	PRL 72 324	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
AVERY	90	PR D41 774	P. Avery, D. Besson	(CLEO Collab.)
ALBRECHT	89B	PL B221 422	H. Albrecht <i>et al.</i>	(ARGUS Collab.) JP
ALBRECHT	89H	PL B232 398	H. Albrecht <i>et al.</i>	(ARGUS Collab.) JP
ANJOS	89C	PRL 62 1717	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)