$N(1675)\ 5/2^-$

$I(J^P) = \frac{1}{2}(\frac{5}{2}^-)$ Status: ****

Most of the results published before 1975 were last included in our 1982 edition, Physics Letters **111B** 1 (1982). Some further obsolete results published before 1984 were last included in our 2006 edition, Journal of Physics **G33** 1 (2006).

N(1675) BREIT-WIGNER MASS

VALUE (MeV)	*DOCUMENT ID*		*TECN*	*COMMENT*
1670 to 1680 (≈ 1675) OUR ESTIMATE				
1666 ± 2	SHKLYAR	13	DPWA	Multichannel
1664 ± 5	ANISOVICH	12A	DPWA	Multichannel
1674.1± 0.2	ARNDT	06	DPWA	$\pi N \to \pi N,\ \eta N$
1675 ±10	CUTKOSKY	80	IPWA	$\pi N \to \pi N$
1679 ± 8	HOEHLER	79	IPWA	$\pi N \to \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1679 ± 1	SHRESTHA	12A	DPWA	Multichannel
1678 ± 5	ANISOVICH	10	DPWA	Multichannel
1679 ± 9	BATINIC	10	DPWA	$\pi N \to N\pi,\ N\eta$
1678 ±15	THOMA	08	DPWA	Multichannel
1676.2± 0.6	ARNDT	04	DPWA	$\pi N \to \pi N,\ \eta N$
1685 ± 4	VRANA	00	DPWA	Multichannel
1673 ± 5	ARNDT	96	IPWA	$\gamma N \to \pi N$
1673	ARNDT	95	DPWA	$\pi N \to N\pi$
1666	LI	93	IPWA	$\gamma N \to \pi N$
1676 ± 2	MANLEY	92	IPWA	$\pi N \to \pi N$ & $N\pi\pi$
1670	SAXON	80	DPWA	$\pi^- p \to \Lambda K^0$
1650	[1] LONGACRE	77	IPWA	$\pi N \to N\pi\pi$
1660	[2] LONGACRE	75	IPWA	$\pi N \to N\pi\pi$

N(1675) BREIT-WIGNER WIDTH

VALUE (MeV)	*DOCUMENT ID*		*TECN*	*COMMENT*
130 to 165 (≈ 150) OUR ESTIMATE				
148 ± 1	SHKLYAR	13	DPWA	Multichannel
152 ± 7	ANISOVICH	12A	DPWA	Multichannel
146.5± 1.0	ARNDT	06	DPWA	$\pi N \to \pi N,\ \eta N$
160 ±20	CUTKOSKY	80	IPWA	$\pi N \to \pi N$
120 ±15	HOEHLER	79	IPWA	$\pi N \to \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
145 ± 4	SHRESTHA	12A	DPWA	Multichannel
177 ±15	ANISOVICH	10	DPWA	Multichannel
152 ± 8	BATINIC	10	DPWA	$\pi N \to N\pi,\ N\eta$
220 ±25	THOMA	08	DPWA	Multichannel
151.8± 3.0	ARNDT	04	DPWA	$\pi N \to \pi N,\ \eta N$
131 ±10	VRANA	00	DPWA	Multichannel
154 ± 7	ARNDT	96	IPWA	$\gamma N \to \pi N$

154		ARNDT	95	DPWA	$\pi N \rightarrow N\pi$
136		LI	93	IPWA	$\gamma N \rightarrow \pi N$
159	± 7	MANLEY	92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$
40		SAXON	80	DPWA	$\pi^- p \rightarrow \Lambda K^0$
130		[1] LONGACRE	77	IPWA	$\pi N \rightarrow N\pi\pi$
150		[2] LONGACRE	75	IPWA	$\pi N \rightarrow N\pi\pi$

N(1675) POLE POSITION

REAL PART

VALUE (MeV)	*DOCUMENT ID*		*TECN*	*COMMENT*
1655 to 1665 (≈ 1660) OUR ESTIMATE				
1654± 2	[3] SVARC	14	MLS	$\pi N \rightarrow \pi N$
1654± 4	ANISOVICH	12A	DPWA	Multichannel
1657	ARNDT	06	DPWA	$\pi N \rightarrow \pi N$, ηN
1656	[4] HOEHLER	93	ARGD	$\pi N \rightarrow \pi N$
1660±10	CUTKOSKY	80	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1640	SHKLYAR	13	DPWA	Multichannel
1656	SHRESTHA	12A	DPWA	Multichannel
1650± 5	ANISOVICH	10	DPWA	Multichannel
1658± 9	BATINIC	10	DPWA	$\pi N \rightarrow N\pi$, $N\eta$
1639±10	THOMA	08	DPWA	Multichannel
1659	ARNDT	04	DPWA	$\pi N \rightarrow \pi N$, ηN
1674	VRANA	00	DPWA	Multichannel
1663	ARNDT	95	DPWA	$\pi N \rightarrow N\pi$
1655	ARNDT	91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
1663 or 1668	[5] LONGACRE	78	IPWA	$\pi N \rightarrow N\pi\pi$
1649 or 1650	[1] LONGACRE	77	IPWA	$\pi N \rightarrow N\pi\pi$

−2×IMAGINARY PART

VALUE (MeV)	*DOCUMENT ID*		*TECN*	*COMMENT*
125 to 150 (≈ 135) OUR ESTIMATE				
125± 3±1	[3] SVARC	14	MLS	$\pi N \rightarrow \pi N$
151± 5	ANISOVICH	12A	DPWA	Multichannel
139	ARNDT	06	DPWA	$\pi N \rightarrow \pi N$, ηN
126	[4] HOEHLER	93	ARGD	$\pi N \rightarrow \pi N$
140±10	CUTKOSKY	80	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
108	SHKLYAR	13	DPWA	Multichannel
128	SHRESTHA	12A	DPWA	Multichannel
143± 7	ANISOVICH	10	DPWA	Multichannel
137± 7	BATINIC	10	DPWA	$\pi N \rightarrow N\pi$, $N\eta$
180±20	THOMA	08	DPWA	Multichannel
146	ARNDT	04	DPWA	$\pi N \rightarrow \pi N$, ηN
120	VRANA	00	DPWA	Multichannel
152	ARNDT	95	DPWA	$\pi N \rightarrow N\pi$
124	ARNDT	91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
146 or 171	[5] LONGACRE	78	IPWA	$\pi N \rightarrow N\pi\pi$
127 or 127	[1] LONGACRE	77	IPWA	$\pi N \rightarrow N\pi\pi$

N(1675) ELASTIC POLE RESIDUE

MODULUS $|r|$

VALUE (MeV)	*DOCUMENT ID*		*TECN*	*COMMENT*
27±5 OUR ESTIMATE				
23±1	[3] SVARC	14	MLS	$\pi N \rightarrow \pi N$
28±1	ANISOVICH	12A	DPWA	Multichannel
27	ARNDT	06	DPWA	$\pi N \rightarrow \pi N, \eta N$
23	HOEHLER	93	ARGD	$\pi N \rightarrow \pi N$
31±5	CUTKOSKY	80	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
20	SHKLYAR	13	DPWA	Multichannel
25	BATINIC	10	DPWA	$\pi N \rightarrow N\pi, N\eta$
29	ARNDT	04	DPWA	$\pi N \rightarrow \pi N, \eta N$
29	ARNDT	95	DPWA	$\pi N \rightarrow N\pi$
28	ARNDT	91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90

PHASE θ

VALUE (°)	*DOCUMENT ID*		*TECN*	*COMMENT*
−25± 6 OUR ESTIMATE				
−25± 2	[3] SVARC	14	MLS	$\pi N \rightarrow \pi N$
−26± 4	ANISOVICH	12A	DPWA	Multichannel
−21	ARNDT	06	DPWA	$\pi N \rightarrow \pi N, \eta N$
−22	HOEHLER	93	ARGD	$\pi N \rightarrow \pi N$
−30±10	CUTKOSKY	80	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
−49	SHKLYAR	13	DPWA	Multichannel
−16	BATINIC	10	DPWA	$\pi N \rightarrow N\pi, N\eta$
−22	ARNDT	04	DPWA	$\pi N \rightarrow \pi N, \eta N$
− 6	ARNDT	95	DPWA	$\pi N \rightarrow N\pi$
−17	ARNDT	91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90

N(1675) INELASTIC POLE RESIDUE

The "normalized residue" is the residue divided by $\Gamma_{pole}/2$.

Normalized residue in $N\pi \rightarrow N(1675) \rightarrow \Delta\pi$, *D*-wave

MODULUS (%)	*PHASE* (°)	*DOCUMENT ID*		*TECN*	*COMMENT*
33±5	**82 ± 10**	ANISOVICH	12A	DPWA	Multichannel

Normalized residue in $N\pi \rightarrow N(1675) \rightarrow N\sigma$

MODULUS (%)	*PHASE* (°)	*DOCUMENT ID*		*TECN*	*COMMENT*
15±4	**132 ± 18**	ANISOVICH	12A	DPWA	Multichannel

N(1675) DECAY MODES

The following branching fractions are our estimates, not fits or averages.

	Mode	Fraction (Γ_i/Γ)
Γ_1	$N\pi$	35–45 %
Γ_2	$N\eta$	(0 ± 7) × 10^{-3}
Γ_3	ΛK	<1 %
Γ_4	ΣK	
Γ_5	$N\pi\pi$	50–60 %
Γ_6	$\Delta\pi$	50–60 %
Γ_7	$\Delta(1232)\pi$, *D*-wave	(50 ±15) %
Γ_8	$\Delta(1232)\pi$, *G*-wave	
Γ_9	$N\rho$	< 1–3 %
Γ_{10}	$N\rho$, *S*=1/2, *D*-wave	(0.0± 1.0) %
Γ_{11}	$N\rho$, *S*=3/2, *D*-wave	(1.0± 1.0) %
Γ_{12}	$N\rho$, *S*=3/2, *G*-wave	
Γ_{13}	$N(\pi\pi)^{I=0}_{S\text{-wave}}$	(7.0± 3.0) %
Γ_{14}	$p\gamma$	0–0.02 %
Γ_{15}	$p\gamma$, helicity=1/2	0–0.01 %
Γ_{16}	$p\gamma$, helicity=3/2	0–0.01 %
Γ_{17}	$n\gamma$	0–0.15 %
Γ_{18}	$n\gamma$, helicity=1/2	0–0.05 %
Γ_{19}	$n\gamma$, helicity=3/2	0–0.10 %

N(1675) BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{total}$ Γ_1/Γ

VALUE (%)	*DOCUMENT ID*		*TECN*	*COMMENT*
35 to 45 OUR ESTIMATE				
41 ±1	SHKLYAR	13	DPWA	Multichannel
40 ±3	ANISOVICH	12A	DPWA	Multichannel
39.3±0.1	ARNDT	06	DPWA	$\pi N \to \pi N, \eta N$
38 ±5	CUTKOSKY	80	IPWA	$\pi N \to \pi N$
38 ±3	HOEHLER	79	IPWA	$\pi N \to \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
38.6±0.6	SHRESTHA	12A	DPWA	Multichannel
37 ±5	ANISOVICH	10	DPWA	Multichannel
35 ±4	BATINIC	10	DPWA	$\pi N \to N\pi, N\eta$
30 ±8	THOMA	08	DPWA	Multichannel
40.0±0.2	ARNDT	04	DPWA	$\pi N \to \pi N, \eta N$
35 ±1	VRANA	00	DPWA	Multichannel
38	ARNDT	95	DPWA	$\pi N \to N\pi$
47 ±2	MANLEY	92	IPWA	$\pi N \to \pi N$ & $N\pi\pi$

$\Gamma(N\eta)/\Gamma_{total}$ Γ_2/Γ

VALUE (%)	DOCUMENT ID		TECN	COMMENT
0.0±0.7 OUR AVERAGE				
0 ±1	SHKLYAR	13	DPWA	Multichannel
0 ±1	VRANA	00	DPWA	Multichannel
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<1	SHRESTHA	12A	DPWA	Multichannel
0.1±0.1	BATINIC	10	DPWA	$\pi N \to N\pi$, $N\eta$
3 ±3	THOMA	08	DPWA	Multichannel

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{total}$ in $N\pi \to N(1675) \to \Lambda K$ $(\Gamma_1\Gamma_3)^{1/2}/\Gamma$

VALUE	DOCUMENT ID		TECN	COMMENT
±0.04 to ±0.08 OUR ESTIMATE				
−0.01	BELL	83	DPWA	$\pi^- p \to \Lambda K^0$
+0.036	[6] SAXON	80	DPWA	$\pi^- p \to \Lambda K^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
−0.03 ±0.01	SHRESTHA	12A	DPWA	Multichannel

Note: Signs of couplings from $\pi N \to N\pi\pi$ analyses were changed in the 1986 edition to agree with the baryon-first convention; the overall phase ambiguity is resolved by choosing a negative sign for the $\Delta(1620)$ S_{31} coupling to $\Delta(1232)\pi$.

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{total}$ in $N\pi \to N(1675) \to \Delta(1232)\pi$, D-wave $(\Gamma_1\Gamma_7)^{1/2}/\Gamma$

VALUE	DOCUMENT ID		TECN	COMMENT
+0.46 to +0.50 OUR ESTIMATE				
+0.46	[1,7] LONGACRE	77	IPWA	$\pi N \to N\pi\pi$
+0.50	[2] LONGACRE	75	IPWA	$\pi N \to N\pi\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
+0.496±0.003	MANLEY	92	IPWA	$\pi N \to \pi N$ & $N\pi\pi$

$\Gamma(\Delta(1232)\pi, D\text{-wave})/\Gamma_{total}$ Γ_7/Γ

VALUE (%)	DOCUMENT ID		TECN	COMMENT
50±15 OUR ESTIMATE				
33± 8	ANISOVICH	12A	DPWA	Multichannel
63± 2	VRANA	00	DPWA	Multichannel
• • • We do not use the following data for averages, fits, limits, etc. • • •				
46± 1	SHRESTHA	12A	DPWA	Multichannel
24± 8	THOMA	08	DPWA	Multichannel

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{total}$ in $N\pi \to N(1675) \to N\rho$, S=1/2, D-wave $(\Gamma_1\Gamma_{10})^{1/2}/\Gamma$

VALUE	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
+0.04±0.02	MANLEY	92	IPWA	$\pi N \to \pi N$ & $N\pi\pi$

$\Gamma(N\rho, S{=}1/2, D\text{-wave})/\Gamma_{total}$ Γ_{10}/Γ

VALUE (%)	DOCUMENT ID		TECN	COMMENT
0±1	VRANA	00	DPWA	Multichannel
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<1	SHRESTHA	12A	DPWA	Multichannel

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{total}$ in $N\pi \to N(1675) \to N\rho, S{=}3/2, D\text{-wave}$ $(\Gamma_1\Gamma_{11})^{1/2}/\Gamma$

VALUE	DOCUMENT ID		TECN	COMMENT
−0.12 to −0.06 OUR ESTIMATE				
−0.15	[1,7] LONGACRE	77	IPWA	$\pi N \to N\pi\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
−0.03±0.02	MANLEY	92	IPWA	$\pi N \to \pi N$ & $N\pi\pi$

$\Gamma(N\rho, S{=}3/2, D\text{-wave})/\Gamma_{total}$ Γ_{11}/Γ

VALUE (%)	DOCUMENT ID		TECN	COMMENT
1±1	VRANA	00	DPWA	Multichannel
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<1	SHRESTHA	12A	DPWA	Multichannel

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{total}$ in $N\pi \to N(1675) \to N(\pi\pi)^{I=0}_{S\text{-wave}}$ $(\Gamma_1\Gamma_{13})^{1/2}/\Gamma$

VALUE	DOCUMENT ID		TECN	COMMENT
+0.03	[1,7] LONGACRE	77	IPWA	$\pi N \to N\pi\pi$

$\Gamma(N(\pi\pi)^{I=0}_{S\text{-wave}})/\Gamma_{total}$ Γ_{13}/Γ

VALUE (%)	DOCUMENT ID		TECN	COMMENT
7±3	ANISOVICH	12A	DPWA	Multichannel

$N(1675)$ PHOTON DECAY AMPLITUDES

Papers on γN amplitudes predating 1981 may be found in our 2006 edition, Journal of Physics **G33** 1 (2006).

$N(1675) \to p\gamma$, helicity-1/2 amplitude $A_{1/2}$

VALUE ($GeV^{-1/2}$)	DOCUMENT ID		TECN	COMMENT
+0.019±0.008 OUR ESTIMATE				
0.024±0.003	ANISOVICH	12A	DPWA	Multichannel
0.013±0.001	WORKMAN	12A	DPWA	$\gamma N \to N\pi$
0.018±0.002	DUGGER	07	DPWA	$\gamma N \to \pi N$
0.021±0.011	CRAWFORD	83	IPWA	$\gamma N \to \pi N$
0.034±0.005	AWAJI	81	DPWA	$\gamma N \to \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.009±0.001	SHKLYAR	13	DPWA	Multichannel
0.011±0.001	SHRESTHA	12A	DPWA	Multichannel
0.021±0.004	ANISOVICH	10	DPWA	Multichannel
0.015	DRECHSEL	07	DPWA	$\gamma N \to \pi N$
0.015±0.010	ARNDT	96	IPWA	$\gamma N \to \pi N$
0.012±0.002	LI	93	IPWA	$\gamma N \to \pi N$

$N(1675) \rightarrow p\gamma$, helicity-3/2 amplitude $A_{3/2}$

VALUE ($GeV^{-1/2}$)	DOCUMENT ID		TECN	COMMENT
+0.020±0.005 OUR ESTIMATE				
0.025±0.007	ANISOVICH	12A	DPWA	Multichannel
0.016±0.001	WORKMAN	12A	DPWA	$\gamma N \rightarrow N\pi$
0.021±0.001	DUGGER	07	DPWA	$\gamma N \rightarrow \pi N$
0.015±0.009	CRAWFORD	83	IPWA	$\gamma N \rightarrow \pi N$
0.024±0.008	AWAJI	81	DPWA	$\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.021±0.001	SHKLYAR	13	DPWA	Multichannel
0.020±0.001	SHRESTHA	12A	DPWA	Multichannel
0.024±0.008	ANISOVICH	10	DPWA	Multichannel
0.022	DRECHSEL	07	DPWA	$\gamma N \rightarrow \pi N$
0.010±0.007	ARNDT	96	IPWA	$\gamma N \rightarrow \pi N$
0.021±0.002	LI	93	IPWA	$\gamma N \rightarrow \pi N$

$N(1675) \rightarrow n\gamma$, helicity-1/2 amplitude $A_{1/2}$

VALUE ($GeV^{-1/2}$)	DOCUMENT ID		TECN	COMMENT
−0.060±0.005 OUR ESTIMATE				
−0.060±0.007	ANISOVICH	13B	DPWA	Multichannel
−0.058±0.002	CHEN	12A	DPWA	$\gamma N \rightarrow \pi N$
−0.057±0.024	AWAJI	81	DPWA	$\gamma N \rightarrow \pi N$
−0.033±0.004	FUJII	81	DPWA	$\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
−0.040±0.004	SHRESTHA	12A	DPWA	Multichannel
−0.062	DRECHSEL	07	DPWA	$\gamma N \rightarrow \pi N$
−0.049±0.010	ARNDT	96	IPWA	$\gamma N \rightarrow \pi N$
−0.060±0.003	LI	93	IPWA	$\gamma N \rightarrow \pi N$

$N(1675) \rightarrow n\gamma$, helicity-3/2 amplitude $A_{3/2}$

VALUE ($GeV^{-1/2}$)	DOCUMENT ID		TECN	COMMENT
−0.085±0.010 OUR ESTIMATE				
−0.088±0.010	ANISOVICH	13B	DPWA	Multichannel
−0.080±0.005	CHEN	12A	DPWA	$\gamma N \rightarrow \pi N$
−0.077±0.018	AWAJI	81	DPWA	$\gamma N \rightarrow \pi N$
−0.069±0.004	FUJII	81	DPWA	$\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
−0.068±0.004	SHRESTHA	12A	DPWA	Multichannel
−0.084	DRECHSEL	07	DPWA	$\gamma N \rightarrow \pi N$
−0.051±0.010	ARNDT	96	IPWA	$\gamma N \rightarrow \pi N$
−0.074±0.003	LI	93	IPWA	$\gamma N \rightarrow \pi N$

N(1675) FOOTNOTES

[1] LONGACRE 77 pole positions are from a search for poles in the unitarized T-matrix; the first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis. The other LONGACRE 77 values are from eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.

[2] From method II of LONGACRE 75: eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.

[3] Fit to the amplitudes of HOEHLER 79.

[4] See HOEHLER 93 for a detailed discussion of the evidence for and the pole parameters of N and Δ resonances as determined from Argand diagrams of πN elastic partial-wave amplitudes and from plots of the speeds with which the amplitudes traverse the diagrams.

[5] LONGACRE 78 values are from a search for poles in the unitarized T-matrix. The first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis.

[6] SAXON 80 finds the coupling phase is near 90°.

[7] LONGACRE 77 considers this coupling to be well determined.

N(1675) REFERENCES

For early references, see Physics Letters **111B** 1 (1982).

SVARC	14	PR C89 045205	A. Svarc *et al.*	
ANISOVICH	13B	EPJ A49 67	A.V. Anisovich *et al.*	
SHKLYAR	13	PR C87 015201	V. Shklyar, H. Lenske, U. Mosel	(GIES)
ANISOVICH	12A	EPJ A48 15	A.V. Anisovich *et al.*	(BONN, PNPI)
CHEN	12A	PR C86 015206	W. Chen *et al.*	(DUKE, GWU, MSST, ITEP+)
SHRESTHA	12A	PR C86 055203	M. Shrestha, D.M. Manley	(KSU)
WORKMAN	12A	PR C86 015202	R. Workman *et al.*	(GWU)
ANISOVICH	10	EPJ A44 203	A.V. Anisovich *et al.*	(BONN, PNPI)
BATINIC	10	PR C82 038203	M. Batinic *et al.*	(ZAGR)
THOMA	08	PL B659 87	U. Thoma *et al.*	(CB-ELSA Collab.)
DRECHSEL	07	EPJ A34 69	D. Drechsel, S.S. Kamalov, L. Tiator	(MAINZ, JINR)
DUGGER	07	PR C76 025211	M. Dugger *et al.*	(Jefferson Lab CLAS Collab.)
ARNDT	06	PR C74 045205	R.A. Arndt *et al.*	(GWU)
PDG	06	JP G33 1	W.-M. Yao *et al.*	(PDG Collab.)
ARNDT	04	PR C69 035213	R.A. Arndt *et al.*	(GWU, TRIU)
VRANA	00	PRPL 328 181	T.P. Vrana, S.A. Dytman,, T.-S.H. Lee	(PITT+)
ARNDT	96	PR C53 430	R.A. Arndt, I.I. Strakovsky, R.L. Workman	(VPI)
ARNDT	95	PR C52 2120	R.A. Arndt *et al.*	(VPI, BRCO)
HOEHLER	93	πN Newsletter 9 1	G. Hohler	(KARL)
LI	93	PR C47 2759	Z.J. Li *et al.*	(VPI)
MANLEY	92	PR D45 4002	D.M. Manley, E.M. Saleski	(KSA) IJP
Also		PR D30 904	D.M. Manley *et al.*	(VPI)
ARNDT	91	PR D43 2131	R.A. Arndt *et al.*	(VPI, TELE) IJP
BELL	83	NP B222 389	K.W. Bell *et al.*	(RL) IJP
CRAWFORD	83	NP B211 1	R.L. Crawford, W.T. Morton	(GLAS)
PDG	82	PL 111B 1	M. Roos *et al.*	(HELS, CIT, CERN)
AWAJI	81	Bonn Conf. 352	N. Awaji, R. Kajikawa	(NAGO)
Also		NP B197 365	K. Fujii *et al.*	(NAGO)
FUJII	81	NP B187 53	K. Fujii *et al.*	(NAGO, OSAK)
CUTKOSKY	80	Toronto Conf. 19	R.E. Cutkosky *et al.*	(CMU, LBL) IJP
Also		PR D20 2839	R.E. Cutkosky *et al.*	(CMU, LBL) IJP
SAXON	80	NP B162 522	D.H. Saxon *et al.*	(RHEL, BRIS) IJP
HOEHLER	79	PDAT 12-1	G. Hohler *et al.*	(KARLT) IJP
Also		Toronto Conf. 3	R. Koch	(KARLT) IJP
LONGACRE	78	PR D17 1795	R.S. Longacre *et al.*	(LBL, SLAC)
LONGACRE	77	NP B122 493	R.S. Longacre, J. Dolbeau	(SACL) IJP
Also		NP B108 365	J. Dolbeau *et al.*	(SACL) IJP
LONGACRE	75	PL 55B 415	R.S. Longacre *et al.*	(LBL, SLAC) IJP