

# $X(4230)$

$I^G(J^{PC}) = ?^?(1^{--})$

## OMITTED FROM SUMMARY TABLE

Enhancement reported by ABLIKIM 15C in  $e^+ e^- \rightarrow \omega \chi_{c0}$  at  $\sqrt{s} = 4.23\text{--}4.26$  GeV at  $9\sigma$  significance. Lineshape found to be inconsistent with origination from  $X(4260)$ . NEEDS CONFIRMATION.

## $X(4230)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4230±8±6</b>	180	<sup>1</sup> ABLIKIM	15C BES3	$e^+ e^- \rightarrow \omega \chi_{c0}$
<sup>1</sup> From a 3-parameter fit of measured cross sections from $\sqrt{s} = 4.21\text{--}4.42$ GeV to a phase-space modified Breit-Wigner function, using the decays $\chi_{c0} \rightarrow \pi^+ \pi^-$ , $\chi_{c0} \rightarrow K^+ K^-$ , and $\omega \rightarrow \pi^+ \pi^- \pi^0$ .				

## $X(4230)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>38±12±2</b>	180	<sup>1</sup> ABLIKIM	15C BES3	$e^+ e^- \rightarrow \omega \chi_{c0}$
<sup>1</sup> From a 3-parameter fit of measured cross sections from $\sqrt{s} = 4.21\text{--}4.42$ GeV to a phase-space modified Breit-Wigner function, using the decays $\chi_{c0} \rightarrow \pi^+ \pi^-$ , $\chi_{c0} \rightarrow K^+ K^-$ , and $\omega \rightarrow \pi^+ \pi^- \pi^0$ .				

## $X(4230)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 \quad e^+ e^-$	
$\Gamma_2 \quad \omega \chi_{c0}$	seen

## $X(4230) \Gamma(i) \Gamma(e^+ e^-)/\Gamma(\text{total})$

$\Gamma(\omega \chi_{c0}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_2 \Gamma_1 / \Gamma$
<b>2.7±0.5±0.4</b>	<sup>1</sup> ABLIKIM 15C BES3 $e^+ e^- \rightarrow \omega \chi_{c0}$
<sup>1</sup> From a 3-parameter fit of measured cross sections from $\sqrt{s} = 4.21\text{--}4.42$ GeV to a phase-space modified Breit-Wigner function, using the decays $\chi_{c0} \rightarrow \pi^+ \pi^-$ , $\chi_{c0} \rightarrow K^+ K^-$ , and $\omega \rightarrow \pi^+ \pi^- \pi^0$ .	

## $X(4230)$ BRANCHING RATIOS

$\Gamma(\omega \chi_{c0})/\Gamma_{\text{total}}$	$\Gamma_2/\Gamma$
<b>seen</b>	<sup>1</sup> ABLIKIM 15C BES3 $e^+ e^- \rightarrow \omega \chi_{c0}$

<sup>1</sup> From a 3-parameter fit of measured cross sections from  $\sqrt{s} = 4.21\text{--}4.42$  GeV to a phase-space modified Breit-Wigner function, using the decays  $\chi_{c0} \rightarrow \pi^+ \pi^-$ ,  $\chi_{c0} \rightarrow K^+ K^-$ , and  $\omega \rightarrow \pi^+ \pi^- \pi^0$ .

---

## X(4230) REFERENCES

---

ABLIKIM

15C PRL 114 092003

M. Ablikim *et al.*

(BES III Collab.)

---