

$\eta_b(1S)$

$I^G(J^{PC}) = 0^+(0^{-+})$

OMMITTED FROM SUMMARY TABLE

Quantum numbers shown are quark-model predictions. Observed in radiative decay of the $\Upsilon(3S)$, therefore $C = +$.


$\eta_b(1S)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
9398.0 \pm 3.2 OUR AVERAGE	Error includes scale factor of 1.8. See the ideogram below.			
9402.4 \pm 1.5 \pm 1.8	34k	¹ MIZUK	12	BELL $e^+ e^- \rightarrow \gamma \pi^+ \pi^-$ + hadrons
9391.8 \pm 6.6 \pm 2.0	$2.3 \pm 0.5k$	² BONVICINI	10	CLEO $\Upsilon(3S) \rightarrow \gamma X$
$9394.2^{+4.8}_{-4.9} \pm 2.0$	$13 \pm 5k$	² AUBERT	09AQ BABR	$\Upsilon(2S) \rightarrow \gamma X$
$9388.9^{+3.1}_{-2.3} \pm 2.7$	$19 \pm 3k$	² AUBERT	08V BABR	$\Upsilon(3S) \rightarrow \gamma X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
9393.2 \pm 3.4 \pm 2.3	10^{+5}_{-4}	^{2,3} DOBBS	12	$\Upsilon(2S) \rightarrow \gamma$ hadrons
9300 \pm 20 \pm 20		HEISTER	02D ALEP	181–209 $e^+ e^-$

¹ With floating width. Not independent of the corresponding mass difference measurement.

² Assuming $\Gamma_{\eta_b(1S)} = 10$ MeV. Not independent of the corresponding γ energy or mass difference measurements.

³ Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.




$m\gamma(1S) - m_{\eta_b}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
62.3±3.2 OUR AVERAGE	Error includes scale factor of 1.8. See the ideogram below.			
57.9±1.5±1.8	34k	⁴ MIZUK	12 BELL	$e^+ e^- \rightarrow \gamma\pi^+\pi^-$ + hadrons
68.5±6.6±2.0	2.3 ± 0.5k	⁵ BONVICINI	10 CLEO	$\gamma(3S) \rightarrow \gamma X$
66.1 ^{+4.8} _{-4.9} ±2.0	13 ± 5k	⁵ AUBERT	09AQ BABR	$\gamma(2S) \rightarrow \gamma X$
71.4 ^{+2.3} _{-3.1} ±2.7	19 ± 3k	⁵ AUBERT	08V BABR	$\gamma(3S) \rightarrow \gamma X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
67.1±3.4±2.3	10^{+5}_{-4}	^{5,6} DOBBS	12	$\gamma(2S) \rightarrow \gamma$ hadrons

⁴ With floating width. Not independent of the corresponding mass measurement.

⁵ Assuming $\Gamma_{\eta_b}(1S) = 10$ MeV. Not independent of the corresponding γ energy or mass measurements.

⁶ Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.



γ ENERGY IN $\Upsilon(3S)$ DECAY

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
920.6^{+2.8}_{-3.2} OUR AVERAGE				
918.6 \pm 6.0 \pm 1.9	2.3 \pm 0.5k	⁷ BONVICINI	10	CLEO $\Upsilon(3S) \rightarrow \gamma X$
921.2 ^{+2.1} _{-2.8} \pm 2.4	19 \pm 3k	⁷ AUBERT	08V	BABR $\Upsilon(3S) \rightarrow \gamma X$
⁷ Assuming $\Gamma_{\eta_b(1S)} = 10$ MeV. Not independent of the corresponding mass or mass difference measurements.				

γ ENERGY IN $\Upsilon(2S)$ DECAY

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
609.3^{+4.6}_{-4.5} \pm 1.9	13 \pm 5k	⁸ AUBERT	09AQ	BABR $\Upsilon(2S) \rightarrow \gamma X$
⁸ Assuming $\Gamma_{\eta_b(1S)} = 10$ MeV. Not independent of the corresponding mass or mass difference measurements.				

$\eta_b(1S)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
10.8^{+4.0+4.5}_{-3.7-2.0}	34k	⁹ MIZUK	12	BELL $e^+ e^- \rightarrow \gamma \pi^+ \pi^- +$ hadrons

⁹ With floating mass.

$\eta_b(1S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 hadrons	seen	
Γ_2 $3h^+ 3h^-$	not seen	
Γ_3 $2h^+ 2h^-$	not seen	
Γ_4 $4h^+ 4h^-$		
Γ_5 $\gamma\gamma$	not seen	
Γ_6 $\mu^+ \mu^-$	$<9 \times 10^{-3}$	90%
Γ_7 $\tau^+ \tau^-$	$<8\%$	90%

$\eta_b(1S) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_2\Gamma_5/\Gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<470	95	ABDALLAH	06	DLPH 161–209 $e^+ e^-$	
<132	95	HEISTER	02D	ALEP 181–209 $e^+ e^-$	

