

$f_1(1285)$

$I^G(J^{PC}) = 0^+(1^{++})$

$f_1(1285)$ MASS

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|------|--------------|-----------|--|
| 1281.9 ± 0.5 OUR AVERAGE | | | | Error includes scale factor of 1.8. See the ideogram below. |
| 1281.16 ± 0.39 ± 0.45 | | 1 LEES | 12X BABR | $\tau^- \rightarrow \pi^- f_1(1285) \nu_\tau$ |
| 1285.1 ± 1.0 ± 1.6 | | 2 ABLIKIM | 11J BES3 | $J/\psi \rightarrow \omega(\eta\pi^+\pi^-)$ |
| 1281 ± 2 ± 1 | | AUBERT | 07AU BABR | $10.6 e^+ e^- \rightarrow f_1(1285)\pi^+\pi^-\gamma$ |
| 1276.1 ± 8.1 ± 8.0 | 203 | BAI | 04J BES2 | $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$ |
| 1274 ± 6 | 237 | ABDALLAH | 03H DLPH | $91.2 e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp + X$ |
| 1280 ± 4 | | ACCIARRI | 01G L3 | |
| 1288 ± 4 ± 5 | 20k | ADAMS | 01B B852 | $18 \text{ GeV } \pi^- p \rightarrow K^+ K^- \pi^0 n$ |
| 1284 ± 6 | 1400 | ALDE | 97B GAM4 | $100 \pi^- p \rightarrow \eta\pi^0\pi^0 n$ |
| 1281 ± 1 | | BARBERIS | 97B OMEG | $450 pp \rightarrow pp2(\pi^+\pi^-)$ |
| 1281 ± 1 | | BARBERIS | 97C OMEG | $450 pp \rightarrow ppK_S^0 K^\pm \pi^\mp$ |
| 1280 ± 2 | | 3 ANTINORI | 95 OMEG | $300,450 pp \rightarrow pp2(\pi^+\pi^-)$ |
| 1282.2 ± 1.5 | | LEE | 94 MPS2 | $18 \pi^- p \rightarrow K^+ \bar{K}^0 2\pi^- p$ |
| 1279 ± 5 | | FUKUI | 91C SPEC | $8.95 \pi^- p \rightarrow \eta\pi^+\pi^- n$ |
| 1278 ± 2 | 140 | ARMSTRONG | 89 OMEG | $300 pp \rightarrow K\bar{K}\pi pp$ |
| 1278 ± 2 | | ARMSTRONG | 89G OMEG | $85 \pi^+ p \rightarrow 4\pi\pi p, pp \rightarrow 4\pi pp$ |
| 1280.1 ± 2.1 | 60 | RATH | 89 MPS | $21.4 \pi^- p \rightarrow K_S^0 K_S^0 \pi^0 n$ |
| 1285 ± 1 | 4750 | 4 BIRMAN | 88 MPS | $8 \pi^- p \rightarrow K^+ \bar{K}^0 \pi^- n$ |
| 1280 ± 1 | 504 | BITYUKOV | 88 SPEC | $32.5 \pi^- p \rightarrow K^+ K^- \pi^0 n$ |
| 1280 ± 4 | | ANDO | 86 SPEC | $8 \pi^- p \rightarrow \eta\pi^+\pi^- n$ |
| 1277 ± 2 | 420 | REEVES | 86 SPEC | $6.6 p\bar{p} \rightarrow K\bar{K}\pi X$ |
| 1285 ± 2 | | CHUNG | 85 SPEC | $8 \pi^- p \rightarrow N\bar{K}\bar{K}\pi$ |
| 1279 ± 2 | 604 | ARMSTRONG | 84 OMEG | $85 \pi^+ p \rightarrow K\bar{K}\pi\pi p, pp \rightarrow K\bar{K}\pi pp$ |
| 1286 ± 1 | | CHAUVAT | 84 SPEC | ISR 31.5 pp |
| 1278 ± 4 | | EVANGELIS... | 81 OMEG | $12 \pi^- p \rightarrow \eta\pi^+\pi^-\pi^- p$ |
| 1283 ± 3 | 103 | DIONISI | 80 HBC | $4 \pi^- p \rightarrow K\bar{K}\pi n$ |
| 1282 ± 2 | 320 | NACASCH | 78 HBC | $0.7, 0.76 \bar{p}p \rightarrow K\bar{K}3\pi$ |
| 1279 ± 5 | 210 | GRASSLER | 77 HBC | $16 \pi^\mp p$ |
| 1286 ± 3 | 180 | DUBOC | 72 HBC | $1.2 \bar{p}p \rightarrow 2K4\pi$ |
| 1283 ± 5 | | DAHL | 67 HBC | $1.6-4.2 \pi^- p$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-----------------------|------------------------|----------|--|--|
| 1284.2 \pm 2.2 | ⁵ AAIJ | 14Y LHCb | $\overline{B}_{(s)}^0 \rightarrow J/\psi 2(\pi^+ \pi^-)$ | |
| 1281.9 \pm 0.5 | ⁵ SOSA | 99 SPEC | $p\bar{p} \rightarrow p_{\text{slow}} (K_S^0 K^+ \pi^-) p_{\text{fast}}$ | |
| 1282.8 \pm 0.6 | ⁵ SOSA | 99 SPEC | $p\bar{p} \rightarrow p_{\text{slow}} (K_S^0 K^- \pi^+) p_{\text{fast}}$ | |
| 1270 \pm 10 | AMELIN | 95 VES | $37 \pi^- N \rightarrow \pi^- \pi^+ \pi^- \gamma N$ | |
| 1280 \pm 2 | ABATZIS | 94 OMEG | $450 p\bar{p} \rightarrow pp2(\pi^+ \pi^-)$ | |
| 1282 \pm 4 | ARMSTRONG | 93C E760 | $\bar{p}p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$ | |
| 1270 \pm 6 \pm 10 | ARMSTRONG | 92C OMEG | $300 p\bar{p} \rightarrow pp\pi^+ \pi^- \gamma$ | |
| 1281 \pm 1 | ARMSTRONG | 89E OMEG | $300 p\bar{p} \rightarrow pp2(\pi^+ \pi^-)$ | |
| 1279 \pm 6 \pm 10 | BECKER | 87 MRK3 | $e^+ e^- \rightarrow \phi K\bar{K}\pi$ | |
| 1286 \pm 9 | GIDAL | 87 MRK2 | $e^+ e^- \rightarrow e^+ e^- \eta \pi^+ \pi^-$ | |
| 1287 \pm 5 | BITYUKOV | 84B SPEC | $32 \pi^- p \rightarrow K^+ K^- \pi^0 n$ | |
| ~ 1279 | ⁶ TORNQVIST | 82B RVUE | | |
| 1275 \pm 6 | BROMBERG | 80 SPEC | $100 \pi^- p \rightarrow K\bar{K}\pi X$ | |
| 1288 \pm 9 | GURTU | 79 HBC | $4.2 K^- p \rightarrow n\eta 2\pi$ | |
| ~ 1275.0 | ⁷ STANTON | 79 CNTR | $8.5 \pi^- p \rightarrow n2\gamma 2\pi$ | |
| 1271 \pm 10 | CORDEN | 78 OMEG | $12-15 \pi^- p \rightarrow K^+ K^- \pi n$ | |
| 1295 \pm 12 | CORDEN | 78 OMEG | $12-15 \pi^- p \rightarrow n5\pi$ | |
| 1292 \pm 10 | DEFOIX | 72 HBC | $0.7 \bar{p}p \rightarrow 7\pi$ | |
| 1280 \pm 3 | 500 ⁸ THUN | 72 MMS | $13.4 \pi^- p$ | |
| 1303 \pm 8 | BARDADIN-... | 71 HBC | $8 \pi^+ p \rightarrow p6\pi$ | |
| 1283 \pm 6 | BOESEBECK | 71 HBC | $16.0 \pi p \rightarrow p5\pi$ | |
| 1270 \pm 10 | CAMPBELL | 69 DBC | $2.7 \pi^+ d$ | |
| 1285 \pm 7 | LORSTAD | 69 HBC | $0.7 \bar{p}p, 4,5\text{-body}$ | |
| 1290 \pm 7 | D'ANDLAU | 68 HBC | $1.2 \bar{p}p, 5-6 \text{ body}$ | |

¹ Using the $2\pi^+ 2\pi^-$ and $\pi^+ \pi^- \eta$ modes of $f_1(1285)$ decay.

² The selected process is $J/\psi \rightarrow \omega a_0(980)\pi$.

³ Supersedes ABATZIS 94, ARMSTRONG 89E.


⁴ From partial wave analysis of $K^+ \bar{K}^0 \pi^-$ system.

⁵ No systematic error given.

⁶ From a unitarized quark-model calculation.

⁷ From phase shift analysis of $\eta \pi^+ \pi^-$ system.

⁸ Seen in the missing mass spectrum.



$f_1(1285)$ mass (MeV)

$f_1(1285)$ WIDTH

Only experiments giving width error less than 20 MeV are kept for averaging.

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------|------|-------------|-----------|---|
| 24.2± 1.1 OUR AVERAGE | | | | Error includes scale factor of 1.3. See the ideogram below. |
| $22.0 \pm 3.1 \pm 2.0$ | 1 | ABLIKIM | 11J BES3 | $J/\psi \rightarrow \omega(\eta\pi^+\pi^-)$ |
| $35 \pm 6 \pm 4$ | | AUBERT | 07AU BABR | $10.6 e^+ e^- \rightarrow f_1(1285)\pi^+\pi^-\gamma$ |
| $40.0 \pm 8.6 \pm 9.3$ | 203 | BAI | 04J BES2 | $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$ |
| 29 ± 12 | 237 | ABDALLAH | 03H DLPH | $91.2 e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp + X$ |
| $45 \pm 9 \pm 7$ | 20k | ADAMS | 01B B852 | $18 \text{ GeV } \pi^- p \rightarrow K^+ K^- \pi^0 n$ |

| | | | | | |
|----------------|------|-----------------------|-----|------|---|
| 55 \pm 18 | 1400 | ALDE | 97B | GAM4 | $100 \pi^- p \rightarrow \eta \pi^0 \pi^0 n$ |
| 24 \pm 3 | | BARBERIS | 97B | OMEG | $450 pp \rightarrow pp2(\pi^+ \pi^-)$ |
| 20 \pm 2 | | BARBERIS | 97C | OMEG | $450 pp \rightarrow ppK_S^0 K^\pm \pi^\mp$ |
| 36 \pm 5 | | ² ANTINORI | 95 | OMEG | $300,450 pp \rightarrow pp2(\pi^+ \pi^-)$ |
| 29.0 \pm 4.1 | | LEE | 94 | MPS2 | $18 \pi^- p \rightarrow K^+ \bar{K}^0 2\pi^- p$ |
| 25 \pm 4 | 140 | ARMSTRONG | 89 | OMEG | $300 pp \rightarrow K\bar{K}\pi pp$ |
| 22 \pm 2 | 4750 | ³ BIRMAN | 88 | MPS | $8 \pi^- p \rightarrow K^+ \bar{K}^0 \pi^- n$ |
| 25 \pm 4 | 504 | BITYUKOV | 88 | SPEC | $32.5 \pi^- p \rightarrow K^+ K^- \pi^0 n$ |
| 19 \pm 5 | | ANDO | 86 | SPEC | $8 \pi^- p \rightarrow \eta \pi^+ \pi^- n$ |
| 32 \pm 8 | 420 | REEVES | 86 | SPEC | $6.6 p\bar{p} \rightarrow K K \pi X$ |
| 22 \pm 2 | | CHUNG | 85 | SPEC | $8 \pi^- p \rightarrow N K \bar{K} \pi$ |
| 32 \pm 3 | 604 | ARMSTRONG | 84 | OMEG | $85 \pi^+ p \rightarrow K\bar{K}\pi\pi p,$ $pp \rightarrow K\bar{K}\pi pp$ |
| 24 \pm 3 | | CHAUVAT | 84 | SPEC | ISR 31.5 pp |
| 29 \pm 10 | 103 | DIONISI | 80 | HBC | $4 \pi^- p \rightarrow K\bar{K}\pi n$ |
| 28.3 \pm 6.7 | 320 | NACASCH | 78 | HBC | $0.7, 0.76 \bar{p}p \rightarrow K\bar{K}3\pi$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-----------------|----------|----------------------|-----|------|--|
| 32.4 \pm 5.8 | | ⁴ AAIJ | 14Y | LHCb | $\bar{B}_{(s)}^0 \rightarrow J/\psi 2(\pi^+ \pi^-)$ |
| 18.2 \pm 1.2 | | ⁴ SOSA | 99 | SPEC | $pp \rightarrow p_{\text{slow}} (K_S^0 K^+ \pi^-)$ |
| 19.4 \pm 1.5 | | ⁴ SOSA | 99 | SPEC | $pp \rightarrow p_{\text{slow}} (K_S^0 K^- \pi^+)$ |
| | | | | | p_{fast} |
| 40 \pm 5 | | ABATZIS | 94 | OMEG | $450 pp \rightarrow pp2(\pi^+ \pi^-)$ |
| 31 \pm 5 | | ARMSTRONG | 89E | OMEG | $300 pp \rightarrow pp2(\pi^+ \pi^-)$ |
| 41 \pm 12 | | ARMSTRONG | 89G | OMEG | $85 \pi^+ p \rightarrow 4\pi\pi p, pp \rightarrow 4\pi pp$ |
| 17.9 \pm 10.9 | 60 | RATH | 89 | MPS | $21.4 \pi^- p \rightarrow K_S^0 K_S^0 \pi^0 n$ |
| 14 \pm 10 | \pm 10 | BECKER | 87 | MRK3 | $e^+ e^- \rightarrow \phi K\bar{K}\pi$ |
| 26 \pm 12 | | EVANGELIS... | 81 | OMEG | $12 \pi^- p \rightarrow \eta \pi^+ \pi^- \pi^- p$ |
| 25 \pm 15 | 200 | GURTU | 79 | HBC | $4.2 K^- p \rightarrow n\eta 2\pi$ |
| \sim 10 | | ⁵ STANTON | 79 | CNTR | $8.5 \pi^- p \rightarrow n 2\gamma 2\pi$ |
| 24 \pm 18 | 210 | GRASSLER | 77 | HBC | $16 \pi^\mp p$ |
| 28 \pm 5 | 150 | ⁶ DEFOIX | 72 | HBC | $0.7 \bar{p}p \rightarrow 7\pi$ |
| 46 \pm 9 | 180 | ⁶ DUBOC | 72 | HBC | $1.2 \bar{p}p \rightarrow 2K 4\pi$ |
| 37 \pm 5 | 500 | ⁷ THUN | 72 | MMS | $13.4 \pi^- p$ |
| 10 \pm 10 | | BOESEBECK | 71 | HBC | $16.0 \pi p \rightarrow p 5\pi$ |
| 30 \pm 15 | | CAMPBELL | 69 | DBC | $2.7 \pi^+ d$ |
| 60 \pm 15 | | ⁶ LORSTAD | 69 | HBC | $0.7 \bar{p}p, 4,5\text{-body}$ |
| 35 \pm 10 | | ⁶ DAHL | 67 | HBC | $1.6\text{--}4.2 \pi^- p$ |

¹ The selected process is $J/\psi \rightarrow \omega a_0(980) \pi$.

² Supersedes ABATZIS 94, ARMSTRONG 89E.


³ From partial wave analysis of $K^+ \bar{K}^0 \pi^-$ system.

⁴ No systematic error given.

⁵ From phase shift analysis of $\eta \pi^+ \pi^-$ system.

⁶ Resolution is not unfolded.

⁷ Seen in the missing mass spectrum.



$f_1(1285)$ DECAY MODES

| Mode | Fraction (Γ_i/Γ) | Scale factor/ Confidence level |
|--|--------------------------------|-----------------------------------|
| $\Gamma_1 \quad 4\pi$ | $(33.1 \pm 2.1) \%$ | $S=1.3$ |
| $\Gamma_2 \quad \pi^0 \pi^0 \pi^+ \pi^-$ | $(22.0 \pm 1.4) \%$ | $S=1.3$ |
| $\Gamma_3 \quad 2\pi^+ 2\pi^-$ | $(11.0 \pm 0.7) \%$ | $S=1.3$ |
| $\Gamma_4 \quad \rho^0 \pi^+ \pi^-$ | $(11.0 \pm 0.7) \%$ | $S=1.3$ |
| $\Gamma_5 \quad \rho^0 \rho^0$ | seen | |
| $\Gamma_6 \quad 4\pi^0$ | $< 7 \times 10^{-4}$ | $CL=90\%$ |
| $\Gamma_7 \quad \eta \pi^+ \pi^-$ | $(35 \pm 15) \%$ | |
| $\Gamma_8 \quad \eta \pi \pi$ | $(52.4 \pm 1.9) \%$ | $S=1.2$ |
| $\Gamma_9 \quad a_0(980)\pi$ [ignoring $a_0(980) \rightarrow K\bar{K}$] | $(36 \pm 7) \%$ | |
| $\Gamma_{10} \quad \eta \pi \pi$ [excluding $a_0(980)\pi$] | $(16 \pm 7) \%$ | |
| $\Gamma_{11} \quad K\bar{K}\pi$ | $(9.0 \pm 0.4) \%$ | $S=1.1$ |

| | | |
|---------------|-------------------|--------------------------------|
| Γ_{12} | $K\bar{K}^*(892)$ | not seen |
| Γ_{13} | $\pi^+\pi^-\pi^0$ | $(3.0 \pm 0.9) \times 10^{-3}$ |
| Γ_{14} | $\rho^\pm\pi^\mp$ | $< 3.1 \times 10^{-3}$ CL=95% |
| Γ_{15} | $\gamma\rho^0$ | $(5.5 \pm 1.3)\%$ S=2.8 |
| Γ_{16} | $\phi\gamma$ | $(7.4 \pm 2.6) \times 10^{-4}$ |
| Γ_{17} | $\gamma\gamma^*$ | |
| Γ_{18} | $\gamma\gamma$ | |

CONSTRAINED FIT INFORMATION

An overall fit to 7 branching ratios uses 16 measurements and one constraint to determine 5 parameters. The overall fit has a $\chi^2 = 24.7$ for 12 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

| | | | | |
|----------|-------|-------|----------|----------|
| x_9 | -17 | | | |
| x_{10} | -8 | -95 | | |
| x_{11} | 46 | -9 | -4 | |
| x_{15} | -36 | -4 | -2 | -34 |
| | x_1 | x_9 | x_{10} | x_{11} |

$f_1(1285) \Gamma(i) \Gamma(\gamma\gamma) / \Gamma(\text{total})$

| $\Gamma(\eta\pi\pi) \times \Gamma(\gamma\gamma) / \Gamma_{\text{total}}$ | | $\Gamma_8 \Gamma_{18} / \Gamma = (\Gamma_9 + \Gamma_{10}) \Gamma_{18} / \Gamma$ | | |
|--|-----|---|------|--|
| VALUE (keV) | CL% | DOCUMENT ID | TECN | COMMENT |
| <0.62 | 95 | GIDAL | 87 | $e^+ e^- \rightarrow e^+ e^- \eta\pi^+\pi^-$ |

| $\Gamma(\eta\pi\pi) \times \Gamma(\gamma\gamma^*) / \Gamma_{\text{total}}$ | | $\Gamma_8 \Gamma_{17} / \Gamma = (\Gamma_9 + \Gamma_{10}) \Gamma_{17} / \Gamma$ | | |
|---|------|---|---------|--|
| VALUE (keV) | EVTS | DOCUMENT ID | TECN | COMMENT |
| 1.4 ± 0.4 OUR AVERAGE | | Error includes scale factor of 1.4. | | |
| 1.18 ± 0.25 ± 0.20 | 26 | 1,2 AIHARA | 88B TPC | $e^+ e^- \rightarrow e^+ e^- \eta\pi^+\pi^-$ |
| 2.30 ± 0.61 ± 0.42 | | 1,3 GIDAL | 87 MRK2 | $e^+ e^- \rightarrow e^+ e^- \eta\pi^+\pi^-$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 1.8 ± 0.3 ± 0.3 | 420 | 4 ACHARD | 02B L3 | $183-209 e^+ e^- \rightarrow e^+ e^- \eta\pi^+\pi^-$ |

¹ Assuming a ρ -pole form factor.

² Published value multiplied by $\eta\pi\pi$ branching ratio 0.49.

³ Published value divided by 2 and multiplied by the $\eta\pi\pi$ branching ratio 0.49.

⁴ Published value multiplied by the $\eta\pi\pi$ branching ratio 0.52.

$f_1(1285)$ BRANCHING RATIOS **$\Gamma(K\bar{K}\pi)/\Gamma(4\pi)$**

$$\Gamma_{11}/\Gamma_1$$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------------------|---|-------------|----------------|
| 0.271±0.016 OUR FIT | Error includes scale factor of 1.3. | | |
| 0.271±0.016 OUR AVERAGE | Error includes scale factor of 1.2. | | |
| 0.265±0.014 | 1 BARBERIS 97C OMEG 450 $p p \rightarrow p p K_S^0 K^\pm \pi^\mp$ | | |
| 0.28 ± 0.05 | 2 ARMSTRONG 89E OMEG 300 $p p \rightarrow p p f_1(1285)$ | | |
| 0.37 ± 0.03 ± 0.05 | 3 ARMSTRONG 89G OMEG 85 $\pi p \rightarrow 4\pi X$ | | |

¹ Using $2(\pi^+ \pi^-)$ data from BARBERIS 97B.² Assuming $\rho\pi\pi$ and $a_0(980)\pi$ intermediate states.³ 4π consistent with being entirely $\rho\pi\pi$. **$\Gamma(\pi^0\pi^+\pi^-)/\Gamma_{\text{total}}$**

$$\Gamma_2/\Gamma = \frac{2}{3}\Gamma_1/\Gamma$$

| <u>VALUE</u> | <u>DOCUMENT ID</u> |
|--|-------------------------------------|
| 0.220^{+0.014}_{-0.012} OUR FIT | Error includes scale factor of 1.3. |

 $\Gamma(2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$

$$\Gamma_3/\Gamma = \frac{1}{3}\Gamma_1/\Gamma$$

| <u>VALUE</u> | <u>DOCUMENT ID</u> |
|--|-------------------------------------|
| 0.110^{+0.007}_{-0.006} OUR FIT | Error includes scale factor of 1.3. |

 $\Gamma(\rho^0\pi^+\pi^-)/\Gamma_{\text{total}}$

$$\Gamma_4/\Gamma = \frac{1}{3}\Gamma_1/\Gamma$$

| <u>VALUE</u> | <u>DOCUMENT ID</u> |
|--|-------------------------------------|
| 0.110^{+0.007}_{-0.006} OUR FIT | Error includes scale factor of 1.3. |

 $\Gamma(\rho^0\pi^+\pi^-)/\Gamma(2\pi^+ 2\pi^-)$

$$\Gamma_4/\Gamma_3$$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------------------------------|-------------|----------------|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 1.0±0.4 | GRASSLER 77 HBC 16 GeV $\pi^\pm p$ | | |

 $\Gamma(\rho^0\rho^0)/\Gamma_{\text{total}}$

$$\Gamma_5/\Gamma$$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>COMMENT</u> |
|--------------|---|----------------|
| seen | BARBERIS 00C 450 $p p \rightarrow p_f 4\pi p_s$ | |

 $\Gamma(4\pi^0)/\Gamma_{\text{total}}$

$$\Gamma_6/\Gamma$$

| <u>VALUE (units 10^{-4})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|------------------------------------|
| <7 | 90 | ALDE 87 | GAM4 | $100 \pi^- p \rightarrow 4\pi^0 n$ |

 $\Gamma(\pi^+\pi^-\pi^0)/\Gamma(\eta\pi^+\pi^-)$

$$\Gamma_{13}/\Gamma_7$$

| <u>VALUE (%)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------|-------------|--------------------|-------------|---|
| 0.86±0.16±0.20 | 2.3k | 1 DOROFEEV 11 VES | | $\pi^- N \rightarrow \pi^- f_1(1285) N$ |

¹ Value obtained selecting the region corresponding to $f_0(980)$ in the $\pi^+\pi^-$ mass spectrum.

$\Gamma(\eta\pi\pi)/\Gamma_{\text{total}}$

VALUE

0.524^{+0.019}_{-0.022} OUR FIT Error includes scale factor of 1.2.

DOCUMENT ID

$\Gamma_8/\Gamma = (\Gamma_9 + \Gamma_{10})/\Gamma$

$\Gamma(4\pi)/\Gamma(\eta\pi\pi)$

VALUE

DOCUMENT ID

TECN

COMMENT

0.63^{+0.06}_{-0.06} OUR FIT Error includes scale factor of 1.2.

0.41^{+0.14}_{-0.14} OUR AVERAGE

$0.37 \pm 0.11 \pm 0.11$ BOLTON 92 MRK3 $J/\psi \rightarrow \gamma f_1(1285)$

0.64 ± 0.40 GURTU 79 HBC 4.2 $K^- p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.93 ± 0.30 ¹ GRASSLER 77 HBC 16 $\pi^\mp p$

¹ Assuming $\rho\pi\pi$ and $a_0(980)\pi$ intermediate states.

$\Gamma_1/\Gamma_8 = \Gamma_1/(\Gamma_9 + \Gamma_{10})$

$\Gamma(2\pi^+ 2\pi^-)/\Gamma(\eta\pi\pi)$

VALUE

DOCUMENT ID

TECN

COMMENT

0.28^{+0.02}_{-0.02}^{+0.02}_{-0.02}

¹ LEES 12X BABR $\tau^- \rightarrow \pi^- f_1(1285) \nu_\tau$

¹ Assuming $B(f_1(1285) \rightarrow \pi\pi\eta) = 3/2 B(f_1(1285) \rightarrow \pi^+\pi^-\eta)$.

Γ_3/Γ_8

$\Gamma(a_0(980)\pi \text{ [ignoring } a_0(980) \rightarrow K\bar{K}])/\Gamma(\eta\pi\pi)$

VALUE CL% EVTS

DOCUMENT ID

TECN

COMMENT

0.69^{+0.13}_{-0.12} OUR FIT

0.69^{+0.13}_{-0.12} OUR AVERAGE

0.72 ± 0.15 GURTU 79 HBC 4.2 $K^- p$

$0.6^{+0.3}_{-0.2}$ CORDEN 78 OMEG 12–15 $\pi^- p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

>0.69 95 318 ACHARD 02B L3 $183\text{--}209 e^+ e^- \rightarrow e^+ e^- \eta \pi^+ \pi^-$

0.28 ± 0.07 1400 ALDE 97B GAM4 100 $\pi^- p \rightarrow \eta \pi^0 \pi^0 n$

1.0 ± 0.3 GRASSLER 77 HBC 16 $\pi^\mp p$

$\Gamma_{11}/\Gamma_8 = \Gamma_{11}/(\Gamma_9 + \Gamma_{10})$

$\Gamma(K\bar{K}\pi)/\Gamma(\eta\pi\pi)$

VALUE

DOCUMENT ID

TECN

COMMENT

0.171^{+0.013}_{-0.013} OUR FIT Error includes scale factor of 1.1.

0.170^{+0.012}_{-0.012} OUR AVERAGE

$0.166 \pm 0.01 \pm 0.008$ BARBERIS 98C OMEG 450 $p p \rightarrow p_f f_1(1285) p_s$

0.42 ± 0.15 GURTU 79 HBC 4.2 $K^- p$

0.5 ± 0.2 ¹ CORDEN 78 OMEG 12–15 $\pi^- p$

0.20 ± 0.08 ² DEFOIX 72 HBC 0.7 $\bar{p} p \rightarrow 7\pi$

0.16 ± 0.08 CAMPBELL 69 DBC 2.7 $\pi^+ d$

¹ CORDEN 78 assumes low-mass $\eta\pi\pi$ region is dominantly 1^{++} . See BARBERIS 98C and MANAK 00A for discussion.

² $K\bar{K}$ system characterized by the $I = 1$ threshold enhancement. (See under $a_0(980)$).

$\Gamma(K\bar{K}^*(892))/\Gamma_{\text{total}}$ Γ_{12}/Γ

| VALUE | DOCUMENT ID | TECN | COMMENT | Γ_{12}/Γ |
|---|---------------------------|------|---|----------------------|
| not seen | NACASCH 78 | HBC | $0.7, 0.76 \bar{p}p \rightarrow K\bar{K}3\pi$ | |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| seen | ¹ ACHARD 07 L3 | | $183-209 e^+e^- \rightarrow e^+e^- K_S^0 K^\pm \pi^\mp$ | |

¹ A clear signal of 19.8 ± 4.4 events observed at high Q^2 .

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{13}/Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT | Γ_{13}/Γ |
|-------------------------|------|--------------------------|------|---|----------------------|
| 0.30±0.055±0.074 | 2.3k | ¹ DOROFEEV 11 | VES | $\pi^- N \rightarrow \pi^- f_1(1285) N$ | |

¹ Value obtained selecting the region corresponding to $f_0(980)$ in the $\pi^+\pi^-$ mass spectrum. The systematic error includes the uncertainty on the partial width $f_1 \rightarrow \eta\pi\pi$ obtained from PDG 10 data.

$\Gamma(\rho^\pm\pi^\mp)/\Gamma_{\text{total}}$ Γ_{14}/Γ

| VALUE (%) | CL% | DOCUMENT ID | TECN | COMMENT | Γ_{14}/Γ |
|-----------------|-----|-------------|------|---|----------------------|
| <0.31 | 95 | DOROFEEV 11 | VES | $\pi^- N \rightarrow \pi^- f_1(1285) N$ | |

$\Gamma(\gamma\rho^0)/\Gamma_{\text{total}}$ Γ_{15}/Γ

| VALUE (units 10^{-2}) | CL% | DOCUMENT ID | TECN | COMMENT | Γ_{15}/Γ |
|--------------------------|-----|-------------------------------------|------|---------|----------------------|
| 5.5±1.3 OUR FIT | | Error includes scale factor of 2.8. | | | |

2.8±0.7±0.6 AMELIN 95 VES $37 \pi^- N \rightarrow \pi^-\pi^+\pi^-\gamma N$
• • • We do not use the following data for averages, fits, limits, etc. **• • •**

<5 95 BITYUKOV 91B SPEC $32 \pi^- p \rightarrow \pi^+\pi^-\gamma n$

$\Gamma(\gamma\rho^0)/\Gamma(2\pi^+2\pi^-)$ $\Gamma_{15}/\Gamma_3 = \Gamma_{15}/\frac{1}{3}\Gamma_1$

| VALUE | DOCUMENT ID | TECN | COMMENT | $\Gamma_{15}/\Gamma_3 = \Gamma_{15}/\frac{1}{3}\Gamma_1$ |
|--------------------------|-------------------------------------|------|---------|--|
| 0.50±0.13 OUR FIT | Error includes scale factor of 2.5. | | | |

| | |
|------------------|--|
| 0.45±0.18 | ¹ COFFMAN 90 MRK3 $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$ |
|------------------|--|

¹ Using $B(J/\psi \rightarrow \gamma f_1(1285) \rightarrow \gamma\gamma\rho^0) = 0.25 \times 10^{-4}$ and $B(J/\psi \rightarrow \gamma f_1(1285) \rightarrow \gamma 2\pi^+ 2\pi^-) = 0.55 \times 10^{-4}$ given by MIR 88.

$\Gamma(\eta\pi\pi)/\Gamma(\gamma\rho^0)$ $\Gamma_8/\Gamma_{15} = (\Gamma_9 + \Gamma_{10})/\Gamma_{15}$

| VALUE | DOCUMENT ID | TECN | COMMENT | $\Gamma_8/\Gamma_{15} = (\Gamma_9 + \Gamma_{10})/\Gamma_{15}$ |
|------------------------|-------------------------------------|------|---------|---|
| 9.5±2.0 OUR FIT | Error includes scale factor of 2.5. | | | |

7.9±0.9 OUR AVERAGE

| | |
|--------------|---|
| 10.0±1.0±2.0 | BARBERIS 98C OMEG 450 $\bar{p}p \rightarrow p_f f_1(1285) p_s$ |
| 7.5±1.0 | ¹ ARMSTRONG 92C OMEG 300 $\bar{p}p \rightarrow p p \pi^+\pi^-\gamma, p p \eta\pi^+\pi^-$ |

¹ Published value multiplied by 1.5.

$\Gamma(\gamma\rho^0)/\Gamma(K\bar{K}\pi)$ Γ_{15}/Γ_{11}

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT | Γ_{15}/Γ_{11} |
|---|-----|-------------|------|---------|---------------------------|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |

| | | |
|--------|----|--|
| >0.035 | 90 | ¹ COFFMAN 90 MRK3 $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$ |
|--------|----|--|

¹ Using $B(J/\psi \rightarrow \gamma f_1(1285) \rightarrow \gamma\gamma\rho^0) = 0.25 \times 10^{-4}$ and $B(J/\psi \rightarrow \gamma f_1(1285) \rightarrow \gamma K\bar{K}\pi) < 0.72 \times 10^{-3}$.

| $\Gamma(\phi\gamma)/\Gamma(K\bar{K}\pi)$ | | | Γ_{16}/Γ_{11} | | |
|---|-----|------|---------------------------|------|---|
| VALUE (units 10^{-2}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
| 0.82±0.21±0.20 | | 19 | BITYUKOV | 88 | SPEC $32.5 \pi^- p \rightarrow K^+ K^- \pi^0 n$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| <0.50 | 95 | | BARBERIS | 98C | OMEG $450 pp \rightarrow p_f f_1(1285) p_s$ |
| <0.93 | 95 | | AMELIN | 95 | VES $37 \pi^- N \rightarrow \pi^- \pi^+ \pi^- \gamma N$ |

f₁(1285) REFERENCES

| | | | | |
|--------------|------|-----------------------------|------------------------------|-----------------------------|
| AAIJ | 14Y | PRL 112 091802 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| LEES | 12X | PR D86 092010 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| ABLIKIM | 11J | PRL 107 182001 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| DOROFEEV | 11 | EPJ A47 68 | V. Dorofeev <i>et al.</i> | (SERP, MIPT) |
| PDG | 10 | JP G37 075021 | K. Nakamura <i>et al.</i> | (PDG Collab.) |
| ACHARD | 07 | JHEP 0703 018 | P. Achard <i>et al.</i> | (L3 Collab.) |
| AUBERT | 07AU | PR D76 092005 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| BAI | 04J | PL B594 47 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| ABDALLAH | 03H | PL B569 129 | J. Abdallah <i>et al.</i> | (DELPHI Collab.) |
| ACHARD | 02B | PL B526 269 | P. Achard <i>et al.</i> | (L3 Collab.) |
| ACCIARRI | 01G | PL B501 1 | M. Acciari <i>et al.</i> | (L3 Collab.) |
| ADAMS | 01B | PL B516 264 | G.S. Adams <i>et al.</i> | (BNL E852 Collab.) |
| BARBERIS | 00C | PL B471 440 | D. Barberis <i>et al.</i> | (WA 102 Collab.) |
| MANAK | 00A | PR D62 012003 | J.J. Manak <i>et al.</i> | (BNL E852 Collab.) |
| SOSA | 99 | PRL 83 913 | M. Sosa <i>et al.</i> | |
| BARBERIS | 98C | PL B440 225 | D. Barberis <i>et al.</i> | (WA 102 Collab.) |
| ALDE | 97B | PAN 60 386 | D. Alde <i>et al.</i> | (GAMS Collab.) |
| | | Translated from YAF 60 458. | | |
| BARBERIS | 97B | PL B413 217 | D. Barberis <i>et al.</i> | (WA 102 Collab.) |
| BARBERIS | 97C | PL B413 225 | D. Barberis <i>et al.</i> | (WA 102 Collab.) |
| AMELIN | 95 | ZPHY C66 71 | D.V. Amelin <i>et al.</i> | (VES Collab.) |
| ANTINORI | 95 | PL B353 589 | F. Antinori <i>et al.</i> | (ATHU, BARI, BIRM+) |
| ABATZIS | 94 | PL B324 509 | S. Abatzis <i>et al.</i> | (ATHU, BARI, BIRM+) |
| LEE | 94 | PL B323 227 | J.H. Lee <i>et al.</i> | (BNL, IND, KYUN, MASD+) |
| ARMSTRONG | 93C | PL B307 394 | T.A. Armstrong <i>et al.</i> | (FNAL, FERR, GENO+) |
| ARMSTRONG | 92C | ZPHY C54 371 | T.A. Armstrong <i>et al.</i> | (ATHU, BARI, BIRM+) |
| BOLTON | 92 | PL B278 495 | T. Bolton <i>et al.</i> | (Mark III Collab.) |
| BITYUKOV | 91B | SJNP 54 318 | S.I. Bityukov <i>et al.</i> | (SERP) |
| | | Translated from YAF 54 529. | | |
| FUKUI | 91C | PL B267 293 | S. Fukui <i>et al.</i> | (SUGI, NAGO, KEK, KYOT+) |
| COFFMAN | 90 | PR D41 1410 | D.M. Coffman <i>et al.</i> | (Mark III Collab.) |
| ARMSTRONG | 89 | PL B221 216 | T.A. Armstrong <i>et al.</i> | (CERN, CDEF, BIRM+) JPC |
| ARMSTRONG | 89E | PL B228 536 | T.A. Armstrong, M. Benayoun | (ATHU, BARI, BIRM+) |
| ARMSTRONG | 89G | ZPHY C43 55 | T.A. Armstrong <i>et al.</i> | (CERN, BIRM, BARI+) |
| RATH | 89 | PR D40 693 | M.G. Rath <i>et al.</i> | (NDAM, BRAN, BNL, CUNY+) |
| AIHARA | 88B | PL B209 107 | H. Aihara <i>et al.</i> | (TPC-2 γ Collab.) |
| BIRMAN | 88 | PRL 61 1557 | A. Birman <i>et al.</i> | (BNL, FSU, IND, MASD) JP |
| BITYUKOV | 88 | PL B203 327 | S.I. Bityukov <i>et al.</i> | (SERP) |
| MIR | 88 | Photon-Photon 88, 126 | R. Mir | (Mark III Collab.) |
| Conference | | | | |
| ALDE | 87 | PL B198 286 | D.M. Alde <i>et al.</i> | (LANL, BRUX, SERP, LAPP) |
| BECKER | 87 | PRL 59 186 | J.J. Becker <i>et al.</i> | (Mark III Collab.) |
| GIDAL | 87 | PRL 59 2012 | G. Gidal <i>et al.</i> | (LBL, SLAC, HARV) |
| ANDO | 86 | PRL 57 1296 | A. Ando <i>et al.</i> | (KEK, KYOT, NIR, SAGA+) IJP |
| REEVES | 86 | PR D34 1960 | D.F. Reeves <i>et al.</i> | (FLOR, BNL, IND+) JP |
| CHUNG | 85 | PRL 55 779 | S.U. Chung <i>et al.</i> | (BNL, FLOR, IND+) JP |
| ARMSTRONG | 84 | PL 146B 273 | T.A. Armstrong <i>et al.</i> | (ATHU, BARI, BIRM+) JP |
| BITYUKOV | 84B | PL 144B 133 | S.I. Bityukov <i>et al.</i> | (SERP) |
| CHAUVAT | 84 | PL 148B 382 | P. Chauvat <i>et al.</i> | (CERN, CLER, UCLA+) |
| TORNQVIST | 82B | NP B203 268 | N.A. Tornqvist | (HELS) |
| EVANGELIS... | 81 | NP B178 197 | C. Evangelista <i>et al.</i> | (BARI, BONN, CERN+) |
| BROMBERG | 80 | PR D22 1513 | C.M. Bromberg <i>et al.</i> | (CIT, FNAL, ILLC+) |
| DIONISI | 80 | NP B169 1 | C. Dionisi <i>et al.</i> | (CERN, MADR, CDEF+) |

| | | | | |
|--------------|----|-------------|--------------------------------------|---------------------------------|
| GURTU | 79 | NP B151 181 | A. Gurtu <i>et al.</i> | (CERN, ZEEM, NIJM, OXF) |
| STANTON | 79 | PRL 42 346 | N.R. Stanton <i>et al.</i> | (OSU, CARL, MCGI+) JP |
| CORDEN | 78 | NP B144 253 | M.J. Corden <i>et al.</i> | (BIRM, RHEL, TELA+) JP |
| NACASCH | 78 | NP B135 203 | R. Nacasch <i>et al.</i> | (PARIS, MADR, CERN) |
| GRASSLER | 77 | NP B121 189 | H. Grassler <i>et al.</i> | (AACH3, BERL, BONN+) |
| DEFOIX | 72 | NP B44 125 | C. Defoix <i>et al.</i> | (CDEF, CERN) |
| DUBOC | 72 | NP B46 429 | J. Duboc <i>et al.</i> | (PARIS, LIVP) |
| THUN | 72 | PRL 28 1733 | R. Thun <i>et al.</i> | (STON, NEAS) |
| BARDADIN-... | 71 | PR D4 2711 | M. Bardadin-Otwinowska <i>et al.</i> | (WARS) |
| BOESEBECK | 71 | PL 34B 659 | K. Boesebeck | (AACH, BERL, BONN, CERN, CRAC+) |
| CAMPBELL | 69 | PRL 22 1204 | J.H. Campbell <i>et al.</i> | (PURD) |
| LORSTAD | 69 | NP B14 63 | B. Lorstad <i>et al.</i> | (CDEF, CERN) JP |
| D'ANDLAU | 68 | NP B5 693 | C. d'Andlau <i>et al.</i> | (CDEF, CERN, IRAD+) IJP |
| DAHL | 67 | PR 163 1377 | O.I. Dahl <i>et al.</i> | (LRL) IJP |