

$f_2(1810)$

$I^G(J^{PC}) = 0^+(2^{++})$

OMITTED FROM SUMMARY TABLE

Needs confirmation.

$f_2(1810)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1815 ± 12 OUR AVERAGE		Error includes scale factor of 1.4.		See the ideogram below.
$1822^{+29+}_{-24-} 66$	5.5k	¹ ABLIKIM	13N BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \eta \eta$
$1737 \pm 9^{+198}_{-65}$		² UEHARA	10A BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \eta \eta$
1800 ± 30	40	ALDE	88D GAM4	$300 \pi^- p \rightarrow \pi^- p 4\pi^0$
1806 ± 10	1600	ALDE	87 GAM4	$100 \pi^- p \rightarrow 4\pi^0 n$
1870 ± 40		³ ALDE	86D GAM4	$100 \pi^- p \rightarrow \eta \eta n$
1857^{+35}_{-24}		⁴ COSTA...	80 OMEG	$10 \pi^- p \rightarrow K^+ K^- n$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1858^{+18}_{-71}		⁵ LONGACRE	86 RVUE	Compilation
1799 ± 15		⁶ CASON	82 STRC	$8 \pi^+ p \rightarrow \Delta^{++} \pi^0 \pi^0$

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.


² Breit-Wigner mass.

³ Seen in only one solution.

⁴ Error increased by spread of two solutions. Included in LONGACRE 86 global analysis.

⁵ From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.

⁶ From an amplitude analysis of the reaction $\pi^+ \pi^- \rightarrow 2\pi^0$. The resonance in the $2\pi^0$ final state is not confirmed by PROKOSHKIN 97.



$f_2(1810)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
197± 22 OUR AVERAGE		Error includes scale factor of 1.5. See the ideogram below.		
229 ⁺ 52 ⁺ 88 42 - 155	5.5k	7 ABLIKIM	13N BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \eta \eta$
228 ⁺ 21 ⁺ 234 20 - 153		8 UEHARA	10A BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \eta \eta$
160± 30	40	ALDE	88D GAM4	$300 \pi^- p \rightarrow \pi^- p 4\pi^0$
190± 20	1600	ALDE	87 GAM4	$100 \pi^- p \rightarrow 4\pi^0 n$
250± 30		9 ALDE	86D GAM4	$100 \pi^- p \rightarrow \eta \eta n$
185 ⁺ 102 - 139		10 COSTA...	80 OMEG	$10 \pi^- p \rightarrow K^+ K^- n$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
388 ⁺ 15 - 21		11 LONGACRE	86 RVUE	Compilation
280 ⁺ 42 - 35		12 CASON	82 STRC	$8 \pi^+ p \rightarrow \Delta^{++} \pi^0 \pi^0$

7 From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.


8 Breit-Wigner width.

9 Seen in only one solution.

10 Error increased by spread of two solutions. Included in LONGACRE 86 global analysis.

11 From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.

12 From an amplitude analysis of the reaction $\pi^+ \pi^- \rightarrow 2\pi^0$. The resonance in the $2\pi^0$ final state is not confirmed by PROKOSHKIN 97.



$f_2(1810)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \pi\pi$	
$\Gamma_2 \eta\eta$	seen
$\Gamma_3 4\pi^0$	seen
$\Gamma_4 K^+K^-$	
$\Gamma_5 \gamma\gamma$	seen

$f_2(1810) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(\eta\eta) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_2\Gamma_5/\Gamma$
$5.2^{+0.9+37.3}_{-0.8-4.5}$	13 UEHARA 10A BELL 10.6 $e^+e^- \rightarrow e^+e^-\eta\eta$

¹³ Including interference with the $f'_2(1525)$ (parameters fixed to the values from the 2008 edition of this review, PDG 08) and $f_2(1270)$. May also be the $f_0(1500)$.

$f_2(1810)$ BRANCHING RATIOS

$\Gamma(\pi\pi)/\Gamma_{\text{total}}$

Γ_1/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	AMSLER 02	CBAR	$0.9 \bar{p}p \rightarrow \pi^0\eta\eta, \pi^0\pi^0\pi^0$
not seen	PROKOSHKIN 97	GAM2	$38 \pi^- p \rightarrow \pi^0\pi^0n$
$0.21^{+0.02}_{-0.03}$	14 LONGACRE 86	RVUE	Compilation
0.44 ± 0.03	15 CASON 82	STRC	$8 \pi^+ p \rightarrow \Delta^{++}\pi^0\pi^0$

¹⁴ From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.

¹⁵ Included in LONGACRE 86 global analysis.

$\Gamma(\eta\eta)/\Gamma_{\text{total}}$

Γ_2/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	ABLIKIM	13N	BES3 PWA of $J/\psi \rightarrow \gamma\eta\eta$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.008^{+0.028}_{-0.003}$	16 LONGACRE 86	RVUE	Compilation

¹⁶ From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.

$\Gamma(\pi\pi)/\Gamma(4\pi^0)$

Γ_1/Γ_3

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
<0.75	ALDE	87	GAM4 $100 \pi^- p \rightarrow 4\pi^0 n$

$\Gamma(4\pi^0)/\Gamma(\eta\eta)$

Γ_3/Γ_2

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.8 ± 0.3	ALDE	87	GAM4 $100 \pi^- p \rightarrow 4\pi^0 n$

$\Gamma(K^+K^-)/\Gamma_{\text{total}}$

Γ_4/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.003^{+0.019}_{-0.002}$	17 LONGACRE 86	RVUE	Compilation
seen	COSTA...	80	OMEG $10 \pi^- p \rightarrow K^+K^- n$

¹⁷ From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.

$f_2(1810)$ REFERENCES

ABLIKIM	13N	PR D87 092009	Ablikim M. <i>et al.</i>	(BES III Collab.)
UEHARA	10A	PR D82 114031	S. Uehara <i>et al.</i>	(BELLE Collab.)
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
AMSLER	02	EPJ C23 29	C. Amsler <i>et al.</i>	
PROKOSHKIN	97	SPD 42 117	Y.D. Prokoshkin <i>et al.</i>	(SERP)
		Translated from DANS 353 323.		
ALDE	88D	SJNP 47 810	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+)
		Translated from YAF 47 1273.		
ALDE	87	PL B198 286	D.M. Alde <i>et al.</i>	(LANL, BRUX, SERP, LAPP)
ALDE	86D	NP B269 485	D.M. Alde <i>et al.</i>	(BELG, LAPP, SERP, CERN+)
LONGACRE	86	PL B177 223	R.S. Longacre <i>et al.</i>	(BNL, BRAN, CUNY+)
CASON	82	PRL 48 1316	N.M. Cason <i>et al.</i>	(NDAM, ANL)
COSTA...	80	NP B175 402	G. Costa de Beauregard <i>et al.</i>	(BARI, BONN+)