

GAUGE AND HIGGS BOSONS

γ (photon)

$$I(J^{PC}) = 0,1(1^{- -})$$

Mass $m < 1 \times 10^{-18}$ eV

Charge $q < 1 \times 10^{-35}$ e

Mean life $\tau = \text{Stable}$

**g
or gluon**

$$I(J^P) = 0(1^{-})$$

Mass $m = 0$ [a]

SU(3) color octet

graviton

$$J = 2$$

Mass $m < 6 \times 10^{-32}$ eV

W

$$J = 1$$

Charge = ± 1 e

Mass $m = 80.385 \pm 0.015$ GeV

$m_Z - m_W = 10.4 \pm 1.6$ GeV

$m_{W^+} - m_{W^-} = -0.2 \pm 0.6$ GeV

Full width $\Gamma = 2.085 \pm 0.042$ GeV

$\langle N_{\pi^\pm} \rangle = 15.70 \pm 0.35$

$\langle N_{K^\pm} \rangle = 2.20 \pm 0.19$

$\langle N_p \rangle = 0.92 \pm 0.14$

$\langle N_{\text{charged}} \rangle = 19.39 \pm 0.08$

W^- modes are charge conjugates of the modes below.

| W^+ DECAY MODES | Fraction (Γ_i/Γ) | Confidence level | P (MeV/c) |
|-------------------------------------|--------------------------------|------------------|----------------|
| $\ell^+ \nu$ | [b] $(10.86 \pm 0.09) \%$ | | — |
| $e^+ \nu$ | $(10.71 \pm 0.16) \%$ | | 40192 |
| $\mu^+ \nu$ | $(10.63 \pm 0.15) \%$ | | 40192 |
| $\tau^+ \nu$ | $(11.38 \pm 0.21) \%$ | | 40173 |
| hadrons | $(67.41 \pm 0.27) \%$ | | — |
| $\pi^+ \gamma$ | < 7 | $\times 10^{-6}$ | 95% 40192 |
| $D_s^+ \gamma$ | < 1.3 | $\times 10^{-3}$ | 95% 40168 |

| | | |
|------------|------------------------|---|
| cX | $(33.3 \pm 2.6) \%$ | — |
| $c\bar{s}$ | $(31^{+13}_{-11}) \%$ | — |
| invisible | $[c] (1.4 \pm 2.9) \%$ | — |

Z $J = 1$

Charge = 0

Mass $m = 91.1876 \pm 0.0021$ GeV ^[d]Full width $\Gamma = 2.4952 \pm 0.0023$ GeV $\Gamma(\ell^+ \ell^-) = 83.984 \pm 0.086$ MeV ^[b] $\Gamma(\text{invisible}) = 499.0 \pm 1.5$ MeV ^[e] $\Gamma(\text{hadrons}) = 1744.4 \pm 2.0$ MeV $\Gamma(\mu^+ \mu^-) / \Gamma(e^+ e^-) = 1.0009 \pm 0.0028$ $\Gamma(\tau^+ \tau^-) / \Gamma(e^+ e^-) = 1.0019 \pm 0.0032$ ^[f]**Average charged multiplicity** $\langle N_{\text{charged}} \rangle = 20.76 \pm 0.16$ ($S = 2.1$)**Couplings to quarks and leptons** $g_V^\ell = -0.03783 \pm 0.00041$ $g_V^u = 0.25^{+0.07}_{-0.06}$ $g_V^d = -0.33^{+0.05}_{-0.06}$ $g_A^\ell = -0.50123 \pm 0.00026$ $g_A^u = 0.50^{+0.04}_{-0.06}$ $g_A^d = -0.523^{+0.050}_{-0.029}$ $g^{\nu\ell} = 0.5008 \pm 0.0008$ $g^{\nu e} = 0.53 \pm 0.09$ $g^{\nu\mu} = 0.502 \pm 0.017$ **Asymmetry parameters ^[g]** $A_e = 0.1515 \pm 0.0019$ $A_\mu = 0.142 \pm 0.015$ $A_\tau = 0.143 \pm 0.004$ $A_s = 0.90 \pm 0.09$ $A_c = 0.670 \pm 0.027$ $A_b = 0.923 \pm 0.020$ **Charge asymmetry (%) at Z pole** $A_{FB}^{(0\ell)} = 1.71 \pm 0.10$ $A_{FB}^{(0u)} = 4 \pm 7$ $A_{FB}^{(0s)} = 9.8 \pm 1.1$ $A_{FB}^{(0c)} = 7.07 \pm 0.35$ $A_{FB}^{(0b)} = 9.92 \pm 0.16$

| Z DECAY MODES | Fraction (Γ_i/Γ) | Scale factor/ Confidence level | p (MeV/c) |
|--|--|-----------------------------------|----------------|
| e^+e^- | (3.363 \pm 0.004) % | | 45594 |
| $\mu^+\mu^-$ | (3.366 \pm 0.007) % | | 45594 |
| $\tau^+\tau^-$ | (3.370 \pm 0.008) % | | 45559 |
| $\ell^+\ell^-$ | [b] (3.3658 \pm 0.0023) % | | — |
| $\ell^+\ell^-\ell^+\ell^-$ | [h] (3.30 \pm 0.31) $\times 10^{-6}$ | S=1.1 | 45594 |
| invisible | (20.00 \pm 0.06) % | | — |
| hadrons | (69.91 \pm 0.06) % | | — |
| ($u\bar{u} + c\bar{c}$)/2 | (11.6 \pm 0.6) % | | — |
| ($d\bar{d} + s\bar{s} + b\bar{b}$)/3 | (15.6 \pm 0.4) % | | — |
| $c\bar{c}$ | (12.03 \pm 0.21) % | | — |
| $b\bar{b}$ | (15.12 \pm 0.05) % | | — |
| $b\bar{b}b\bar{b}$ | (3.6 \pm 1.3) $\times 10^{-4}$ | | — |
| ggg | < 1.1 | % CL=95% | — |
| $\pi^0\gamma$ | < 2.01 | $\times 10^{-5}$ CL=95% | 45594 |
| $\eta\gamma$ | < 5.1 | $\times 10^{-5}$ CL=95% | 45592 |
| $\omega\gamma$ | < 6.5 | $\times 10^{-4}$ CL=95% | 45590 |
| $\eta'(958)\gamma$ | < 4.2 | $\times 10^{-5}$ CL=95% | 45589 |
| $\gamma\gamma$ | < 1.46 | $\times 10^{-5}$ CL=95% | 45594 |
| $\pi^0\pi^0$ | < 1.52 | $\times 10^{-5}$ CL=95% | 45594 |
| $\gamma\gamma\gamma$ | < 1.0 | $\times 10^{-5}$ CL=95% | 45594 |
| $\pi^\pm W^\mp$ | [i] < 7 | $\times 10^{-5}$ CL=95% | 10162 |
| $\rho^\pm W^\mp$ | [i] < 8.3 | $\times 10^{-5}$ CL=95% | 10136 |
| $J/\psi(1S)X$ | (3.51 $^{+0.23}_{-0.25}$) $\times 10^{-3}$ | S=1.1 | — |
| $\psi(2S)X$ | (1.60 \pm 0.29) $\times 10^{-3}$ | | — |
| $\chi_{c1}(1P)X$ | (2.9 \pm 0.7) $\times 10^{-3}$ | | — |
| $\chi_{c2}(1P)X$ | < 3.2 | $\times 10^{-3}$ CL=90% | — |
| $\Upsilon(1S)X + \Upsilon(2S)X$ $+ \Upsilon(3S)X$ | (1.0 \pm 0.5) $\times 10^{-4}$ | | — |
| $\Upsilon(1S)X$ | < 4.4 | $\times 10^{-5}$ CL=95% | — |
| $\Upsilon(2S)X$ | < 1.39 | $\times 10^{-4}$ CL=95% | — |
| $\Upsilon(3S)X$ | < 9.4 | $\times 10^{-5}$ CL=95% | — |
| (D^0/\bar{D}^0)X | (20.7 \pm 2.0) % | | — |
| $D^\pm X$ | (12.2 \pm 1.7) % | | — |
| $D^*(2010)^\pm X$ | [i] (11.4 \pm 1.3) % | | — |
| $D_{s1}(2536)^\pm X$ | (3.6 \pm 0.8) $\times 10^{-3}$ | | — |
| $D_{sJ}(2573)^\pm X$ | (5.8 \pm 2.2) $\times 10^{-3}$ | | — |
| $D^{*J}(2629)^\pm X$ | searched for | | — |
| B^+X | [j] (6.08 \pm 0.13) % | | — |
| B_s^0X | [j] (1.59 \pm 0.13) % | | — |
| B_c^+X | searched for | | — |
| Λ_c^+X | (1.54 \pm 0.33) % | | — |

| | | | | |
|-------------------------------|-----|-------------------------|-------------------------|-------|
| $\Xi_c^0 X$ | | seen | | — |
| $\Xi_b X$ | | seen | | — |
| b -baryon X | | $[j]$ (1.38 ± 0.22) % | | — |
| anomalous γ + hadrons | | $[k] < 3.2$ | $\times 10^{-3}$ CL=95% | — |
| $e^+ e^- \gamma$ | | $[k] < 5.2$ | $\times 10^{-4}$ CL=95% | 45594 |
| $\mu^+ \mu^- \gamma$ | | $[k] < 5.6$ | $\times 10^{-4}$ CL=95% | 45594 |
| $\tau^+ \tau^- \gamma$ | | $[k] < 7.3$ | $\times 10^{-4}$ CL=95% | 45559 |
| $\ell^+ \ell^- \gamma \gamma$ | | $[l] < 6.8$ | $\times 10^{-6}$ CL=95% | — |
| $q\bar{q}\gamma\gamma$ | | $[l] < 5.5$ | $\times 10^{-6}$ CL=95% | — |
| $\nu\bar{\nu}\gamma\gamma$ | | $[l] < 3.1$ | $\times 10^{-6}$ CL=95% | 45594 |
| $e^\pm \mu^\mp$ | LF | $[i] < 7.5$ | $\times 10^{-7}$ CL=95% | 45594 |
| $e^\pm \tau^\mp$ | LF | $[i] < 9.8$ | $\times 10^{-6}$ CL=95% | 45576 |
| $\mu^\pm \tau^\mp$ | LF | $[i] < 1.2$ | $\times 10^{-5}$ CL=95% | 45576 |
| pe | L,B | < 1.8 | $\times 10^{-6}$ CL=95% | 45589 |
| $p\mu$ | L,B | < 1.8 | $\times 10^{-6}$ CL=95% | 45589 |



$$J = 0$$

$$\text{Mass } m = 125.09 \pm 0.24 \text{ GeV}$$

H^0 Signal Strengths in Different Channels

See Listings for the latest unpublished results.

$$\text{Combined Final States} = 1.17 \pm 0.17 \quad (S = 1.2)$$

$$W W^* = 0.81 \pm 0.16$$

$$Z Z^* = 1.15^{+0.27}_{-0.23} \quad (S = 1.2)$$

$$\gamma\gamma = 1.17^{+0.19}_{-0.17}$$

$$b\bar{b} = 0.85 \pm 0.29$$

$$\mu^+ \mu^- < 7.0, \text{ CL} = 95\%$$

$$\tau^+ \tau^- = 0.79 \pm 0.26$$

$$Z\gamma < 9.5, \text{ CL} = 95\%$$

$$t\bar{t}H^0 \text{ Production} = 2.5^{+0.9}_{-0.8}$$

| H^0 DECAY MODES | Fraction (Γ_i/Γ) | Confidence level | P (MeV/c) |
|-------------------|--------------------------------|------------------|-------------|
| invisible | <58 % | 95% | — |

Neutral Higgs Bosons, Searches for

Searches for a Higgs Boson with Standard Model Couplings

Mass $m > 122$ and none 128–710 GeV, CL = 95%

The limits for H_1^0 and A^0 in supersymmetric models refer to the m_h^{\max} benchmark scenario for the supersymmetric parameters.

H_1^0 in Supersymmetric Models ($m_{H_1^0} < m_{H_2^0}$)

Mass $m > 92.8$ GeV, CL = 95%

A^0 Pseudoscalar Higgs Boson in Supersymmetric Models ^[n]

Mass $m > 93.4$ GeV, CL = 95% $\tan\beta > 0.4$

Charged Higgs Bosons (H^\pm and $H^{\pm\pm}$), Searches for

H^\pm Mass $m > 80$ GeV, CL = 95%

New Heavy Bosons (W' , Z' , leptoquarks, etc.), Searches for

Additional W Bosons

W' with standard couplings

Mass $m > 3.240 \times 10^3$ GeV, CL = 95% (pp direct search)

W_R (Right-handed W Boson)

Mass $m > 715$ GeV, CL = 90% (electroweak fit)

Additional Z Bosons

Z'_{SM} with standard couplings

Mass $m > 2.900 \times 10^3$ GeV, CL = 95% (pp direct search)

Mass $m > 1.500 \times 10^3$ GeV, CL = 95% (electroweak fit)

Z_{LR} of $SU(2)_L \times SU(2)_R \times U(1)$ (with $g_L = g_R$)

Mass $m > 630$ GeV, CL = 95% ($p\bar{p}$ direct search)

Mass $m > 1162$ GeV, CL = 95% (electroweak fit)

Z_χ of $SO(10) \rightarrow SU(5) \times U(1)_\chi$ (with $g_\chi = e/\cos\theta_W$)

Mass $m > 2.620 \times 10^3$ GeV, CL = 95% (pp direct search)

Mass $m > 1.141 \times 10^3$ GeV, CL = 95% (electroweak fit)

Z_ψ of $E_6 \rightarrow SO(10) \times U(1)_\psi$ (with $g_\psi = e/\cos\theta_W$)

Mass $m > 2.510 \times 10^3$ GeV, CL = 95% (pp direct search)

Mass $m > 476$ GeV, CL = 95% (electroweak fit)

Z_η of $E_6 \rightarrow SU(3) \times SU(2) \times U(1) \times U(1)_\eta$ (with $g_\eta = e/\cos\theta_W$)

Mass $m > 1.870 \times 10^3$ GeV, CL = 95% (pp direct search)

Mass $m > 619$ GeV, CL = 95% (electroweak fit)

Scalar Leptoquarks

Mass $m > 830$ GeV, CL = 95% (1st generation, pair prod.)

Mass $m > 304$ GeV, CL = 95% (1st gener., single prod.)

Mass $m > 840$ GeV, CL = 95% (2nd gener., pair prod.)

Mass $m > 73$ GeV, CL = 95% (2nd gener., single prod.)

Mass $m > 740$ GeV, CL = 95% (3rd gener., pair prod.)

(See the Particle Listings for assumptions on leptoquark quantum numbers and branching fractions.)

Diquarks

Mass $m > 3.750 \times 10^3$ GeV, CL = 95%

Axigluon

Mass $m > 3.360 \times 10^3$ GeV, CL = 95%

Axions (A^0) and Other Very Light Bosons, Searches for

The standard Peccei-Quinn axion is ruled out. Variants with reduced couplings or much smaller masses are constrained by various data. The Particle Listings in the full *Review* contain a Note discussing axion searches.

The best limit for the half-life of neutrinoless double beta decay with Majoron emission is $> 7.2 \times 10^{24}$ years (CL = 90%).

NOTES

[a] Theoretical value. A mass as large as a few MeV may not be precluded.

[b] ℓ indicates each type of lepton (e , μ , and τ), not sum over them.

[c] This represents the width for the decay of the W boson into a charged particle with momentum below detectability, $p < 200$ MeV.

[d] The Z -boson mass listed here corresponds to a Breit-Wigner resonance parameter. It lies approximately 34 MeV above the real part of the position of the pole (in the energy-squared plane) in the Z -boson propagator.

[e] This partial width takes into account Z decays into $\nu\bar{\nu}$ and any other possible undetected modes.

[f] This ratio has not been corrected for the τ mass.

[g] Here $A \equiv 2g_V g_A / (g_V^2 + g_A^2)$.

[*h*] Here ℓ indicates e or μ .

[*i*] The value is for the sum of the charge states or particle/antiparticle states indicated.

[*j*] This value is updated using the product of (i) the $Z \rightarrow b\bar{b}$ fraction from this listing and (ii) the b -hadron fraction in an unbiased sample of weakly decaying b -hadrons produced in Z -decays provided by the Heavy Flavor Averaging Group (HFAG, http://www.slac.stanford.edu/xorg/hfag/osc/PDG_2009/#FRACZ).

[*k*] See the Z Particle Listings for the γ energy range used in this measurement.

[*l*] For $m_{\gamma\gamma} = (60 \pm 5)$ GeV.

[*n*] The limits assume no invisible decays.