

K(3100)

$$I^G(J^{PC}) = ??(???)$$

OMITTED FROM SUMMARY TABLE

Narrow peak observed in several ($\Lambda\bar{p}$ + pions) and ($\bar{\Lambda}p$ + pions) states in Σ^- Be reactions by BOURQUIN 86 and in np and nA reactions by ALEEV 93. Not seen by BOEHNLEIN 91. If due to strong decays, this state has exotic quantum numbers ($B=0, Q=+1, S=-1$ for $\Lambda\bar{p}\pi^+\pi^+$ and $I \geq 3/2$ for $\Lambda\bar{p}\pi^-$). Needs confirmation.

K(3100) MASS

VALUE (MeV)
 ≈ 3100 OUR ESTIMATE

DOCUMENT ID**3-BODY DECAYS**

| <u>VALUE (MeV)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-----------------------|-------------|---|
| 3054 ± 11 OUR AVERAGE | | | |
| $3060 \pm 7 \pm 20$ | ¹ ALEEV 93 | BIS2 | $K(3100) \rightarrow \Lambda\bar{p}\pi^+$ |
| $3056 \pm 7 \pm 20$ | ¹ ALEEV 93 | BIS2 | $K(3100) \rightarrow \bar{\Lambda}p\pi^-$ |
| $3055 \pm 8 \pm 20$ | ¹ ALEEV 93 | BIS2 | $K(3100) \rightarrow \Lambda\bar{p}\pi^-$ |
| $3045 \pm 8 \pm 20$ | ¹ ALEEV 93 | BIS2 | $K(3100) \rightarrow \bar{\Lambda}p\pi^+$ |

4-BODY DECAYS

| <u>VALUE (MeV)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-----------------------|-------------|--|
| 3059 ± 11 OUR AVERAGE | | | |
| $3067 \pm 6 \pm 20$ | ¹ ALEEV 93 | BIS2 | $K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^+$ |
| $3060 \pm 8 \pm 20$ | ¹ ALEEV 93 | BIS2 | $K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^-$ |
| $3055 \pm 7 \pm 20$ | ¹ ALEEV 93 | BIS2 | $K(3100) \rightarrow \bar{\Lambda}p\pi^-\pi^-$ |
| $3052 \pm 8 \pm 20$ | ¹ ALEEV 93 | BIS2 | $K(3100) \rightarrow \bar{\Lambda}p\pi^-\pi^+$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| 3105 ± 30 | BOURQUIN 86 | SPEC | $K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^+$ |
| 3115 ± 30 | BOURQUIN 86 | SPEC | $K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^-$ |

5-BODY DECAYS

| <u>VALUE (MeV)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|---|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| 3095 ± 30 | BOURQUIN 86 | SPEC | $K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^+\pi^-$ |
| ¹ Supersedes ALEEV 90. | | | |

K(3100) WIDTH**3-BODY DECAYS**

| <u>VALUE (MeV)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-----------------------|-------------|---|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| 42 ± 16 | ² ALEEV 93 | BIS2 | $K(3100) \rightarrow \Lambda\bar{p}\pi^+$ |
| 36 ± 15 | ² ALEEV 93 | BIS2 | $K(3100) \rightarrow \bar{\Lambda}p\pi^-$ |
| 50 ± 18 | ² ALEEV 93 | BIS2 | $K(3100) \rightarrow \Lambda\bar{p}\pi^-$ |
| 30 ± 15 | ² ALEEV 93 | BIS2 | $K(3100) \rightarrow \bar{\Lambda}p\pi^+$ |

4-BODY DECAYS

| <u>VALUE (MeV)</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|--|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 22 ± 8 | | ² ALEEV | 93 | BIS2 $K(3100) \rightarrow \Lambda \bar{p} \pi^+ \pi^+$ |
| 28 ± 12 | | ² ALEEV | 93 | BIS2 $K(3100) \rightarrow \Lambda \bar{p} \pi^+ \pi^-$ |
| 32 ± 15 | | ² ALEEV | 93 | BIS2 $K(3100) \rightarrow \bar{\Lambda} p \pi^- \pi^-$ |
| 30 ± 15 | | ² ALEEV | 93 | BIS2 $K(3100) \rightarrow \bar{\Lambda} p \pi^- \pi^+$ |
| <30 | 90 | BOURQUIN | 86 | SPEC $K(3100) \rightarrow \Lambda \bar{p} \pi^+ \pi^+$ |
| <80 | 90 | BOURQUIN | 86 | SPEC $K(3100) \rightarrow \Lambda \bar{p} \pi^+ \pi^-$ |

5-BODY DECAYS

| <u>VALUE (MeV)</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|--|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| <30 | 90 | BOURQUIN | 86 | SPEC $K(3100) \rightarrow \Lambda \bar{p} \pi^+ \pi^+ \pi^-$ |
| ² Supersedes ALEEV 90. | | | | |

K(3100) DECAY MODES

| Mode | |
|------------|---|
| Γ_1 | $K(3100)^0 \rightarrow \Lambda \bar{p} \pi^+$ |
| Γ_2 | $K(3100)^{-} \rightarrow \Lambda \bar{p} \pi^-$ |
| Γ_3 | $K(3100)^- \rightarrow \Lambda \bar{p} \pi^+ \pi^-$ |
| Γ_4 | $K(3100)^+ \rightarrow \Lambda \bar{p} \pi^+ \pi^+$ |
| Γ_5 | $K(3100)^0 \rightarrow \Lambda \bar{p} \pi^+ \pi^+ \pi^-$ |
| Γ_6 | $K(3100)^0 \rightarrow \Sigma(1385)^+ \bar{p}$ |

 $\Gamma(\Sigma(1385)^+ \bar{p})/\Gamma(\Lambda \bar{p} \pi^+)$ **Γ_6/Γ_1**

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|------------|--------------------|-------------|---|
| <0.04 | 90 | ALEEV | 93 | BIS2 $K(3100)^0 \rightarrow \Sigma(1385)^+ \bar{p}$ |

K(3100) REFERENCES

| | | | | |
|-----------|----|--|-----------------------------|---------------------|
| ALEEV | 93 | PAN 56 1358 Translated from YAF 56 100. | A.N. Aleev <i>et al.</i> | (BIS-2 Collab.) |
| BOEHNLEIN | 91 | NPBPS B21 174 | A. Boehnlein <i>et al.</i> | (FLOR, BNL, IND+) |
| ALEEV | 90 | ZPHY C47 533 | A.N. Aleev <i>et al.</i> | (BIS-2 Collab.) |
| BOURQUIN | 86 | PL B172 113 | M.H. Bourquin <i>et al.</i> | (GEVA, RAL, HEIDP+) |