

$K_0^*(1430)$

$$I(J^P) = \frac{1}{2}(0^+)$$

See our minireview in the 1994 edition and in this edition under the $f_0(500)$.

 $K_0^*(1430)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1425 ±50	OUR ESTIMATE			
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1438 ± 8 ± 4	5.4k	¹ LEES	14E BABR	$\eta_c(1S) \rightarrow K^+ K^- \eta/\pi^0$
1427 ± 4 ±13		² BUGG	10 RVUE	S-matrix pole
1466.6 ± 0.7 ± 3.4	141k	³ BONVICINI	08A CLEO	$D^+ \rightarrow K^- \pi^+ \pi^+$
~ 1412		⁴ LINK	07 FOCS	$D^+ \rightarrow K^- K^+ \pi^+$
1461.0 ± 4.0 ± 2.1	54k	⁵ LINK	07B FOCS	$D^+ \rightarrow K^- \pi^+ \pi^+$
1406 ±29		⁶ BUGG	06 RVUE	
1435 ± 6		⁷ ZHOU	06 RVUE	$K p \rightarrow K^- \pi^+ n$
1455 ±20 ±15		ABLIKIM	05Q BES2	$\psi(2S) \rightarrow$ $\gamma \pi^+ \pi^- K^+ K^-$
1456 ± 8		⁸ ZHENG	04 RVUE	$K^- p \rightarrow K^- \pi^+ n$
~ 1419		⁹ BUGG	03 RVUE	$11 K^- p \rightarrow K^- \pi^+ n$
~ 1440		¹⁰ LI	03 RVUE	$11 K^- p \rightarrow K^- \pi^+ n$
1459 ± 9	15k	¹¹ AITALA	02 E791	$D^+ \rightarrow K^- \pi^+ \pi^+$
~ 1440		¹² JAMIN	00 RVUE	$K p \rightarrow K p$
1436 ± 8		¹³ BARBERIS	98E OMEG	$450 p p \rightarrow$ $p_f p_s K^+ K^- \pi^+ \pi^-$
1415 ±25		⁹ ANISOVICH	97C RVUE	$11 K^- p \rightarrow K^- \pi^+ n$
~ 1450		¹⁴ TORNQVIST	96 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi$
1412 ± 6		¹⁵ ASTON	88 LASS	$11 K^- p \rightarrow K^- \pi^+ n$
~ 1430		BAUBILLIER	84B HBC	$8.25 K^- p \rightarrow \bar{K}^0 \pi^- p$
~ 1425		¹⁶ ESTABROOKS	78 ASPK	$13 K^\pm p \rightarrow K^\pm \pi^\pm (n, \Delta)$
~ 1450.0		MARTIN	78 SPEC	$10 K^\pm p \rightarrow K_S^0 \pi p$

¹ Using both $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow \pi^+ \pi^- \pi^0$. From a likelihood scan in the presence of several interfering scalar-meson resonances with fixed width $\Gamma(K_0^*(1430)) = 210$ MeV.

² S-Matrix pole. Supersedes BUGG 06. Combined analysis of ASTON 88, ABLIKIM 06C, AITALA 06, and LINK 09 using an s-dependent width with couplings to $K\pi$ and $K\eta'$, and the Adler zero near thresholds.

³ From the isobar model with a complex pole for the κ .

⁴ From a non-parametric analysis.

⁵ A Breit-Wigner mass and width.

⁶ S-matrix pole. Reanalysis of ASTON 88, AITALA 02, and ABLIKIM 06C including the κ with an s-dependent width and an Adler zero near threshold.

⁷ S-matrix pole. Using ASTON 88 and assuming $K_0^*(800)$, $K_0^*(1950)$.

⁸ Using ASTON 88 and assuming $K_0^*(800)$.

⁹ T-matrix pole. Reanalysis of ASTON 88 data.

¹⁰ Breit-Wigner fit. Using ASTON 88.

¹¹ Assuming a low-mass scalar $K\pi$ resonance, $\kappa(800)$.

¹² T-matrix pole. Using data from ESTABROOKS 78 and ASTON 88.

¹³ J^P not determined, could be $K_2^*(1430)$.

¹⁴ T-matrix pole.

¹⁵ Uses a model for the background, without this background they get a mass 1340 MeV, where the phase shift passes 90° .

¹⁶ Mass defined by pole position. From elastic $K\pi$ partial-wave analysis.

$K_0^*(1430)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
270 ±80	OUR ESTIMATE			
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
210 ±20 ±12	5.4k	¹ LEES	14E BABR	$\eta_c(1S) \rightarrow K^+ K^- \eta/\pi^0$
270 ±10 ±40		² BUGG	10 RVUE	S-matrix pole
174.2 ± 1.9 ± 3.2	141k	³ BONVICINI	08A CLEO	$D^+ \rightarrow K^- \pi^+ \pi^+$
~ 500		⁴ LINK	07 FOCS	$D^+ \rightarrow K^- K^+ \pi^+$
177.0 ± 8.0 ± 3.4	54k	⁵ LINK	07B FOCS	$D^+ \rightarrow K^- \pi^+ \pi^+$
350 ±40		⁶ BUGG	06 RVUE	
288 ±22		⁷ ZHOU	06 RVUE	$Kp \rightarrow K^- \pi^+ n$
270 ±45 ⁺³⁰ -35		ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$
217 ±31		⁸ ZHENG	04 RVUE	$K^- p \rightarrow K^- \pi^+ n$
~ 316		⁹ BUGG	03 RVUE	11 $K^- p \rightarrow K^- \pi^+ n$
~ 350		¹⁰ LI	03 RVUE	11 $K^- p \rightarrow K^- \pi^+ n$
175 ±17	15k	¹¹ AITALA	02 E791	$D^+ \rightarrow K^- \pi^+ \pi^+$
~ 300		¹² JAMIN	00 RVUE	$Kp \rightarrow Kp$
196 ±45		¹³ BARBERIS	98E OMEG	450 $pp \rightarrow$ $p_f p_s K^+ K^- \pi^+ \pi^-$
330 ±50		⁹ ANISOVICH	97C RVUE	11 $K^- p \rightarrow K^- \pi^+ n$
~ 320		¹⁴ TORNQVIST	96 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi$
294 ±23		ASTON	88 LASS	11 $K^- p \rightarrow K^- \pi^+ n$
~ 200		BAUBILLIER	84B HBC	8.25 $K^- p \rightarrow \bar{K}^0 \pi^- p$
200 to 300		¹⁵ ESTABROOKS	78 ASPK	13 $K^\pm p \rightarrow K^\pm \pi^\pm (n, \Delta)$

¹ Using both $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow \pi^+ \pi^- \pi^0$. From a likelihood scan in the presence of several interfering scalar-meson resonances with fixed mass $M(K_0^*(1430)) = 1435$ MeV.

² S-Matrix pole. Supersedes BUGG 06. Combined analysis of ASTON 88, ABLIKIM 06C, AITALA 06, and LINK 09 using an s-dependent width with couplings to $K\pi$ and $K\eta'$, and the Adler zero near thresholds.

³ From the isobar model with a complex pole for the κ .

⁴ From a non-parametric analysis.

⁵ A Breit-Wigner mass and width.

⁶ S-matrix pole. Reanalysis of ASTON 88, AITALA 02, and ABLIKIM 06C including the κ with an s-dependent width and an Adler zero near threshold.

⁷ S-matrix pole. Using ASTON 88 and assuming $K_0^*(800)$, $K_0^*(1950)$.

⁸ Using ASTON 88 and assuming $K_0^*(800)$.

⁹ T-matrix pole. Reanalysis of ASTON 88 data.

¹⁰ Breit-Wigner fit. Using ASTON 88.

¹¹ Assuming a low-mass scalar $K\pi$ resonance, $\kappa(800)$.

¹² T-matrix pole. Using data from ESTABROOKS 78 and ASTON 88.

¹³ J^P not determined, could be $K_2^*(1430)$.

¹⁴ T-matrix pole.

¹⁵ From elastic $K\pi$ partial-wave analysis.

$K_0^*(1430)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K\pi$	(93 \pm 10) %
Γ_2 $K\eta$	(8.6 $^{+2.7}_{-3.4}$) %
Γ_3 $K\eta'(958)$	seen

 $K_0^*(1430)$ BRANCHING RATIOS

$\Gamma(K\pi)/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
0.93\pm0.04\pm0.09	ASTON	88	LASS	0 11 $K^- p \rightarrow K^- \pi^+ n$

$\Gamma(K\eta)/\Gamma(K\pi)$ Γ_2/Γ_1

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
9.2\pm2.5$^{+1.0}_{-2.5}$	5.4k	¹ LEES	14E	BABR $\eta_c(1S) \rightarrow K^+ K^- \eta/\pi^0$

¹ Using both $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow \pi^+ \pi^- \pi^0$. From a Dalitz analysis in the presence of several interfering scalar-meson resonances.

$\Gamma(K\eta'(958))/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	ABLIKIM	14J	BES3 $\psi(2S) \rightarrow \gamma K^+ K^- \eta'(958)$

 $K_0^*(1430)$ REFERENCES

ABLIKIM	14J	PR D89 074030	M. Ablikim <i>et al.</i>	(BES III Collab.)
LEES	14E	PR D89 112004	J.P. Lees <i>et al.</i>	(BABAR Collab.)
BUGG	10	PR D81 014002	D.V. Bugg	(LOQM)
LINK	09	PL B681 14	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
BONVICINI	08A	PR D78 052001	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
LINK	07	PL B648 156	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	07B	PL B653 1	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
ABLIKIM	06C	PL B633 681	M. Ablikim <i>et al.</i>	(BES Collab.)
AITALA	06	PR D73 032004	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
Also		PR D74 059901 (err.)	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
BUGG	06	PL B632 471	D.V. Bugg	(LOQM)
ZHOU	06	NP A775 212	Z.Y. Zhou, H.Q. Zheng	
ABLIKIM	05Q	PR D72 092002	M. Ablikim <i>et al.</i>	(BES Collab.)
ZHENG	04	NP A733 235	H.Q. Zheng <i>et al.</i>	
BUGG	03	PL B572 1	D.V. Bugg	
LI	03	PR D67 034025	L. Li, B. Zou, G. Li	
AITALA	02	PRL 89 121801	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
JAMIN	00	NP B587 331	M. Jamin <i>et al.</i>	
BARBERIS	98E	PL B436 204	D. Barberis <i>et al.</i>	(Omega Expt.)
ANISOVICH	97C	PL B413 137	A.V. Anisovich, A.V. Sarantsev	
TORNQVIST	96	PRL 76 1575	N.A. Tornqvist, M. Roos	(HELS)
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
BAUBILLIER	84B	ZPHY C26 37	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)
ESTABROOKS	78	NP B133 490	P.G. Estabrooks <i>et al.</i>	(MCGI, CARL, DURH+)
MARTIN	78	NP B134 392	A.D. Martin <i>et al.</i>	(DURH, GEVA)