

X(4020)

$$I(J^P) = 1(?^?)$$

Charged X(4020) seen by ABLIKIM 13X from $e^+e^- \rightarrow \pi^+\pi^-h_c(1P)$ at c.m. energy from 3.90 to 4.42 GeV as a peak in the invariant mass distribution of the $\pi^\pm h_c(1P)$ system, and by ABLIKIM 14B from $e^+e^- \rightarrow (D^*\bar{D}^*)^\pm\pi^\mp$ events in $(D^*\bar{D}^*)^\pm$ mass. A neutral X(4020) seen by ABLIKIM 14P at three c.m. energies in the same range in $e^+e^- \rightarrow \pi^0\pi^0h_c(1P)$ as a peak in the larger of the two masses recoiling against a π^0 . ABLIKIM 15AA observes a 5.9σ signal in $(D^*\bar{D}^*)^0$ in $e^+e^- \rightarrow (D^*\bar{D}^*)^0\pi^0$ events using collisions at two c.m. energies. Production rates and mass values support grouping neutral and charged X(4020) together as manifestations of a single $I = 1$ particle.

X(4020) MASS

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
4024.1±1.9 OUR AVERAGE					
4025.5 ^{+2.0} _{-4.7} ±3.1	116	¹ ABLIKIM 15AA	BES3	0	$e^+e^- \rightarrow (D^*\bar{D}^*)^0\pi^0$
4026.3±2.6±3.7	401	¹ ABLIKIM 14B	BES3	±	$e^+e^- \rightarrow (D^*\bar{D}^*)^\pm\pi^\mp$
4023.9±2.2±3.8	61	^{1,2} ABLIKIM 14P	BES3	0	$e^+e^- \rightarrow \pi^0\pi^0h_c$
4022.9±0.8±2.7	253	¹ ABLIKIM 13X	BES3	±	$e^+e^- \rightarrow \pi^+\pi^-h_c$

¹ Neglecting interference between the X(4020) and non-resonant continuum.


² Assuming $J^P = 1^+$ and width of 7.9 ± 2.6 MeV.

X(4020) WIDTH

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
13 ±5 OUR AVERAGE					
Error includes scale factor of 1.7. See the ideogram below.					
23.0±6.0±1.0	116	¹ ABLIKIM 15AA	BES3	0	$e^+e^- \rightarrow (D^*\bar{D}^*)^0\pi^0$
24.8±5.6±7.7	401	¹ ABLIKIM 14B	BES3	±	$e^+e^- \rightarrow (D^*\bar{D}^*)^\pm\pi^\mp$
7.9±2.7±2.6	253	¹ ABLIKIM 13X	BES3	±	$e^+e^- \rightarrow \pi^+\pi^-h_c$

¹ Neglecting interference between the X(4020) and non-resonant continuum.

WEIGHTED AVERAGE
 13 ± 5 (Error scaled by 1.7)



X(4020) DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $h_c(1P)\pi$	seen
Γ_2 $D^*\bar{D}^*$	seen
Γ_3 $D\bar{D}^* + \text{c.c.}$	not seen
Γ_4 $\eta_c\pi^+\pi^-$	not seen

X(4020) BRANCHING RATIOS

$\Gamma(h_c(1P)\pi)/\Gamma_{\text{total}}$						Γ_1/Γ
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
seen	61	ABLIKIM 14P	BES3	0	$e^+e^- \rightarrow \pi^0\pi^0 h_c$	
seen	253	ABLIKIM 13X	BES3	\pm	$e^+e^- \rightarrow \pi^+\pi^- h_c$	

$\Gamma(D^*\bar{D}^*)/\Gamma_{\text{total}}$						Γ_2/Γ
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
seen	116	¹ ABLIKIM 15AA	BES3	0	$e^+e^- \rightarrow (D^*\bar{D}^*)^0\pi^0$	
seen	401	¹ ABLIKIM 14B	BES3	\pm	$e^+e^- \rightarrow (D^*\bar{D}^*)^\pm\pi^\mp$	

¹ Neglecting interference between the X(4020) and non-resonant continuum.

$\Gamma(D\bar{D}^* + \text{c.c.})/\Gamma_{\text{total}}$						Γ_3/Γ
VALUE	DOCUMENT ID	TECN	CHG	COMMENT		
not seen	ABLIKIM 15AC	BES3	\pm	$e^+e^- \rightarrow \pi^\pm(D\bar{D}^*)^\mp$		

$\Gamma(\eta_c \pi^+ \pi^-)/\Gamma_{\text{total}}$			Γ_4/Γ
<i>VALUE</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
not seen	¹ VINOKUROVA 15	BELL	$B^+ \rightarrow K^+ \eta_c \pi^+ \pi^-$
¹ VINOKUROVA 15 reports $B(B^+ \rightarrow K^+ X(4020)^0) \times B(X \rightarrow \eta_c \pi^+ \pi^-) < 1.6 \times 10^{-5}$ at 90% CL.			

X(4020) REFERENCES

ABLIKIM	15AA	PRL 115 182002	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	15AC	PR D92 092006	M. Ablikim <i>et al.</i>	(BES III Collab.)
VINOKUROVA	15	JHEP 1506 132	A. Vinokurova <i>et al.</i>	(BELLE Collab.)
ABLIKIM	14B	PRL 112 132001	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	14P	PRL 113 212002	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	13X	PRL 111 242001	M. Ablikim <i>et al.</i>	(BES III Collab.)