

X(4050)[±]

$$I(J^P) = ?(??)$$

OMITTED FROM SUMMARY TABLE

Observed by MIZUK 08 in the $\pi^+ \chi_{c1}(1P)$ invariant mass distribution in $\bar{B}^0 \rightarrow K^- \pi^+ \chi_{c1}(1P)$ decays. Not seen by LEES 12B in this same mode after accounting for $K\pi$ resonant mass and angular structure.

X(4050)[±] MASS

| <u>VALUE (MeV)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|--------------------|-------------|--|
| 4051 ± 14⁺²⁰₋₄₁ | ¹ MIZUK | 08 | BELL $\bar{B}^0 \rightarrow K^- \pi^+ \chi_{c1}(1P)$ |

¹ From a Dalitz plot analysis with two Breit-Wigner amplitudes.

X(4050)[±] WIDTH

| <u>VALUE (MeV)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|--|
| 82⁺²¹⁺⁴⁷₋₁₇₋₂₂ | ¹ MIZUK | 08 | BELL $\bar{B}^0 \rightarrow K^- \pi^+ \chi_{c1}(1P)$ |

¹ From a Dalitz plot analysis with two Breit-Wigner amplitudes.

X(4050)[±] DECAY MODES

| <u>Mode</u> | <u>Fraction (Γ_i/Γ)</u> |
|--------------------------------------|--|
| $\Gamma_1 \quad \pi^+ \chi_{c1}(1P)$ | seen |

X(4050)[±] BRANCHING RATIOS

| <u>$\Gamma(\pi^+ \chi_{c1}(1P))/\Gamma_{\text{total}}$</u> | <u>$\Gamma_1/\Gamma$</u> |
|---|---|
| seen | |
| ¹ MIZUK | 08 |
| BELL | $\bar{B}^0 \rightarrow K^- \pi^+ \chi_{c1}(1P)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen ² LEES 12B BABR $B \rightarrow K\pi \chi_{c1}(1P)$

¹ With a product branching fraction measurement of $B(\bar{B}^0 \rightarrow K^- X(4050)^+) \times B(X(4050)^+ \rightarrow \pi^+ \chi_{c1}(1P)) = (3.0^{+1.5+3.7}_{-0.8-1.6}) \times 10^{-5}$.

² With a product branching fraction limit of $B(\bar{B}^0 \rightarrow X(4050)^+ K^-) \times B(X(4050)^+ \rightarrow \chi_{c1} \pi^+) < 1.8 \times 10^{-5}$ at 90% CL.

X(4050)[±] REFERENCES

| | | | | |
|-------|-----|---------------|-------------------------|-----------------|
| LEES | 12B | PR D85 052003 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| MIZUK | 08 | PR D78 072004 | R. Mizuk <i>et al.</i> | (BELLE Collab.) |