

$a_1(1420)$

$$I^G(J^{PC}) = 1^-(1^{++})$$

OMITTED FROM SUMMARY TABLE

 $a_1(1420)$ MASS

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1414^{+15}_{-13}	¹ ADOLPH	15C	COMP 190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$

¹ Using the isobar model and partial-wave analysis with 88 waves. **$a_1(1420)$ WIDTH**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
153^{+8}_{-23}	¹ ADOLPH	15C	COMP 190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$

¹ Using the isobar model and partial-wave analysis with 88 waves. **$a_1(1420)$ DECAY MODES**

Mode	Fraction (Γ_j/Γ)
Γ_1 $f_0(980)\pi$	seen

 $a_1(1420)$ BRANCHING RATIOS

$\Gamma(f_0(980)\pi)/\Gamma_{\text{total}}$	Γ_1/Γ		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
seen	¹ ADOLPH	15C	COMP 190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$

¹ Using the isobar model and partial-wave analysis with 88 waves. **$a_1(1420)$ REFERENCES**

ADOLPH	15C	PRL 115 082001	C. Adolph <i>et al.</i>	(COMPASS Collab.)
--------	-----	----------------	-------------------------	-------------------