

$\eta_c(2S)$ 

$$I^G(J^{PC}) = 0^+(0^{-+})$$

Quantum numbers are quark model predictions.

### $\eta_c(2S)$ MASS

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3639.2<math>\pm</math>1.2 OUR AVERAGE</b>					
3637.0 $\pm$ 5.7 $\pm$ 3.4		178	<sup>1,2</sup> LEES	14E BABR	$\gamma\gamma \rightarrow K^+ K^- \pi^0$
3635.1 $\pm$ 5.8 $\pm$ 2.1		47	<sup>1,3</sup> LEES	14E BABR	$\gamma\gamma \rightarrow K^+ K^- \eta$
3646.9 $\pm$ 1.6 $\pm$ 3.6		57 $\pm$ 17	ABLIKIM	13K BES3	$\psi(2S) \rightarrow$ $\gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$
3637.6 $\pm$ 2.9 $\pm$ 1.6		127 $\pm$ 18	<sup>4</sup> ABLIKIM	12G BES3	$\psi(2S) \rightarrow \gamma K^0 K \pi,$ $K K \pi^0$
3638.5 $\pm$ 1.5 $\pm$ 0.8		624	<sup>1</sup> DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$
3640.5 $\pm$ 3.2 $\pm$ 2.5		1201	<sup>1</sup> DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
3636.1 $^{+3.9+0.7}_{-4.2-2.0}$		128	<sup>5</sup> VINOKUROVA 11	BELL	$B^\pm \rightarrow K^\pm (K_S^0 K^\pm \pi^\mp)$
3626 $\pm$ 5 $\pm$ 6		311	<sup>6</sup> ABE	07 BELL	$e^+ e^- \rightarrow J/\psi(c\bar{c})$
3645.0 $\pm$ 5.5 $^{+4.9}_{-7.8}$		121 $\pm$ 27	AUBERT	05C BABR	$e^+ e^- \rightarrow J/\psi c\bar{c}$
3642.9 $\pm$ 3.1 $\pm$ 1.5		61	ASNER	04 CLEO	$\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0 K^\pm \pi^\mp$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
3639 $\pm$ 7		98 $\pm$ 52	<sup>7</sup> AUBERT	06E BABR	$B^\pm \rightarrow K^\pm X_{c\bar{c}}$
3630.8 $\pm$ 3.4 $\pm$ 1.0		112 $\pm$ 24	<sup>8</sup> AUBERT	04D BABR	$\gamma\gamma \rightarrow \eta_c(2S) \rightarrow K\bar{K}\pi$
3654 $\pm$ 6 $\pm$ 8		39 $\pm$ 11	<sup>9</sup> CHOI	02 BELL	$B \rightarrow K K_S K^- \pi^+$
3594 $\pm$ 5			<sup>10</sup> EDWARDS	82C CBAL	$e^+ e^- \rightarrow \gamma X$

<sup>1</sup> Ignoring possible interference with continuum.<sup>2</sup> With a width fixed to 11.3 MeV.<sup>3</sup> With a width fixed to 11.3 MeV. Using both  $\eta \rightarrow \gamma\gamma$  and  $\eta \rightarrow \pi^+ \pi^- \pi^0$  decays.<sup>4</sup> From a simultaneous fit to  $K_S^0 K^\pm \pi^\mp$  and  $K^+ K^- \pi^0$  decay modes.<sup>5</sup> Accounts for interference with non-resonant continuum.<sup>6</sup> From a fit of the  $J/\psi$  recoil mass spectrum. Supersedes ABE,K 02 and ABE 04G.<sup>7</sup> From the fit of the kaon momentum spectrum. Systematic errors not evaluated.<sup>8</sup> Superseded by DEL-AMO-SANCHEZ 11M.<sup>9</sup> Superseded by VINOKUROVA 11.<sup>10</sup> Assuming mass of  $\psi(2S) = 3686$  MeV.

### $\eta_c(2S)$ WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>11.3<math>^{+3.2}_{-2.9}</math> OUR AVERAGE</b>					
9.9 $\pm$ 4.8 $\pm$ 2.9		57 $\pm$ 17	ABLIKIM	13K BES3	$\psi(2S) \rightarrow$ $\gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$
16.9 $\pm$ 6.4 $\pm$ 4.8		127 $\pm$ 18	<sup>11</sup> ABLIKIM	12G BES3	$\psi(2S) \rightarrow \gamma K^0 K \pi,$ $K K \pi^0$
13.4 $\pm$ 4.6 $\pm$ 3.2		624	<sup>12</sup> DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$

$6.6^{+8.4+2.6}_{-5.1-0.9}$	128	<sup>13</sup> VINOKUROVA 11	BELL	$B^{\pm} \rightarrow K^{\pm}(K_S^0 K^{\pm} \pi^{\mp})$
$6.3 \pm 12.4 \pm 4.0$	61	ASNER	04 CLEO	$\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0 K^{\pm} \pi^{\mp}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 23	90	$98 \pm 52$	<sup>14</sup> AUBERT	06E BABR	$B^{\pm} \rightarrow K^{\pm} X_{c\bar{c}}$
$22 \pm 14$		$121 \pm 27$	AUBERT	05C BABR	$e^+ e^- \rightarrow J/\psi c\bar{c}$
$17.0 \pm 8.3 \pm 2.5$		$112 \pm 24$	<sup>15</sup> AUBERT	04D BABR	$\gamma\gamma \rightarrow \eta_c(2S) \rightarrow K\bar{K}\pi$
<55	90	$39 \pm 11$	<sup>16</sup> CHOI	02 BELL	$B \rightarrow K K_S K^- \pi^+$
<8.0	95		<sup>17</sup> EDWARDS	82C CBAL	$e^+ e^- \rightarrow \gamma X$

<sup>11</sup> From a simultaneous fit to  $K_S^0 K^{\pm} \pi^{\mp}$  and  $K^+ K^- \pi^0$  decay modes.

<sup>12</sup> Ignoring possible interference with continuum.

<sup>13</sup> Accounts for interference with non-resonant continuum.

<sup>14</sup> From the fit of the kaon momentum spectrum. Systematic errors not evaluated.

<sup>15</sup> Superseded by DEL-AMO-SANCHEZ 11M.

<sup>16</sup> For a mass value of  $3654 \pm 6$  MeV. Superseded by VINOKUROVA 11.

<sup>17</sup> For a mass value of  $3594 \pm 5$  MeV

### $\eta_c(2S)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1$ hadrons	not seen	
$\Gamma_2$ $K\bar{K}\pi$	( $1.9 \pm 1.2$ ) %	
$\Gamma_3$ $K\bar{K}\eta$	( $5 \pm 4$ ) $\times 10^{-3}$	
$\Gamma_4$ $2\pi^+ 2\pi^-$	not seen	
$\Gamma_5$ $\rho^0 \rho^0$	not seen	
$\Gamma_6$ $3\pi^+ 3\pi^-$	not seen	
$\Gamma_7$ $K^+ K^- \pi^+ \pi^-$	not seen	
$\Gamma_8$ $K^{*0} \bar{K}^{*0}$	not seen	
$\Gamma_9$ $K^+ K^- \pi^+ \pi^- \pi^0$	( $1.4 \pm 1.0$ ) %	
$\Gamma_{10}$ $K^+ K^- 2\pi^+ 2\pi^-$	not seen	
$\Gamma_{11}$ $K_S^0 K^- 2\pi^+ \pi^- + c.c.$	seen	
$\Gamma_{12}$ $2K^+ 2K^-$	not seen	
$\Gamma_{13}$ $\phi\phi$	not seen	
$\Gamma_{14}$ $p\bar{p}$	< 2.0 $\times 10^{-3}$	90%
$\Gamma_{15}$ $\gamma\gamma$	( $1.9 \pm 1.3$ ) $\times 10^{-4}$	
$\Gamma_{16}$ $\pi^+ \pi^- \eta$	not seen	
$\Gamma_{17}$ $\pi^+ \pi^- \eta'$	not seen	
$\Gamma_{18}$ $\pi^+ \pi^- \eta_c(1S)$	< 25 %	90%

$\eta_c(2S)$  PARTIAL WIDTHS $\Gamma(\gamma\gamma)$  $\Gamma_{15}$ 

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.3 \pm 0.6$	<sup>18</sup> ASNER	04	CLEO $\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0 K^\pm \pi^\mp$
---------------	---------------------	----	--

<sup>18</sup> They measure  $\Gamma(\eta_c(2S)\gamma\gamma) B(\eta_c(2S) \rightarrow K\bar{K}\pi) = (0.18 \pm 0.05 \pm 0.02) \Gamma(\eta_c(1S)\gamma\gamma) B(\eta_c(1S) \rightarrow K\bar{K}\pi)$ . The value for  $\Gamma(\eta_c(2S) \rightarrow \gamma\gamma)$  is derived assuming that the branching fractions for  $\eta_c(2S)$  and  $\eta_c(1S)$  decays to  $K_S K\pi$  are equal and using  $\Gamma(\eta_c(1S) \rightarrow \gamma\gamma) = 7.4 \pm 0.4 \pm 2.3$  keV.

 $\eta_c(2S) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$ 

$\Gamma(2\pi^+ 2\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$				$\Gamma_4 \Gamma_{15}/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT

<b>&lt;6.5</b>	90	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S) \rightarrow 2(\pi^+ \pi^-)$
----------------	----	--------	----	---

$\Gamma(K\bar{K}\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$				$\Gamma_2 \Gamma_{15}/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT

<b>41 ± 4 ± 6</b>	624	<sup>19</sup> DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$
-------------------	-----	-------------------------------	------	--

<sup>19</sup> Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.

$\Gamma(K^+ K^- \pi^+ \pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$				$\Gamma_7 \Gamma_{15}/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT

<b>&lt;5.0</b>	90	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S) \rightarrow K^+ K^- \pi^+ \pi^-$
----------------	----	--------	----	--

$\Gamma(K^+ K^- \pi^+ \pi^- \pi^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$				$\Gamma_9 \Gamma_{15}/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT

<b>30 ± 6 ± 5</b>	1201	<sup>20</sup> DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
-------------------	------	-------------------------------	------	--

<sup>20</sup> Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.

$\Gamma(2K^+ 2K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$				$\Gamma_{12} \Gamma_{15}/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT

<b>&lt;2.9</b>	90	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S) \rightarrow 2(K^+ K^-)$
----------------	----	--------	----	---

$\Gamma(\pi^+ \pi^- \eta_c(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$				$\Gamma_{18} \Gamma_{15}/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT

<b>&lt;133</b>	90	LEES	12AE	BABR $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \eta_c$
----------------	----	------	------	---

 $\eta_c(2S) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma^2(\text{total})$ 

$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$				$\Gamma_{14}/\Gamma \times \Gamma_{15}/\Gamma$
VALUE (units $10^{-8}$ )	CL%	DOCUMENT ID	TECN	COMMENT

<b>&lt; 5.6</b>	90 <sup>21,22,23</sup>	AMBROGIANI 01	E835	$\bar{p}p \rightarrow \gamma\gamma$
-----------------	------------------------	---------------	------	-------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 8.0	90 <sup>21,22,24</sup>	AMBROGIANI 01	E835	$\bar{p}p \rightarrow \gamma\gamma$
-------	------------------------	---------------	------	-------------------------------------

<12.0	90 <sup>22,24</sup>	AMBROGIANI 01	E835	$\bar{p}p \rightarrow \gamma\gamma$
-------	---------------------	---------------	------	-------------------------------------

<sup>21</sup> Including the measurements of of ARMSTRONG 95F in the AMBROGIANI 01 analysis.

<sup>22</sup> For a total width  $\Gamma=5$  MeV.

<sup>23</sup> For the resonance mass region 3589–3599 MeV/ $c^2$ .

<sup>24</sup> For the resonance mass region 3575–3660 MeV/ $c^2$ .

## $\eta_c(2S)$ BRANCHING RATIOS

### $\Gamma(\text{hadrons})/\Gamma_{\text{total}}$

$\Gamma_1/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>not seen</b>	ABREU	98O	DLPH $e^+e^- \rightarrow e^+e^- + \text{hadrons}$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
seen	<sup>25</sup> EDWARDS	82C	CBAL $e^+e^- \rightarrow \gamma X$
<sup>25</sup> For a mass value of $3594 \pm 5$ MeV			

### $\Gamma(K\bar{K}\pi)/\Gamma_{\text{total}}$

$\Gamma_2/\Gamma$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>1.9 \pm 0.4 \pm 1.1</math></b>	$59 \pm 12$	<sup>26</sup> AUBERT	08AB	BABR $B \rightarrow \eta_c(2S)K \rightarrow K\bar{K}\pi K$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	$127 \pm 18$	ABLIKIM	13K	BES3 $\psi(2S) \rightarrow \gamma K\bar{K}\pi$
seen	$39 \pm 11$	<sup>27</sup> CHOI	02	BELL $B \rightarrow K K_S K^- \pi^+$

<sup>26</sup> Derived from a measurement of  $[B(B^+ \rightarrow \eta_c(2S)K^+) \times B(\eta_c(2S) \rightarrow K\bar{K}\pi)] / [B(B^+ \rightarrow \eta_c K^+) \times B(\eta_c \rightarrow K\bar{K}\pi)] = (9.6^{+2.0}_{-1.9} \pm 2.5)\%$  and using  $B(B^+ \rightarrow \eta_c(2S)K^+) = (3.4 \pm 1.8) \times 10^{-4}$ , and  $[B(B^+ \rightarrow \eta_c K^+) \times B(\eta_c \rightarrow K\bar{K}\pi)] = (6.88 \pm 0.77^{+0.55}_{-0.66}) \times 10^{-5}$ .

<sup>27</sup> For a mass value of  $3654 \pm 6$  MeV

### $\Gamma(K\bar{K}\eta)/\Gamma(K\bar{K}\pi)$

$\Gamma_3/\Gamma_2$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>27.3 \pm 7.0 \pm 9.0</math></b>	225	<sup>28</sup> LEES	14E	BABR $\gamma\gamma \rightarrow K^+ K^- \gamma\gamma$

<sup>28</sup> LEES 14E reports  $B(\eta_c(2S) \rightarrow K^+ K^- \eta)/B(\eta_c(2S) \rightarrow K^+ K^- \pi^0) = 0.82 \pm 0.21 \pm 0.27$ , which we divide by 3 to account for isospin symmetry.

### $\Gamma(2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$

$\Gamma_4/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>not seen</b>	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S)$

### $\Gamma(\rho^0 \rho^0)/\Gamma_{\text{total}}$

$\Gamma_5/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>not seen</b>	ABLIKIM	11H	BES3 $\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$

### $\Gamma(K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$

$\Gamma_7/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>not seen</b>	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S)$

$\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K\bar{K}\pi)$   $\Gamma_9/\Gamma_2$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.73±0.17±0.17</b>	1201	<sup>29</sup> DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

<sup>29</sup> We have multiplied the value of  $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K_S^0 K^\pm \pi^\mp)$  reported in DEL-AMO-SANCHEZ 11M by a factor 1/3 to obtain  $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K\bar{K}\pi)$ . Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.

$\Gamma(K^*0\bar{K}^*0)/\Gamma_{total}$   $\Gamma_8/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>not seen</b>	ABLIKIM	11H BES3	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$

$\Gamma(K_S^0 K^- 2\pi^+ \pi^- + c.c.)/\Gamma_{total}$   $\Gamma_{11}/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	57±17	ABLIKIM	13K BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$

$\Gamma(2K^+ 2K^-)/\Gamma_{total}$   $\Gamma_{12}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>not seen</b>	UEHARA	08 BELL	$\gamma\gamma \rightarrow \eta_c(2S)$

$\Gamma(\phi\phi)/\Gamma_{total}$   $\Gamma_{13}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>not seen</b>	ABLIKIM	11H BES3	$\psi(2S) \rightarrow \gamma K^+ K^- K^+ K^-$

$\Gamma(\gamma\gamma)/\Gamma_{total}$   $\Gamma_{15}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • •				We do not use the following data for averages, fits, limits, etc. • • •
<5 × 10 <sup>-4</sup>	90	<sup>30</sup> WICHT	08 BELL	$B^\pm \rightarrow K^\pm \gamma\gamma$
not seen		AMBROGIANI	01 E835	$\bar{p}p \rightarrow \gamma\gamma$
<0.01	90	LEE	85 CBAL	$\psi' \rightarrow \text{photons}$

<sup>30</sup> WICHT 08 reports  $[\Gamma(\eta_c(2S) \rightarrow \gamma\gamma)/\Gamma_{total}] \times [B(B^+ \rightarrow \eta_c(2S) K^+)] < 0.18 \times 10^{-6}$  which we divide by our best value  $B(B^+ \rightarrow \eta_c(2S) K^+) = 3.4 \times 10^{-4}$ .

$\Gamma(\pi^+ \pi^- \eta_c(1S))/\Gamma(K\bar{K}\pi)$   $\Gamma_{18}/\Gamma_2$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;3.33</b>	90	<sup>31</sup> LEES	12AE BABR	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \eta_c$

<sup>31</sup> We divided the reported limit by 3 to take into account isospin relations.

**$\eta_c(2S)$  CROSS-PARTICLE BRANCHING RATIOS**

$\Gamma(\eta_c(2S) \rightarrow 2\pi^+ 2\pi^-)/\Gamma_{total} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{total}$   
 $\Gamma_4/\Gamma \times \Gamma_{136}^{\psi(2S)}/\Gamma\psi(2S)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;14.6 × 10<sup>-6</sup></b>	90	<sup>32</sup> CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$

<sup>32</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow \rho^0 \rho^0) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}}$$

$$\Gamma_5 / \Gamma \times \Gamma_{136}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<12.7 \times 10^{-7}$	90	ABLIKIM	11H BES3	$\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$

$$\Gamma(\eta_c(2S) \rightarrow 3\pi^+ 3\pi^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}}$$

$$\Gamma_6 / \Gamma \times \Gamma_{136}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<13.2 \times 10^{-6}$	90	33 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma 3\pi^+ 3\pi^-$

<sup>33</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow K^+ K^- \pi^+ \pi^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}}$$

$$\Gamma_7 / \Gamma \times \Gamma_{136}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<9.6 \times 10^{-6}$	90	34 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$

<sup>34</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow K^{*0} \bar{K}^{*0}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}}$$

$$\Gamma_8 / \Gamma \times \Gamma_{136}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<19.6 \times 10^{-7}$	90	ABLIKIM	11H BES3	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$

$$\Gamma(\eta_c(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}}$$

$$\Gamma_9 / \Gamma \times \Gamma_{136}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<43.0 \times 10^{-6}$	90	35 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^- \pi^0$

<sup>35</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow K^+ K^- 2\pi^+ 2\pi^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}}$$

$$\Gamma_{10} / \Gamma \times \Gamma_{136}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<9.7 \times 10^{-6}$	90	36 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- 2\pi^+ 2\pi^-$

<sup>36</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}}$$

$$\Gamma_{11} / \Gamma \times \Gamma_{136}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$7.03 \pm 2.10 \pm 0.7$	60		ABLIKIM	13K BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 15.2$	90	37 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}$
----------	----	-------------------	------	--

<sup>37</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow \phi\phi)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}} \quad \Gamma_{13}/\Gamma \times \Gamma_{136}^{\psi(2S)}/\Gamma\psi(2S)$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.8 \times 10^{-7}$	90	ABLIKIM	11H BES3	$\psi(2S) \rightarrow \gamma K^+ K^- K^+ K^-$

$$\Gamma(\eta_c(2S) \rightarrow \pi^+\pi^-\eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}} \quad \Gamma_{16}/\Gamma \times \Gamma_{136}^{\psi(2S)}/\Gamma\psi(2S)$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.3 \times 10^{-6}$	90	<sup>38</sup> CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma\pi^+\pi^-\eta$

<sup>38</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow \pi^+\pi^-\eta')/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}} \quad \Gamma_{17}/\Gamma \times \Gamma_{136}^{\psi(2S)}/\Gamma\psi(2S)$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<14.2 \times 10^{-6}$	90	<sup>39</sup> CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma\pi^+\pi^-\eta'$

<sup>39</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow K\bar{K}\eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}} \quad \Gamma_3/\Gamma \times \Gamma_{136}^{\psi(2S)}/\Gamma\psi(2S)$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<11.8 \times 10^{-6}$	90	<sup>40</sup> CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- \eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>40</sup> CRONIN-HENNESSY 10 reports a limit of  $< 5.9 \times 10^{-6}$  for the decay  $\eta_c(2S) \rightarrow K^+ K^- \eta$  which we multiply by 2 account for isospin symmetry. It assumes  $\Gamma(\eta_c(2S)) = 14$  MeV. It also gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow \pi^+\pi^-\eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}} \quad \Gamma_{18}/\Gamma \times \Gamma_{136}^{\psi(2S)}/\Gamma\psi(2S)$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.7 \times 10^{-4}$	90	<sup>41</sup> CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma\pi^+\pi^-\eta_c(1S)$

<sup>41</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}} \quad \Gamma_{14}/\Gamma \times \Gamma_{136}^{\psi(2S)}/\Gamma\psi(2S)$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.4 \times 10^{-6}$	90	ABLIKIM	13V BES3	$\psi(2S) \rightarrow \gamma p\bar{p}$

$\eta_c(2S)$  REFERENCES

LEES	14E	PR D89 112004	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	13K	PR D87 052005	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	13V	PR D88 112001	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	12G	PRL 109 042003	M. Ablikim <i>et al.</i>	(BES III Collab.)
LEES	12AE	PR D86 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	11H	PR D84 091102	M. Ablikim <i>et al.</i>	(BES III Collab.)
DEL-AMO-SA...	11M	PR D84 012004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
VINOKUROVA	11	PL B706 139	A. Vinokurova <i>et al.</i>	(BELLE Collab.)
CRONIN-HEN...	10	PR D81 052002	D. Cronin-Hennessey <i>et al.</i>	(CLEO Collab.)
AUBERT	08AB	PR D78 012006	B. Aubert <i>et al.</i>	(BABAR Collab.)
UEHARA	08	EPJ C53 1	S. Uehara <i>et al.</i>	(BELLE Collab.)
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)
ABE	07	PRL 98 082001	K. Abe <i>et al.</i>	(BELLE Collab.)
AUBERT	06E	PRL 96 052002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	05C	PR D72 031101	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABE	04G	PR D70 071102	K. Abe <i>et al.</i>	(BELLE Collab.)
ASNER	04	PRL 92 142001	D.M. Asner <i>et al.</i>	(CLEO Collab.)
AUBERT	04D	PRL 92 142002	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABE,K	02	PRL 89 142001	K. Abe <i>et al.</i>	(BELLE Collab.)
CHOI	02	PRL 89 102001	S.-K. Choi <i>et al.</i>	(BELLE Collab.)
AMBROGIANI	01	PR D64 052003	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)
ABREU	98O	PL B441 479	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ARMSTRONG	95F	PR D52 4839	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
LEE	85	SLAC 282	R.A. Lee	(SLAC)
EDWARDS	82C	PRL 48 70	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)

---