

$h_c(1P)$

$$I^G(J^{PC}) = ??(1^{+-})$$

Quantum numbers are quark model prediction, $C = -$ established by $\eta_c \gamma$ decay.

$h_c(1P)$ MASS

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3525.38 ± 0.11 OUR AVERAGE				
3525.31 ± 0.11 ± 0.14	832	¹ ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma$ hadrons
3525.40 ± 0.13 ± 0.18	3679	ABLIKIM	10B BES3	$\psi(2S) \rightarrow \pi^0 \gamma \eta_c$
3525.20 ± 0.18 ± 0.12	1282	² DOBBS	08A CLEO	$\psi(2S) \rightarrow \pi^0 \eta_c \gamma$
3525.8 ± 0.2 ± 0.2	13	ANDREOTTI	05B E835	$\bar{p} p \rightarrow \eta_c \gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3525.6 ± 0.5	92 ⁺²³ ₋₂₂	ADAMS	09 CLEO	$\psi(2S) \rightarrow 2(\pi^+ \pi^- \pi^0)$
3524.4 ± 0.6 ± 0.4	168 ± 40	³ ROSNER	05 CLEO	$\psi(2S) \rightarrow \pi^0 \eta_c \gamma$
3527 ± 8	42	ANTONIAZZI	94 E705	300 $\pi^\pm, p\text{Li} \rightarrow J/\psi \pi^0 X$
3526.28 ± 0.18 ± 0.19	59	⁴ ARMSTRONG	92D E760	$\bar{p} p \rightarrow J/\psi \pi^0$
3525.4 ± 0.8 ± 0.4	5	BAGLIN	86 SPEC	$\bar{p} p \rightarrow J/\psi X$

¹ With floating width.

² Combination of exclusive and inclusive analyses for the reaction $\psi(2S) \rightarrow \pi^0 h_c \rightarrow \pi^0 \eta_c \gamma$. This result is the average of DOBBS 08A and ROSNER 05.

³ Superseded by DOBBS 08A.

⁴ Mass central value and systematic error recalculated by us according to Eq. (16) in ARMSTRONG 93B, using the value for the $\psi(2S)$ mass from AULCHENKO 03.

$h_c(1P)$ WIDTH

<u>VALUE (MeV)</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.70 ± 0.28 ± 0.22					
		832	⁵ ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 1.44	90	3679	⁶ ABLIKIM	10B BES3	$\psi(2S) \rightarrow \pi^0 \gamma \eta_c$
< 1		13	ANDREOTTI	05B E835	$\bar{p} p \rightarrow \eta_c \gamma$
< 1.1	90	59	ARMSTRONG	92D E760	$\bar{p} p \rightarrow J/\psi \pi^0$

⁵ With floating mass.

⁶ The central value is $\Gamma = 0.73 \pm 0.45 \pm 0.28$ MeV.

$h_c(1P)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $J/\psi(1S) \pi^0$		
Γ_2 $J/\psi(1S) \pi \pi$	not seen	
Γ_3 $p \bar{p}$	< 1.5 × 10 ⁻⁴	90%

Γ_4	$\eta_c(1S)\gamma$	$(51 \pm 6) \%$
Γ_5	$\pi^+\pi^-\pi^0$	$< 2.2 \times 10^{-3}$
Γ_6	$2\pi^+2\pi^-\pi^0$	$(2.2^{+0.8}_{-0.7}) \%$
Γ_7	$3\pi^+3\pi^-\pi^0$	$< 2.9 \%$

$h_c(1P)$ PARTIAL WIDTHS

$h_c(1P) \Gamma(i)\Gamma(\bar{p}p)/\Gamma(\text{total})$

$\Gamma(\eta_c(1S)\gamma) \times \Gamma(p\bar{p})/\Gamma_{\text{total}}$ $\Gamma_4\Gamma_3/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
12.0 ± 4.5	13	⁷ ANDREOTTI	05B E835	$\bar{p}p \rightarrow \eta_c\gamma$
⁷ Assuming $\Gamma = 1$ MeV.				

$h_c(1P)$ BRANCHING RATIOS

$\Gamma(J/\psi(1S)\pi\pi)/\Gamma(J/\psi(1S)\pi^0)$ Γ_2/Γ_1

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.18	90	ARMSTRONG	92D E760	$\bar{p}p \rightarrow J/\psi\pi^0$

$\Gamma(\eta_c(1S)\gamma)/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
51 ± 6 OUR AVERAGE				
$54.3 \pm 6.7 \pm 5.2$	3679	ABLIKIM	10B BES3	$\psi(2S) \rightarrow \pi^0\gamma\eta_c$
$48 \pm 6 \pm 7$		⁸ DOBBS	08A CLEO	$\psi(2S) \rightarrow \pi^0\eta_c\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$48 \pm 6 \pm 7$	1282	⁹ DOBBS	08A CLEO	$\psi(2S) \rightarrow \pi^0\eta_c\gamma$
$46 \pm 12 \pm 7$	168	¹⁰ ROSNER	05 CLEO	$\psi(2S) \rightarrow \pi^0\eta_c\gamma$

⁸ Average of DOBBS 08A and ROSNER 05. DOBBS 08A reports $[\Gamma(h_c(1P) \rightarrow \eta_c(1S)\gamma)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \pi^0 h_c(1P))]$ = $(4.16 \pm 0.30 \pm 0.37) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \pi^0 h_c(1P)) = (8.6 \pm 1.3) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁹ DOBBS 08A reports $[\Gamma(h_c(1P) \rightarrow \eta_c(1S)\gamma)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \pi^0 h_c(1P))]$ = $(4.19 \pm 0.32 \pm 0.45) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \pi^0 h_c(1P)) = (8.6 \pm 1.3) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

¹⁰ ROSNER 05 reports $[\Gamma(h_c(1P) \rightarrow \eta_c(1S)\gamma)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \pi^0 h_c(1P))]$ = $(4.0 \pm 0.8 \pm 0.7) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \pi^0 h_c(1P)) = (8.6 \pm 1.3) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
<2.2	¹¹ ADAMS	09 CLEO	$\psi(2S) \rightarrow \pi^0\gamma\eta_c$

¹¹ ADAMS 09 reports $[\Gamma(h_c(1P) \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \pi^0 h_c(1P))]$ = $< 0.19 \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \pi^0 h_c(1P)) = 8.6 \times 10^{-4}$.

$\Gamma(2\pi^+2\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.2^{+0.8}_{-0.6} \pm 0.3$	92	¹² ADAMS	09	CLEO $\psi(2S) \rightarrow \pi^0 \gamma \eta_c$

¹² ADAMS 09 reports $[\Gamma(h_c(1P) \rightarrow 2\pi^+2\pi^-\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \pi^0 h_c(1P))] = (1.88^{+0.48}_{-0.45} + 0.47_{-0.30}) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \pi^0 h_c(1P)) = (8.6 \pm 1.3) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(3\pi^+3\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
<2.9	¹³ ADAMS	09	CLEO $\psi(2S) \rightarrow \pi^0 \gamma \eta_c$

¹³ ADAMS 09 reports $[\Gamma(h_c(1P) \rightarrow 3\pi^+3\pi^-\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \pi^0 h_c(1P))] < 2.5 \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \pi^0 h_c(1P)) = 8.6 \times 10^{-4}$.

$\Gamma(h_c(1P) \rightarrow \eta_c(1S)\gamma)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}$ $\Gamma_4/\Gamma \times \Gamma_{15}^{\psi(2S)}/\Gamma_{\psi(2S)}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
4.3 ± 0.4 OUR AVERAGE				
$4.58 \pm 0.40 \pm 0.50$	3679	¹⁴ ABLIKIM	10B	BES3 $\psi(2S) \rightarrow \pi^0 \gamma X$
$4.16 \pm 0.30 \pm 0.37$	1430	¹⁵ DOBBS	08A	CLEO $\psi(2S) \rightarrow \pi^0 \gamma \eta_c$

¹⁴ Not independent of other branching fractions in ABLIKIM 10B.

¹⁵ Not independent of other branching fractions in DOBBS 08A.

$\Gamma(h_c(1P) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}$ $\Gamma_3/\Gamma \times \Gamma_{15}^{\psi(2S)}/\Gamma_{\psi(2S)}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.3 × 10⁻⁷	90	ABLIKIM	13V	BES3 $\psi(2S) \rightarrow \gamma p\bar{p}$

$h_c(1P)$ REFERENCES

ABLIKIM	13V	PR D88 112001	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	12N	PR D86 092009	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	10B	PRL 104 132002	M. Ablikim <i>et al.</i>	(BES III Collab.)
ADAMS	09	PR D80 051106	G.S. Adams <i>et al.</i>	(CLEO Collab.)
DOBBS	08A	PRL 101 182003	S. Dobbs <i>et al.</i>	(CLEO Collab.)
ANDREOTTI	05B	PR D72 032001	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)
ROSNER	05	PRL 95 102003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)
ANTONIAZZI	94	PR D50 4258	L. Antoniazzi <i>et al.</i>	(E705 Collab.)
ARMSTRONG	93B	PR D47 772	T.A. Armstrong <i>et al.</i>	(FNAL E760 Collab.)
ARMSTRONG	92D	PRL 69 2337	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
BAGLIN	86	PL B171 135	C. Baglin <i>et al.</i>	(LAPP, CERN, TORI, STRB+)