

$\Upsilon(10860)$ 

$$J^{PC} = 0^{-}(1^{-}-)$$

### $\Upsilon(10860)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$10891.1 \pm 3.2^{+1.2}_{-2.0}$	<sup>1</sup> SANTEL	16	BELL $e^+e^- \rightarrow \Upsilon(1S, 2S, 3S)\pi^+\pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$10881.8^{+1.0}_{-1.1} \pm 1.2$	<sup>2,3</sup> SANTEL	16	BELL $e^+e^- \rightarrow$ hadrons
$10879 \pm 3$	<sup>4,5</sup> CHEN	10	BELL $e^+e^- \rightarrow$ hadrons
$10888.4^{+2.7}_{-2.6} \pm 1.2$	<sup>6</sup> CHEN	10	BELL $e^+e^- \rightarrow \Upsilon(1S, 2S, 3S)\pi^+\pi^-$
$10876 \pm 2$	<sup>4</sup> AUBERT	09E	BABR $e^+e^- \rightarrow$ hadrons
$10869 \pm 2$	<sup>7</sup> AUBERT	09E	BABR $e^+e^- \rightarrow$ hadrons
$10868 \pm 6 \pm 5$	<sup>8</sup> BESSON	85	CLEO $e^+e^- \rightarrow$ hadrons
$10845 \pm 20$	<sup>9</sup> LOVELOCK	85	CUSB $e^+e^- \rightarrow$ hadrons

<sup>1</sup> From a simultaneous fit to the  $\Upsilon(nS)\pi^+\pi^-$ ,  $n = 1, 2, 3$  cross sections at 25 energy points within  $\sqrt{s} = 10.6\text{--}11.05$  GeV to a pair of interfering Breit-Wigner amplitudes modified by phase space factors, with fourteen resonance parameters (a mass, width, and three amplitudes for each of  $\Upsilon(10860)$  and  $\Upsilon(11020)$ , a single universal relative phase, and three decoherence coefficients, one for each  $n$ ). Continuum contributions were measured (and therefore fixed) to be zero.

<sup>2</sup> From a fit to the total hadronic cross sections measured at 60 energy points within  $\sqrt{s} = 10.82\text{--}11.05$  GeV to a pair of interfering Breit-Wigner amplitudes and two floating continuum amplitudes with  $1/\sqrt{s}$  dependence, one coherent with the resonances and one incoherent, with six resonance parameters (a mass, width, and an amplitude for each of  $\Upsilon(10860)$  and  $\Upsilon(11020)$ , one relative phase, and one decoherence coefficient).

<sup>3</sup> Not including uncertain and potentially large systematic errors due to assumed continuum amplitude  $1/\sqrt{s}$  dependence and related interference contributions.

<sup>4</sup> In a model where a flat non-resonant  $b\bar{b}$ -continuum is incoherently added to a second flat component interfering with two Breit-Wigner resonances. Systematic uncertainties not estimated.

<sup>5</sup> The parameters of the  $\Upsilon(11020)$  are fixed to those in AUBERT 09E.

<sup>6</sup> In a model where a flat nonresonant  $\Upsilon(1S, 2S, 3S)\pi^+\pi^-$  continuum interferes with a single Breit-Wigner resonance.

<sup>7</sup> In a model where a non-resonant  $b\bar{b}$ -continuum represented by a threshold function at  $\sqrt{s}=2m_B$  is incoherently added to a flat component interfering with two Breit-Wigner resonances. Not independent of other AUBERT 09E results. Systematic uncertainties not estimated.

<sup>8</sup> Assuming four Gaussians with radiative tails and a single step in  $R$ .

<sup>9</sup> In a coupled-channel model with three resonances and a smooth step in  $R$ .

### $\Upsilon(10860)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$53.7^{+7.1+1.3}_{-5.6-5.4}$	<sup>10</sup> SANTEL	16	BELL $e^+e^- \rightarrow \Upsilon(1S, 2S, 3S)\pi^+\pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$48.5^{+1.9+2.0}_{-1.8-2.8}$	<sup>11,12</sup> SANTEL	16	BELL $e^+e^- \rightarrow$ hadrons

46	$\begin{matrix} +9 \\ -7 \end{matrix}$	13,14 CHEN	10	BELL	$e^+e^- \rightarrow \text{hadrons}$
30.7	$\begin{matrix} +8.3 \\ -7.0 \end{matrix} \pm 3.1$	15 CHEN	10	BELL	$e^+e^- \rightarrow \Upsilon(1S, 2S, 3S)\pi^+\pi^-$
43	$\pm 4$	13 AUBERT	09E	BABR	$e^+e^- \rightarrow \text{hadrons}$
74	$\pm 4$	16 AUBERT	09E	BABR	$e^+e^- \rightarrow \text{hadrons}$
112	$\pm 17 \pm 23$	17 BESSON	85	CLEO	$e^+e^- \rightarrow \text{hadrons}$
110	$\pm 15$	18 LOVELOCK	85	CUSB	$e^+e^- \rightarrow \text{hadrons}$

<sup>10</sup> From a simultaneous fit to the  $\Upsilon(nS)\pi^+\pi^-$ ,  $n = 1, 2, 3$  cross sections at 25 energy points within  $\sqrt{s} = 10.6\text{--}11.05$  GeV to a pair of interfering Breit-Wigner amplitudes modified by phase space factors, with fourteen resonance parameters (a mass, width, and three amplitudes for each of  $\Upsilon(10860)$  and  $\Upsilon(11020)$ , a single universal relative phase, and three decoherence coefficients, one for each  $n$ ). Continuum contributions were measured (and therefore fixed) to be zero.

<sup>11</sup> From a fit to the total hadronic cross sections measured at 60 energy points within  $\sqrt{s} = 10.82\text{--}11.05$  GeV to a pair of interfering Breit-Wigner amplitudes and two floating continuum amplitudes with  $1/\sqrt{s}$  dependence, one coherent with the resonances and one incoherent, with six resonance parameters (a mass, width, and an amplitude for each of  $\Upsilon(10860)$  and  $\Upsilon(11020)$ , one relative phase, and one decoherence coefficient).

<sup>12</sup> Not including uncertain and potentially large systematic errors due to assumed continuum amplitude  $1/\sqrt{s}$  dependence and related interference contributions.

<sup>13</sup> In a model where a flat non-resonant  $b\bar{b}$ -continuum is incoherently added to a second flat component interfering with two Breit-Wigner resonances. Systematic uncertainties not estimated.

<sup>14</sup> The parameters of the  $\Upsilon(11020)$  are fixed to those in AUBERT 09E.

<sup>15</sup> In a model where a flat nonresonant  $\Upsilon(1S, 2S, 3S)\pi^+\pi^-$  continuum interferes with a single Breit-Wigner resonance.

<sup>16</sup> In a model where a non-resonant  $b\bar{b}$ -continuum represented by a threshold function at  $\sqrt{s}=2m_B$  is incoherently added to a flat component interfering with two Breit-Wigner resonances. Not independent of other AUBERT 09E results. Systematic uncertainties not estimated.

<sup>17</sup> Assuming four Gaussians with radiative tails and a single step in  $R$ .

<sup>18</sup> In a coupled-channel model with three resonances and a smooth step in  $R$ .

### $\Upsilon(10860)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1$ $B\bar{B}X$	( 76.2 $\begin{matrix} +2.7 \\ -4.0 \end{matrix}$ ) %	
$\Gamma_2$ $B\bar{B}$	( 5.5 $\pm 1.0$ ) %	
$\Gamma_3$ $B\bar{B}^* + \text{c.c.}$	( 13.7 $\pm 1.6$ ) %	
$\Gamma_4$ $B^*\bar{B}^*$	( 38.1 $\pm 3.4$ ) %	
$\Gamma_5$ $B\bar{B}^{(*)}\pi$	< 19.7 %	90%
$\Gamma_6$ $B\bar{B}\pi$	( 0.0 $\pm 1.2$ ) %	
$\Gamma_7$ $B^*\bar{B}\pi + B\bar{B}^*\pi$	( 7.3 $\pm 2.3$ ) %	
$\Gamma_8$ $B^*\bar{B}^*\pi$	( 1.0 $\pm 1.4$ ) %	
$\Gamma_9$ $B\bar{B}\pi\pi$	< 8.9 %	90%
$\Gamma_{10}$ $B_s^{(*)}\bar{B}_s^{(*)}$	( 20.1 $\pm 3.1$ ) %	
$\Gamma_{11}$ $B_s\bar{B}_s$	( 5 $\pm 5$ ) $\times 10^{-3}$	
$\Gamma_{12}$ $B_s\bar{B}_s^* + \text{c.c.}$	( 1.35 $\pm 0.32$ ) %	
$\Gamma_{13}$ $B_s^*\bar{B}_s^*$	( 17.6 $\pm 2.7$ ) %	

$\Gamma_{14}$	no open-bottom	( 3.8 $^{+5.0}_{-0.5}$ ) %	
$\Gamma_{15}$	$e^+ e^-$	( 5.7 $\pm 1.5$ ) $\times 10^{-6}$	
$\Gamma_{16}$	$K^*(892)^0 \bar{K}^0$	< 1.0 $\times 10^{-5}$	90%
$\Gamma_{17}$	$\Upsilon(1S) \pi^+ \pi^-$	( 5.3 $\pm 0.6$ ) $\times 10^{-3}$	
$\Gamma_{18}$	$\Upsilon(2S) \pi^+ \pi^-$	( 7.8 $\pm 1.3$ ) $\times 10^{-3}$	
$\Gamma_{19}$	$\Upsilon(3S) \pi^+ \pi^-$	( 4.8 $^{+1.9}_{-1.7}$ ) $\times 10^{-3}$	
$\Gamma_{20}$	$\Upsilon(1S) K^+ K^-$	( 6.1 $\pm 1.8$ ) $\times 10^{-4}$	
$\Gamma_{21}$	$h_b(1P) \pi^+ \pi^-$	( 3.5 $^{+1.0}_{-1.3}$ ) $\times 10^{-3}$	
$\Gamma_{22}$	$h_b(2P) \pi^+ \pi^-$	( 6.0 $^{+2.1}_{-1.8}$ ) $\times 10^{-3}$	
$\Gamma_{23}$	$\chi_{b0}(1P) \pi^+ \pi^- \pi^0$	< 6.3 $\times 10^{-3}$	90%
$\Gamma_{24}$	$\chi_{b0}(1P) \omega$	< 3.9 $\times 10^{-3}$	90%
$\Gamma_{25}$	$\chi_{b0}(1P) (\pi^+ \pi^- \pi^0)_{\text{non-}\omega}$	< 4.8 $\times 10^{-3}$	90%
$\Gamma_{26}$	$\chi_{b1}(1P) \pi^+ \pi^- \pi^0$	( 1.85 $\pm 0.33$ ) $\times 10^{-3}$	
$\Gamma_{27}$	$\chi_{b1}(1P) \omega$	( 1.57 $\pm 0.30$ ) $\times 10^{-3}$	
$\Gamma_{28}$	$\chi_{b1}(1P) (\pi^+ \pi^- \pi^0)_{\text{non-}\omega}$	( 5.2 $\pm 1.9$ ) $\times 10^{-4}$	
$\Gamma_{29}$	$\chi_{b2}(1P) \pi^+ \pi^- \pi^0$	( 1.17 $\pm 0.30$ ) $\times 10^{-3}$	
$\Gamma_{30}$	$\chi_{b2}(1P) \omega$	( 6.0 $\pm 2.7$ ) $\times 10^{-4}$	
$\Gamma_{31}$	$\chi_{b2}(1P) (\pi^+ \pi^- \pi^0)_{\text{non-}\omega}$	( 6 $\pm 4$ ) $\times 10^{-4}$	
$\Gamma_{32}$	$\gamma X_b \rightarrow \gamma \Upsilon(1S) \omega$	< 3.8 $\times 10^{-5}$	90%

### Inclusive Decays.

These decay modes are submodes of one or more of the decay modes above.

$\Gamma_{33}$	$\phi$ anything	( 13.8 $^{+2.4}_{-1.7}$ ) %
$\Gamma_{34}$	$D^0$ anything + c.c.	( 108 $\pm 8$ ) %
$\Gamma_{35}$	$D_s$ anything + c.c.	( 46 $\pm 6$ ) %
$\Gamma_{36}$	$J/\psi$ anything	( 2.06 $\pm 0.21$ ) %
$\Gamma_{37}$	$B^0$ anything + c.c.	( 77 $\pm 8$ ) %
$\Gamma_{38}$	$B^+$ anything + c.c.	( 72 $\pm 6$ ) %

### $\Upsilon(10860)$ PARTIAL WIDTHS

$\Gamma(e^+ e^-)$				$\Gamma_{15}$
VALUE (keV)	DOCUMENT ID	TECN	COMMENT	
<b>0.31 <math>\pm 0.07</math> OUR AVERAGE</b>	Error includes scale factor of 1.3.			
0.22 $\pm 0.05$ $\pm 0.07$	BESSION	85	CLEO $e^+ e^- \rightarrow$ hadrons	
0.365 $\pm 0.070$	LOVELOCK	85	CUSB $e^+ e^- \rightarrow$ hadrons	

**$\Upsilon(10860)$  BRANCHING RATIOS**

“OUR EVALUATION” is obtained based on averages of rescaled data listed below. The averages and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>.

 **$\Gamma(B\bar{B}X)/\Gamma_{\text{total}}$   $\Gamma_1/\Gamma$** 

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.762^{+0.027}_{-0.043}</math> OUR EVALUATION</b>				
<b><math>0.71 \pm 0.06</math> OUR AVERAGE</b>				
$0.737 \pm 0.032 \pm 0.051$	1063	<sup>19</sup> DRUTSKOY	10 BELL	$\Upsilon(5S) \rightarrow B^+ X, B^0 X$
$0.589 \pm 0.100 \pm 0.092$		<sup>20</sup> HUANG	07 CLEO	$\Upsilon(5S) \rightarrow \text{hadrons}$

 **$\Gamma(B\bar{B})/\Gamma_{\text{total}}$   $\Gamma_2/\Gamma$** 

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>5.5^{+1.0}_{-0.9} \pm 0.4</math></b>		<sup>21</sup> DRUTSKOY	10 BELL	$\Upsilon(5S) \rightarrow B^+ X, B^0 X$
••• We do not use the following data for averages, fits, limits, etc. •••				
<13.8	90	<sup>20</sup> HUANG	07 CLEO	$\Upsilon(5S) \rightarrow \text{hadrons}$

 **$\Gamma(B\bar{B})/\Gamma(B\bar{B}X)$   $\Gamma_2/\Gamma_1$** 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;0.22</b>	90	AQUINES	06 CLE3	$\Upsilon(5S) \rightarrow \text{hadrons}$

 **$\Gamma(B\bar{B}^* + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_3/\Gamma$** 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.137 \pm 0.016</math> OUR AVERAGE</b>			
$0.137 \pm 0.013 \pm 0.011$	<sup>21</sup> DRUTSKOY	10 BELL	$\Upsilon(5S) \rightarrow B^+ X, B^0 X$
$0.143 \pm 0.053 \pm 0.027$	<sup>20</sup> HUANG	07 CLEO	$\Upsilon(5S) \rightarrow \text{hadrons}$

 **$\Gamma(B\bar{B}^* + \text{c.c.})/\Gamma(B\bar{B}X)$   $\Gamma_3/\Gamma_1$** 

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.24 \pm 0.09 \pm 0.03</math></b>	10	AQUINES	06 CLE3	$\Upsilon(5S) \rightarrow \text{hadrons}$

 **$\Gamma(B^*\bar{B}^*)/\Gamma_{\text{total}}$   $\Gamma_4/\Gamma$** 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.381 \pm 0.034</math> OUR AVERAGE</b>			
$0.375^{+0.021}_{-0.019} \pm 0.030$	<sup>21</sup> DRUTSKOY	10 BELL	$\Upsilon(5S) \rightarrow B^+ X, B^0 X$
$0.436 \pm 0.083 \pm 0.072$	<sup>20</sup> HUANG	07 CLEO	$\Upsilon(5S) \rightarrow \text{hadrons}$

 **$\Gamma(B^*\bar{B}^*)/\Gamma(B\bar{B}X)$   $\Gamma_4/\Gamma_1$** 

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.74 \pm 0.15 \pm 0.08</math></b>	31	AQUINES	06 CLE3	$\Upsilon(5S) \rightarrow \text{hadrons}$

 **$\Gamma(B\bar{B}^{(*)}\pi)/\Gamma_{\text{total}}$   $\Gamma_5/\Gamma$** 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;0.197</b>	90	<sup>20</sup> HUANG	07 CLEO	$\Upsilon(5S) \rightarrow \text{hadrons}$

$\Gamma(B\bar{B}^{(*)}\pi)/\Gamma(B\bar{B}X)$   $\Gamma_5/\Gamma_1$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;0.32</b>	90	AQUINES	06	CLE3 $\Upsilon(5S) \rightarrow$ hadrons

$\Gamma(B\bar{B}\pi)/\Gamma_{total}$   $\Gamma_6/\Gamma$

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.0 \pm 1.2 \pm 0.3</math></b>	0	<sup>21</sup> DRUTSKOY	10	BELL $\Upsilon(5S) \rightarrow B^{+,0}\pi^- X$

$[\Gamma(B^*\bar{B}\pi) + \Gamma(B\bar{B}^*\pi)]/\Gamma_{total}$   $\Gamma_7/\Gamma$

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>7.3^{+2.3}_{-2.1} \pm 0.8</math></b>	38	<sup>21</sup> DRUTSKOY	10	BELL $\Upsilon(5S) \rightarrow B^{+,0}\pi^- X$

$\Gamma(B^*\bar{B}^*\pi)/\Gamma_{total}$   $\Gamma_8/\Gamma$

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>1.0^{+1.4}_{-1.3} \pm 0.4</math></b>	5	<sup>21</sup> DRUTSKOY	10	BELL $\Upsilon(5S) \rightarrow B^{+,0}\pi^- X$

$\Gamma(B\bar{B}\pi\pi)/\Gamma_{total}$   $\Gamma_9/\Gamma$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;0.089</b>	90	<sup>20</sup> HUANG	07	CLEO $\Upsilon(5S) \rightarrow$ hadrons

$\Gamma(B\bar{B}\pi\pi)/\Gamma(B\bar{B}X)$   $\Gamma_9/\Gamma_1$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;0.14</b>	90	AQUINES	06	CLE3 $\Upsilon(5S) \rightarrow$ hadrons

$\Gamma(B_s^{(*)}\bar{B}_s^{(*)})/\Gamma_{total}$   $\Gamma_{10}/\Gamma = (\Gamma_{11} + \Gamma_{12} + \Gamma_{13})/\Gamma$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

**$0.201^{+0.030}_{-0.031}$  OUR EVALUATION**

**$0.189^{+0.027}_{-0.021}$  OUR AVERAGE**

0.172 ± 0.030 <sup>22</sup> ESEN 13 BELL  $\Upsilon(5S) \rightarrow D^0 X, D_s X$

0.21  $^{+0.06}_{-0.03}$  <sup>23</sup> HUANG 07 CLEO  $\Upsilon(5S) \rightarrow D_s X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.180 ± 0.013 ± 0.032 <sup>24</sup> DRUTSKOY 07 BELL  $\Upsilon(5S) \rightarrow D^0 X, D_s X$

0.160 ± 0.026 ± 0.058 <sup>25</sup> ARTUSO 05B CLEO  $e^+e^- \rightarrow D_X X$

$\Gamma(B_s^{(*)}\bar{B}_s^{(*)})/\Gamma(B\bar{B}X)$   $\Gamma_{10}/\Gamma_1$

<u>VALUE</u>	<u>DOCUMENT ID</u>
--------------	--------------------

**$0.264^{+0.052}_{-0.045}$  OUR EVALUATION**

$$\Gamma(B_s^* \bar{B}_s^*) / \Gamma(B_s^{(*)} \bar{B}_s^{(*)}) \qquad \Gamma_{13} / \Gamma_{10} = \Gamma_{13} / (\Gamma_{11} + \Gamma_{12} + \Gamma_{13})$$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>87.8 ± 1.5 OUR AVERAGE</b>				
87.0 ± 1.7	26,27	ESEN	13	BELL $B_s^0 \rightarrow D_s^- \pi^+$
90.5 ± 3.2 ± 0.1	227 27,28	LI	12	BELL $B_s^0 \rightarrow J/\psi \eta^{(\prime)}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
90.1 <sup>+3.8</sup> <sub>-4.0</sub> ± 0.2	29	LOUVOT	09	BELL 10.86 $e^+ e^- \rightarrow B_s^{(*)} \bar{B}_s^{(*)}$
93 <sup>+7</sup> <sub>-9</sub> ± 1	29	DRUTSKOY	07A	BELL Superseded by LOUVOT 09

$$\Gamma(B_s \bar{B}_s) / \Gamma(B_s^{(*)} \bar{B}_s^{(*)}) \qquad \Gamma_{11} / \Gamma_{10} = \Gamma_{11} / (\Gamma_{11} + \Gamma_{12} + \Gamma_{13})$$

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>2.6<sup>+2.6</sup></b> <b>-2.5</b>	LOUVOT	09	BELL 10.86 $e^+ e^- \rightarrow B_s^{(*)} \bar{B}_s^{(*)}$

$$\Gamma(B_s \bar{B}_s) / \Gamma(B_s^* \bar{B}_s^*) \qquad \Gamma_{11} / \Gamma_{13}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.16</b>	90	BONVICINI	06	CLE3 $e^+ e^-$

$$\Gamma(B_s \bar{B}_s^* + \text{c.c.}) / \Gamma(B_s^{(*)} \bar{B}_s^{(*)}) \qquad \Gamma_{12} / \Gamma_{10} = \Gamma_{12} / (\Gamma_{11} + \Gamma_{12} + \Gamma_{13})$$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.7 ± 1.2 OUR AVERAGE</b>				
7.3 ± 1.4	26,27	ESEN	13	BELL $B_s^0 \rightarrow D_s^- \pi^+$
4.9 ± 2.5 ± 0.0	227 27,28	LI	12	BELL $B_s^0 \rightarrow J/\psi \eta^{(\prime)}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
7.3 <sup>+3.3</sup> <sub>-3.0</sub> ± 0.1		LOUVOT	09	BELL 10.86 $e^+ e^- \rightarrow B_s^{(*)} \bar{B}_s^{(*)}$

$$\Gamma(B_s \bar{B}_s^* + \text{c.c.}) / \Gamma(B_s^* \bar{B}_s^*) \qquad \Gamma_{12} / \Gamma_{13}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.16</b>	90	BONVICINI	06	CLE3 $e^+ e^-$

$$\Gamma(\text{no open-bottom}) / \Gamma_{\text{total}} \qquad \Gamma_{14} / \Gamma$$

VALUE	DOCUMENT ID
<b>0.038<sup>+0.051</sup></b> <b>-0.005</b> OUR EVALUATION	

$$\Gamma(K^*(892)^0 \bar{K}^0) / \Gamma_{\text{total}} \qquad \Gamma_{16} / \Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;1.0 × 10<sup>-5</sup></b>	90	SHEN	13A	BELL $e^+ e^- \rightarrow K^*(892)^0 \bar{K}^0$

$$\Gamma(\Upsilon(1S) \pi^+ \pi^-) / \Gamma_{\text{total}} \qquad \Gamma_{17} / \Gamma$$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.3 ± 0.3 ± 0.5</b>	325 30	CHEN	08	BELL 10.87 $e^+ e^- \rightarrow \Upsilon(1S) \pi^+ \pi^-$

$\Gamma(\Upsilon(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$						$\Gamma_{18}/\Gamma$
VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT		
$7.8 \pm 0.6 \pm 1.1$	186	<sup>30</sup> CHEN	08	BELL	$10.87 e^+ e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$	
$\Gamma(\Upsilon(3S)\pi^+\pi^-)/\Gamma_{\text{total}}$						$\Gamma_{19}/\Gamma$
VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT		
$4.8^{+1.8}_{-1.5} \pm 0.7$	10	<sup>30</sup> CHEN	08	BELL	$10.87 e^+ e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$	
$\Gamma(\Upsilon(1S)K^+K^-)/\Gamma_{\text{total}}$						$\Gamma_{20}/\Gamma$
VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT		
$6.1^{+1.6}_{-1.4} \pm 1.0$	20	<sup>30</sup> CHEN	08	BELL	$10.87 e^+ e^- \rightarrow \Upsilon(1S)K^+K^-$	
$\Gamma(h_b(1P)\pi^+\pi^-)/\Gamma(\Upsilon(2S)\pi^+\pi^-)$						$\Gamma_{21}/\Gamma_{18}$
VALUE	DOCUMENT ID	TECN	COMMENT			
$0.45 \pm 0.08^{+0.07}_{-0.12}$	ADACHI	12	BELL	$10.86 e^+ e^- \rightarrow \text{hadrons}$		
$\Gamma(h_b(2P)\pi^+\pi^-)/\Gamma(\Upsilon(2S)\pi^+\pi^-)$						$\Gamma_{22}/\Gamma_{18}$
VALUE	DOCUMENT ID	TECN	COMMENT			
$0.77 \pm 0.08^{+0.22}_{-0.17}$	ADACHI	12	BELL	$10.86 e^+ e^- \rightarrow \text{hadrons}$		
$\Gamma(\chi_{b0}(1P)\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$						$\Gamma_{23}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$< 6.3 \times 10^{-3}$	90	<sup>31</sup> HE	14	BELL	$\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$	
$\Gamma(\chi_{b0}(1P)\omega)/\Gamma_{\text{total}}$						$\Gamma_{24}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$< 3.9 \times 10^{-3}$	90	<sup>31</sup> HE	14	BELL	$\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$	
$\Gamma(\chi_{b0}(1P)(\pi^+\pi^-\pi^0)_{\text{non-}\omega})/\Gamma_{\text{total}}$						$\Gamma_{25}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$< 4.8 \times 10^{-3}$	90	<sup>31</sup> HE	14	BELL	$\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$	
$\Gamma(\chi_{b1}(1P)\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$						$\Gamma_{26}/\Gamma$
VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT		
$1.85 \pm 0.23 \pm 0.23$	80	<sup>31</sup> HE	14	BELL	$\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$	
$\Gamma(\chi_{b1}(1P)\omega)/\Gamma_{\text{total}}$						$\Gamma_{27}/\Gamma$
VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT		
$1.57 \pm 0.22 \pm 0.21$	60	<sup>31</sup> HE	14	BELL	$\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$	
$\Gamma(\chi_{b1}(1P)(\pi^+\pi^-\pi^0)_{\text{non-}\omega})/\Gamma_{\text{total}}$						$\Gamma_{28}/\Gamma$
VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT		
$0.52 \pm 0.15 \pm 0.11$	24	<sup>31</sup> HE	14	BELL	$\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$	

$\Gamma(\chi_{b2}(1P)\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{29}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.17±0.27±0.14</b>	29	31 HE	14 BELL	$\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma \Upsilon(1S)$

$\Gamma(\chi_{b2}(1P)\omega)/\Gamma_{\text{total}}$   $\Gamma_{30}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.60±0.23±0.15</b>	13	31 HE	14 BELL	$\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma \Upsilon(1S)$

$\Gamma(\chi_{b2}(1P)\omega)/\Gamma(\chi_{b1}(1P)\omega)$   $\Gamma_{30}/\Gamma_{27}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.38±0.16±0.09	32 HE	14 BELL	$\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma \Upsilon(1S)$
----------------	-------	---------	---

$\Gamma(\chi_{b2}(1P)(\pi^+\pi^-\pi^0)_{\text{non-}\omega})/\Gamma_{\text{total}}$   $\Gamma_{31}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.61±0.22±0.28</b>	16	31 HE	14 BELL	$\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma \Upsilon(1S)$

$\Gamma(\chi_{b2}(1P)(\pi^+\pi^-\pi^0)_{\text{non-}\omega})/\Gamma(\chi_{b1}(1P)(\pi^+\pi^-\pi^0)_{\text{non-}\omega})$   $\Gamma_{31}/\Gamma_{28}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.20±0.55±0.65	32 HE	14 BELL	$\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma \Upsilon(1S)$
----------------	-------	---------	---

$\Gamma(\gamma X_b \rightarrow \gamma \Upsilon(1S)\omega)/\Gamma_{\text{total}}$   $\Gamma_{32}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;3.8 × 10<sup>-5</sup></b>	90	33 HE	14 BELL	$\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma \Upsilon(1S)$

$\Gamma(\phi \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_{33}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

<b>0.138±0.007<sup>+0.023</sup><sub>-0.015</sub></b>	HUANG	07 CLEO	$\Upsilon(5S) \rightarrow \phi X$
--	-------	---------	-----------------------------------

$\Gamma(D^0 \text{ anything} + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{34}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

<b>1.076±0.040±0.068</b>	DRUTSKOY	07 BELL	$\Upsilon(5S) \rightarrow D^0 X$
--------------------------	----------	---------	----------------------------------

$\Gamma(D_s \text{ anything} + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{35}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

**0.46 ±0.06 OUR AVERAGE**

0.472±0.024±0.072	24 DRUTSKOY	07 BELL	$\Upsilon(5S) \rightarrow D_s X$
-------------------	-------------	---------	----------------------------------

0.44 ±0.09 ±0.04	34 ARTUSO	05B CLE3	$e^+e^- \rightarrow D_s X$
------------------	-----------	----------	----------------------------

$\Gamma(J/\psi \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_{36}/\Gamma$

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

<b>2.060±0.160±0.134</b>	DRUTSKOY	07 BELL	$\Upsilon(5S) \rightarrow J/\psi X$
--------------------------	----------	---------	-------------------------------------

$\Gamma(B^0 \text{ anything} + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{37}/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

<b>0.770<sup>+0.058</sup><sub>-0.056</sub> ±0.061</b>	352	DRUTSKOY	10 BELL	$\Upsilon(5S) \rightarrow B^0 X$
---	-----	----------	---------	----------------------------------



$\Gamma(B^+ \text{ anything} + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{38}/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$0.721^{+0.039}_{-0.038} \pm 0.050$	711	DRUTSKOY 10	BELL	$\Upsilon(5S) \rightarrow B^+ X$
<p><sup>19</sup> Not independent of DRUTSKOY 10 values for <math>\Upsilon(5S) \rightarrow B^{\pm,0}</math> anything.  <sup>20</sup> Using measurements or limits from AQUINES 06.  <sup>21</sup> Assuming isospin conservation.  <sup>22</sup> Supersedes DRUTSKOY 07.  <sup>23</sup> Supersedes ARTUSO 05B. Combining inclusive <math>\phi</math>, <math>D_s</math>, and <math>B</math> measurements. Using <math>B(D_s^+ \rightarrow \phi\pi^+) = 4.4 \pm 0.6\%</math> from PDG 06.  <sup>24</sup> Using <math>B(D_s^+ \rightarrow \phi\pi^+) = (4.4 \pm 0.6)\%</math> from PDG 06.  <sup>25</sup> Uses a model-dependent estimate <math>B(B_s \rightarrow D_s X) = (92 \pm 11)\%</math>.  <sup>26</sup> Supersedes LOUVOT 09.  <sup>27</sup> With <math>N(B_s^{(*)}\bar{B}_s^{(*)}) = (7.11 \pm 1.30) \times 10^6</math>.  <sup>28</sup> The ratios <math>N(B_s^*\bar{B}_s^*) / N(B_s^{(*)}\bar{B}_s^{(*)})</math> and <math>N(B_s^*\bar{B}_s^0) / N(B_s^{(*)}\bar{B}_s^{(*)})</math> are measured with a correlation coefficient of <math>-0.72</math>.  <sup>29</sup> From a measurement of <math>\sigma(e^+e^- \rightarrow B_s^*\bar{B}_s^*) / \sigma(e^+e^- \rightarrow B_s^{(*)}\bar{B}_s^{(*)})</math> at <math>\sqrt{s} = 10.86</math> GeV.  <sup>30</sup> Assuming that the observed events are solely due to the <math>\Upsilon(5S)</math> resonance.  <sup>31</sup> Assuming that all the <math>b\bar{b}</math> events are from <math>\Upsilon(5S)</math> resonance decays and using <math>\sigma(e^+e^- \rightarrow b\bar{b}) = 0.340 \pm 0.016</math> nb from ESEN 13. Correlated with other results from HE 14.  <sup>32</sup> Accounting for correlated systematics.  <sup>33</sup> Assuming that all the <math>b\bar{b}</math> events are from <math>\Upsilon(5S)</math> resonance decays and using <math>\sigma(e^+e^- \rightarrow b\bar{b}) = 0.340 \pm 0.016</math> nb from ESEN 13. Correlated with other results from HE 14. For a state <math>X_b</math> with mass between <math>10.55 \text{ GeV}/c^2</math> and <math>10.65 \text{ GeV}/c^2</math>, the obtained 90% upper limit as a function of <math>m_{X_b}</math> varies from <math>2.6 \times 10^{-5}</math> to <math>3.8 \times 10^{-5}</math>.  <sup>34</sup> ARTUSO 05B reports <math>[\Gamma(\Upsilon(10860) \rightarrow D_s \text{ anything} + \text{c.c.})/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)] = 0.0198 \pm 0.0019 \pm 0.0038</math> which we divide by our best value <math>B(D_s^+ \rightarrow \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}</math>. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p>				

**$\Upsilon(10860)$  REFERENCES**

SANTEL	16	PR D93 011101	D. Santel <i>et al.</i>	(BELLE Collab.)
HE	14	PRL 113 142001	X.H. He <i>et al.</i>	(BELLE Collab.)
ESEN	13	PR D87 031101	S. Esen <i>et al.</i>	(BELLE Collab.)
SHEN	13A	PR D88 052019	C.P. Shen <i>et al.</i>	(BELLE Collab.)
ADACHI	12	PRL 108 032001	I. Adachi <i>et al.</i>	(BELLE Collab.)
LI	12	PRL 108 181808	J. Li <i>et al.</i>	(BELLE Collab.)
CHEN	10	PR D82 091106	K.-F. Chen <i>et al.</i>	(BELLE Collab.)
DRUTSKOY	10	PR D81 112003	A. Drutskoy <i>et al.</i>	(BELLE Collab.)
AUBERT	09E	PRL 102 012001	B. Aubert <i>et al.</i>	(BABAR Collab.)
LOUVOT	09	PRL 102 021801	R. Louvot <i>et al.</i>	(BELLE Collab.)
CHEN	08	PRL 100 112001	K.-F. Chen <i>et al.</i>	(BELLE Collab.)
DRUTSKOY	07	PRL 98 052001	A. Drutskoy <i>et al.</i>	(BELLE Collab.)
DRUTSKOY	07A	PR D76 012002	A. Drutskoy <i>et al.</i>	(BELLE Collab.)
HUANG	07	PR D75 012002	G.S. Huang <i>et al.</i>	(CLEO Collab.)
AQUINES	06	PRL 96 152001	O. Aquines <i>et al.</i>	(CLEO Collab.)
BONVICINI	06	PRL 96 022002	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
PDG	06	JP G33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)
ARTUSO	05B	PRL 95 261801	M. Artuso <i>et al.</i>	(CLEO Collab.)
BESSION	85	PRL 54 381	D. Besson <i>et al.</i>	(CLEO Collab.)
LOVELLOCK	85	PRL 54 377	D.M.J. Lovelock <i>et al.</i>	(CUSB Collab.)