

$\rho(770)$ $I^G(J^{PC}) = 1^+(1^{--})$

A REVIEW GOES HERE – Check our WWW List of Reviews

 $\rho(770)$ MASS

We no longer list S -wave Breit-Wigner fits, or data with high combinatorial background.

NEUTRAL ONLY, e^+e^-

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
775.26 ± 0.25 OUR AVERAGE				
775.02 ± 0.35		1 LEES	12G	BABR $e^+e^- \rightarrow \pi^+\pi^-\gamma$
775.97 $\pm 0.46 \pm 0.70$	900k	2 AKHMETSHIN 07		$e^+e^- \rightarrow \pi^+\pi^-$
774.6 $\pm 0.4 \pm 0.5$	800k	3,4 ACHASOV 06	SND	$e^+e^- \rightarrow \pi^+\pi^-$
775.65 $\pm 0.64 \pm 0.50$	114k	5,6 AKHMETSHIN 04	CMD2	$e^+e^- \rightarrow \pi^+\pi^-$
775.9 $\pm 0.5 \pm 0.5$	1.98M	7 ALOISIO 03	KLOE	$1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
775.8 $\pm 0.9 \pm 2.0$	500k	7 ACHASOV 02	SND	$1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
775.9 ± 1.1		8 BARKOV 85	OLYA	$e^+e^- \rightarrow \pi^+\pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
775.8 $\pm 0.5 \pm 0.3$	1.98M	9 ALOISIO 03	KLOE	$1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
775.9 $\pm 0.6 \pm 0.5$	1.98M	10 ALOISIO 03	KLOE	$1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
775.0 $\pm 0.6 \pm 1.1$	500k	11 ACHASOV 02	SND	$1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
775.1 $\pm 0.7 \pm 5.3$		12 BENAYOUN 98	RVUE	$e^+e^- \rightarrow \pi^+\pi^-, \mu^+\mu^-$
770.5 $\pm 1.9 \pm 5.1$		13 GARDNER 98	RVUE	$0.28-0.92 e^+e^- \rightarrow \pi^+\pi^-$
764.1 ± 0.7		14 O'CONNELL 97	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
757.5 ± 1.5		15 BERNICHA 94	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
768 ± 1		16 GESHKEN...	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$

CHARGED ONLY, τ DECAYS and e^+e^-

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
775.11 ± 0.34 OUR AVERAGE					
774.6 $\pm 0.2 \pm 0.5$	5.4M	17,18 FUJIKAWA 08	BELL	\pm	$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
775.5 ± 0.7		18,19 SCHABEL 05C	ALEP		$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
775.5 $\pm 0.5 \pm 0.4$	1.98M	7 ALOISIO 03	KLOE		$1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
775.1 $\pm 1.1 \pm 0.5$	87k	20,21 ANDERSON 00A	CLE2		$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
774.8 $\pm 0.6 \pm 0.4$	1.98M	10 ALOISIO 03	KLOE	–	$1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
776.3 $\pm 0.6 \pm 0.7$	1.98M	10 ALOISIO 03	KLOE	+	$1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
773.9 $\pm 2.0 \begin{matrix} +0.3 \\ -1.0 \end{matrix}$		22 SANZ-CILLERO 03	RVUE		$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
774.5 $\pm 0.7 \pm 1.5$	500k	7 ACHASOV 02	SND	\pm	$1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
775.1 ± 0.5		23 PICH 01	RVUE		$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$

MIXED CHARGES, OTHER REACTIONS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
763.0±0.3±1.2	600k	24 ABELE	99E CBAR	0±	0.0 $\bar{p}p \rightarrow \pi^+ \pi^- \pi^0$

CHARGED ONLY, HADROPRODUCED

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
766.5±1.1 OUR AVERAGE					
763.7±3.2		ABELE	97	CBAR	$\bar{p}n \rightarrow \pi^- \pi^0 \pi^0$
768 ± 9		AGUILAR-...	91	EHS	400 $p\bar{p}$
767 ± 3	2935	25 CAPRARO	87	SPEC	—
761 ± 5	967	25 CAPRARO	87	SPEC	—
771 ± 4		HUSTON	86	SPEC	+
766 ± 7	6500	26 BYERLY	73	OSPK	—
766.8±1.5	9650	27 PISUT	68	RVUE	—
767 ± 6	900	25 EISNER	67	HBC	—

NEUTRAL ONLY, PHOTOPRODUCED


VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
769.0± 1.0 OUR AVERAGE				
771 ± 2 +2 -1	63.5k	28 ABRAMOWICZ12	ZEUS	$e p \rightarrow e \pi^+ \pi^- p$
770 ± 2 ±1	79k	29 BREITWEG	98B ZEUS	50–100 γp
767.6± 2.7		BARTALUCCI	78 CNTR	$\gamma p \rightarrow e^+ e^- p$
775 ± 5		GLADDING	73 CNTR	2.9–4.7 γp
767 ± 4	1930	BALLAM	72 HBC	2.8 γp
770 ± 4	2430	BALLAM	72 HBC	4.7 γp
765 ± 10		ALVENSLEB...	70 CNTR	$\gamma A, t < 0.01$
767.7± 1.9	140k	BIGGS	70 CNTR	<4.1 $\gamma C \rightarrow \pi^+ \pi^- C$
765 ± 5	4000	ASBURY	67B CNTR	$\gamma + Pb$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
771 ± 2	79k	30 BREITWEG	98B ZEUS	50–100 γp

NEUTRAL ONLY, OTHER REACTIONS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
769.0±0.9 OUR AVERAGE					
Error includes scale factor of 1.4. See the ideogram below.					
765 ± 6		BERTIN	97C OBLX	0.0	$\bar{p}p \rightarrow \pi^+ \pi^- \pi^0$
773 ± 1.6		WEIDENAUER	93 ASTE		$\bar{p}p \rightarrow \pi^+ \pi^- \omega$
762.6±2.6		AGUILAR-...	91 EHS		400 $p\bar{p}$
770 ± 2	31 HEYN	81 RVUE			Pion form factor
768 ± 4	32,33 BOHACIK	80 RVUE	0		
769 ± 3	26 WICKLUND	78 ASPK	0		3,4,6 $\pi^\pm N$
768 ± 1	76000	DEUTSCH...	76 HBC	0	16 $\pi^+ p$
767 ± 4	4100	ENGLER	74 DBC	0	6 $\pi^+ n \rightarrow \pi^+ \pi^- p$
775 ± 4	32000	32 PROTOPOP...	73 HBC	0	7.1 $\pi^+ p, t < 0.4$
764 ± 3	6800	RATCLIFF	72 ASPK	0	15 $\pi^- p, t < 0.3$
774 ± 3	1700	REYNOLDS	69 HBC	0	2.26 $\pi^- p$
769.2±1.5	13300	34 PISUT	68 RVUE	0	1.7–3.2 $\pi^- p, t < 10$

• • • We do not use the following data for averages, fits, limits, etc. • • •

773.5 ± 2.5		35	COLANGELO 01	RVUE	$\pi\pi \rightarrow \pi\pi$
$762.3 \pm 0.5 \pm 1.2$	600k	36	ABELE 99E	CBAR 0	$0.0 \bar{p}p \rightarrow \pi^+ \pi^- \pi^0$
777 ± 2	4943	37	ADAMS 97	E665	$470 \mu p \rightarrow \mu XB$
770 ± 2		38	BOGOLYUB... 97	MIRA	$32 \bar{p}p \rightarrow \pi^+ \pi^- X$
768 ± 8		38	BOGOLYUB... 97	MIRA	$32 pp \rightarrow \pi^+ \pi^- X$
761.1 ± 2.9			DUBNICKA 89	RVUE	π form factor
777.4 ± 2.0		39	CHABAUD 83	ASPK 0	$17 \pi^- p$ polarized
769.5 ± 0.7		32,33	LANG 79	RVUE 0	
770 ± 9		33	ESTABROOKS 74	RVUE 0	$17 \pi^- p \rightarrow \pi^+ \pi^- n$
773.5 ± 1.7	11200	25	JACOBS 72	HBC 0	$2.8 \pi^- p$
775 ± 3	2250		HYAMS 68	OSPK 0	$11.2 \pi^- p$



¹ Using the GOUNARIS 68 parametrization with the complex phase of the $\rho - \omega$ interference and leaving the masses and widths of the $\rho(1450)$, $\rho(1700)$, and $\rho(2150)$ resonances as free parameters of the fit.

² A combined fit of AKHMETSHIN 07, AULCHENKO 06, and AULCHENKO 05.

³ Supersedes ACHASOV 05A.

⁴ A fit of the SND data from 400 to 1000 MeV using parameters of the $\rho(1450)$ and $\rho(1700)$ from a fit of the data of BARKOV 85, BISELLO 89 and ANDERSON 00A.

⁵ Using the GOUNARIS 68 parametrization with the complex phase of the $\rho - \omega$ interference.

⁶ Update of AKHMETSHIN 02.

⁷ Assuming $m_{\rho^+} = m_{\rho^-}$, $\Gamma_{\rho^+} = \Gamma_{\rho^-}$.

⁸ From the GOUNARIS 68 parametrization of the pion form factor.

⁹ Assuming $m_{\rho^+} = m_{\rho^-} = m_{\rho^0}$, $\Gamma_{\rho^+} = \Gamma_{\rho^-} = \Gamma_{\rho^0}$.

¹⁰ Without limitations on masses and widths.

- 11 Assuming $m_{\rho^0} = m_{\rho^\pm}$, $g_{\rho^0 \pi\pi} = g_{\rho^\pm \pi\pi}$.
- 12 Using the data of BARKOV 85 in the hidden local symmetry model.
- 13 From the fit to $e^+ e^- \rightarrow \pi^+ \pi^-$ data from the compilations of HEYN 81 and BARKOV 85, including the GOUNARIS 68 parametrization of the pion form factor.
- 14 A fit of BARKOV 85 data assuming the direct $\omega \pi\pi$ coupling.
- 15 Applying the S-matrix formalism to the BARKOV 85 data.
- 16 Includes BARKOV 85 data. Model-dependent width definition.
- 17 $|F_\pi(0)|^2$ fixed to 1.
- 18 From the GOUNARIS 68 parametrization of the pion form factor.
- 19 The error combines statistical and systematic uncertainties. Supersedes BARATE 97M.
- 20 $\rho(1700)$ mass and width fixed at 1700 MeV and 235 MeV respectively.
- 21 From the GOUNARIS 68 parametrization of the pion form factor. The second error is a model error taking into account different parametrizations of the pion form factor.
- 22 Using the data of BARATE 97M and the effective chiral Lagrangian.
- 23 From a fit of the model-independent parameterization of the pion form factor to the data of BARATE 97M.
- 24 Assuming the equality of ρ^+ and ρ^- masses and widths.
- 25 Mass errors enlarged by us to Γ/\sqrt{N} ; see the note with the $K^*(892)$ mass.
- 26 Phase shift analysis. Systematic errors added corresponding to spread of different fits.
- 27 From fit of 3-parameter relativistic P -wave Breit-Wigner to total mass distribution. Includes BATON 68, MILLER 67B, ALFF-STEINBERGER 66, HAGOPIAN 66, HAGOPIAN 66B, JACOBS 66B, JAMES 66, WEST 66, BLIEDEN 65 and CARMONY 64.
- 28 Using the KUHN 90 parametrization of the pion form factor, neglecting $\rho-\omega$ interference.
- 29 From the parametrization according to SOEDING 66.
- 30 From the parametrization according to ROSS 66.
- 31 HEYN 81 includes all spacelike and timelike F_π values until 1978.
- 32 From pole extrapolation.
- 33 From phase shift analysis of GRAYER 74 data.
- 34 Includes MALAMUD 69, ARMENISE 68, BACON 67, HUWE 67, MILLER 67B, ALFF-STEINBERGER 66, HAGOPIAN 66, HAGOPIAN 66B, JACOBS 66B, JAMES 66, WEST 66, GOLDHABER 64, ABOLINS 63.
- 35 Breit-Wigner mass from a phase-shift analysis of HYAMS 73 and PROTOPOPESCU 73 data.
- 36 Using relativistic Breit-Wigner and taking into account $\rho-\omega$ interference.
- 37 Systematic errors not evaluated.
- 38 Systematic effects not studied.
- 39 From fit of 3-parameter relativistic Breit-Wigner to helicity-zero part of P-wave intensity. CHABAUD 83 includes data of GRAYER 74.

$m_{\rho(770)^0} - m_{\rho(770)^\pm}$


VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-0.7 ± 0.8 OUR AVERAGE	Error includes scale factor of 1.5. See the ideogram below.				
-2.4 ± 0.8		¹ SCHAEL	05C	ALEP	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
$0.4 \pm 0.7 \pm 0.6$	1.98M	² ALOISIO	03	KLOE	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
$1.3 \pm 1.1 \pm 2.0$	500k	² ACHASOV	02	SND	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
$1.6 \pm 0.6 \pm 1.7$	600k	ABELE	99E	CBAR	$0 \pm 0.0 \bar{p}p \rightarrow \pi^+ \pi^- \pi^0$
-4 ± 4	3000	³ REYNOLDS	69	HBC	$-0 \quad 2.26 \pi^- p$
-5 ± 5	3600	³ FOSTER	68	HBC	$\pm 0 \quad 0.0 \bar{p}p$
2.4 ± 2.1	22950	⁴ PISUT	68	RVUE	$\pi N \rightarrow \rho N$

¹ From the combined fit of the τ^- data from ANDERSON 00A and SCHAEEL 05C and $e^+ e^-$ data from the compilation of BARKOV 85, AKHMETSHIN 04, and ALOISIO 05. Supersedes BARATE 97M.

² Assuming $m_{\rho^+} = m_{\rho^-}$, $\Gamma_{\rho^+} = \Gamma_{\rho^-}$.

³ From quoted masses of charged and neutral modes.

⁴ Includes MALAMUD 69, ARMENISE 68, BATON 68, BACON 67, HUWE 67, MILLER 67B, ALFF-STEINBERGER 66, HAGOPIAN 66, HAGOPIAN 66B, JACOBS 66B, JAMES 66, WEST 66, BLIEDEN 65, CARMONY 64, GOLDHABER 64, ABOLINS 63.



$m_{\rho(770)^+} - m_{\rho(770)^-}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.5±0.8±0.7 1.98M 1 ALOISIO 03 KLOE 1.02 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$

1 Without limitations on masses and widths.

$\rho(770)$ RANGE PARAMETER

The range parameter R enters an energy-dependent correction to the width, of the form $(1 + q_r^2 R^2)/(1 + q^2 R^2)$, where q is the momentum of one of the pions in the $\pi\pi$ rest system. At resonance, $q = q_r$.

VALUE (GeV^{-1})	DOCUMENT ID	TECN	CHG	COMMENT
-----------------------------	-------------	------	-----	---------

5.3^{+0.9}_{-0.7}	CHABAUD	83	ASPK	0 17 $\pi^- p$ polarized
--	---------	----	------	-------------------------------


$\rho(770)$ WIDTH

We no longer list S -wave Breit-Wigner fits, or data with high combinatorial background.

NEUTRAL ONLY, $e^+ e^-$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
147.8 ± 0.9 OUR AVERAGE		Error includes scale factor of 2.0. See the ideogram below.			
149.59 ± 0.67		¹ LEES 12G BABR			$e^+ e^- \rightarrow \pi^+ \pi^- \gamma$
145.98 ± 0.75 ± 0.50	900k	² AKHMETSHIN 07			$e^+ e^- \rightarrow \pi^+ \pi^-$
146.1 ± 0.8 ± 1.5	800k	^{3,4} ACHASOV 06	SND		$e^+ e^- \rightarrow \pi^+ \pi^-$
143.85 ± 1.33 ± 0.80	114k	^{5,6} AKHMETSHIN 04	CMD2		$e^+ e^- \rightarrow \pi^+ \pi^-$
147.3 ± 1.5 ± 0.7	1.98M	⁷ ALOISIO 03	KLOE		$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
151.1 ± 2.6 ± 3.0	500k	⁷ ACHASOV 02	SND	0	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
150.5 ± 3.0		⁸ BARKOV 85	OLYA	0	$e^+ e^- \rightarrow \pi^+ \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
143.9 ± 1.3 ± 1.1	1.98M	⁹ ALOISIO 03	KLOE		$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
147.4 ± 1.5 ± 0.7	1.98M	¹⁰ ALOISIO 03	KLOE		$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
149.8 ± 2.2 ± 2.0	500k	¹¹ ACHASOV 02	SND		$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
147.9 ± 1.5 ± 7.5		¹² BENAYOUN 98	RVUE		$e^+ e^- \rightarrow \pi^+ \pi^- , \mu^+ \mu^-$
153.5 ± 1.3 ± 4.6		¹³ GARDNER 98	RVUE		$0.28-0.92 e^+ e^- \rightarrow \pi^+ \pi^-$
145.0 ± 1.7		¹⁴ O'CONNELL 97	RVUE		$e^+ e^- \rightarrow \pi^+ \pi^-$
142.5 ± 3.5		¹⁵ BERNICHA 94	RVUE		$e^+ e^- \rightarrow \pi^+ \pi^-$
138 ± 1		¹⁶ GESHKEN... 89	RVUE		$e^+ e^- \rightarrow \pi^+ \pi^-$

WEIGHTED AVERAGE
147.8±0.9 (Error scaled by 2.0)



CHARGED ONLY, τ DECAYS and $e^+ e^-$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
149.1±0.8 OUR FIT					
149.1±0.8 OUR AVERAGE					
148.1±0.4±1.7	5.4M	17,18 FUJIKAWA	08 BELL	±	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
149.0±1.2		18,19 SCHABEL	05C ALEP		$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
149.9±2.3±2.0	500k	7 ACHASOV	02 SND	±	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
150.4±1.4±1.4	87k	20,21 ANDERSON	00A CLE2		$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
143.7±1.3±1.2	1.98M	7 ALOISIO	03 KLOE	±	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
142.9±1.3±1.4	1.98M	10 ALOISIO	03 KLOE	-	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
144.7±1.4±1.2	1.98M	10 ALOISIO	03 KLOE	+	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
150.2±2.0 ^{+0.7} _{-1.6}		22 SANZ-CILLERO03	RVUE		$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
150.9±2.2±2.0	500k	11 ACHASOV	02 SND		$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$

MIXED CHARGES, OTHER REACTIONS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
149.5±1.3					
	600k	23 ABELE	99E CBAR	0±	$0.0 \bar{p}p \rightarrow \pi^+ \pi^- \pi^0$

CHARGED ONLY, HADROPRODUCED

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
150.2± 2.4 OUR FIT					
150.2± 2.4 OUR AVERAGE					
152.8± 4.3		ABELE	97 CBAR		$\bar{p}n \rightarrow \pi^- \pi^0 \pi^0$
155 ±11	2935	24 CAPRARO	87 SPEC	-	$200 \pi^- Cu \rightarrow \pi^- \pi^0 Cu$
154 ±20	967	24 CAPRARO	87 SPEC	-	$200 \pi^- Pb \rightarrow \pi^- \pi^0 Pb$
150 ± 5		HUSTON	86 SPEC	+	$202 \pi^+ A \rightarrow \pi^+ \pi^0 A$
146 ±12	6500	25 BYERLY	73 OSPK	-	$5 \pi^- p$
148.2± 4.1	9650	26 PISUT	68 RVUE	-	$1.7\text{--}3.2 \pi^- p, t < 10$
146 ±13	900	EISNER	67 HBC	-	$4.2 \pi^- p, t < 10$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
137.0± 0.4		27 ABLIKIM	17 BES3		$J/\psi \rightarrow \gamma 3\pi$

NEUTRAL ONLY, PHOTOPRODUCED

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
151.7± 2.6 OUR AVERAGE				
155 ± 5 ± 2	63.5k	28 ABRAMOWICZ12	ZEUS	$ep \rightarrow e\pi^+ \pi^- p$
146 ± 3 ± 13	79k	29 BREITWEG	98B ZEUS	$50\text{--}100 \gamma p$
150.9± 3.0		BARTALUCCI	78 CNTR	$\gamma p \rightarrow e^+ e^- p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

138	\pm 3	79k	30 BREITWEG	98B ZEUS	50–100 γp
147	\pm 11		GLADDING	73 CNTR	2.9–4.7 γp
155	\pm 12	2430	BALLAM	72 HBC	4.7 γp
145	\pm 13	1930	BALLAM	72 HBC	2.8 γp
140	\pm 5		ALVENSLEB...	70 CNTR	$\gamma A, t < 0.01$
146.1 \pm 2.9		140k	BIGGS	70 CNTR	$< 4.1 \gamma C \rightarrow \pi^+ \pi^- C$
160	\pm 10		LANZEROTTI	68 CNTR	γp
130	\pm 5	4000	ASBURY	67B CNTR	$\gamma + Pb$

NEUTRAL ONLY, OTHER REACTIONS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
150.9 \pm 1.7 OUR AVERAGE		Error includes scale factor of 1.1.			
122 \pm 20		BERTIN	97C OBLX	0	$0.0 \bar{p}p \rightarrow \pi^+ \pi^- \pi^0$
145.7 \pm 5.3		WEIDENAUER	93 ASTE		$\bar{p}p \rightarrow \pi^+ \pi^- \omega$
144.9 \pm 3.7		DUBNICKA	89 RVUE		π form factor
148 \pm 6	31,32 BOHACIK		80 RVUE	0	
152 \pm 9	25 WICKLUND		78 ASPK	0	$3,4,6 \pi^\pm p N$
154 \pm 2	76000 DEUTSCH...		76 HBC	0	$16 \pi^+ p$
157 \pm 8	6800 RATCLIFF		72 ASPK	0	$15 \pi^- p, t < 0.3$
143 \pm 8	1700 REYNOLDS		69 HBC	0	$2.26 \pi^- p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

147.0 \pm 2.5	600k	33 ABELE	99E CBAR	0	$0.0 \bar{p}p \rightarrow \pi^+ \pi^- \pi^0$
146 \pm 3	4943	34 ADAMS	97 E665		$470 \mu p \rightarrow \mu XB$
160.0 \pm 4.1		35 CHABAUD	83 ASPK	0	$17 \pi^- p$ polarized
155 \pm 1	36 HEYN		81 RVUE	0	π form factor
148.0 \pm 1.3	31,32 LANG		79 RVUE	0	
146 \pm 14	4100 ENGLER		74 DBC	0	$6 \pi^+ n \rightarrow \pi^+ \pi^- p$
143 \pm 13	32 ESTABROOKS	74 RVUE	0		$17 \pi^- p \rightarrow \pi^+ \pi^- n$
160 \pm 10	32000 31 PROTOPOP...	73 HBC	0		$7.1 \pi^+ p, t < 0.4$
145 \pm 12	2250 HYAMS		68 OSPK	0	$11.2 \pi^- p$
163 \pm 15	13300 PISUT		68 RVUE	0	$1.7\text{--}3.2 \pi^- p, t < 10$

¹ Using the GOUNARIS 68 parametrization with the complex phase of the $\rho-\omega$ interference and leaving the masses and widths of the $\rho(1450)$, $\rho(1700)$, and $\rho(2150)$ resonances as free parameters of the fit.

² A combined fit of AKHMETSHIN 07, AULCHENKO 06, and AULCHENKO 05.

³ Supersedes ACHASOV 05A.

⁴ A fit of the SND data from 400 to 1000 MeV using parameters of the $\rho(1450)$ and $\rho(1700)$ from a fit of the data of BARKOV 85, BISELLO 89 and ANDERSON 00A.

⁵ Using the GOUNARIS 68 parametrization with the complex phase of the $\rho-\omega$ interference.

⁶ From a fit in the energy range 0.61 to 0.96 GeV. Update of AKHMETSHIN 02.

⁷ Assuming $m_{\rho^+} = m_{\rho^-}$, $\Gamma_{\rho^+} = \Gamma_{\rho^-}$.

⁸ From the GOUNARIS 68 parametrization of the pion form factor.

⁹ Assuming $m_{\rho^+} = m_{\rho^-} = m_{\rho^0}$, $\Gamma_{\rho^+} = \Gamma_{\rho^-} = \Gamma_{\rho^0}$.

¹⁰ Without limitations on masses and widths.

¹¹ Assuming $m_{\rho^0} = m_{\rho^\pm}$, $g_{\rho^0 \pi\pi} = g_{\rho^\pm \pi\pi}$.

¹² Using the data of BARKOV 85 in the hidden local symmetry model.

¹³ From the fit to $e^+ e^- \rightarrow \pi^+ \pi^-$ data from the compilations of HEYN 81 and BARKOV 85, including the GOUNARIS 68 parametrization of the pion form factor.

- ¹⁴ A fit of BARKOV 85 data assuming the direct $\omega\pi\pi$ coupling.
¹⁵ Applying the S-matrix formalism to the BARKOV 85 data.
¹⁶ Includes BARKOV 85 data. Model-dependent width definition.
¹⁷ $|F_\pi(0)|^2$ fixed to 1.
¹⁸ From the GOUNARIS 68 parametrization of the pion form factor.
¹⁹ The error combines statistical and systematic uncertainties. Supersedes BARATE 97M.
²⁰ $\rho(1700)$ mass and width fixed at 1700 MeV and 235 MeV respectively.
²¹ From the GOUNARIS 68 parametrization of the pion form factor. The second error is a model error taking into account different parametrizations of the pion form factor.
²² Using the data of BARATE 97M and the effective chiral Lagrangian.
²³ Assuming the equality of ρ^+ and ρ^- masses and widths.
²⁴ Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.
²⁵ Phase shift analysis. Systematic errors added corresponding to spread of different fits.
²⁶ From fit of 3-parameter relativistic P -wave Breit-Wigner to total mass distribution. Includes BATON 68, MILLER 67B, ALFF-STEINBERGER 66, HAGOPIAN 66, HAGOPIAN 66B, JACOBS 66B, JAMES 66, WEST 66, BLIEDEN 65 and CARMONY 64.
²⁷ S-matrix pole at a fixed ρ meson mass of 775.49 MeV.
²⁸ Using the KUHN 90 parametrization of the pion form factor, neglecting $\rho-\omega$ interference.
²⁹ From the parametrization according to SOEDING 66.
³⁰ From the parametrization according to ROSS 66.
³¹ From pole extrapolation.
³² From phase shift analysis of GRAYER 74 data.
³³ Using relativistic Breit-Wigner and taking into account $\rho-\omega$ interference.
³⁴ Systematic errors not evaluated.
³⁵ From fit of 3-parameter relativistic Breit-Wigner to helicity-zero part of P -wave intensity. CHABAUD 83 includes data of GRAYER 74.
³⁶ HEYN 81 includes all spacelike and timelike F_π values until 1978.
³⁷ Includes MALAMUD 69, ARMENISE 68, BACON 67, HUWE 67, MILLER 67B, ALFF-STEINBERGER 66, HAGOPIAN 66, HAGOPIAN 66B, JACOBS 66B, JAMES 66, WEST 66, GOLDHABER 64, ABOLINS 63.

 $\Gamma_{\rho(770)^0} - \Gamma_{\rho(770)^\pm}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.3±1.3 OUR AVERAGE		Error includes scale factor of 1.4.		
-0.2±1.0		¹ SCHAEL	05C ALEP	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
3.6±1.8±1.7	1.98M	² ALOISIO	03 KLOE	$1.02 \frac{e^+ e^-}{\pi^+ \pi^- \pi^0} \rightarrow$

 $\Gamma_{\rho(770)^+} - \Gamma_{\rho(770)^-}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.8±2.0±0.5	1.98M	³ ALOISIO	03 KLOE	$1.02 \frac{e^+ e^-}{\pi^+ \pi^- \pi^0} \rightarrow$

¹ From the combined fit of the τ^- data from ANDERSON 00A and SCHAEL 05C and $e^+ e^-$ data from the compilation of BARKOV 85, AKHMETSHIN 04, and ALOISIO 05. Supersedes BARATE 97M.

² Assuming $m_{\rho^+} = m_{\rho^-}$, $\Gamma_{\rho^+} = \Gamma_{\rho^-}$.

³ Without limitations on masses and widths.

$\rho(770)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
$\Gamma_1 \pi\pi$	~ 100	%
$\rho(770)^{\pm}$ decays		
$\Gamma_2 \pi^{\pm}\pi^0$	~ 100	%
$\Gamma_3 \pi^{\pm}\gamma$	(4.5 ± 0.5)	$\times 10^{-4}$ S=2.2
$\Gamma_4 \pi^{\pm}\eta$	< 6	$\times 10^{-3}$ CL=84%
$\Gamma_5 \pi^{\pm}\pi^+\pi^-\pi^0$	< 2.0	$\times 10^{-3}$ CL=84%
$\rho(770)^0$ decays		
$\Gamma_6 \pi^+\pi^-$	~ 100	%
$\Gamma_7 \pi^+\pi^-\gamma$	(9.9 ± 1.6)	$\times 10^{-3}$
$\Gamma_8 \pi^0\gamma$	(4.7 ± 0.6)	$\times 10^{-4}$ S=1.4
$\Gamma_9 \eta\gamma$	(3.00 ± 0.21)	$\times 10^{-4}$
$\Gamma_{10} \pi^0\pi^0\gamma$	(4.5 ± 0.8)	$\times 10^{-5}$
$\Gamma_{11} \mu^+\mu^-$	[a] (4.55 ± 0.28)	$\times 10^{-5}$
$\Gamma_{12} e^+e^-$	[a] (4.72 ± 0.05)	$\times 10^{-5}$
$\Gamma_{13} \pi^+\pi^-\pi^0$	$(1.01^{+0.54}_{-0.36} \pm 0.34)$	$\times 10^{-4}$
$\Gamma_{14} \pi^+\pi^-\pi^+\pi^-$	(1.8 ± 0.9)	$\times 10^{-5}$
$\Gamma_{15} \pi^+\pi^-\pi^0\pi^0$	(1.6 ± 0.8)	$\times 10^{-5}$
$\Gamma_{16} \pi^0e^+e^-$	< 1.2	$\times 10^{-5}$ CL=90%
$\Gamma_{17} \eta e^+e^-$		

[a] The $\omega\rho$ interference is then due to $\omega\rho$ mixing only, and is expected to be small. If $e\mu$ universality holds, $\Gamma(\rho^0 \rightarrow \mu^+\mu^-) = \Gamma(\rho^0 \rightarrow e^+e^-) \times 0.99785$.

CONSTRAINED FIT INFORMATION

An overall fit to the total width and a partial width uses 10 measurements and one constraint to determine 3 parameters. The overall fit has a $\chi^2 = 10.7$ for 8 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i/\Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

$$\begin{array}{c|cc} x_3 & -100 & \\ \hline \Gamma & 15 & -15 \\ \hline x_2 & x_3 \end{array}$$

Mode	Rate (MeV)	Scale factor
$\Gamma_2 \pi^\pm \pi^0$	150.2 \pm 2.4	
$\Gamma_3 \pi^\pm \gamma$	0.068 \pm 0.007	2.3

CONSTRAINED FIT INFORMATION

An overall fit to the total width, a partial width, and 7 branching ratios uses 22 measurements and one constraint to determine 9 parameters. The overall fit has a $\chi^2 = 9.5$ for 14 degrees of freedom.


The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_7	-100							
x_8	-4	0						
x_9	-1	0	1					
x_{10}	-1	0	0	0				
x_{11}	2	-3	0	0	0			
x_{12}	0	0	-8	-9	0	0		
x_{14}	-1	0	0	0	0	0	0	
Γ	0	0	4	5	0	0	-54	
	x_6	x_7	x_8	x_9	x_{10}	x_{11}	x_{12}	x_{14}

Mode	Rate (MeV)	Scale factor
$\Gamma_6 \pi^+ \pi^-$	147.5 \pm 0.9	
$\Gamma_7 \pi^+ \pi^- \gamma$	1.48 \pm 0.24	
$\Gamma_8 \pi^0 \gamma$	0.070 \pm 0.009	1.4
$\Gamma_9 \eta \gamma$	0.0447 \pm 0.0032	
$\Gamma_{10} \pi^0 \pi^0 \gamma$	0.0066 \pm 0.0012	
$\Gamma_{11} \mu^+ \mu^-$	[a] 0.0068 \pm 0.0004	
$\Gamma_{12} e^+ e^-$	[a] 0.00704 \pm 0.00006	
$\Gamma_{14} \pi^+ \pi^- \pi^+ \pi^-$	0.0027 \pm 0.0014	

$\rho(770)$ PARTIAL WIDTHS

$\Gamma(\pi^\pm \gamma)$		Γ_3
VALUE (keV)	DOCUMENT ID	TECN CHG COMMENT
68 \pm 7 OUR FIT	Error includes scale factor of 2.3.	
68 \pm 7 OUR AVERAGE	Error includes scale factor of 2.2. See the ideogram below.	
81 \pm 4 \pm 4	CAPRARO	87 SPEC - $200 \pi^- A \rightarrow \pi^- \pi^0 A$
59.8 \pm 4.0	HUSTON	86 SPEC + $202 \pi^+ A \rightarrow \pi^+ \pi^0 A$
71 \pm 7	JENSEN	83 SPEC - $156-260 \pi^- A \rightarrow \pi^- \pi^0 A$



$\Gamma(e^+e^-)$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{12}
7.04 ±0.06 OUR FIT					
7.04 ±0.06 OUR AVERAGE					
7.048±0.057±0.050	900k	¹ AKHMETSHIN 07		$e^+e^- \rightarrow \pi^+\pi^-$	
7.06 ±0.11 ±0.05	114k	2,3 AKHMETSHIN 04	CMD2	$e^+e^- \rightarrow \pi^+\pi^-$	
6.77 ±0.10 ±0.30		BARKOV 85	OLYA	$e^+e^- \rightarrow \pi^+\pi^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
7.12 ±0.02 ±0.11	800k	⁴ ACHASOV 06	SND	$e^+e^- \rightarrow \pi^+\pi^-$	
6.3 ±0.1		⁵ BENAYOUN 98	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$, $\mu^+\mu^-$	

$\Gamma(\pi^0\gamma)$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_8
• • • We do not use the following data for averages, fits, limits, etc. • • •					
77±17±11	36500	⁶ ACHASOV 03	SND	$0.60\text{--}0.97 e^+e^- \rightarrow \pi^0\gamma$	
121±31		DOLINSKY 89	ND	$e^+e^- \rightarrow \pi^0\gamma$	

$\Gamma(\eta\gamma)$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT	Γ_9
• • • We do not use the following data for averages, fits, limits, etc. • • •				
62±17	⁷ DOLINSKY 89	ND	$e^+e^- \rightarrow \eta\gamma$	

$\Gamma(\pi^+\pi^-\pi^+\pi^-)$ Γ_{14}

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$2.8 \pm 1.4 \pm 0.5$	153	AKHMETSHIN 00	CMD2	$0.6-0.97 e^+ e^- \rightarrow \pi^+ \pi^- \pi^+ \pi^-$
¹ A combined fit of AKHMETSHIN 07, AULCHENKO 06, and AULCHENKO 05.				
² Using the GOUNARIS 68 parametrization with the complex phase of the ρ - ω interference.				
³ From a fit in the energy range 0.61 to 0.96 GeV. Update of AKHMETSHIN 02.				
⁴ Supersedes ACHASOV 05A.				
⁵ Using the data of BARKOV 85 in the hidden local symmetry model.				
⁶ Using $\Gamma_{\text{total}} = 147.9 \pm 1.3$ MeV and $B(\rho \rightarrow \pi^0 \gamma)$ from ACHASOV 03.				
⁷ Solution corresponding to constructive ω - ρ interference.				

$$\rho(770) \Gamma(e^+ e^-) \Gamma(i) / \Gamma^2(\text{total})$$

 $\Gamma(e^+ e^-) / \Gamma_{\text{total}} \times \Gamma(\pi^+ \pi^-) / \Gamma_{\text{total}}$ $\Gamma_{12}/\Gamma \times \Gamma_6/\Gamma$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$4.876 \pm 0.023 \pm 0.064$	800k	1,2 ACHASOV	06	$e^+ e^- \rightarrow \pi^+ \pi^-$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
4.72 ± 0.02	³ BENAYOUN	10	RVUE	0.4–1.05 $e^+ e^-$

¹ Supersedes ACHASOV 05A.² A fit of the SND data from 400 to 1000 MeV using parameters of the $\rho(1450)$ and $\rho(1700)$ from a fit of the data of BARKOV 85, BISELLO 89 and ANDERSON 00A.³ A simultaneous fit of $e^+ e^- \rightarrow \pi^+ \pi^-, \pi^+ \pi^- \pi^0, \pi^0 \gamma, \eta \gamma$ data. $\Gamma(e^+ e^-) / \Gamma_{\text{total}} \times \Gamma(\eta \gamma) / \Gamma_{\text{total}}$ $\Gamma_{12}/\Gamma \times \Gamma_9/\Gamma$

VALUE (units 10^{-8})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.42 ± 0.10 OUR FIT**1.45 ± 0.12 OUR AVERAGE**

$1.32 \pm 0.14 \pm 0.08$	33k	¹ ACHASOV 07B	SND	$0.6-1.38 e^+ e^- \rightarrow \eta \gamma$
$1.50 \pm 0.65 \pm 0.09$	17.4k	² AKHMETSHIN 05	CMD2	$0.60-1.38 e^+ e^- \rightarrow \eta \gamma$
$1.61 \pm 0.20 \pm 0.11$	23k	^{3,4} AKHMETSHIN 01B	CMD2	$e^+ e^- \rightarrow \eta \gamma$
1.85 ± 0.49		⁵ DOLINSKY 89	ND	$e^+ e^- \rightarrow \eta \gamma$

 $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$

1.05 ± 0.02	⁶ BENAYOUN	10	RVUE	0.4–1.05 $e^+ e^-$
-----------------	-----------------------	----	------	--------------------

¹ From a combined fit of $\sigma(e^+ e^- \rightarrow \eta \gamma)$ with $\eta \rightarrow 3\pi^0$ and $\eta \rightarrow \pi^+ \pi^- \pi^0$, and fixing $B(\eta \rightarrow 3\pi^0) / B(\eta \rightarrow \pi^+ \pi^- \pi^0) = 1.44 \pm 0.04$. Recalculated by us from the cross section at the peak. Supersedes ACHASOV 00D and ACHASOV 06A.² From the $\eta \rightarrow 2\gamma$ decay and using $B(\eta \rightarrow \gamma\gamma) = 39.43 \pm 0.26\%$.³ From the $\eta \rightarrow 3\pi^0$ decay and using $B(\eta \rightarrow 3\pi^0) = (32.24 \pm 0.29) \times 10^{-2}$.⁴ The combined fit from 600 to 1380 MeV taking into account $\rho(770)$, $\omega(782)$, $\phi(1020)$, and $\rho(1450)$ (mass and width fixed at 1450 MeV and 310 MeV respectively).⁵ Recalculated by us from the cross section in the peak.⁶ A simultaneous fit of $e^+ e^- \rightarrow \pi^+ \pi^-, \pi^+ \pi^- \pi^0, \pi^0 \gamma, \eta \gamma$ data.

$\Gamma(e^+ e^-)/\Gamma_{\text{total}} \times \Gamma(\pi^0 \gamma)/\Gamma_{\text{total}}$

$\Gamma_{12}/\Gamma \times \Gamma_8/\Gamma$

VALUE (units 10^{-8})	EVTS	DOCUMENT ID	TECN	COMMENT
2.22 ± 0.29 OUR FIT		Error includes scale factor of 1.4.		
2.22 ± 0.26 OUR AVERAGE		Error includes scale factor of 1.3. See the ideogram below.		
1.98 ± 0.22 ± 0.10	1 ACHASOV	16A SND	0.60-1.38 $e^+ e^- \rightarrow \pi^0 \gamma$	
2.90 ± 0.60 ± 0.55	18k AKHMETSHIN	05 CMD2	0.60-1.38 $e^+ e^- \rightarrow \pi^0 \gamma$	
2.37 ± 0.53 ± 0.33	36k ACHASOV	03 SND	0.60-0.97 $e^+ e^- \rightarrow \pi^0 \gamma$	
3.61 ± 0.74 ± 0.49	10k DOLINSKY	89 ND	$e^+ e^- \rightarrow \pi^0 \gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.875 ± 0.026	4 BENAYOUN	10 RVUE	0.4-1.05 $e^+ e^-$	


¹ From the VMD model with the rho(770), omega(782), phi(1020) resonances, and an additional resonance describing the total contribution of the rho(1450) and omega(1420) states. Supersedes ACHASOV 03.

² Using $\sigma_{\phi \rightarrow \pi^0 \gamma}$ from ACHASOV 00 and $m_\rho = 775.97$ MeV in the model with the energy-independent phase of rho-omega interference equal to $(-10.2 \pm 7.0)^\circ$.

³ Recalculated by us from the cross section in the peak.

⁴ A simultaneous fit of $e^+ e^- \rightarrow \pi^+ \pi^-, \pi^+ \pi^- \pi^0, \pi^0 \gamma, \eta \gamma$ data.

WEIGHTED AVERAGE
2.22±0.26 (Error scaled by 1.3)



$\Gamma(e^+ e^-)/\Gamma_{\text{total}} \times \Gamma(\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$

$\Gamma_{12}/\Gamma \times \Gamma_{13}/\Gamma$

VALUE (units 10^{-9})	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.903 ± 0.076	1 BENAYOUN	10 RVUE	0.4-1.05 $e^+ e^-$	
4.58 ± 2.46 ± 1.56	1.2M ACHASOV	03D RVUE	0.44-2.00 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$	

¹A simultaneous fit of $e^+ e^- \rightarrow \pi^+ \pi^-$, $\pi^+ \pi^- \pi^0$, $\pi^0 \gamma$, $\eta \gamma$ data.²Statistical significance is less than 3σ .

$\rho(770)$ BRANCHING RATIOS

$\Gamma(\pi^\pm \eta)/\Gamma(\pi\pi)$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	Γ_4/Γ_1
<60	84	FERBEL	66	HBC	\pm	$\pi^\pm p$ above 2.5

$\Gamma(\pi^\pm \pi^+ \pi^- \pi^0)/\Gamma(\pi\pi)$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	Γ_5/Γ_1
<20	84	FERBEL	66	HBC	\pm	$\pi^\pm p$ above 2.5

• • • We do not use the following data for averages, fits, limits, etc. • • •

35 \pm 40JAMES 66 HBC + 2.1 $\pi^+ p$

$\Gamma(\mu^+ \mu^-)/\Gamma(\pi^+ \pi^-)$

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{11}/Γ_6
---	--------------------	-------------	----------------	------------------------

4.60 \pm 0.28 OUR FIT

4.6 \pm 0.2 \pm 0.2ANTIPOV 89 SIGM $\pi^- \text{Cu} \rightarrow \mu^+ \mu^- \pi^- \text{Cu}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

8.2 $^{+1.6}_{-3.6}$ ¹ ROTHWELL 69 CNTR Photoproduction5.6 \pm 1.5² WEHMANN 69 OSPK 12 π^- C, Fe9.7 $^{+3.1}_{-3.3}$ ³ HYAMS 67 OSPK 11 π^- Li, H

$\Gamma(e^+ e^-)/\Gamma(\pi\pi)$

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{12}/Γ_1
---	--------------------	-------------	----------------	------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.40 \pm 0.05⁴ BENAKSAS 72 OSPK $e^+ e^- \rightarrow \pi^+ \pi^-$

$\Gamma(\eta\gamma)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	Γ_9/Γ
---	-------------	--------------------	-------------	------------	----------------	-------------------

3.00 \pm 0.21 OUR FIT

2.90 \pm 0.32 OUR AVERAGE

2.79 \pm 0.34 \pm 0.03 33k ⁵ ACHASOV 07B SND 0.6–1.38 $e^+ e^- \rightarrow \eta\gamma$ 3.6 \pm 0.9 ⁶ ANDREWS 77 CNTR 0 6.7–10 γ Cu

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.21 \pm 1.39 \pm 0.20 17.4k ^{7,8} AKHMETSHIN 05 CMD2 0.60–1.38 $e^+ e^- \rightarrow \eta\gamma$ 3.39 \pm 0.42 \pm 0.23 ^{6,9,10} AKHMETSHIN 01B CMD2 $e^+ e^- \rightarrow \eta\gamma$ 1.9 $^{+0.6}_{-0.8}$ ¹¹ BENAYOUN 96 RVUE 0.54–1.04 $e^+ e^- \rightarrow \eta\gamma$ 4.0 \pm 1.1 ^{6,8} DOLINSKY 89 ND $e^+ e^- \rightarrow \eta\gamma$

$\Gamma(\pi^+\pi^-\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{14}/Γ

<u>VALUE</u> (units 10^{-5})	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.8 ± 0.9 OUR FIT					
$1.8 \pm 0.9 \pm 0.3$	153		AKHMETSHIN 00	CMD2	$0.6\text{--}0.97 e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<20	90	KURDADZE	88	OLYA	$e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$
-----	----	----------	----	------	---

 $\Gamma(\pi^+\pi^-\pi^+\pi^-)/\Gamma(\pi\pi)$ Γ_{14}/Γ_1

<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<15	90	ERBE	69	HBC	$0 2.5\text{--}5.8 \gamma p$
<20		CHUNG	68	HBC	$0 3.2, 4.2 \pi^- p$
<20	90	HUSON	68	HLBC	$0 16.0 \pi^- p$
<80		JAMES	66	HBC	$0 2.1 \pi^+ p$

 $\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{13}/Γ

<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$1.01^{+0.54}_{-0.36} \pm 0.34$	1.2M	¹² ACHASOV	03D	RVUE	$0.44\text{--}2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
<1.2	90	VASSERMAN	88B	ND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$

 $\Gamma(\pi^+\pi^-\pi^0)/\Gamma(\pi\pi)$ Γ_{13}/Γ_1

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •					
~ 0.01		BRAMON	86	RVUE	$0 J/\psi \rightarrow \omega\pi^0$
<0.01	84	¹³ ABRAMS	71	HBC	$0 3.7 \pi^+ p$

 $\Gamma(\pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_{15}/Γ

<u>VALUE</u> (units 10^{-5})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.60 \pm 0.74 \pm 0.18$		¹⁴ ACHASOV	09A	$SND e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 4	90	AULCHENKO	87C	ND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$
<20	90	KURDADZE	86	OLYA	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$

 $\Gamma(\pi^+\pi^-\gamma)/\Gamma_{\text{total}}$ Γ_7/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0099 ± 0.0016 OUR FIT				
0.0099 ± 0.0016		¹⁵ DOLINSKY	91	ND $e^+e^- \rightarrow \pi^+\pi^-\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.0111 ± 0.0014		¹⁶ VASSERMAN	88	ND $e^+e^- \rightarrow \pi^+\pi^-\gamma$
<0.005	90	¹⁷ VASSERMAN	88	ND $e^+e^- \rightarrow \pi^+\pi^-\gamma$

$\Gamma(\pi^0\gamma)/\Gamma_{\text{total}}$ Γ_8/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<u>Γ_8/Γ</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •					
4.20 \pm 0.52	18	ACHASOV	16A	SND 0.60–1.38 $e^+e^- \rightarrow \pi^0\gamma$	■
6.21 $^{+1.28}_{-1.18} \pm 0.39$	18k	AKHMETSHIN 05	CMD2	0.60–1.38 $e^+e^- \rightarrow \pi^0\gamma$	
5.22 $\pm 1.17 \pm 0.75$	36k	ACHASOV	03	SND 0.60–0.97 $e^+e^- \rightarrow \pi^0\gamma$	
6.8 ± 1.7	22	BENAYOUN	96	RVUE 0.54–1.04 $e^+e^- \rightarrow \pi^0\gamma$	
7.9 ± 2.0	20	DOLINSKY	89	ND $e^+e^- \rightarrow \pi^0\gamma$	

 $\Gamma(\pi^0e^+e^-)/\Gamma_{\text{total}}$ Γ_{16}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<u>Γ_{16}/Γ</u>
<1.2	90	ACHASOV	08	SND 0.36–0.97 $e^+e^- \rightarrow \pi^0e^+e^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<1.6		AKHMETSHIN 05A	CMD2	0.72–0.84 e^+e^-	

 $\Gamma(\eta e^+e^-)/\Gamma_{\text{total}}$ Γ_{17}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<u>Γ_{17}/Γ</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.7	AKHMETSHIN 05A	CMD2	0.72–0.84 e^+e^-	

 $\Gamma(\pi^0\pi^0\gamma)/\Gamma_{\text{total}}$ Γ_{10}/Γ

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<u>Γ_{10}/Γ</u>
4.5 \pm 0.8 OUR FIT					
4.5 $^{+0.9}_{-0.8}$ OUR AVERAGE					
5.2 $^{+1.5}_{-1.3} \pm 0.6$	190	23 AKHMETSHIN 04B	CMD2	$0.6\text{--}0.97 e^+e^- \rightarrow \pi^0\pi^0\gamma$	
4.1 $^{+1.0}_{-0.9} \pm 0.3$	295	24 ACHASOV	02F	SND 0.36–0.97 $e^+e^- \rightarrow \pi^0\pi^0\gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
4.8 $^{+3.4}_{-1.8} \pm 0.5$	63	25 ACHASOV	00G	SND $e^+e^- \rightarrow \pi^0\pi^0\gamma$	

1 Possibly large ρ - ω interference leads us to increase the minus error.2 Result contains $11 \pm 11\%$ correction using SU(3) for central value. The error on the correction takes account of possible ρ - ω interference and the upper limit agrees with the upper limit of $\omega \rightarrow \mu^+\mu^-$ from this experiment.3 HYAMS 67's mass resolution is 20 MeV. The ω region was excluded.4 The ρ' contribution is not taken into account.5 ACHASOV 07B reports $[\Gamma(\rho(770) \rightarrow \eta\gamma)/\Gamma_{\text{total}}] \times [B(\rho(770) \rightarrow e^+e^-)] = (1.32 \pm 0.14 \pm 0.08) \times 10^{-8}$ which we divide by our best value $B(\rho(770) \rightarrow e^+e^-) = (4.72 \pm 0.05) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Supersedes ACHASOV 00D and ACHASOV 06A.6 Solution corresponding to constructive ω - ρ interference.7 Using $B(\rho \rightarrow e^+e^-) = (4.67 \pm 0.09) \times 10^{-5}$ and $B(\eta \rightarrow \gamma\gamma) = 39.43 \pm 0.26\%$.8 Not independent of the corresponding $\Gamma(e^+e^-) \times \Gamma(\eta\gamma)/\Gamma_{\text{total}}^2$.

- ⁹ The combined fit from 600 to 1380 MeV taking into account $\rho(770)$, $\omega(782)$, $\phi(1020)$, and $\rho(1450)$ (mass and width fixed at 1450 MeV and 310 MeV respectively).
- ¹⁰ Using $B(\rho \rightarrow e^+ e^-) = (4.75 \pm 0.10) \times 10^{-5}$ from AKHMETSHIN 02 and $B(\eta \rightarrow 3\pi^0) = (32.24 \pm 0.29) \times 10^{-2}$.
- ¹¹ Reanalysis of DRUZHININ 84, DOLINSKY 89, and DOLINSKY 91 taking into account a triangle anomaly contribution. Constructive ρ - ω interference solution.
- ¹² Statistical significance is less than 3σ .
- ¹³ Model dependent, assumes $I = 1, 2$, or 3 for the 3π system.
- ¹⁴ Assuming no interference between the ρ and ω contributions.
- ¹⁵ Bremsstrahlung from a decay pion and for photon energy above 50 MeV.
- ¹⁶ Superseded by DOLINSKY 91.
- ¹⁷ Structure radiation due to quark rearrangement in the decay.
- ¹⁸ Using $B(\rho \rightarrow e^+ e^-)$ from PDG 15. Supersedes ACHASOV 03.
- ¹⁹ Using $B(\rho \rightarrow e^+ e^-) = (4.67 \pm 0.09) \times 10^{-5}$.
- ²⁰ Not independent of the corresponding $\Gamma(e^+ e^-) \times \Gamma(\pi^0 \gamma) / \Gamma_{\text{total}}^2$.
- ²¹ Using $B(\rho \rightarrow e^+ e^-) = (4.54 \pm 0.10) \times 10^{-5}$.
- ²² Reanalysis of DRUZHININ 84, DOLINSKY 89, and DOLINSKY 91 taking into account a triangle anomaly contribution.
- ²³ This branching ratio includes the conventional VMD mechanism $\rho \rightarrow \omega \pi^0$, $\omega \rightarrow \pi^0 \gamma$, and the new decay mode $\rho \rightarrow f_0(500)\gamma$, $f_0(500) \rightarrow \pi^0 \pi^0$ with a branching ratio $(2.0^{+1.1}_{-0.9} \pm 0.3) \times 10^{-5}$ differing from zero by 2.0 standard deviations.
- ²⁴ This branching ratio includes the conventional VMD mechanism $\rho \rightarrow \omega \pi^0$, $\omega \rightarrow \pi^0 \gamma$ and the new decay mode $\rho \rightarrow f_0(500)\gamma$, $f_0(500) \rightarrow \pi^0 \pi^0$ with a branching ratio $(1.9^{+0.9}_{-0.8} \pm 0.4) \times 10^{-5}$ differing from zero by 2.4 standard deviations. Supersedes ACHASOV 00G.
- ²⁵ Superseded by ACHASOV 02F.

$\rho(770)$ REFERENCES

ABLIKIM	17	PRL 118 012001	M. Ablikim <i>et al.</i>	(BES III Collab.)
ACHASOV	16A	PR D93 092001	M.N. Achasov <i>et al.</i>	(SND Collab.)
PDG	15	RPP 2015 at pdg.lbl.gov		(PDG Collab.)
ABRAMOWICZ	12	EPJ C72 1869	H. Abramowicz <i>et al.</i>	(ZEUS Collab.)
LEES	12G	PR D86 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)
BENAYOUN	10	EPJ C65 211	M. Benayoun <i>et al.</i>	
ACHASOV	09A	JETP 109 379	M.N. Achasov <i>et al.</i>	(SND Collab.)
		Translated from ZETF 136 442.		
ACHASOV	08	JETP 107 61	M.N. Achasov <i>et al.</i>	(SND Collab.)
		Translated from ZETF 134 80.		
FUJKAWA	08	PR D78 072006	M. Fujikawa <i>et al.</i>	(BELLE Collab.)
ACHASOV	07B	PR D76 077101	M.N. Achasov <i>et al.</i>	(SND Collab.)
AKHMETSHIN	07	PL B648 28	R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
ACHASOV	06	JETP 103 380	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
		Translated from ZETF 130 437.		
ACHASOV	06A	PR D74 014016	M.N. Achasov <i>et al.</i>	(SND Collab.)
AULCHENKO	06	JETPL 84 413	V.M. Aulchenko <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
		Translated from ZETFP 84 491.		
ACHASOV	05A	JETP 101 1053	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
		Translated from ZETF 128 1201.		
AKHMETSHIN	05	PL B605 26	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
AKHMETSHIN	05A	PL B613 29	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
ALOISIO	05	PL B606 12	A. Aloisio <i>et al.</i>	(KLOE Collab.)
AULCHENKO	05	JETPL 82 743	V.M. Aulchenko <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
		Translated from ZETFP 82 841.		

SCHAEL	05C	PRPL 421 191	S. Schael <i>et al.</i>	(ALEPH Collab.)
AKHMETSHIN	04	PL B578 285	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
AKHMETSHIN	04B	PL B580 119	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
ACHASOV	03	PL B559 171	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	03D	PR D68 052006	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ALOISIO	03	PL B561 55	A. Aloisio <i>et al.</i>	(KLOE Collab.)
SANZ-CILLERO	03	EPJ C27 587	J.J. Sanz-Cillero, A. Pich	
ACHASOV	02	PR D65 032002	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	02F	PL B537 201	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
AKHMETSHIN	02	PL B527 161	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
AKHMETSHIN	01B	PL B509 217	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
COLANGELO	01	NP B603 125	G. Colangelo, J. Gasser, H. Leytwyler	
PICH	01	PR D63 093005	A. Pich, J. Portoles	
ACHASOV	00	EPJ C12 25	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	00D	JETPL 72 282	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	00G	Translated from ZETFP 72 411.	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	00G	JETPL 71 355	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	Translated from ZETFP 71 519.			
AKHMETSHIN	00	PL B475 190	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
ANDERSON	00A	PR D61 112002	S. Anderson <i>et al.</i>	(CLEO Collab.)
ABELE	99E	PL B469 270	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
BENAYOUN	98	EPJ C2 269	M. Benayoun <i>et al.</i>	(IPNP, NOVO, ADLD+)
BREITWEG	98B	EPJ C2 247	J. Breitweg <i>et al.</i>	(ZEUS Collab.)
GARDNER	98	PR D57 2716	S. Gardner, H.B. O'Connell	
Also		PR D62 019903 (errat.)	S. Gardner, H.B. O'Connell	
ABELE	97	PL B391 191	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ADAMS	97	ZPHY C74 237	M.R. Adams <i>et al.</i>	(E665 Collab.)
BARATE	97M	ZPHY C76 15	R. Barate <i>et al.</i>	(ALEPH Collab.)
BERTIN	97C	PL B408 476	A. Bertin <i>et al.</i>	(OBELIX Collab.)
BOGOLYUB...	97	PAN 60 46	M.Y. Bogolyubsky <i>et al.</i>	(MOSU, SERP)
Translated from YAF 60 53.				
O'CONNELL	97	NP A623 559	H.B. O'Connell <i>et al.</i>	(ADLD)
BENAYOUN	96	ZPHY C72 221	M. Benayoun <i>et al.</i>	(IPNP, NOVO)
BERNICA	94	PR D50 4454	A. Bernicha, G. Lopez Castro, J. Pestieau	(LOUV+)
WEIDENAUER	93	ZPHY C59 387	P. Weidenauer <i>et al.</i>	(ASTERIX Collab.)
AGUILAR-...	91	ZPHY C50 405	M. Aguilar-Benitez <i>et al.</i>	(LEBC-EHS Collab.)
DOLINSKY	91	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)
KUHN	90	ZPHY C48 445	J.H. Kuhn <i>et al.</i>	(MPIM)
ANTIROV	89	ZPHY C42 185	Y.M. Antipov <i>et al.</i>	(SERP, JINR, BGNA+)
BISELLO	89	PL B220 321	D. Bisello <i>et al.</i>	(DM2 Collab.)
DOLINSKY	89	ZPHY C42 511	S.I. Dolinsky <i>et al.</i>	(NOVO)
DUBNICKA	89	JP G15 1349	S. Dubnicka <i>et al.</i>	(JINR, SLOV)
GESHKEN...	89	ZPHY C45 351	B.V. Geshkenbein	(ITEP)
KURDADZE	88	JETPL 47 512	L.M. Kurdadze <i>et al.</i>	(NOVO)
Translated from ZETFP 47 432.				
VASSERMAN	88	SJNP 47 1035	I.B. Vasserman <i>et al.</i>	(NOVO)
Translated from YAF 47 1635.				
VASSERMAN	88B	SJNP 48 480	I.B. Vasserman <i>et al.</i>	(NOVO)
Translated from YAF 48 753.				
AULCHENKO	87C	IYF 87-90 Preprint	V.M. Aulchenko <i>et al.</i>	(NOVO)
CAPRARO	87	NP B288 659	L. Capraro <i>et al.</i>	(CLER, FRAS, MILA+)
BRAMON	86	PL B173 97	A. Bramon, J. Casulleras	(BARC)
HUSTON	86	PR D33 3199	J. Huston <i>et al.</i>	(ROCH, FNAL, MINN)
KURDADZE	86	JETPL 43 643	L.M. Kurdadze <i>et al.</i>	(NOVO)
Translated from ZETFP 43 497.				
BARKOV	85	NP B256 365	L.M. Barkov <i>et al.</i>	(NOVO)
DRUZHININ	84	PL 144B 136	V.P. Druzhinin <i>et al.</i>	(NOVO)
CHABAUD	83	NP B223 1	V. Chabaud <i>et al.</i>	(CERN, CRAC, MPIM)
JENSEN	83	PR D27 26	T. Jensen <i>et al.</i>	(ROCH, FNAL, MINN)
HEYN	81	ZPHY C7 169	M.F. Heyn, C.B. Lang	(GRAZ)
BOHACIK	80	PR D21 1342	J. Bohacik, H. Kuhnel	(SLOV, WIEN)
LANG	79	PR D19 956	C.B. Lang, A. Mas-Parareda	(GRAZ)
BARTALUCCI	78	NC 44A 587	S. Bartalucci <i>et al.</i>	(DESY, FRAS)
WICKLUND	78	PR D17 1197	A.B. Wicklund <i>et al.</i>	(ANL)
ANDREWS	77	PRL 38 198	D.E. Andrews <i>et al.</i>	(ROCH)
DEUTSCH...	76	NP B103 426	M. Deutschmann <i>et al.</i>	(AACH3, BERL, BONN+)
ENGLER	74	PR D10 2070	A. Engler <i>et al.</i>	(CMU, CASE)
ESTABROOKS	74	NP B79 301	P.G. Estabrooks, A.D. Martin	(DURH)
GRAYER	74	NP B75 189	G. Grayer <i>et al.</i>	(CERN, MPIM)
BYERLY	73	PR D7 637	W.L. Byerly <i>et al.</i>	(MICH)
GLADDING	73	PR D8 3721	G.E. Gladding <i>et al.</i>	(HARV)
HYAMS	73	NP B64 134	B.D. Hyams <i>et al.</i>	(CERN, MPIM)

PROTOPOP...	73	PR D7 1279	S.D. Protopopescu <i>et al.</i>	(LBL)
BALLAM	72	PR D5 545	J. Ballam <i>et al.</i>	(SLAC, LBL, TUFTS)
BENAKSAS	72	PL 39B 289	D. Benaksas <i>et al.</i>	(ORSAY)
JACOBS	72	PR D6 1291	L.D. Jacobs	(SACL)
RATCLIFF	72	PL 38B 345	B.N. Ratcliff <i>et al.</i>	(SLAC)
ABRAMS	71	PR D4 653	G.S. Abrams <i>et al.</i>	(LBL)
ALVENSLEB...	70	PRL 24 786	H. Alvensleben <i>et al.</i>	(DESY)
BIGGS	70	PRL 24 1197	P.J. Biggs <i>et al.</i>	(DARE)
ERBE	69	PR 188 2060	R. Erbe <i>et al.</i>	(German Bubble Chamber Collab.)
MALAMUD	69	Argonne Conf. 93	E.I. Malamud, P.E. Schlein	(UCLA)
REYNOLDS	69	PR 184 1424	B.G. Reynolds <i>et al.</i>	(FSU)
ROTHWELL	69	PRL 23 1521	P.L. Rothwell <i>et al.</i>	(NEAS)
WEHMANN	69	PR 178 2095	A.A. Wehmann <i>et al.</i>	(HARV, CASE, SLAC+)
ARMENISE	68	NC 54A 999	N. Armenise <i>et al.</i>	(BARI, BGNA, FIRZ+)
BATON	68	PR 176 1574	J.P. Baton, G. Laurens	(SACL)
CHUNG	68	PR 165 1491	S.U. Chung <i>et al.</i>	(LRL)
FOSTER	68	NP B6 107	M. Foster <i>et al.</i>	(CERN, CDEF)
GOUNARIS	68	PRL 21 244	G.J. Gounaris, J.J. Sakurai	
HUSON	68	PL 28B 208	R. Huson <i>et al.</i>	(ORSAY, MILA, UCLA)
HYAMS	68	NP B7 1	B.D. Hyams <i>et al.</i>	(CERN, MPIM)
LANZEROTTI	68	PR 166 1365	L.J. Lanzerotti <i>et al.</i>	(HARV)
PISUT	68	NP B6 325	J. Pisut, M. Roos	(CERN)
ASBURY	67B	PRL 19 865	J.G. Asbury <i>et al.</i>	(DESY, COLU)
BACON	67	PR 157 1263	T.C. Bacon <i>et al.</i>	(BNL)
EISNER	67	PR 164 1699	R.L. Eisner <i>et al.</i>	(PURD)
HUWE	67	PL 24B 252	D.O. Huwe <i>et al.</i>	(COLU)
HYAMS	67	PL 24B 634	B.D. Hyams <i>et al.</i>	(CERN, MPIM)
MILLER	67B	PR 153 1423	D.H. Miller <i>et al.</i>	(PURD)
ALFF-...	66	PR 145 1072	C. Alff-Steinberger <i>et al.</i>	(COLU, RUTG)
FERBEL	66	PL 21 111	T. Ferbel	(ROCH)
HAGOPIAN	66	PR 145 1128	V. Hagopian <i>et al.</i>	(PENN, SACL)
HAGOPIAN	66B	PR 152 1183	V. Hagopian, Y.L. Pan	(PENN, LRL)
JACOBS	66B	UCRL 16877	L.D. Jacobs	(LRL)
JAMES	66	PR 142 896	F.E. James, H.L. Kraybill	(YALE, BNL)
ROSS	66	PR 149 1172	M. Ross, L. Stodolsky	
SOEDING	66	PL B19 702	P. Soeding	
WEST	66	PR 149 1089	E. West <i>et al.</i>	(WISC)
BLIEDEN	65	PL 19 444	H.R. Blieden <i>et al.</i>	(CERN MMS Collab.)
CARMONY	64	PRL 12 254	D.D. Carmony <i>et al.</i>	(UCB)
GOLDHABER	64	PRL 12 336	G. Goldhaber <i>et al.</i>	(LRL, UCB)
ABOLINS	63	PRL 11 381	M.A. Abolins <i>et al.</i>	(UCSD)
