45. Clebsch-Gordan Coefficients, Spherical Harmonics, and d Functions

Note: A square-root sign is to be understood over *every* coefficient, *e.g.*, for $-8/15$ read $-\sqrt{8/15}$.

Notation:

		J	J	...	
		M	M	...	
m_1	m_2				
m_1	m_2	Coefficients			
.	.				
.	.				
.	.				

$$Y_1^0 = \sqrt{\frac{3}{4\pi}}\cos\theta$$

$$Y_1^1 = -\sqrt{\frac{3}{8\pi}}\sin\theta\, e^{i\phi}$$

$$Y_2^0 = \sqrt{\frac{5}{4\pi}}\Big(\frac{3}{2}\cos^2\theta - \frac{1}{2}\Big)$$

$$Y_2^1 = -\sqrt{\frac{15}{8\pi}}\sin\theta\cos\theta\, e^{i\phi}$$

$$Y_2^2 = \frac{1}{4}\sqrt{\frac{15}{2\pi}}\sin^2\theta\, e^{2i\phi}$$

$$Y_\ell^{-m} = (-1)^m Y_\ell^{m*}$$

$$d^{\,\ell}_{m,0} = \sqrt{\frac{4\pi}{2\ell+1}}\, Y_\ell^m\, e^{-im\phi}$$

$$\langle j_1 j_2 m_1 m_2 | j_1 j_2 J M\rangle = (-1)^{J-j_1-j_2}\langle j_2 j_1 m_2 m_1 | j_2 j_1 J M\rangle$$

$$d^{\,j}_{m',m} = (-1)^{m-m'} d^{\,j}_{m,m'} = d^{\,j}_{-m,-m'}$$

$$d^{\,1}_{0,0} = \cos\theta \qquad d^{\,1/2}_{1/2,1/2} = \cos\frac{\theta}{2} \qquad d^{\,1/2}_{1/2,-1/2} = -\sin\frac{\theta}{2}$$

$$d^{\,1}_{1,1} = \frac{1+\cos\theta}{2} \qquad d^{\,1}_{1,0} = -\frac{\sin\theta}{\sqrt{2}} \qquad d^{\,1}_{1,-1} = \frac{1-\cos\theta}{2}$$

$$d^{\,3/2}_{3/2,3/2} = \frac{1+\cos\theta}{2}\cos\frac{\theta}{2}$$

$$d^{\,3/2}_{3/2,1/2} = -\sqrt{3}\frac{1+\cos\theta}{2}\sin\frac{\theta}{2} \qquad d^{\,2}_{2,2} = \Big(\frac{1+\cos\theta}{2}\Big)^2$$

$$d^{\,3/2}_{3/2,-1/2} = \sqrt{3}\frac{1-\cos\theta}{2}\cos\frac{\theta}{2} \qquad d^{\,2}_{2,1} = -\frac{1+\cos\theta}{2}\sin\theta$$

$$d^{\,3/2}_{3/2,-3/2} = -\frac{1-\cos\theta}{2}\sin\frac{\theta}{2} \qquad d^{\,2}_{2,0} = \frac{\sqrt{6}}{4}\sin^2\theta \qquad d^{\,2}_{1,1} = \frac{1+\cos\theta}{2}(2\cos\theta - 1)$$

$$d^{\,3/2}_{1/2,1/2} = \frac{3\cos\theta - 1}{2}\cos\frac{\theta}{2} \qquad d^{\,2}_{2,-1} = -\frac{1-\cos\theta}{2}\sin\theta \qquad d^{\,2}_{1,0} = -\sqrt{\frac{3}{2}}\sin\theta\cos\theta$$

$$d^{\,3/2}_{1/2,-1/2} = -\frac{3\cos\theta + 1}{2}\sin\frac{\theta}{2} \qquad d^{\,2}_{2,-2} = \Big(\frac{1-\cos\theta}{2}\Big)^2 \qquad d^{\,2}_{1,-1} = \frac{1-\cos\theta}{2}(2\cos\theta + 1) \qquad d^{\,2}_{0,0} = \Big(\frac{3}{2}\cos^2\theta - \frac{1}{2}\Big)$$

Figure 45.1: The sign convention is that of Wigner (*Group Theory*, Academic Press, New York, 1959), also used by Condon and Shortley (*The Theory of Atomic Spectra*, Cambridge Univ. Press, New York, 1953), Rose (*Elementary Theory of Angular Momentum*, Wiley, New York, 1957), and Cohen (*Tables of the Clebsch-Gordan Coefficients*, North American Rockwell Science Center, Thousand Oaks, Calif., 1974).