

**K(1460)**

$$I(J^P) = \frac{1}{2}(0^-)$$

OMITTED FROM SUMMARY TABLE

Observed in  $K\pi\pi$  partial-wave analysis.**K(1460) MASS**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
~ 1460	DAUM	81C	CNTR	– 63 $K^- p \rightarrow K^- 2\pi p$
~ 1400	<sup>1</sup> BRANDENB...	76B	ASPK	± 13 $K^\pm p \rightarrow K^\pm 2\pi p$
<sup>1</sup> Coupled mainly to $K f_0(1370)$ . Decay into $K^*(892)\pi$ seen.				

**K(1460) WIDTH**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
~ 260	DAUM	81C	CNTR	– 63 $K^- p \rightarrow K^- 2\pi p$
~ 250	<sup>2</sup> BRANDENB...	76B	ASPK	± 13 $K^\pm p \rightarrow K^\pm 2\pi p$
<sup>2</sup> Coupled mainly to $K f_0(1370)$ . Decay into $K^*(892)\pi$ seen.				

**K(1460) DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $K^*(892)\pi$	seen
$\Gamma_2$ $K\rho$	seen
$\Gamma_3$ $K_0^*(1430)\pi$	seen

**K(1460) PARTIAL WIDTHS** **$\Gamma(K^*(892)\pi)$   $\Gamma_1$** 

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
~ 109	DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$

 **$\Gamma(K\rho)$   $\Gamma_2$** 

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
~ 34	DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$

 **$\Gamma(K_0^*(1430)\pi)$   $\Gamma_3$** 

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
~ 117	DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$

## ***K*(1460) REFERENCES**

DAUM            81C   NP B187 1            C. Daum *et al.*            (AMST, CERN, CRAC, MPIM+)  
BRANDENB... 76B   PRL 36 1239           G.W. Brandenburg *et al.*            (SLAC) JP

---