

# $b$ -baryon ADMIXTURE ( $\Lambda_b$ , $\Xi_b$ , $\Sigma_b$ , $\Omega_b$ )

## $b$ -baryon ADMIXTURE MEAN LIFE

Each measurement of the  $b$ -baryon mean life is an average over an admixture of various  $b$  baryons which decay weakly. Different techniques emphasize different admixtures of produced particles, which could result in a different  $b$ -baryon mean life. More  $b$ -baryon flavor specific channels are not included in the measurement.

VALUE ( $10^{-12}$ s)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
1.218 <sup>+0.130</sup> <sub>-0.115</sub> ± 0.042		<sup>1</sup> ABAZOV	07S D0	Repl. by ABAZOV 12U
1.22 <sup>+0.22</sup> <sub>-0.18</sub> ± 0.04		<sup>1</sup> ABAZOV	05C D0	Repl. by ABAZOV 07S
1.16 ± 0.20 ± 0.08		<sup>2</sup> ABREU	99W DLPH	$e^+ e^- \rightarrow Z$
1.19 ± 0.14 ± 0.07		<sup>3</sup> ABREU	99W DLPH	$e^+ e^- \rightarrow Z$
1.14 ± 0.08 ± 0.04		<sup>4</sup> ABREU	99W DLPH	$e^+ e^- \rightarrow Z$
1.11 <sup>+0.19</sup> <sub>-0.18</sub> ± 0.05		<sup>5</sup> ABREU	99W DLPH	$e^+ e^- \rightarrow Z$
1.29 <sup>+0.24</sup> <sub>-0.22</sub> ± 0.06		<sup>5</sup> ACKERSTAFF	98G OPAL	$e^+ e^- \rightarrow Z$
1.20 ± 0.08 ± 0.06		<sup>6</sup> BARATE	98D ALEP	$e^+ e^- \rightarrow Z$
1.21 ± 0.11		<sup>5</sup> BARATE	98D ALEP	$e^+ e^- \rightarrow Z$
1.32 ± 0.15 ± 0.07		<sup>7</sup> ABE	96M CDF	$p\bar{p}$ at 1.8 TeV
1.46 <sup>+0.22</sup> <sub>-0.21</sub> ± 0.07		ABREU	96D DLPH	Repl. by ABREU 99W
1.10 <sup>+0.19</sup> <sub>-0.17</sub> ± 0.09		<sup>5</sup> ABREU	96D DLPH	$e^+ e^- \rightarrow Z$
1.16 ± 0.11 ± 0.06		<sup>5</sup> AKERS	96 OPAL	$e^+ e^- \rightarrow Z$
1.27 <sup>+0.35</sup> <sub>-0.29</sub> ± 0.09		ABREU	95S DLPH	Repl. by ABREU 99W
1.05 <sup>+0.12</sup> <sub>-0.11</sub> ± 0.09	290	BUSKULIC	95L ALEP	Repl. by BARATE 98D
1.04 <sup>+0.48</sup> <sub>-0.38</sub> ± 0.10	11	<sup>8</sup> ABREU	93F DLPH	Excess $\Lambda\mu^-$ , decay lengths
1.05 <sup>+0.23</sup> <sub>-0.20</sub> ± 0.08	157	<sup>9</sup> AKERS	93 OPAL	Excess $\Lambda\ell^-$ , decay lengths
1.12 <sup>+0.32</sup> <sub>-0.29</sub> ± 0.16	101	<sup>10</sup> BUSKULIC	92I ALEP	Excess $\Lambda\ell^-$ , impact parameters

<sup>1</sup> Measured mean life using fully reconstructed  $\Lambda_b^0 \rightarrow J/\psi \Lambda$  decays.

<sup>2</sup> Measured using  $\Lambda\ell^-$  decay length.

<sup>3</sup> Measured using  $p\ell^-$  decay length.

<sup>4</sup> This ABREU 99W result is the combined result of the  $\Lambda\ell^-$ ,  $p\ell^-$ , and excess  $\Lambda\mu^-$  impact parameter measurements.

<sup>5</sup> Measured using  $\Lambda_c\ell^-$  and  $\Lambda\ell^+\ell^-$ .

<sup>6</sup> Measured using the excess of  $\Lambda\ell^-$ , lepton impact parameter.

<sup>7</sup> Measured using  $\Lambda_c\ell^-$ .

<sup>8</sup> ABREU 93F superseded by ABREU 96D.

<sup>9</sup> AKERS 93 superseded by AKERS 96.<sup>10</sup> BUSKULIC 92I superseded by BUSKULIC 95L.

## **$b$ -baryon ADMIXTURE DECAY MODES ( $\Lambda_b, \Xi_b, \Sigma_b, \Omega_b$ )**

These branching fractions are actually an average over weakly decaying  $b$ -baryons weighted by their production rates at the LHC, LEP, and Tevatron, branching ratios, and detection efficiencies. They scale with the  $b$ -baryon production fraction  $B(b \rightarrow b\text{-baryon})$ .

The branching fractions  $B(b\text{-baryon} \rightarrow \Lambda \ell^- \bar{\nu}_\ell \text{anything})$  and  $B(\Lambda_b^0 \rightarrow \Lambda_c^+ \ell^- \bar{\nu}_\ell \text{anything})$  are not pure measurements because the underlying measured products of these with  $B(b \rightarrow b\text{-baryon})$  were used to determine  $B(b \rightarrow b\text{-baryon})$ , as described in the note “Production and Decay of  $b$ -Flavored Hadrons.”

For inclusive branching fractions, e.g.,  $B \rightarrow D^\pm \text{anything}$ , the values usually are multiplicities, not branching fractions. They can be greater than one.

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 p \mu^- \bar{\nu} \text{anything}$	( 5.5 $\pm$ 2.2 ) %
$\Gamma_2 p \ell \bar{\nu}_\ell \text{anything}$	( 5.3 $\pm$ 1.1 ) %
$\Gamma_3 p \text{anything}$	( 66 $\pm$ 21 ) %
$\Gamma_4 \Lambda \ell^- \bar{\nu}_\ell \text{anything}$	( 3.6 $\pm$ 0.6 ) %
$\Gamma_5 \Lambda \ell^+ \nu_\ell \text{anything}$	( 3.0 $\pm$ 0.8 ) %
$\Gamma_6 \Lambda \text{anything}$	( 37 $\pm$ 7 ) %
$\Gamma_7 \Xi^- \ell^- \bar{\nu}_\ell \text{anything}$	( 6.2 $\pm$ 1.6 ) $\times 10^{-3}$

## **$b$ -baryon ADMIXTURE ( $\Lambda_b, \Xi_b, \Sigma_b, \Omega_b$ ) BRANCHING RATIOS**

$\Gamma(p \mu^- \bar{\nu} \text{anything})/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma$
$5.5^{+2.1}_{-1.7} \pm 0.7$	

<sup>11</sup> ABREU 95S reports  $[\Gamma(b\text{-baryon} \rightarrow p \mu^- \bar{\nu} \text{anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})]$

$= 0.0049 \pm 0.0011^{+0.0015}_{-0.0011}$  which we divide by our best value  $B(\bar{b} \rightarrow b\text{-baryon}) = (8.9 \pm 1.2) \times 10^{-2}$ . Our first error is their experiment’s error and our second error is the systematic error from using our best value.

$\Gamma(p \ell \bar{\nu}_\ell \text{anything})/\Gamma_{\text{total}}$	$\Gamma_2/\Gamma$
$5.3 \pm 0.9 \pm 0.7$	

<sup>12</sup> BARATE 98V reports  $[\Gamma(b\text{-baryon} \rightarrow p \ell \bar{\nu}_\ell \text{anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})]$   
 $= (4.72 \pm 0.66 \pm 0.44) \times 10^{-3}$  which we divide by our best value  $B(\bar{b} \rightarrow b\text{-baryon}) = (8.9 \pm 1.2) \times 10^{-2}$ . Our first error is their experiment’s error and our second error is the systematic error from using our best value.

$\Gamma(p\ell\bar{\nu}_\ell \text{anything})/\Gamma(p\text{anything})$   $\Gamma_2/\Gamma_3$ 

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b>8.0±1.2±1.4</b>	BARATE 98V	ALEP	$e^+ e^- \rightarrow Z$

 $\Gamma(\Lambda\ell^-\bar{\nu}_\ell \text{anything})/\Gamma_{\text{total}}$   $\Gamma_4/\Gamma$ 

The values and averages in this section serve only to show what values result if one assumes our  $B(b \rightarrow b\text{-baryon})$ . They cannot be thought of as measurements since the underlying product branching fractions were also used to determine  $B(b \rightarrow b\text{-baryon})$  as described in the note on “Production and Decay of  $b$ -Flavored Hadrons.”

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.6±0.6 OUR AVERAGE</b>				

3.7±0.5±0.5	13	BARATE	98D	ALEP $e^+ e^- \rightarrow Z$
3.3±0.4±0.4	14	AKERS	96	OPAL Excess of $\Lambda\ell^-$ over $\Lambda\ell^+$
3.4±0.8±0.5	262	ABREU	95S	DLPH Excess of $\Lambda\ell^-$ over $\Lambda\ell^+$
6.9±1.3±0.9	290	BUSKULIC	95L	ALEP Excess of $\Lambda\ell^-$ over $\Lambda\ell^+$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	157	17 AKERS	93	OPAL Excess of $\Lambda\ell^-$ over $\Lambda\ell^+$
7.9±2.3±1.1	101	18 BUSKULIC	92I	ALEP Excess of $\Lambda\ell^-$ over $\Lambda\ell^+$

<sup>13</sup> BARATE 98D reports  $[\Gamma(b\text{-baryon} \rightarrow \Lambda\ell^-\bar{\nu}_\ell \text{anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})] = 0.00326 \pm 0.00016 \pm 0.00039$  which we divide by our best value  $B(\bar{b} \rightarrow b\text{-baryon}) = (8.9 \pm 1.2) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. Measured using the excess of  $\Lambda\ell^-$ , lepton impact parameter.

<sup>14</sup> AKERS 96 reports  $[\Gamma(b\text{-baryon} \rightarrow \Lambda\ell^-\bar{\nu}_\ell \text{anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})] = 0.00291 \pm 0.00023 \pm 0.00025$  which we divide by our best value  $B(\bar{b} \rightarrow b\text{-baryon}) = (8.9 \pm 1.2) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>15</sup> ABREU 95S reports  $[\Gamma(b\text{-baryon} \rightarrow \Lambda\ell^-\bar{\nu}_\ell \text{anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})] = 0.0030 \pm 0.0006 \pm 0.0004$  which we divide by our best value  $B(\bar{b} \rightarrow b\text{-baryon}) = (8.9 \pm 1.2) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>16</sup> BUSKULIC 95L reports  $[\Gamma(b\text{-baryon} \rightarrow \Lambda\ell^-\bar{\nu}_\ell \text{anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})] = 0.0061 \pm 0.0006 \pm 0.0010$  which we divide by our best value  $B(\bar{b} \rightarrow b\text{-baryon}) = (8.9 \pm 1.2) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>17</sup> AKERS 93 superseded by AKERS 96.

<sup>18</sup> BUSKULIC 92I reports  $[\Gamma(b\text{-baryon} \rightarrow \Lambda\ell^-\bar{\nu}_\ell \text{anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})] = 0.0070 \pm 0.0010 \pm 0.0018$  which we divide by our best value  $B(\bar{b} \rightarrow b\text{-baryon}) = (8.9 \pm 1.2) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. Superseded by BUSKULIC 95L.

 $\Gamma(\Lambda\ell^+\nu_\ell \text{anything})/\Gamma(\Lambda \text{anything})$   $\Gamma_5/\Gamma_6$ 

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>8.0±1.2±0.8</b>	ABBIENDI 99L	OPAL	$e^+ e^- \rightarrow Z$

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.0±1.2±0.7	ACKERSTAFF 97N	OPAL	Repl. by ABBIENDI 99L
-------------	----------------	------	-----------------------

$\Gamma(\Lambda\text{anything})/\Gamma_{\text{total}}$  $\Gamma_6/\Gamma$ 

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b>37±7 OUR AVERAGE</b>			
39±5±5	19 ABBIENDI	99L OPAL	$e^+ e^- \rightarrow Z$
$25^{+14}_{-9} \pm 3$	20 ABREU	95C DLPH	$e^+ e^- \rightarrow Z$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
44±7±6	21 ACKERSTAFF	97N OPAL	Repl. by ABBIENDI 99L
19 ABBIENDI 99L reports $[\Gamma(b\text{-baryon} \rightarrow \Lambda\text{anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})]$ = $0.035 \pm 0.0032 \pm 0.0035$ which we divide by our best value $B(\bar{b} \rightarrow b\text{-baryon}) = (8.9 \pm 1.2) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.			
20 ABREU 95C reports $0.28^{+0.17}_{-0.12}$ from a measurement of $[\Gamma(b\text{-baryon} \rightarrow \Lambda\text{anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})]$ assuming $B(\bar{b} \rightarrow b\text{-baryon}) = 0.08 \pm 0.02$ , which we rescale to our best value $B(\bar{b} \rightarrow b\text{-baryon}) = (8.9 \pm 1.2) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.			
21 ACKERSTAFF 97N reports $[\Gamma(b\text{-baryon} \rightarrow \Lambda\text{anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})]$ = $0.0393 \pm 0.0046 \pm 0.0037$ which we divide by our best value $B(\bar{b} \rightarrow b\text{-baryon}) = (8.9 \pm 1.2) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.			

 $\Gamma(\Xi^-\ell^-\bar{\nu}_\ell\text{anything})/\Gamma_{\text{total}}$  $\Gamma_7/\Gamma$ 

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>6.2±1.6 OUR AVERAGE</b>			
6.1±1.5±0.8	22 BUSKULIC	96T ALEP	Excess $\Xi^-\ell^-$ over $\Xi^-\ell^+$
6.6±2.6±0.9	23 ABREU	95V DLPH	Excess $\Xi^-\ell^-$ over $\Xi^-\ell^+$
22 BUSKULIC 96T reports $[\Gamma(b\text{-baryon} \rightarrow \Xi^-\ell^-\bar{\nu}_\ell\text{anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})]$ = $0.00054 \pm 0.00011 \pm 0.00008$ which we divide by our best value $B(\bar{b} \rightarrow b\text{-baryon}) = (8.9 \pm 1.2) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.			
23 ABREU 95V reports $[\Gamma(b\text{-baryon} \rightarrow \Xi^-\ell^-\bar{\nu}_\ell\text{anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})]$ = $0.00059 \pm 0.00021 \pm 0.0001$ which we divide by our best value $B(\bar{b} \rightarrow b\text{-baryon}) = (8.9 \pm 1.2) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.			

**b-baryon ADMIXTURE ( $\Lambda_b$ ,  $\Xi_b$ ,  $\Sigma_b$ ,  $\Omega_b$ ) REFERENCES**

ABAZOV	12U	PR D85 112003	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	07S	PRL 99 142001	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	05C	PRL 94 102001	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABBIENDI	99L	EPJ C9 1	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABREU	99W	EPJ C10 185	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACKERSTAFF	98G	PL B426 161	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
BARATE	98D	EPJ C2 197	R. Barate <i>et al.</i>	(ALEPH Collab.)
BARATE	98V	EPJ C5 205	R. Barate <i>et al.</i>	(ALEPH Collab.)
ACKERSTAFF	97N	ZPHY C74 423	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
ABE	96M	PRL 77 1439	F. Abe <i>et al.</i>	(CDF Collab.)
ABREU	96D	ZPHY C71 199	P. Abreu <i>et al.</i>	(DELPHI Collab.)
AKERS	96	ZPHY C69 195	R. Akers <i>et al.</i>	(OPAL Collab.)
BUSKULIC	96T	PL B384 449	D. Buskulic <i>et al.</i>	(ALEPH Collab.)

ABREU	95C	PL B347 447	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	95S	ZPHY C68 375	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	95V	ZPHY C68 541	P. Abreu <i>et al.</i>	(DELPHI Collab.)
BUSKULIC	95L	PL B357 685	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
ABREU	93F	PL B311 379	P. Abreu <i>et al.</i>	(DELPHI Collab.)
AKERS	93	PL B316 435	R. Akers <i>et al.</i>	(OPAL Collab.)
BUSKULIC	92I	PL B297 449	D. Buskulic <i>et al.</i>	(ALEPH Collab.)