

$\chi_{c1}(3872)$

$$I^G(J^{PC}) = 0^+(1^{++})$$

also known as $X(3872)$

This state shows properties different from a conventional $q\bar{q}$ state. A candidate for an exotic structure. See the review on non- $q\bar{q}$ states.

First observed by CHOI 03 in $B \rightarrow K\pi^+\pi^- J/\psi(1S)$ decays as a narrow peak in the invariant mass distribution of the $\pi^+\pi^- J/\psi(1S)$ final state. Isovector hypothesis excluded by AUBERT 05B and CHOI 11.

AAIJ 13Q perform a full five-dimensional amplitude analysis of the angular correlations between the decay products in $B^+ \rightarrow \chi_{c1}(3872)K^+$ decays, where $\chi_{c1}(3872) \rightarrow J/\psi\pi^+\pi^-$ and $J/\psi \rightarrow \mu^+\mu^-$, which unambiguously gives the $J^{PC} = 1^{++}$ assignment under the assumption that the $\pi^+\pi^-$ and J/ψ are in an S -wave. AAIJ 15AO extend this analysis with more data to limit D -wave contributions to $< 4\%$ at 95% CL.

See our note on “Developments in Heavy Quarkonium Spectroscopy”.

$\chi_{c1}(3872)$ MASS FROM $J/\psi X$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3871.69 ± 0.17 OUR AVERAGE				
$3871.9 \pm 0.7 \pm 0.2$	20 ± 5	ABLIKIM	14 BES3	$e^+e^- \rightarrow J/\psi\pi^+\pi^-\gamma$
$3871.95 \pm 0.48 \pm 0.12$	$0.6k$	AAIJ	12H LHCb	$pp \rightarrow J/\psi\pi^+\pi^- X$
$3871.85 \pm 0.27 \pm 0.19$	~ 170	¹ CHOI	11 BELL	$B \rightarrow K\pi^+\pi^- J/\psi$
$3873 \begin{smallmatrix} +1.8 \\ -1.6 \end{smallmatrix} \pm 1.3$	27 ± 8	² DEL-AMO-SA.10B	BABR	$B \rightarrow \omega J/\psi K$
$3871.61 \pm 0.16 \pm 0.19$	$6k$	^{2,3} AALTONEN	09AU CDF2	$p\bar{p} \rightarrow J/\psi\pi^+\pi^- X$
$3871.4 \pm 0.6 \pm 0.1$	93.4	AUBERT	08Y BABR	$B^+ \rightarrow K^+ J/\psi\pi^+\pi^-$
$3868.7 \pm 1.5 \pm 0.4$	9.4	AUBERT	08Y BABR	$B^0 \rightarrow K_S^0 J/\psi\pi^+\pi^-$
$3871.8 \pm 3.1 \pm 3.0$	522	^{2,4} ABAZOV	04F D0	$p\bar{p} \rightarrow J/\psi\pi^+\pi^- X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$3868.6 \pm 1.2 \pm 0.2$	8	⁵ AUBERT	06 BABR	$B^0 \rightarrow K_S^0 J/\psi\pi^+\pi^-$
$3871.3 \pm 0.6 \pm 0.1$	61	⁵ AUBERT	06 BABR	$B^- \rightarrow K^- J/\psi\pi^+\pi^-$
3873.4 ± 1.4	25	⁶ AUBERT	05R BABR	$B^+ \rightarrow K^+ J/\psi\pi^+\pi^-$
$3871.3 \pm 0.7 \pm 0.4$	730	^{2,7} ACOSTA	04 CDF2	$p\bar{p} \rightarrow J/\psi\pi^+\pi^- X$
$3872.0 \pm 0.6 \pm 0.5$	36	⁸ CHOI	03 BELL	$B \rightarrow K\pi^+\pi^- J/\psi$
3836 ± 13	58	^{2,9} ANTONIAZZI	94 E705	$300 \pi^\pm Li \rightarrow J/\psi\pi^+\pi^- X$

¹ The mass difference for the $\chi_{c1}(3872)$ produced in B^+ and B^0 decays is $(-0.71 \pm 0.96 \pm 0.19)$ MeV.

² Width consistent with detector resolution.

³ A possible equal mixture of two states with a mass difference greater than $3.6 \text{ MeV}/c^2$ is excluded at 95% CL.

- ⁴ Calculated from the corresponding $m_{\chi_{c1}(3872)} - m_{J/\psi}$ using $m_{J/\psi} = 3096.916$ MeV.
⁵ Calculated from the corresponding $m_{\chi_{c1}(3872)} - m_{\psi(2S)}$ using $m_{\psi(2S)} = 3686.093$ MeV. Superseded by AUBERT 08Y.
⁶ Calculated from the corresponding $m_{\chi_{c1}(3872)} - m_{\psi(2S)}$ using $m_{\psi(2S)} = 3685.96$ MeV. Superseded by AUBERT 06.
⁷ Superseded by AALTONEN 09AU.
⁸ Superseded by CHOI 11.
⁹ A lower mass value can be due to an incorrect momentum scale for soft pions.

$\chi_{c1}(3872)$ MASS FROM $\bar{D}^{*0} D^0$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$3872.9^{+0.6+0.4}_{-0.4-0.5}$	50	1,2 AUSHEV	10 BELL	$B \rightarrow \bar{D}^{*0} D^0 K$
$3875.1^{+0.7}_{-0.5} \pm 0.5$	33 ± 6	² AUBERT	08B BABR	$B \rightarrow \bar{D}^{*0} D^0 K$
$3875.2 \pm 0.7^{+0.9}_{-1.8}$	24 ± 6	^{2,3} GOKHROO	06 BELL	$B \rightarrow D^0 \bar{D}^0 \pi^0 K$

- ¹ Calculated from the measured $m_{\chi_{c1}(3872)} - m_{D^{*0}} - m_{\bar{D}^0} = 1.1^{+0.6+0.1}_{-0.4-0.3}$ MeV.
² Experiments report $D^{*0} \bar{D}^0$ invariant mass above $D^{*0} \bar{D}^0$ threshold because D^{*0} decay products are kinematically constrained to the D^{*0} mass, even though the D^{*0} may decay off-shell.
³ Superseded by AUSHEV 10.

$m_{\chi_{c1}(3872)} - m_{J/\psi}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$774.9 \pm 3.1 \pm 3.0$	522	ABAZOV	04F D0	$p\bar{p} \rightarrow J/\psi \pi^+ \pi^- X$

$m_{\chi_{c1}(3872)} - m_{\psi(2S)}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
187.4 ± 1.4	25	¹ AUBERT	05R BABR	$B^+ \rightarrow K^+ J/\psi \pi^+ \pi^-$

- ¹ Superseded by AUBERT 06.

$\chi_{c1}(3872)$ WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<1.2	90		CHOI	11 BELL	$B \rightarrow K \pi^+ \pi^- J/\psi$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<2.4	90		ABLIKIM	14 BES3	$e^+ e^- \rightarrow J/\psi \pi^+ \pi^- \gamma$
<3.3	90		AUBERT	08Y BABR	$B^+ \rightarrow K^+ J/\psi \pi^+ \pi^-$
<4.1	90	69	AUBERT	06 BABR	$B \rightarrow K \pi^+ \pi^- J/\psi$
<2.3	90	36	¹ CHOI	03 BELL	$B \rightarrow K \pi^+ \pi^- J/\psi$

- ¹ Superseded by CHOI 11.

$\chi_{c1}(3872)$ WIDTH FROM $\bar{D}^{*0} D^0$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$3.9^{+2.8+0.2}_{-1.4-1.1}$	50	¹ AUSHEV	10	BELL $B \rightarrow \bar{D}^{*0} D^0 K$
$3.0^{+1.9}_{-1.4} \pm 0.9$	33 ± 6	AUBERT	08B	BABR $B \rightarrow \bar{D}^{*0} D^0 K$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹With a measured value of $B(B \rightarrow \chi_{c1}(3872) K) \times B(\chi_{c1}(3872) \rightarrow D^{*0} \bar{D}^0) = (0.80 \pm 0.20 \pm 0.10) \times 10^{-4}$, assumed to be equal for both charged and neutral modes.

$\chi_{c1}(3872)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $e^+ e^-$	
Γ_2 $\pi^+ \pi^- J/\psi(1S)$	$> 3.2 \%$
Γ_3 $\rho^0 J/\psi(1S)$	
Γ_4 $\omega J/\psi(1S)$	$> 2.3 \%$
Γ_5 $D^0 \bar{D}^0 \pi^0$	$> 40 \%$
Γ_6 $\bar{D}^{*0} D^0$	$> 30 \%$
Γ_7 $\gamma\gamma$	
Γ_8 $D^0 \bar{D}^0$	
Γ_9 $D^+ D^-$	
Γ_{10} $\gamma \chi_{c1}$	
Γ_{11} $\gamma \chi_{c2}$	
Γ_{12} $\gamma J/\psi$	$> 7 \times 10^{-3}$
Γ_{13} $\gamma \psi(2S)$	$> 4 \%$
Γ_{14} $\pi^+ \pi^- \eta_c(1S)$	not seen
Γ_{15} $\pi^+ \pi^- \chi_{c1}$	not seen
Γ_{16} $\rho \bar{\rho}$	not seen

C-violating decays

Γ_{17} $\eta J/\psi$

$\chi_{c1}(3872)$ PARTIAL WIDTHS

$\Gamma(e^+ e^-)$					Γ_1
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 4.3	90	¹ ABLIKIM	15V	BES3	$4.0-4.4 e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$
< 280	90	² YUAN	04	RVUE	$e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$

¹ABLIKIM 15V reports this limit from the measurement of $\Gamma(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S)) \times \Gamma(\chi_{c1}(3872) \rightarrow e^+ e^-) / \Gamma < 0.13$ eV using $\Gamma(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S)) / \Gamma = 3\%$.

²Using BAI 98E data on $e^+ e^- \rightarrow \pi^+ \pi^- \ell^+ \ell^-$. Assuming that $\Gamma(\pi^+ \pi^- J/\psi)$ of $\chi_{c1}(3872)$ is the same as that of $\psi(2S)$ (85.4 keV).

$\chi_{c1}(3872) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$ $\Gamma(\pi^+\pi^- J/\psi(1S)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_2\Gamma_1/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
< 0.13	90	ABLIKIM	15V BES3	4.0–4.4 $e^+e^- \rightarrow \pi^+\pi^- J/\psi$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
< 6.2	90	^{1,2} AUBERT	05D BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
< 8.3	90	² DOBBS	05 CLE3	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$
< 10	90	³ YUAN	04 RVUE	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$

¹ Using $B(\chi_{c1}(3872) \rightarrow J/\psi\pi^+\pi^-) \cdot B(J/\psi \rightarrow \mu^+\mu^-) \cdot \Gamma(\chi_{c1}(3872) \rightarrow e^+e^-) < 0.37$ eV from AUBERT 05D and $B(J/\psi \rightarrow \mu^+\mu^-) = 0.0588 \pm 0.0010$ from the PDG 04.

² Assuming $\chi_{c1}(3872)$ has $J^{PC} = 1^{--}$.

³ Using BAI 98E data on $e^+e^- \rightarrow \pi^+\pi^-\ell^+\ell^-$. From theoretical calculation of the production cross section and using $B(J/\psi \rightarrow \mu^+\mu^-) = (5.88 \pm 0.10)\%$.

 $\chi_{c1}(3872) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$ $\Gamma(\pi^+\pi^- J/\psi(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_2\Gamma_7/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
< 12.9	90	¹ DOBBS	05 CLE3	$e^+e^- \rightarrow \pi^+\pi^- J/\psi\gamma$

¹ Assuming $\chi_{c1}(3872)$ has positive C parity and spin 0.

 $\Gamma(\omega J/\psi(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_4\Gamma_7/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
< 1.7	90	¹ LEES	12AD BABR	$e^+e^- \rightarrow e^+e^-\omega J/\psi$

¹ Assuming $\chi_{c1}(3872)$ has spin 2.

 $\Gamma(\pi^+\pi^-\eta_c(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{14}\Gamma_7/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
< 11.1	90	LEES	12AE BABR	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\eta_c$

 $\chi_{c1}(3872)$ BRANCHING RATIOS $\Gamma(\pi^+\pi^- J/\psi(1S))/\Gamma_{\text{total}} \quad \Gamma_2/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
> 0.032	93 ± 17	¹ AUBERT	08Y BABR	$B \rightarrow \chi_{c1}(3872)K$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
seen	151	² BALA	15 BELL	$B \rightarrow \chi_{c1}(3872)K\pi$
> 0.05	30	³ AUBERT	05R BABR	$B^+ \rightarrow K^+\pi^+\pi^- J/\psi$
> 0.05	36 ± 7	⁴ CHOI	03 BELL	$B^+ \rightarrow K^+\pi^+\pi^- J/\psi$

¹ AUBERT 08Y reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (8.4 \pm 1.5 \pm 0.7) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) < 2.6 \times 10^{-4}$.

² BALA 15 reports $B(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi) \times B(B^0 \rightarrow \chi_{c1}(3872) K^+ \pi^-) = (7.9 \pm 1.3 \pm 0.4) \times 10^{-6}$ and $B(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi) \times B(B^+ \rightarrow \chi_{c1}(3872) K^0 \pi^+) = (10.6 \pm 3.0 \pm 0.9) \times 10^{-6}$.

³ Superseded by AUBERT 08Y. AUBERT 05R reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (1.28 \pm 0.41) \times 10^{-5}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) < 2.6 \times 10^{-4}$.

⁴ CHOI 03 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] / [B(B^+ \rightarrow \psi(2S) K^+)] / [B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)] = 0.063 \pm 0.012 \pm 0.007$ which we multiply or divide by our best values $B(B^+ \rightarrow \chi_{c1}(3872) K^+) < 2.6 \times 10^{-4}$, $B(B^+ \rightarrow \psi(2S) K^+) = (6.21 \pm 0.23) \times 10^{-4}$, $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (34.67 \pm 0.30) \times 10^{-2}$.

$\Gamma(\omega J/\psi(1S))/\Gamma_{\text{total}}$

Γ_4/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
>0.023	21 ± 7	¹ DEL-AMO-SA..10B	BABR	$B^+ \rightarrow \omega J/\psi K^+$

¹ DEL-AMO-SANCHEZ 10B reports $[\Gamma(\chi_{c1}(3872) \rightarrow \omega J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (6 \pm 2 \pm 1) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) < 2.6 \times 10^{-4}$. DEL-AMO-SANCHEZ 10B also reports $B(B^0 \rightarrow \chi_{c1}(3872) K^0) \times B(\chi_{c1}(3872) \rightarrow J/\psi \omega) = (6 \pm 3 \pm 1) \times 10^{-6}$.

$\Gamma(\omega J/\psi(1S))/\Gamma(\pi^+ \pi^- J/\psi(1S))$

Γ_4/Γ_2

VALUE	DOCUMENT ID	TECN	COMMENT
0.8 ± 0.3	¹ DEL-AMO-SA..10B	BABR	$B \rightarrow \omega J/\psi K$

¹ Statistical and systematic errors added in quadrature. Uses the values of $B(B \rightarrow \chi_{c1}(3872) K) \times B(\chi_{c1}(3872) \rightarrow J/\psi \pi^+ \pi^-)$ reported in AUBERT 08Y, taking into account the common systematics.

$\Gamma(D^0 \bar{D}^0 \pi^0)/\Gamma_{\text{total}}$

Γ_5/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
>0.4	17 ± 5	¹ GOKHROO 06	BELL	$B^+ \rightarrow D^0 \bar{D}^0 \pi^0 K^+$

¹ GOKHROO 06 reports $[\Gamma(\chi_{c1}(3872) \rightarrow D^0 \bar{D}^0 \pi^0)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (1.02 \pm 0.31^{+0.21}_{-0.29}) \times 10^{-4}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) < 2.6 \times 10^{-4}$.

$\Gamma(D^0 \bar{D}^0 \pi^0)/\Gamma(\pi^+ \pi^- J/\psi(1S))$

Γ_5/Γ_2

VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ GOKHROO 06	BELL	$B \rightarrow D^0 \bar{D}^0 \pi^0 K$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	AUSHEV 10	BELL	$B \rightarrow D^0 \bar{D}^0 \pi^0 K$
------	-----------	------	---------------------------------------

¹ May not necessarily be the same state as that observed in the $J/\psi \pi^+ \pi^-$ mode. Supersedes CHISTOV 04.

$\Gamma(\bar{D}^{*0} D^0)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
>0.30	41^{+9}_{-8}	¹ AUSHEV	10	BELL $B^+ \rightarrow D^{*0} \bar{D}^0 K^+$

• • • We do not use the following data for averages, fits, limits, etc. • • •

>0.6	27 ± 6	² AUBERT	08B	BABR $B^+ \rightarrow \bar{D}^{*0} D^0 K^+$
¹ AUSHEV 10 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \bar{D}^{*0} D^0)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (0.77 \pm 0.16 \pm 0.10) \times 10^{-4}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) < 2.6 \times 10^{-4}$.				
² AUBERT 08B reports $[\Gamma(\chi_{c1}(3872) \rightarrow \bar{D}^{*0} D^0)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (1.67 \pm 0.36 \pm 0.47) \times 10^{-4}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) < 2.6 \times 10^{-4}$.				

 $\Gamma(D^0 \bar{D}^0)/\Gamma(\pi^+ \pi^- J/\psi(1S))$ Γ_8/Γ_2

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	CHISTOV	04	BELL $B \rightarrow K D^0 \bar{D}^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $\Gamma(D^+ D^-)/\Gamma(\pi^+ \pi^- J/\psi(1S))$ Γ_9/Γ_2

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	CHISTOV	04	BELL $B \rightarrow K D^+ D^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $\Gamma(\gamma \chi_{c1})/\Gamma(\pi^+ \pi^- J/\psi(1S))$ Γ_{10}/Γ_2

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
not seen		¹ BHARDWAJ	13	BELL $B^+ \rightarrow \chi_{c1} \gamma K^+$
<0.89	90	CHOI	03	BELL $B \rightarrow K \pi^+ \pi^- J/\psi$

¹ Reported $B(B^\pm \rightarrow \chi_{c1}(3872) K^\pm) \times B(\chi_{c1}(3872) \rightarrow \gamma \chi_{c1}) < 1.9 \times 10^{-6}$ at 90% CL.

 $\Gamma(\gamma \chi_{c2})/\Gamma(\pi^+ \pi^- J/\psi(1S))$ Γ_{11}/Γ_2

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	¹ BHARDWAJ	13	BELL $B^\pm \rightarrow \chi_{c2} \gamma K^\pm$

¹ Reported $B(B^\pm \rightarrow \chi_{c1}(3872) K^\pm) \times B(\chi_{c1}(3872) \rightarrow \gamma \chi_{c2}) < 6.7 \times 10^{-6}$ at 90% CL.

 $\Gamma(\gamma J/\psi)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
>7 $\times 10^{-3}$		¹ BHARDWAJ	11	BELL $B^\pm \rightarrow \gamma J/\psi K^\pm$

• • • We do not use the following data for averages, fits, limits, etc. • • •

>0.011	20	² AUBERT	09B	BABR $B^+ \rightarrow \gamma J/\psi K^+$
>0.013	19	³ AUBERT,BE	06M	BABR $B^+ \rightarrow \gamma J/\psi K^+$
¹ BHARDWAJ 11 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma J/\psi)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (1.78^{+0.48}_{-0.44} \pm 0.12) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) < 2.6 \times 10^{-4}$.				

² AUBERT 09B reports $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma J/\psi)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (2.8 \pm 0.8 \pm 0.1) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) < 2.6 \times 10^{-4}$.

³ Superseded by AUBERT 09B. AUBERT, BE 06M reports $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma J/\psi)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (3.3 \pm 1.0 \pm 0.3) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) < 2.6 \times 10^{-4}$.

 $\Gamma(\gamma\psi(2S))/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	36 ± 9	¹ AAIJ	14AH LHCB	$B^+ \rightarrow \gamma\psi(2S) K^+$
>0.04	25 ± 7	² AUBERT	09B BABR	$B^+ \rightarrow \gamma\psi(2S) K^+$

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen ³ BHARDWAJ 11 BELL $B^+ \rightarrow \gamma\psi(2S) K^+$

¹ From 36.4 ± 9.0 events of $\chi_{c1}(3872) \rightarrow J/\psi\gamma$ decays with a statistical significance of 4.4σ .

² AUBERT 09B reports $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma\psi(2S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (9.5 \pm 2.7 \pm 0.6) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) < 2.6 \times 10^{-4}$.

³ BHARDWAJ 11 reports $B(B^+ \rightarrow K^+ \chi_{c1}(3872)) \times B(\chi_{c1} \rightarrow \gamma\psi(2S)) < 3.45 \times 10^{-6}$ at 90% CL.

 $\Gamma(\gamma\psi(2S))/\Gamma(\gamma J/\psi)$ Γ_{13}/Γ_{12}

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.6 ± 0.6 OUR AVERAGE					
$2.46 \pm 0.64 \pm 0.29$		36 ± 9	¹ AAIJ	14AH LHCB	$B^+ \rightarrow \gamma\psi(2S) K^+$
3.4 ± 1.4			AUBERT	09B BABR	$B^+ \rightarrow \gamma c\bar{c} K^+$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.1 90 BHARDWAJ 11 BELL $B^+ \rightarrow \gamma\psi(2S) K^+$

¹ From 36.4 ± 9.0 events of $\chi_{c1}(3872) \rightarrow J/\psi\gamma$ decays with a statistical significance of 4.4σ .

 $\Gamma(\pi^+ \pi^- \chi_{c1})/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	¹ BHARDWAJ 16	BELL	$B^+ \rightarrow \pi^+ \pi^- \chi_{c1} K^+$

¹ BHARDWAJ 16 quotes $B(B^+ \rightarrow \chi_{c1}(3872) K^+) \cdot B(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- \chi_{c1}) < 1.5 \times 10^{-6}$ at 90% CL.

 $\Gamma(p\bar{p})/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	¹ AAIJ	17AD LHCB	$pp \rightarrow B^+ X \rightarrow p\bar{p} K^+ X$

¹ AAIJ 17AD reports $B(B^+ \rightarrow \chi_{c1}(3872) K^+ \rightarrow p\bar{p} K^+)/B(B^+ \rightarrow J/\psi K^+ \rightarrow p\bar{p} K^+) < 2.0 (2.5) \times 10^{-3}$ at 90% (95%) CL.

 $\Gamma(p\bar{p})/\Gamma(\pi^+ \pi^- J/\psi(1S))$ Γ_{16}/Γ_2

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.0 $\times 10^{-3}$	95	¹ AAIJ	13S LHCB	$B^+ \rightarrow p\bar{p} K^+$

¹ AAIJ 13S reports $[\Gamma(\chi_{c1}(3872) \rightarrow p\bar{p})/\Gamma(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S))] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+, \chi_{c1} \rightarrow J/\psi \pi^+ \pi^-)] < 1.7 \times 10^{-8}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+, \chi_{c1} \rightarrow J/\psi \pi^+ \pi^-) = 8.6 \times 10^{-6}$.

C-violating decays

 $\Gamma(\eta J/\psi)/\Gamma(\pi^+\pi^- J/\psi(1S))$ Γ_{17}/Γ_2

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.4	90	1, ² IWASHITA	14	BELL $B \rightarrow K\eta J/\psi$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<0.6	90	AUBERT	04Y	BABR $B \rightarrow K\eta J/\psi$
¹ IWASHITA 14 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \eta J/\psi)/\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi(1S))] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+, \chi_{c1} \rightarrow J/\psi\pi^+\pi^-)] < 3.8 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+, \chi_{c1} \rightarrow J/\psi\pi^+\pi^-) = 8.6 \times 10^{-6}$.				
² IWASHITA 14 also scans the $\eta J/\psi$ mass range 3.8–4.75 GeV and sets upper limits for $B(B^\pm \rightarrow \chi_{c1}(3872)K^\pm) \times B(\chi_{c1}(3872) \rightarrow \eta J/\psi)$ in 5 MeV intervals.				

 $\chi_{c1}(3872)$ REFERENCES

AAIJ	17AD	PL B769 305	R. Aaij <i>et al.</i>	(LHCb Collab.)
BHARDWAJ	16	PR D93 052016	V. Bhardwaj <i>et al.</i>	(BELLE Collab.)
AAIJ	15AO	PR D92 011102	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	15V	PL B749 414	M. Ablikim <i>et al.</i>	(BES III Collab.)
BALA	15	PR D91 051101	A. Bala <i>et al.</i>	(BELLE Collab.)
AAIJ	14AH	NP B886 665	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	14	PRL 112 092001	M. Ablikim <i>et al.</i>	(BES III Collab.)
IWASHITA	14	PTEP 2014 043C01	T. Iwashita <i>et al.</i>	(BELLE Collab.)
AAIJ	13Q	PRL 110 222001	R. Aaij <i>et al.</i>	(LHCb Collab.) JP
AAIJ	13S	EPJ C73 2462	R. Aaij <i>et al.</i>	(LHCb Collab.)
BHARDWAJ	13	PRL 111 032001	V. Bhardwaj <i>et al.</i>	(BELLE Collab.)
AAIJ	12H	EPJ C72 1972	R. Aaij <i>et al.</i>	(LHCb Collab.)
LEES	12AD	PR D86 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	12AE	PR D86 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
BHARDWAJ	11	PRL 107 091803	V. Bhardwaj <i>et al.</i>	(BELLE Collab.)
CHOI	11	PR D84 052004	S.-K. Choi <i>et al.</i>	(BELLE Collab.)
AUSHEV	10	PR D81 031103	T. Aushev <i>et al.</i>	(BELLE Collab.)
DEL-AMO-SA...	10B	PR D82 011101	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
AALTONEN	09AU	PRL 103 152001	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AUBERT	09B	PRL 102 132001	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08B	PR D77 011102	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08Y	PR D77 111101	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	06	PR D73 011101	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	06M	PR D74 071101	B. Aubert <i>et al.</i>	(BABAR Collab.)
GOKHROO	06	PRL 97 162002	G. Gokhroo <i>et al.</i>	(BELLE Collab.)
AUBERT	05B	PR D71 031501	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	05D	PR D71 052001	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	05R	PR D71 071103	B. Aubert <i>et al.</i>	(BABAR Collab.)
DOBBS	05	PRL 94 032004	S. Dobbs <i>et al.</i>	(CLEO Collab.)
ABAZOV	04F	PRL 93 162002	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ACOSTA	04	PRL 93 072001	D. Acosta <i>et al.</i>	(CDF Collab.)
AUBERT	04Y	PRL 93 041801	B. Aubert <i>et al.</i>	(BABAR Collab.)
CHISTOV	04	PRL 93 051803	R. Chistov <i>et al.</i>	(BELLE Collab.)
PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)
YUAN	04	PL B579 74	C.Z. Yuan <i>et al.</i>	
CHOI	03	PRL 91 262001	S.-K. Choi <i>et al.</i>	(BELLE Collab.)
BAI	98E	PR D57 3854	J.Z. Bai <i>et al.</i>	(BES Collab.)
ANTONIAZZI	94	PR D50 4258	L. Antoniazzi <i>et al.</i>	(E705 Collab.)
