

$\omega(1650)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

 $\omega(1650)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1670 ± 30 OUR ESTIMATE				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1671 ± 6 ± 10	824	¹ AKHMETSHIN 17A	CMD3	1.4–2.0 $e^+e^- \rightarrow \omega\eta$
1660 ± 10	898	² ACHASOV 16B	SND	1.34–2.00 $e^+e^- \rightarrow \omega\eta$
1680 ± 10	13.1k	³ AULCHENKO 15A	SND	1.05–1.80 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
1667 ± 13 ± 6		AUBERT 07AU	BABR	10.6 $e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$
1645 ± 8	13	AUBERT 06D	BABR	10.6 $e^+e^- \rightarrow \omega\eta\gamma$
1660 ± 10 ± 2		AUBERT,B 04N	BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
1770 ± 50 ± 60	1.2M	⁴ ACHASOV 03D	RVUE	0.44–2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
1619 ± 5		⁵ HENNER 02	RVUE	1.2–2.0 $e^+e^- \rightarrow \rho\pi,$ $\omega\pi\pi$
1700 ± 20		EUGENIO 01	SPEC	18 $\pi^-p \rightarrow \omega\eta n$
1705 ± 26	612	⁶ AKHMETSHIN 00D	CMD2	$e^+e^- \rightarrow \omega\pi^+\pi^-$
1820 ⁺¹⁹⁰ _{–150}		⁷ ACHASOV 98H	RVUE	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
1840 ⁺¹⁰⁰ _{–70}		⁸ ACHASOV 98H	RVUE	$e^+e^- \rightarrow \omega\pi^+\pi^-$
1780 ⁺¹⁷⁰ _{–300}		⁹ ACHASOV 98H	RVUE	$e^+e^- \rightarrow K^+K^-$
~ 2100		¹⁰ ACHASOV 98H	RVUE	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$
1606 ± 9		¹¹ CLEGG 94	RVUE	
1662 ± 13	750	¹² ANTONELLI 92	DM2	1.34–2.4 $e^+e^- \rightarrow \rho\pi,$ $\omega\pi\pi$
1670 ± 20		ATKINSON 83B	OMEG	20–70 $\gamma p \rightarrow 3\pi X$
1657 ± 13		CORDIER 81	DM1	$e^+e^- \rightarrow \omega 2\pi$
1679 ± 34	21	ESPOSITO 80	FRAM	$e^+e^- \rightarrow 3\pi$
1652 ± 17		COSME 79	OSPK	$e^+e^- \rightarrow 3\pi$

¹ From a fit of the interfering $\omega(1420)$ and $\omega(1650)$ with a relative phase of π and other parameters floating.

² From a fit with contributions from $\omega(1420)$, $\omega(1650)$, and $\phi(1680)$.

³ From a fit with contributions from $\omega(782)$, $\phi(1020)$, $\omega(1420)$, and $\omega(1650)$.

⁴ From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+\pi^-\pi^0$ and ANTONELLI 92 on the $\omega\pi^+\pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.

⁵ Using results of CORDIER 81 and preliminary data of DOLINSKY 91 and ANTONELLI 92.

⁶ Using the data of AKHMETSHIN 00D and ANTONELLI 92. The $\rho\pi$ dominance for the energy dependence of the $\omega(1420)$ and $\omega(1650)$ width assumed.

⁷ Using data from BARKOV 87, DOLINSKY 91, and ANTONELLI 92.

⁸ Using the data from ANTONELLI 92.

⁹ Using the data from IVANOV 81 and BISELLO 88B.

¹⁰ Using the data from BISELLO 91C.

¹¹ From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.

¹²From the combined fit of the $\rho\pi$ and $\omega\pi\pi$ final states. **$\omega(1650)$ WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
315 ± 35 OUR ESTIMATE				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
113 ± 9 ± 10	824	¹ AKHMETSHIN 17A	CMD3	1.4–2.0 $e^+e^- \rightarrow \omega\eta$
110 ± 20	898	² ACHASOV 16B	SND	1.34–2.00 $e^+e^- \rightarrow \omega\eta$
310 ± 30	13.1k	³ AULCHENKO 15A	SND	1.05–1.80 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
222 ± 25 ± 20		AUBERT 07AU	BABR	10.6 $e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$
114 ± 14	13	AUBERT 06D	BABR	10.6 $e^+e^- \rightarrow \omega\eta\gamma$
230 ± 30 ± 20		AUBERT,B 04N	BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
490 ⁺²⁰⁰ ₋₁₅₀ ± 130	1.2M	⁴ ACHASOV 03D	RVUE	0.44–2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
250 ± 14		⁵ HENNER 02	RVUE	1.2–2.0 $e^+e^- \rightarrow \rho\pi, \omega\pi\pi$
250 ± 50		EUGENIO 01	SPEC	18 $\pi^-p \rightarrow \omega\eta n$
370 ± 25	612	⁶ AKHMETSHIN 00D	CMD2	$e^+e^- \rightarrow \omega\pi^+\pi^-$
113 ± 20		⁷ CLEGG 94	RVUE	
280 ± 24	750	⁸ ANTONELLI 92	DM2	1.34–2.4 $e^+e^- \rightarrow \rho\pi, \omega\pi\pi$
160 ± 20		ATKINSON 83B	OMEG	20–70 $\gamma p \rightarrow 3\pi X$
136 ± 46		CORDIER 81	DM1	$e^+e^- \rightarrow \omega 2\pi$
99 ± 49	21	ESPOSITO 80	FRAM	$e^+e^- \rightarrow 3\pi$
42 ± 17		COSME 79	OSPK	$e^+e^- \rightarrow 3\pi$

¹From a fit of the interfering $\omega(1420)$ and $\omega(1650)$ with a relative phase of π and other parameters floating.²From a fit with contributions from $\omega(1420)$, $\omega(1650)$, and $\phi(1680)$.³From a fit with contributions from $\omega(782)$, $\phi(1020)$, $\omega(1420)$, and $\omega(1650)$.⁴From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+\pi^-\pi^0$ and ANTONELLI 92 on the $\omega\pi^+\pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.⁵Using results of CORDIER 81 and preliminary data of DOLINSKY 91 and ANTONELLI 92.⁶Using the data of AKHMETSHIN 00D and ANTONELLI 92. The $\rho\pi$ dominance for the energy dependence of the $\omega(1420)$ and $\omega(1650)$ width assumed.⁷From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.⁸From the combined fit of the $\rho\pi$ and $\omega\pi\pi$ final states. **$\omega(1650)$ DECAY MODES**

Mode	Fraction (Γ_i/Γ)
Γ_1 $\rho\pi$	seen
Γ_2 $\omega\pi\pi$	seen
Γ_3 $\omega\eta$	seen
Γ_4 e^+e^-	seen

$\omega(1650) \Gamma(i)\Gamma(e^+e^-)/\Gamma^2(\text{total})$ $\Gamma(\rho\pi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_1/\Gamma \times \Gamma_4/\Gamma$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1.56 ± 0.23	13.1k	¹ AULCHENKO	15A SND	$1.05\text{--}1.80 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
$1.3 \pm 0.1 \pm 0.1$		AUBERT,B	04N BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
$1.2 \begin{smallmatrix} +0.4 \\ -0.1 \end{smallmatrix} \pm 0.8$	1.2M	^{2,3} ACHASOV	03D RVUE	$0.44\text{--}2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.921 ± 0.230		^{4,5} CLEGG	94 RVUE	
0.479 ± 0.050	750	^{6,7} ANTONELLI	92 DM2	$1.34\text{--}2.4 e^+e^- \rightarrow \rho\pi, \omega\pi\pi$

 $\Gamma(\omega\pi\pi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_2/\Gamma \times \Gamma_4/\Gamma$

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
7.0 ± 0.5		AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$
$4.1 \pm 0.9 \pm 1.3$	1.2M	^{2,3} ACHASOV	03D RVUE	$0.44\text{--}2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
5.40 ± 0.95		⁸ AKHMETSHIN	00D CMD2	$1.2\text{--}1.38 e^+e^- \rightarrow \omega\pi^+\pi^-$
3.18 ± 0.80		^{4,5} CLEGG	94 RVUE	
6.07 ± 0.61	750	^{6,7} ANTONELLI	92 DM2	$1.34\text{--}2.4 e^+e^- \rightarrow \rho\pi, \omega\pi\pi$

 $\Gamma(\omega\eta)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_3/\Gamma \times \Gamma_4/\Gamma$

VALUE (units 10^{-7})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
$4.5 \pm 0.3 \pm 0.3$		824	⁹ AKHMETSHIN	17A CMD3	$1.4\text{--}2.0 e^+e^- \rightarrow \omega\eta$
4.4 ± 0.5		898	¹⁰ ACHASOV	16B SND	$1.34\text{--}2.00 e^+e^- \rightarrow \omega\eta$
5.7 ± 0.6		13	AUBERT	06D BABR	$10.6 e^+e^- \rightarrow \omega\eta\gamma$
<60	90		¹¹ AKHMETSHIN	03B CMD2	$e^+e^- \rightarrow \eta\pi^0\gamma$

¹ From a fit with contributions from $\omega(782)$, $\phi(1020)$, $\omega(1420)$, and $\omega(1650)$.² Calculated by us from the cross section at the peak.³ From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+\pi^-\pi^0$ and ANTONELLI 92 on the $\omega\pi^+\pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.⁴ From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.⁵ From the partial and leptonic width given by the authors.⁶ From the combined fit of the $\rho\pi$ and $\omega\pi\pi$ final states.⁷ From the product of the leptonic width and partial branching ratio given by the authors.⁸ Using the data of AKHMETSHIN 00D and ANTONELLI 92. The $\rho\pi$ dominance for the energy dependence of the $\omega(1420)$ and $\omega(1650)$ width assumed.⁹ From a fit of the interfering $\omega(1420)$ and $\omega(1650)$ with a relative phase of π and other parameters floating. From an alternative fit $\Gamma(\omega(1650) \rightarrow \omega\eta)/\Gamma_{\text{total}} \times \Gamma(\omega(1650) \rightarrow e^+e^-) = 51 \pm 3$ eV.¹⁰ From a fit with contributions from $\omega(1420)$, $\omega(1650)$, and $\phi(1680)$.¹¹ $\omega(1650)$ mass and width fixed at 1700 MeV and 250 MeV, respectively.

$\omega(1650)$ BRANCHING RATIOS **$\Gamma(\omega\pi\pi)/\Gamma_{\text{total}}$ Γ_2/Γ**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
~ 0.35	1.2M	¹ ACHASOV	03D	RVUE 0.44–2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.620 ± 0.014		² HENNER	02	RVUE 1.2–2.0 $e^+e^- \rightarrow \rho\pi, \omega\pi\pi$

 $\Gamma(\rho\pi)/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
~ 0.65	1.2M	¹ ACHASOV	03D	RVUE 0.44–2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.380 ± 0.014		² HENNER	02	RVUE 1.2–2.0 $e^+e^- \rightarrow \rho\pi, \omega\pi\pi$

 $\Gamma(e^+e^-)/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
~ 18	1.2M	^{2,3} ACHASOV	03D	RVUE 0.44–2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
32 ± 1		² HENNER	02	RVUE 1.2–2.0 $e^+e^- \rightarrow \rho\pi, \omega\pi\pi$

¹From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+\pi^-\pi^0$ and ANTONELLI 92 on the $\omega\pi^+\pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.

²Assuming that the $\omega(1650)$ decays into $\rho\pi$ and $\omega\pi\pi$ only.

³Calculated by us from the cross section at the peak.

 $\omega(1650)$ REFERENCES

AKHMETSHIN 17A	PL B773 150	R.R. Akhmetshin <i>et al.</i>	(CMD-3 Collab.)
ACHASOV 16B	PR D94 092002	M.N. Achasov <i>et al.</i>	(SND Collab.)
AULCHENKO 15A	JETP 121 27	V.M. Aulchenko <i>et al.</i>	(SND Collab.)
	Translated from ZETF 148 34.		
AUBERT 07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT 06D	PR D73 052003	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B 04N	PR D70 072004	B. Aubert <i>et al.</i>	(BABAR Collab.)
ACHASOV 03D	PR D68 052006	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
AKHMETSHIN 03B	PL B562 173	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
ACHASOV 02E	PR D66 032001	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
HENNER 02	EPJ C26 3	V.K. Henner <i>et al.</i>	
ACHASOV 01E	PR D63 072002	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
EUGENIO 01	PL B497 190	P. Eugenio <i>et al.</i>	
AKHMETSHIN 00D	PL B489 125	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
ACHASOV 99E	PL B462 365	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV 98H	PR D57 4334	N.N. Achasov, A.A. Kozhevnikov	
CLEGG 94	ZPHY C62 455	A.B. Clegg, A. Donnachie	(LANC, MCHS)
ANTONELLI 92	ZPHY C56 15	A. Antonelli <i>et al.</i>	(DM2 Collab.)
BISELLO 91C	ZPHY C52 227	D. Bisello <i>et al.</i>	(DM2 Collab.)
DOLINSKY 91	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)
BISELLO 88B	ZPHY C39 13	D. Bisello <i>et al.</i>	(PADO, CLER, FRAS+)
BARKOV 87	JETPL 46 164	L.M. Barkov <i>et al.</i>	(NOVO)
	Translated from ZETFP 46 132.		
ATKINSON 83B	PL 127B 132	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
CORDIER 81	PL 106B 155	A. Cordier <i>et al.</i>	(ORSAY)
IVANOV 81	PL 107B 297	P.M. Ivanov <i>et al.</i>	(NOVO)
ESPOSITO 80	LNC 28 195	B. Esposito <i>et al.</i>	(FRAS, NAPL, PADO+)
COSME 79	NP B152 215	G. Cosme <i>et al.</i>	(IPN)