

$D_{s0}^*(2317)^\pm$
 $I(J^P) = 0(0^+)$
 J, P need confirmation.

AUBERT 06P and CHOI 15A do not observe neutral and doubly charged partners of the $D_{s0}^*(2317)^+$.

 $D_{s0}^*(2317)^\pm$ MASS

The fit includes $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2317.8±0.5 OUR FIT				
2318.0±0.7 OUR AVERAGE				
2318.3±1.2±1.2	115	1 ABLIKIM	18J BES3	$4.6 e^+ e^- \rightarrow D_s^{*\pm} D_{s0}^*(2317)^\mp$
2319.6±0.2±1.4	3.1k	AUBERT	06P BABR	$10.6 e^+ e^- \rightarrow D_s^+ \pi^0 X$
2317.3±0.4±0.8	1.0k	2 AUBERT	04E BABR	$10.6 e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2317.2±1.3	88	3 AUBERT,B	04S BABR	$B \rightarrow D_{s0}^{(*)}(2317)^+ \bar{D}^{(*)}$
2317.2±0.5±0.9	761	4 MIKAMI	04 BELL	$10.6 e^+ e^-$
2316.8±0.4±3.0	1.2k	4,5 AUBERT	03G BABR	$10.6 e^+ e^-$
2317.6±1.3	273	4,6 AUBERT	03G BABR	$10.6 e^+ e^-$
2319.8±2.1±2.0	24	4 KROKOVNY	03B BELL	$10.6 e^+ e^-$

¹ From a fit of the D_s^* recoil mass where the $D_{s0}^*(2317)$ signal is described with a Crystal Ball function convolved with a Gaussian function.

² Supersedes AUBERT 03G.

³ Systematic errors not evaluated.

⁴ Not independent of the corresponding $m_{D_{s0}^*(2317)} - m_{D_s}$.

⁵ From $D_s^+ \rightarrow K^+ K^- \pi^+$ decay.

⁶ From $D_s^+ \rightarrow K^+ K^- \pi^+ \pi^0$ decay.

 $m_{D_{s0}^*(2317)^\pm} - m_{D_s^\pm}$

The fit includes $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
349.4±0.5 OUR FIT				
349.2±0.7 OUR AVERAGE				
348.7±0.5±0.7	761	MIKAMI	04 BELL	$10.6 e^+ e^-$
350.0±1.2±1.0	135	BESSION	03 CLE2	$10.6 e^+ e^-$
351.3±2.1±1.9	24	7 KROKOVNY	03B BELL	$10.6 e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
349.6±0.4±3.0	1267	8,9 AUBERT	03G BABR	$10.6 e^+ e^-$
350.2±1.3	273	10,11 AUBERT	03G BABR	$10.6 e^+ e^-$

⁷ Recalculated by us using $m_{D_s^+} = 1968.5 \pm 0.6$ MeV.

⁸ From $D_s^+ \rightarrow K^+ K^- \pi^+$ decay.

⁹ Recalculated by us using $m_{D_s^+} = 1967.20 \pm 0.03$ MeV.

¹⁰ From $D_s^+ \rightarrow K^+ K^- \pi^+ \pi^0$ decay.

¹¹ Recalculated by us using $m_{D_s^+} = 1967.4 \pm 0.2$ MeV. Systematic errors not estimated.

$D_{s0}^*(2317)^\pm$ WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 3.8	95	3180	AUBERT	06P	BABR $10.6 e^+ e^- \rightarrow D_s^+ \pi^0 X$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
< 4.6	90	761	MIKAMI	04	BELL $10.6 e^+ e^-$
< 10			AUBERT	03G	BABR $10.6 e^+ e^-$
< 7	90	135	BESSON	03	CLE2 $10.6 e^+ e^-$

$D_{s0}^*(2317)^\pm$ DECAY MODES

$D_{s0}^*(2317)^-$ modes are charge conjugates of modes below.

Mode	Fraction (Γ_i/Γ)	Confidence level
$\Gamma_1 D_s^+ \pi^0$	(100^{+0}_{-20}) %	
$\Gamma_2 D_s^+ \gamma$	< 5 %	90%
$\Gamma_3 D_s^*(2112)^+ \gamma$	< 6 %	90%
$\Gamma_4 D_s^+ \gamma\gamma$	< 18 %	95%
$\Gamma_5 D_s^*(2112)^+ \pi^0$	< 11 %	90%
$\Gamma_6 D_s^+ \pi^+ \pi^-$	< 4 $\times 10^{-3}$	90%
$\Gamma_7 D_s^+ \pi^0 \pi^0$	not seen	

$D_{s0}^*(2317)^\pm$ BRANCHING RATIOS

$\Gamma(D_s^+ \pi^0)/\Gamma_{\text{total}}$	EVTS	DOCUMENT ID	TECN	Γ_1/Γ
$1.00^{+0.00}_{-0.14} {}^{+0.00}_{-0.14}$	47	ABLIKIM	18J	BES3 $4.6 e^+ e^- \rightarrow D_s^* \pm D_{s0}^*(2317)^\mp$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
seen	1.5k	AUBERT	03G	BABR $10.6 e^+ e^-$

$\Gamma(D_s^+ \gamma)/\Gamma(D_s^+ \pi^0)$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_2/Γ_1
<0.05	90	MIKAMI	04	BELL 10.6 e ⁺ e ⁻	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.14	95	AUBERT	06P	BABR 10.6 e ⁺ e ⁻	
<0.052	90	BESSION	03	CLE2 10.6 e ⁺ e ⁻	

 $\Gamma(D_s^*(2112)^+ \gamma)/\Gamma(D_s^+ \pi^0)$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_3/Γ_1
<0.059	90	BESSION	03	CLE2 10.6 e ⁺ e ⁻	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.16	95	AUBERT	06P	BABR 10.6 e ⁺ e ⁻	
<0.18	90	MIKAMI	04	BELL 10.6 e ⁺ e ⁻	

 $\Gamma(D_s^+ \gamma\gamma)/\Gamma(D_s^+ \pi^0)$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_4/Γ_1
<0.18	95	AUBERT	06P	BABR 10.6 e ⁺ e ⁻	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
not seen		AUBERT	03G	BABR 10.6 e ⁺ e ⁻	

 $\Gamma(D_s^*(2112)^+ \pi^0)/\Gamma(D_s^+ \pi^0)$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_5/Γ_1
<0.11	90	BESSION	03	CLE2 10.6 e ⁺ e ⁻	

 $\Gamma(D_s^+ \pi^+ \pi^-)/\Gamma(D_s^+ \pi^0)$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_6/Γ_1
<0.004	90	MIKAMI	04	BELL 10.6 e ⁺ e ⁻	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.005	95	AUBERT	06P	BABR 10.6 e ⁺ e ⁻	
<0.019	90	BESSION	03	CLE2 10.6 e ⁺ e ⁻	

 $\Gamma(D_s^+ \pi^0 \pi^0)/\Gamma(D_s^+ \pi^0)$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_7/Γ_1
<0.25	95	AUBERT	06P	BABR 10.6 e ⁺ e ⁻	

 $D_{s0}^*(2317)^\pm$ REFERENCES

ABLIKIM	18J	PR D97 051103	M. Ablikim <i>et al.</i>	(BESIII Collab.)
CHOI	15A	PR D91 092011	S.-K. Choi <i>et al.</i>	(BELLE Collab.)
AUBERT	06P	PR D74 032007	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	04E	PR D69 031101	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04S	PRL 93 181801	B. Aubert <i>et al.</i>	(BABAR Collab.)
MIKAMI	04	PRL 92 012002	Y. Mikami <i>et al.</i>	(BELLE Collab.)
AUBERT	03G	PRL 90 242001	B. Aubert <i>et al.</i>	(BABAR Collab.)
BESSION	03	PR D68 032002	D. Besson <i>et al.</i>	(CLEO Collab.)
KROKOVNY	03B	PRL 91 262002	P. Krokovny <i>et al.</i>	(BELLE Collab.)