Heavy Charged Lepton Searches

Charged Heavy Lepton MASS LIMITS

Sequential Charged Heavy Lepton (L[±]) MASS LIMITS

These experiments assumed that a fourth generation L^\pm decayed to a fourth generation ν_L (or L^0) where ν_L was stable, or that L^\pm decays to a light ν_ℓ via mixing.

See the "Quark and Lepton Compositeness, Searches for" Listings for limits on radiatively decaying excited leptons, i.e. $\ell^* \to \ell \gamma$. See the "WIMPs and other Particle Searches" section for heavy charged particle search limits in which the charged particle could be a lepton.

| VALUE (GeV) | CL% | DOCUMENT ID | DOCUMENT ID TECN | | COMMENT | |
|---|-----|-----------------------|------------------|------|---|--|
| >100.8 | 95 | ACHARD | 01 B | L3 | Decay to νW | |
| >101.9 | 95 | ACHARD | 01 B | L3 | $m_L - m_{I0} > 15 \text{ GeV}$ | |
| • • • We do not use the following data for averages, fits, limits, etc. • • | | | | | _ | |
| > 81.5 | 95 | ACKERSTAFF | 98 C | OPAL | Assumed $m_{L^{\pm}} - m_{L^{0}} > 8.4$ GeV | |
| > 80.2 | 95 | ACKERSTAFF | 98C | OPAL | $m_{L^0}^{\rm GeV} > m_{L^\pm}^{\rm and} L^\pm \rightarrow \nu W$ | |
| < 48 or $>$ 61 | 95 | ¹ ACCIARRI | 96G | L3 | | |
| > 63.9 | 95 | ALEXANDER | 96 P | OPAL | Decay to massless $ u$'s | |
| > 63.5 | 95 | BUSKULIC | 96 S | ALEP | $m_L - m_{I0} > 7 \text{ GeV}$ | |
| > 65 | 95 | BUSKULIC | 96 S | ALEP | Decay to massless ν 's | |
| none 10-225 | | ² AHMED | 94 | CNTR | H1 Collab. at HERA | |
| none 12.6-29.6 | 95 | KIM | 91 B | AMY | Massless $ u$ assumed | |
| > 44.3 | 95 | AKRAWY | 90 G | OPAL | | |
| none 0.5–10 | 95 | ³ RILES | 90 | MRK2 | For $(m_{10} - m_{10}) > 0.25 - 0.4 \text{GeV}$ | |
| > 8 | | ⁴ STOKER | 89 | | For $(m_{I^+} - m_{I^0}) = 0.4 \text{ GeV}$ | |
| > 12 | | ⁴ STOKER | 89 | | For $m_{10} = 0.9 \text{ GeV}$ | |
| none 18.4-27.6 | 95 | ⁵ ABE | 88 | VNS | L | |
| > 25.5 | 95 | ⁶ ADACHI | 88 B | TOPZ | | |
| none 1.5-22.0 | 95 | BEHREND | 88C | CELL | | |
| > 41 | 90 | ⁷ ALBAJAR | 87 B | UA1 | | |
| > 22.5 | 95 | ⁸ ADEVA | 85 | MRKJ | | |
| > 18.0 | 95 | ⁹ BARTEL | 83 | JADE | | |
| none 4–14.5 | 95 | ¹⁰ BERGER | 81 B | PLUT | | |
| > 15.5 | 95 | 11 BRANDELIK | 81 | TASS | | |
| > 13. | | 12 AZIMOV | 80 | | | |
| > 16. | 95 | ¹³ BARBER | 80 B | CNTR | | |
| > 0.490 | | ¹⁴ ROTHE | 69 | RVUE | | |

 $^{^1}$ ACCIARRI 96G assumes LEP result that the associated neutral heavy lepton mass > 40 GeV.

Created: 6/1/2020 08:33

² The AHMED 94 limits are from a search for neutral and charged sequential heavy leptons at HERA via the decay channels $L^- \to e \gamma$, $L^- \to \nu W^-$, $L^- \to e Z$; and $L^0 \to \nu \gamma$, $L^0 \to e^- W^+$, $L^- \to \nu Z$, where the W decays to $\ell \nu_\ell$, or to jets, and Z decays to $\ell^+ \ell^-$ or jets.

³ RILES 90 limits were the result of a special analysis of the data in the case where the mass difference $m_{L^-} - m_{L^0}$ was allowed to be quite small, where L^0 denotes the neutrino

into which the sequential charged lepton decays. With a slightly reduced m_{L^\pm} range, the mass difference extends to about 4 GeV.

- ⁴STOKER 89 (Mark II at PEP) gives bounds on charged heavy lepton (L^+) mass for the generalized case in which the corresponding neutral heavy lepton (L^0) in the SU(2) doublet is not of negligible mass.
- 5 ABE 88 search for L^+ and $L^- o hadrons looking for acoplanar jets. The bound is valid for <math>m_
 u < 10$ GeV.
- 6 ADACHI 88B search for hadronic decays giving acoplanar events with large missing energy. ${\sf E_{cm}}^{ee}=$ 52 GeV.
- ⁷ Assumes associated neutrino is approximately massless.
- 8 ADEVA 85 analyze one-isolated-muon data and sensitive to τ $\,<\!10$ nanosec. Assume B(lepton) = 0.30. $E_{\rm cm}$ = 40–47 GeV.
- ⁹BARTEL 83 limit is from PETRA e^+e^- experiment with average $E_{\rm cm}=34.2$ GeV.
- 10 BERGER 81B is DESY DORIS and PETRA experiment. Looking for $e^+e^- \rightarrow L^+L^-$.
- ¹¹BRANDELIK 81 is DESY-PETRA experiment. Looking for $e^+e^- \rightarrow L^+L^-$.
- ¹² AZIMOV 80 estimated probabilities for M+N type events in $e^+e^- \rightarrow L^+L^-$ deducing semi-hadronic decay multiplicities of L from e^+e^- annihilation data at $E_{\rm cm}=(2/3)m_L$. Obtained above limit comparing these with e^+e^- data (BRANDELIK 80).
- 13 BARBER 80B looked for $e^+e^-
 ightarrow ~L^+L^-$, $L
 ightarrow ~\nu_L^+$ X with MARK-J at DESY-PETRA.

Stable Charged Heavy Lepton (L^{\pm}) MASS LIMITS

| <i>VALUE</i> (GeV) | CL% | DOCUMENT ID | | TECN |
|-------------------------|-------------|---------------------|-------------|------------------------------|
| >102.6 | 95 | ACHARD | 01 B | L3 |
| • • • We do not use the | following o | data for averages | , fits, | limits, etc. $ullet$ $ullet$ |
| > 28.2 | 95 1 | ⁵ ADACHI | 90c | TOPZ |
| none 18.5-42.8 | 95 | AKRAWY | 900 | OPAL |
| > 26.5 | 95 | DECAMP | 90F | ALEP |
| none m_{μ} –36.3 | 95 | SODERSTROM | Л 90 | MRK2 |

 $^{^{15}}$ ADACHI 90C put lower limits on the mass of stable charged particles with electric charge Q satisfying 2/3 < Q/e < 4/3 and with spin 0 or 1/2. We list here the special case for a stable charged heavy lepton.

Charged Long-Lived Heavy Lepton MASS LIMITS

| VALUE (GeV) | CL% | DOCUMENT ID | | TECN | CHG | COMMENT | |
|---|-----|-----------------------|-------------|------|-----|---------------------------|--|
| • • We do not use the following data for averages, fits, limits, etc. | | | | | | | |
| >574 | 95 | CHATRCHYA | N 13AE | CMS | | Leptons singlet model | |
| >102.0 | 95 | ABBIENDI | 03L | OPAL | | pair produced in e^+e^- | |
| > 0.1 | | ¹⁶ ANSORGE | 73 B | HBC | _ | Long-lived | |
| none 0.55-4.5 | | ¹⁷ BUSHNIN | 73 | CNTR | _ | Long-lived | |
| none 0.2-0.92 | | ¹⁸ BARNA | 68 | CNTR | _ | Long-lived | |
| none 0.97-1.03 | | ¹⁸ BARNA | 68 | CNTR | _ | Long-lived | |

¹⁶ ANSORGE 73B looks for electron pair production and electron-like Bremsstrahlung.

Created: 6/1/2020 08:33

 $^{^{14}\,\}mathrm{ROTHE}$ 69 examines previous data on μ pair production and π and K decays.

 $^{^{17}}$ BUSHNIN 73 is SERPUKHOV 70 GeV p experiment. Masses assume mean life above 7×10^{-10} and 3×10^{-8} respectively. Calculated from cross section (see "Charged Quasi-Stable Lepton Production Differential Cross Section" below) and 30 GeV muon pair production data.

¹⁸BARNA 68 is SLAC photoproduction experiment.

Doubly-Charged Heavy Lepton MASS LIMITS

<u>VALUE (GeV)</u> <u>CL%</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>CHG</u>

• • • We do not use the following data for averages, fits, limits, etc. • • • none 1–9 GeV 90 19 CLARK 81 SPEC ++

Doubly-Charged Lepton Production Cross Section $(\mu N \text{ Scattering})$

<u>VALUE (cm²)</u> <u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>CHG</u>

• • • We do not use the following data for averages, fits, limits, etc. • • • $<6. \times 10^{-38}$ 0 ²⁰ CLARK 81 SPEC ++

REFERENCES FOR Heavy Charged Lepton Searches

| CHATROHNAN | 1040 | UIED 1007 100 | 6 6 | (CNAC C II I) |
|------------|------|-----------------|----------------------------|-------------------|
| | - | JHEP 1307 122 | | (CMS Collab.) |
| ABBIENDI | 03L | PL B572 8 | G. Abbiendi <i>et al.</i> | (OPAL Collab.) |
| ACHARD | 01B | PL B517 75 | P. Achard <i>et al.</i> | (L3 Collab.) |
| ACKERSTAFF | 98C | EPJ C1 45 | K. Ackerstaff et al. | (OPAL Collab.) |
| ACCIARRI | 96G | PL B377 304 | M. Acciarri et al. | (L3 Collab.) |
| ALEXANDER | 96P | PL B385 433 | G. Alexander et al. | (OPAL Collab.) |
| BUSKULIC | 96S | PL B384 439 | D. Buskulic <i>et al.</i> | (ALEPH Collab.) |
| AHMED | 94 | PL B340 205 | T. Ahmed <i>et al.</i> | (H1 Collab.) |
| KIM | 91B | IJMP A6 2583 | G.N. Kim <i>et al.</i> | (AMY Collab.) |
| ADACHI | 90C | PL B244 352 | I. Adachi <i>et al.</i> | (TOPAZ Collab.) |
| AKRAWY | 90G | PL B240 250 | M.Z. Akrawy <i>et al.</i> | (OPAL Collab.) |
| AKRAWY | 90O | PL B252 290 | M.Z. Akrawy <i>et al.</i> | (OPAL Collab.) |
| DECAMP | 90F | PL B236 511 | D. Decamp et al. | (ALEPH Collab.) |
| RILES | 90 | PR D42 1 | K. Riles et al. | (Mark II Collab.) |
| SODERSTROM | 1 90 | PRL 64 2980 | E. Soderstrom et al. | (Mark II Collab.) |
| STOKER | 89 | PR D39 1811 | D.P. Stoker et al. | (Mark II Collab.) |
| ABE | 88 | PRL 61 915 | K. Abe <i>et al</i> . | (VENUS Collab.) |
| ADACHI | 88B | PR D37 1339 | I. Adachi <i>et al.</i> | (TOPAZ Collab.) |
| BEHREND | 88C | ZPHY C41 7 | H.J. Behrend et al. | (CELLO Collab.) |
| ALBAJAR | 87B | PL B185 241 | C. Albajar <i>et al.</i> | ` (UA1 Collab.) |
| ADEVA | 85 | PL 152B 439 | B. Adeva <i>et al.</i> | (Mark-J Collab.) |
| Also | | PRPL 109 131 | B. Adeva et al. | (Mark-J Collab.) |
| BARTEL | 83 | PL 123B 353 | W. Bartel et al. | (JADE Collab.) |
| BERGER | 81B | PL 99B 489 | C. Berger et al. | (PLUTO Collab.) |
| BRANDELIK | 81 | PL 99B 163 | R. Brandelik <i>et al.</i> | (TASSO Collab.) |
| CLARK | 81 | PRL 46 299 | A.R. Clark et al. | (UCB, LBL, FNAL+) |
| Also | - | PR D25 2762 | W.H. Smith et al. | (LBL, FNAL, PRIN) |
| AZIMOV | 80 | JETPL 32 664 | Y.I. Azimov, V.A. Khoze | (PNPI) |
| 7.2 | 00 | Translated from | ZETFP 32 677. | () |
| BARBER | 80B | PRL 45 1904 | D.P. Barber et al. | (Mark-J Collab.) |
| BRANDELIK | 80 | PL 92B 199 | R. Brandelik <i>et al.</i> | (TASSO Collab.) |
| ANSORGE | 73B | PR D7 26 | R.E. Ansorge et al. | (CAVE) |
| BUSHNIN | 73 | NP B58 476 | Y.B. Bushnin et al. | (SERP) |
| Also | | PL 42B 136 | S.V. Golovkin et al. | (SERP) |
| ROTHE | 69 | NP B10 241 | K.W. Rothe, A.M. Wolsky | (PENN) |
| BARNA | 68 | PR 173 1391 | A. Barna et al. | (SLAC, STAN) |
| | | | | , |

- OTHER RELATED PAPERS ———

Created: 6/1/2020 08:33

PERL 81 SLAC-PUB-2752 M.L. Perl (SLAC) Physics in Collision Conference.

 $^{^{19}}$ CLARK 81 is FNAL experiment with 209 GeV muons. Bounds apply to μ_P which couples with full weak strength to muon. See also section on "Doubly-Charged Lepton Production Cross Section."

 $^{^{20}}$ CLARK 81 is FNAL experiment with 209 GeV muon. Looked for μ^+ nucleon $\to ~\overline{\mu}^0_P$ X, $\overline{\mu}^0_P \to ~\mu^+\mu^-\overline{\nu}_\mu$ and $\mu^+\,n \to ~\mu^{++}_P$ X, $\mu^{++}_P \to ~2\mu^+\nu_\mu$. Above limits are for $\sigma\times BR$ taken from their mass-dependence plot figure 2.