

**$K^*(1410)$**  $I(J^P) = \frac{1}{2}(1^-)$  **$K^*(1410)$  T-MATRIX POLE  $\sqrt{s}$** 

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>			
(1368 ± 38) – $i$ (106 <sup>+48</sup> <sub>-59</sub> )	<sup>1</sup> PELAEZ	17	RVUE $\pi K \rightarrow \pi K$
<b>1</b> Reanalysis of ESTABROOKS 78 and ASTON 88 satisfying Forward Dispersion Relations and using sequences of Pade approximants.			

 **$K^*(1410)$  MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>1414±15 OUR AVERAGE</b> Error includes scale factor of 1.3.					
1380 ± 21 ± 19		ASTON	88	LASS 0	11 $K^- p \rightarrow K^- \pi^+ n$
1420 ± 7 ± 10		ASTON	87	LASS 0	11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>					
1437 ± 8 ± 16	190k	<sup>1</sup> AAIJ	16N	LHCb	$D^0 \rightarrow (K_S^0 \pi^\mp) K^\pm$
1426 ± 8 ± 24	190k	<sup>2</sup> AAIJ	16N	LHCb	$D^0 \rightarrow K_S^0 (K^\pm \pi^\mp)$
1276 <sup>+72</sup> <sub>-77</sub>		<sup>3,4</sup> BOITO	09	RVUE	$\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$
1367 ± 54		BIRD	89	LASS	–
1474 ± 25		BAUBILLIER	82B	HBC	0
1500 ± 30		ETKIN	80	MPS	0
1 Using a parametrization for the $K\pi$ S-wave similar to ASTON 88 with fixed resonance width.					
2 Using a $K\pi$ S-wave parametrization with resonant and non-resonant contributions.					
3 From the pole position of the $K\pi$ vector form factor in the complex $s$ -plane and using EPIFANOV 07 data.					
4 Systematic uncertainties not estimated.					

 **$K^*(1410)$  WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>232± 21 OUR AVERAGE</b> Error includes scale factor of 1.1.					
176 ± 52 ± 22		ASTON	88	LASS 0	11 $K^- p \rightarrow K^- \pi^+ n$
240 ± 18 ± 12		ASTON	87	LASS 0	11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>					
210 ± 20 ± 60	190k	<sup>1</sup> AAIJ	16N	LHCb	$D^0 \rightarrow (K_S^0 \pi^\mp) K^\pm$
270 ± 20 ± 40	190k	<sup>1</sup> AAIJ	16N	LHCb	$D^0 \rightarrow K_S^0 (K^\pm \pi^\mp)$
198 <sup>+61</sup> <sub>-87</sub>		<sup>2,3</sup> BOITO	09	RVUE	$\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$
114 ± 101		BIRD	89	LASS	–
275 ± 65		BAUBILLIER	82B	HBC	0
500 ± 100		ETKIN	80	MPS	0
11 $K^- p \rightarrow \bar{K}^0 \pi^- p$					
8.25 $K^- p \rightarrow \bar{K}^0 2\pi n$					
6 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$					

<sup>1</sup> Using a  $K\pi$  S-wave parametrization with resonant and non-resonant contributions.<sup>2</sup> From the pole position of the  $K\pi$  vector form factor in the complex  $s$ -plane and using EPIFANOV 07 data.<sup>3</sup> Systematic uncertainties not estimated.

## $K^*(1410)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1 K^*(892)\pi$	> 40 %	95%
$\Gamma_2 K\pi$	( $6.6 \pm 1.3$ ) %	
$\Gamma_3 K\rho$	< 7 %	95%
$\Gamma_4 \gamma K^0$	< $2.3 \times 10^{-4}$	90%

## $K^*(1410)$ PARTIAL WIDTHS

$\Gamma(\gamma K^0)$	$\Gamma_4$
$\text{VALUE (keV)}$	$\text{CL \%}$

$\Gamma(K\rho)/\Gamma(K^*(892)\pi)$	$\Gamma_3/\Gamma_1$
$\text{VALUE}$	$\text{CL \%}$

$\Gamma(K\pi)/\Gamma(K^*(892)\pi)$	$\Gamma_2/\Gamma_1$
$\text{VALUE}$	$\text{CL \%}$

$\Gamma(K\pi)/\Gamma_{\text{total}}$	$\Gamma_2/\Gamma$
$\text{VALUE}$	$\text{DOCUMENT ID}$

## $K^*(1410)$ REFERENCES

PELAEZ	17	EPJ C77 91	J.R. Pelaez, A.Rodas, J.R. de Elvira
AAIJ	16N	PR D93 052018	R. Aaij <i>et al.</i> (LHCb Collab.)
BOITO	09	EPJ C59 821	D.R. Boito, R. Escribano, M. Jamin
EPIFANOV	07	PL B654 65	D. Epifanov <i>et al.</i> (BELLE Collab.)
ALAVI-HARATI	02B	PRL 89 072001	A. Alavi-Harati <i>et al.</i> (FNAL KTeV Collab.)
BIRD	89	SLAC-332	P.F. Bird (SLAC)
ASTON	88	NP B296 493	D. Aston <i>et al.</i> (SLAC, NAGO, CINC, INUS)
ASTON	87	NP B292 693	D. Aston <i>et al.</i> (SLAC, NAGO, CINC, INUS)
ASTON	84	PL 149B 258	D. Aston <i>et al.</i> (SLAC, CARL, OTTA) JP
BAUBILLIER	82B	NP B202 21	M. Baubillier <i>et al.</i> (BIRM, CERN, GLAS+) JP
ETKIN	80	PR D22 42	A. Etkin <i>et al.</i> (BNL, CUNY) JP
ESTABROOKS	78	NP B133 490	P.G. Estabrooks <i>et al.</i> (MCGI, CARL, DURH+)