

**$\phi(1680)$**

$I^G(J^{PC}) = 0^-(1^{--})$

## **$\phi(1680)$ MASS**

### e<sup>+</sup> e<sup>-</sup> PRODUCTION

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

#### **1680±20 OUR ESTIMATE**

• • • We do not use the following data for averages, fits, limits, etc. • • •

1680 <sup>+12</sup> <sub>-13</sub> ± 21	1.8k	1 ABLIKIM	20F BES3	$\psi(2S) \rightarrow K^+ K^- \eta$
1662 ± 20		2 ACHASOV	20C SND	$1.3\text{--}2.0 e^+ e^- \rightarrow K^+ K^- \pi^0$
1641 <sup>+24</sup> <sub>-18</sub>		ACHASOV	19 SND	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \eta$
1667 ± 5 ± 11	3k	3 IVANOV	19A CMD3	$1.59\text{--}2.007 e^+ e^- \rightarrow K^+ K^- \eta$
1700 ± 23	2k	4 ACHASOV	18A SND	$1.3\text{--}2.0 e^+ e^- \rightarrow K_S^0 K_L^0 \pi^0$
1674 ± 12 ± 6	6.2k	5 LEES	14H BABR	$e^+ e^- \rightarrow K_S^0 K_L^0 \gamma$
1733 ± 10 ± 10		6 LEES	12F BABR	$10.6 e^+ e^- \rightarrow \phi \pi^+ \pi^- \gamma$
1689 ± 7 ± 10	4.8k	7 SHEN	09 BELL	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$
1709 ± 20 ± 43		8 AUBERT	08S BABR	$10.6 e^+ e^- \rightarrow \text{hadrons}$
1623 ± 20	948	9 AKHMETSHIN 03	CMD2	$1.05\text{--}1.38 e^+ e^- \rightarrow K_L^0 K_S^0$
~ 1500		10 ACHASOV	98H RVUE	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0, \omega \pi^+ \pi^-, K^+ K^-$
~ 1900		11 ACHASOV	98H RVUE	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp$
1700 ± 20		12 CLEGG	94 RVUE	$e^+ e^- \rightarrow K^+ K^-, K_S^0 K \pi$
1657 ± 27	367	13 BISELLO	91C DM2	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp$
1655 ± 17		13 BISELLO	88B DM2	$e^+ e^- \rightarrow K^+ K^-$
1680 ± 10		14 BUON	82 DM1	$e^+ e^- \rightarrow \text{hadrons}$
1677 ± 12		15 MANE	82 DM1	$e^+ e^- \rightarrow K_S^0 K \pi$

<sup>1</sup> Seen in  $\psi(2S)$  decay with branching ratio  $\psi(2S) \rightarrow X \eta \rightarrow K^+ K^- \eta = (12.0 \pm 1.3^{+6.5}_{-6.9}) \times 10^{-6}$ .

<sup>2</sup> From a fit using a vector meson dominance model with contribution from  $\rho(770)$ ,  $\omega(782)$ ,  $\phi(1020)$ ,  $\omega(1420)$ ,  $\rho(1450)$ .

<sup>3</sup> From a fit with coherent interference of the  $\phi(1680)$  with a non-resonant contribution.

<sup>4</sup> Assuming the  $K \bar{K}^*(892) + \text{c.c.}$  dynamics. Systematic uncertainties not estimated.

<sup>5</sup> Using a vector meson dominance model with contribution from  $\phi(1020)$ ,  $\phi(1680)$ , and higher mass excitations of  $\rho(770)$  and  $\omega(782)$ .

<sup>6</sup> Using events with  $\pi\pi$  invariant mass less than 0.85 GeV.

<sup>7</sup> From a fit with two incoherent Breit-Wigners.

<sup>8</sup> From the simultaneous fit to the  $K \bar{K}^*(892) + \text{c.c.}$  and  $\phi \eta$  data from AUBERT 08S using the results of AUBERT 07AK.

<sup>9</sup> From the combined fit of AKHMETSHIN 03 and MANE 81 also including  $\rho$ ,  $\omega$ , and  $\phi$ . Neither isospin nor flavor structure known.

<sup>10</sup> Using data from IVANOV 81, BARKOV 87, BISELLO 88B, DOLINSKY 91, and ANTONELLI 92.

<sup>11</sup> Using the data from BISELLO 91C.

<sup>12</sup> Using BISELLO 88B and MANE 82 data.

<sup>13</sup> From global fit including  $\rho$ ,  $\omega$ ,  $\phi$  and  $\rho(1700)$  assume mass 1570 MeV and width 510 MeV for  $\rho$  radial excitation.

- <sup>14</sup> From global fit of  $\rho$ ,  $\omega$ ,  $\phi$  and their radial excitations to channels  $\omega\pi^+\pi^-$ ,  $K^+K^-$ ,  $K_S^0 K_L^0$ ,  $K_S^0 K^\pm \pi^\mp$ . Assume mass 1570 MeV and width 510 MeV for  $\rho$  radial excitations, mass 1570 and width 500 MeV for  $\omega$  radial excitation.  
<sup>15</sup> Fit to one channel only, neglecting interference with  $\omega$ ,  $\rho(1700)$ .

## PHOTOPRODUCTION

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>			
1753 $\pm$ 3	<sup>1</sup> LINK	02K FOCS	20–160 $\gamma p \rightarrow K^+ K^- p$
1726 $\pm$ 22	<sup>1</sup> BUSENITZ	89 TPS	$\gamma p \rightarrow K^+ K^- X$
1760 $\pm$ 20	<sup>1</sup> ATKINSON	85C OMEG	20–70 $\gamma p \rightarrow K\bar{K}X$
1690 $\pm$ 10	<sup>1</sup> ASTON	81F OMEG	25–70 $\gamma p \rightarrow K^+ K^- X$

<sup>1</sup> We list here a state decaying into  $K^+ K^-$  possibly different from  $\phi(1680)$ .

## p $\bar{p}$ ANNIHILATION

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>			
1700 $\pm$ 8	<sup>1</sup> AMSLER	06 CBAR	$0.9 \bar{p}p \rightarrow K^+ K^- \pi^0$
<sup>1</sup> Could also be $\rho(1700)$ .			

## $\phi(1680)$ WIDTH

### e<sup>+</sup> e<sup>-</sup> PRODUCTION

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>150 <math>\pm</math> 50 OUR ESTIMATE</b>				This is only an educated guess; the error given is larger than the error on the average of the published values.
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
185 $\pm$ 30 $\pm$ 25	1.8k	<sup>1</sup> ABLIKIM	20F BES3	$\psi(2S) \rightarrow K^+ K^- \eta$
159 $\pm$ 32		<sup>2</sup> ACHASOV	20C SND	$1.3\text{--}2.0 e^+ e^- \rightarrow K^+ K^- \pi^0$
103 $\pm$ 26		ACHASOV	19 SND	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \eta$
176 $\pm$ 23 $\pm$ 38	3k	<sup>3</sup> IVANOV	19A CMD3	$1.59\text{--}2.007 e^+ e^- \rightarrow K^+ K^- \eta$
300 $\pm$ 50	2k	<sup>4</sup> ACHASOV	18A SND	$1.3\text{--}2.0 e^+ e^- \rightarrow K_S^0 K_L^0 \pi^0$
165 $\pm$ 38 $\pm$ 70	6.2k	<sup>5</sup> LEES	14H BABR	$e^+ e^- \rightarrow K_S^0 K_L^0 \gamma$
300 $\pm$ 15 $\pm$ 37		<sup>6</sup> LEES	12F BABR	$10.6 e^+ e^- \rightarrow \phi \pi^+ \pi^- \gamma$
211 $\pm$ 14 $\pm$ 19	4.8k	<sup>7</sup> SHEN	09 BELL	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$
322 $\pm$ 77 $\pm$ 160		<sup>8</sup> AUBERT	08S BABR	$10.6 e^+ e^- \rightarrow \text{hadrons}$
139 $\pm$ 60	948	<sup>9</sup> AKHMETSHIN	03 CMD2	$1.05\text{--}1.38 e^+ e^- \rightarrow K_L^0 K_S^0$
300 $\pm$ 60		<sup>10</sup> CLEGG	94 RVUE	$e^+ e^- \rightarrow K^+ K^-, K_S^0 K\pi$
146 $\pm$ 55	367	BISELLO	91C DM2	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp$
207 $\pm$ 45		<sup>11</sup> BISELLO	88B DM2	$e^+ e^- \rightarrow K^+ K^-$
185 $\pm$ 22		<sup>12</sup> BUON	82 DM1	$e^+ e^- \rightarrow \text{hadrons}$
102 $\pm$ 36		<sup>13</sup> MANE	82 DM1	$e^+ e^- \rightarrow K_S^0 K\pi$

<sup>1</sup> Seen in  $\psi(2S)$  decay with branching ratio  $\psi(2S) \rightarrow X\eta \rightarrow K^+ K^- \eta = (12.0 \pm 1.3^{+6.5}_{-6.9}) \times 10^{-6}$ .

<sup>2</sup> From a fit using a vector meson dominance model with contribution from  $\rho(770)$ ,  $\omega(782)$ ,  $\phi(1020)$ ,  $\omega(1420)$ ,  $\rho(1450)$ .

- <sup>3</sup> From a fit with coherent interference of the  $\phi(1680)$  with a non-resonant contribution.  
<sup>4</sup> Assuming the  $K\bar{K}^*(892) + \text{c.c.}$  dynamics. Systematic uncertainties not estimated.  
<sup>5</sup> Using a vector meson dominance model with contribution from  $\phi(1020)$ ,  $\phi(1680)$ , and higher mass excitations of  $\rho(770)$  and  $\omega(782)$ .  
<sup>6</sup> Using events with  $\pi\pi$  invariant mass less than 0.85 GeV.  
<sup>7</sup> From a fit with two incoherent Breit-Wigners.  
<sup>8</sup> From the simultaneous fit to the  $K\bar{K}^*(892) + \text{c.c.}$  and  $\phi\eta$  data from AUBERT 08S using the results of AUBERT 07AK.  
<sup>9</sup> From the combined fit of AKHMETSHIN 03 and MANE 81 also including  $\rho$ ,  $\omega$ , and  $\phi$ . Neither isospin nor flavor structure known.  
<sup>10</sup> Using BISELLO 88B and MANE 82 data.  
<sup>11</sup> From global fit including  $\rho$ ,  $\omega$ ,  $\phi$  and  $\rho(1700)$   
<sup>12</sup> From global fit of  $\rho$ ,  $\omega$ ,  $\phi$  and their radial excitations to channels  $\omega\pi^+\pi^-$ ,  $K^+K^-$ ,  $K_S^0 K_L^0$ ,  $K_S^0 K^\pm\pi^\mp$ . Assume mass 1570 MeV and width 510 MeV for  $\rho$  radial excitations, mass 1570 and width 500 MeV for  $\omega$  radial excitation.  
<sup>13</sup> Fit to one channel only, neglecting interference with  $\omega$ ,  $\rho(1700)$ .

## PHOTOPRODUCTION

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>			
122 ± 63	<sup>1</sup> LINK	02K FOCS	20–160 $\gamma p \rightarrow K^+K^-p$
121 ± 47	<sup>1</sup> BUSENITZ	89 TPS	$\gamma p \rightarrow K^+K^-X$
80 ± 40	<sup>1</sup> ATKINSON	85C OMEG	20–70 $\gamma p \rightarrow K\bar{K}X$
100 ± 40	<sup>1</sup> ASTON	81F OMEG	25–70 $\gamma p \rightarrow K^+K^-X$

<sup>1</sup> We list here a state decaying into  $K^+K^-$  possibly different from  $\phi(1680)$ .

## p $\bar{p}$ ANNIHILATION

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>			
143 ± 24	<sup>1</sup> AMSLER	06 CBAR	0.9 $\bar{p}p \rightarrow K^+K^-\pi^0$
<sup>1</sup> Could also be $\rho(1700)$ .			

## $\phi(1680)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 K\bar{K}^*(892) + \text{c.c.}$	seen
$\Gamma_2 K_S^0 K\pi$	seen
$\Gamma_3 K\bar{K}$	seen
$\Gamma_4 K_L^0 K_S^0$	
$\Gamma_5 e^+e^-$	seen
$\Gamma_6 \omega\pi\pi$	not seen
$\Gamma_7 \phi\pi\pi$	
$\Gamma_8 K^+K^-\pi^+\pi^-$	seen
$\Gamma_9 \eta\phi$	seen
$\Gamma_{10} K^+K^-\eta$	
$\Gamma_{11} \eta\gamma$	seen
$\Gamma_{12} K^+K^-\pi^0$	

$$\phi(1680) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$$

This combination of a partial width with the partial width into  $e^+ e^-$  and with the total width is obtained from the integrated cross section into channel (I) in  $e^+ e^-$  annihilation. We list only data that have not been used to determine the partial width  $\Gamma(I)$  or the branching ratio  $\Gamma(I)/\text{total}$ .

$$\Gamma(K_L^0 K_S^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} = 4\Gamma_4 \Gamma_5 / \Gamma$$

VALUE (eV)      EVTS      DOCUMENT ID      TECN      COMMENT

- • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup> Using a vector meson dominance model with contribution from  $\phi(1020)$ ,  $\phi(1680)$ , and light-cone contributions of  $(770)$  and  $(700)$ .

$$\Gamma(\phi\pi\pi) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$$

1.000 x 1.000 / total 11151

VALUE ( $10^{-2}$ keV)	DOCUMENT_ID	TECN	COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.2 ± 0.2 ± 0.3                  LEFS                  12F BABR 10.6  $e^+e^- \rightarrow \phi\pi^+\pi^-\gamma$

$$\Gamma(\eta\phi) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} = \Gamma_9\Gamma_5/\Gamma$$

VALUE (eV)      EVTS      DOCUMENT ID      TECN      COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

$94 \pm 13 \pm 15$       3k      <sup>1</sup> IVANOV      19A    CMD3    1.59–2.007     $e^+ e^- \rightarrow K^+ K^- \eta$

<sup>1</sup>From a fit with coherent interference of the  $\phi(1680)$  with a non-resonant contribution.

$$\phi(1680) \Gamma(i)\Gamma(e^+e^-)/\Gamma^2(\text{total})$$

This combination of a branching ratio into channel ( $i$ ) and branching ratio into  $e^+ e^-$  is directly measured and obtained from the cross section at the peak. We list only data that have not been used to determine the branching ratio into ( $i$ ) or  $e^+ e^-$ .

$$\Gamma(K_L^0 K_S^0)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_4/\Gamma \times \Gamma_5/\Gamma$$

VALUE (units  $10^{-6}$ )    EVTS            DOCUMENT ID            TECN    COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.131 \pm 0.059$       948      <sup>1</sup> AKHMETSHIN 03    CMD2 1.05–1.38  $e^+e^- \rightarrow K_L^0 K_S^0$

<sup>1</sup> From the combined fit of AKHMETSHIN 03 and MANE 81 also including  $\rho$ ,  $\omega$ , and  $\phi$ . Neither isospin nor flavor structure known. Recalculated by us.

$$\Gamma(K\bar{K}^*(892)+\text{c.c.})/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_1/\Gamma \times \Gamma_5/\Gamma$$

VALUE (units 10<sup>-6</sup>)    EVTS    DOCUMENT ID    TECN    COMMENT

- • • We do not use the following data for averages, fits, limits, etc. • • •

$1.15 \pm 0.16 \pm 0.01$        $^1$  AUBERT      08s BABR 10.6  $e^+e^- \rightarrow K\bar{K}^*(892)\gamma +$

$3.29 \pm 1.57$       367      <sup>2</sup> BISELLO      91c DM2      <sup>c.c.</sup>  $1.35\text{--}2.40 \text{ } e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp$

<sup>1</sup> From the simultaneous fit to the  $K\bar{K}^*(892) + \text{c.c.}$  and  $\phi\eta$  data from AUBERT 08S using the results of AUBERT 07AK.

<sup>2</sup> Recalculated by us with the published value of  $B(K\bar{K}^*(892) + \text{c.c.}) \times \Gamma(e^+ e^-)$ .

$\Gamma(\phi\pi\pi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$  $\Gamma_7/\Gamma \times \Gamma_5/\Gamma$ 

<u>VALUE</u> (units $10^{-7}$ )	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$1.86 \pm 0.14 \pm 0.21$	4.8k	<sup>1</sup> SHEN	09	BELL $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
<sup>1</sup> Multiplied by 3/2 to take into account the $\phi\pi^0\pi^0$ mode. Using $B(\phi \rightarrow K^+K^-) = (49.2 \pm 0.6)\%$ .				

 $\Gamma(\eta\phi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$  $\Gamma_9/\Gamma \times \Gamma_5/\Gamma$ 

<u>VALUE</u> (units $10^{-7}$ )	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$5.64^{+1.74}_{-1.80}$		ACHASOV	19	SND $e^+e^- \rightarrow \pi^+\pi^-\pi^0\eta$
$5.3 \pm 0.6 \pm 0.9$	3k	<sup>1</sup> IVANOV	19A	CMD3 $1.59-2.007 e^+e^- \rightarrow K^+K^-\eta$
$4.3 \pm 1.0 \pm 0.9$		<sup>2</sup> AUBERT	08S	BABR $10.6 e^+e^- \rightarrow \phi\eta\gamma$
<sup>1</sup> From a fit with coherent interference of the $\phi(1680)$ with a non-resonant contribution.				
<sup>2</sup> From the simultaneous fit to the $K\bar{K}^*(892)+$ c.c. and $\phi\eta$ data from AUBERT 08S using the results of AUBERT 07AK.				

 **$\phi(1680)$  BRANCHING RATIOS** $\Gamma(K\bar{K}^*(892)+\text{c.c.})/\Gamma(K_S^0 K\pi)$  $\Gamma_1/\Gamma_2$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
dominant	MANE	82	$e^+e^- \rightarrow K_S^0 K^\pm\pi^\mp$

 $\Gamma(K\bar{K})/\Gamma(K\bar{K}^*(892)+\text{c.c.})$  $\Gamma_3/\Gamma_1$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
$0.07 \pm 0.01$	BUON	82	DM1 $e^+e^-$

 $\Gamma(\omega\pi\pi)/\Gamma(K\bar{K}^*(892)+\text{c.c.})$  $\Gamma_6/\Gamma_1$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.10	BUON	82	DM1 $e^+e^-$

 $\Gamma(\eta\phi)/\Gamma_{\text{total}}$  $\Gamma_9/\Gamma$ 

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
seen	35	<sup>1</sup> ACHASOV	14	SND $1.15-2.00 e^+e^- \rightarrow \eta\gamma$

<sup>1</sup> From a phenomenological model based on vector meson dominance with  $\rho(1450)$  and  $\phi(1680)$  masses and widths from the PDG 12.

 $\Gamma(\eta\phi)/\Gamma(K\bar{K}^*(892)+\text{c.c.})$  $\Gamma_9/\Gamma_1$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
$\approx 0.37$	<sup>1</sup> AUBERT	08S	BABR $10.6 e^+e^- \rightarrow \text{hadrons}$

<sup>1</sup> From the fit including data from AUBERT 07AK.

$\Gamma(\eta\gamma)/\Gamma_{\text{total}}$	$\Gamma_{11}/\Gamma$			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	35	<sup>1</sup> ACHASOV	14	SND    1.15–2.00 $e^+e^- \rightarrow \eta\gamma$
<sup>1</sup> From a phenomenological model based on vector meson dominance with $\rho(1450)$ and $\phi(1680)$ masses and widths from the PDG 12.				

## $\phi(1680)$ REFERENCES

ABLIKIM	20F	PR D101 032008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ACHASOV	20C	EPJ C80 1139	M.N. Achasov <i>et al.</i>	(SND Collab.)
ACHASOV	19	PR D99 112004	M.N. Achasov <i>et al.</i>	(SND Collab.)
IVANOV	19A	PL B798 134946	V.L. Ivanov <i>et al.</i>	(CMD-3 Collab.)
ACHASOV	18A	PR D97 032011	M.N. Achasov <i>et al.</i>	(SND Collab.)
ACHASOV	14	PR D90 032002	M.N. Achasov <i>et al.</i>	(SND Collab.)
LEES	14H	PR D89 092002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	12F	PR D86 012008	J.P. Lees <i>et al.</i>	(BABAR Collab.)
PDG	12	PR D86 010001	J. Beringer <i>et al.</i>	(PDG Collab.)
SHEN	09	PR D80 031101	C.P. Shen <i>et al.</i>	(BELLE Collab.)
AUBERT	08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)
AMSLER	06	PL B639 165	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AKHMETSHIN	03	PL B551 27	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
Also		PAN 65 1222	E.V. Anashkin, V.M. Aulchenko, R.R. Akhmetshin	
LINK	02K	PL B545 50	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
ACHASOV	98H	PR D57 4334	N.N. Achasov, A.A. Kozhevnikov	
CLEGG	94	ZPHY C62 455	A.B. Clegg, A. Donnachie	(LANC, MCHS)
ANTONELLI	92	ZPHY C56 15	A. Antonelli <i>et al.</i>	(DM2 Collab.)
BISELLO	91C	ZPHY C52 227	D. Bisello <i>et al.</i>	(DM2 Collab.)
DOLINSKY	91	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)
BUSENITZ	89	PR D40 1	J.K. Busenitz <i>et al.</i>	(ILL, FNAL)
BISELLO	88B	ZPHY C39 13	D. Bisello <i>et al.</i>	(PADO, CLÉR, FRAS+)
BARKOV	87	JETPL 46 164	L.M. Barkov <i>et al.</i>	(NOVO)
ATKINSON	85C	ZPHY C27 233	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
BUON	82	PL 118B 221	J. Buon <i>et al.</i>	(LALO, MONP)
MANE	82	PL 112B 178	F. Mane <i>et al.</i>	(LALO)
ASTON	81F	PL 104B 231	D. Aston	(BONN, CERN, EPOL, GLAS, LANC+)
IVANOV	81	PL 107B 297	P.M. Ivanov <i>et al.</i>	(NOVO)
MANE	81	PL 99B 261	F. Mane <i>et al.</i>	(ORsay)