

$\Xi(1820)$ $I(J^P) = \frac{1}{2}(\frac{3}{2}^-)$ Status: ***

The clearest evidence is an 8-standard-deviation peak in ΛK^- seen by GAY 76C. TEODORO 78 favors $J = 3/2$, but cannot make a parity discrimination. BIAGI 87C is consistent with $J = 3/2$ and favors negative parity for this J value.

 $\Xi(1820)$ MASS

We only average the measurements that appear to us to be most significant and best determined.

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | CHG | COMMENT |
|---|------|-----------------------|------|------|------------------------------------|
| 1823 \pm 5 OUR ESTIMATE | | | | | |
| 1823.5 \pm 1.4 OUR AVERAGE | | | | | |
| 1825.5 \pm 4.7 \pm 4.7 | 288 | ABLIKIM | 20C | BES3 | — |
| 1819.4 \pm 3.1 \pm 2.0 | 280 | ¹ BIAGI | 87 | SPEC | 0 |
| 1826 \pm 3 \pm 1 | 54 | BIAGI | 87C | SPEC | 0 |
| 1822 \pm 6 | | JENKINS | 83 | MPS | — |
| 1830 \pm 6 | 300 | BIAGI | 81 | SPEC | — |
| 1823 \pm 2 | 130 | GAY | 76C | HBC | — |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| 1817 \pm 3 | | ADAMOVICH | 99B | WA89 | Σ^- nucleus, 345 GeV |
| 1797 \pm 19 | 74 | BRIEFEL | 77 | HBC | 0 $K^- p$ 2.87 GeV/c |
| 1829 \pm 9 | 68 | BRIEFEL | 77 | HBC | —0 $\Xi(1530)\pi$ |
| 1860 \pm 14 | 39 | BRIEFEL | 77 | HBC | — $\Sigma^-\bar{K}^0$ |
| 1870 \pm 9 | 44 | BRIEFEL | 77 | HBC | 0 $\Lambda\bar{K}^0$ |
| 1813 \pm 4 | 57 | BRIEFEL | 77 | HBC | — ΛK^- |
| 1807 \pm 27 | | DIBIANCA | 75 | DBC | —0 $\Xi\pi\pi, \Xi^*\pi$ |
| 1762 \pm 8 | 28 | ² BADIER | 72 | HBC | —0 $\Xi\pi, \Xi\pi\pi, YK$ |
| 1838 \pm 5 | 38 | ² BADIER | 72 | HBC | —0 $\Xi\pi, \Xi\pi\pi, YK$ |
| 1830 \pm 10 | 25 | ³ CRENNELL | 70B | DBC | —0 3.6, 3.9 GeV/c |
| 1826 \pm 12 | | ⁴ CRENNELL | 70B | DBC | —0 3.6, 3.9 GeV/c |
| 1830 \pm 10 | 40 | ALITTI | 69 | HBC | — $\Lambda, \Sigma\bar{K}$ |
| 1814 \pm 4 | 30 | BADIER | 65 | HBC | 0 $\Lambda\bar{K}^0$ |
| 1817 \pm 7 | 29 | SMITH | 65C | HBC | —0 $\Lambda\bar{K}^0, \Lambda K^-$ |
| 1770 | | HALSTEINSLID63 | FBC | —0 | K^- freon 3.5 GeV/c |


 $\Xi(1820)$ WIDTH

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | CHG | COMMENT |
|--|------|--------------------|------|------|---|
| 24 \pm 15 OUR ESTIMATE | | | | | |
| 24 \pm 5 OUR AVERAGE | | | | | Error includes scale factor of 1.3. See the ideogram below. |
| 17.0 \pm 15.0 \pm 7.9 | 288 | ABLIKIM | 20C | BES3 | — |
| 17.0 \pm 15.0 \pm 7.9 | | | | | $e^+ e^- \rightarrow \Xi(1820)^-\Xi^+$ |
| 24.6 \pm 5.3 | 280 | ¹ BIAGI | 87 | SPEC | 0 $\Xi^- Be \rightarrow (\Lambda K^-) X$ |
| 12 \pm 14 \pm 1.7 | 54 | BIAGI | 87C | SPEC | 0 $\Xi^- Be \rightarrow (\Lambda\bar{K}^0) X$ |

| | | | | | | |
|---------|-----|-------|-----|------|---|-------------------|
| 72 ± 20 | 300 | BIAGI | 81 | SPEC | — | SPS hyperon beam |
| 21 ± 7 | 130 | GAY | 76C | HBC | — | $K^- p$ 4.2 GeV/c |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | | |
|----------|----|----------------|-----|------|-----------------------------|--------------------------|
| 23 ± 13 | | ADAMOVICH | 99B | WA89 | Σ^- nucleus, 345 GeV | |
| 99 ± 57 | 74 | BRIEFEL | 77 | HBC | 0 | $K^- p$ 2.87 GeV/c |
| 52 ± 34 | 68 | BRIEFEL | 77 | HBC | —0 | $\Xi(1530)\pi$ |
| 72 ± 17 | 39 | BRIEFEL | 77 | HBC | — | $\Sigma^-\bar{K}^0$ |
| 44 ± 11 | 44 | BRIEFEL | 77 | HBC | 0 | $\Lambda\bar{K}^0$ |
| 26 ± 11 | 57 | BRIEFEL | 77 | HBC | — | ΛK^- |
| 85 ± 58 | | DIBIANCA | 75 | DBC | —0 | $\Xi\pi\pi, \Xi^*\pi$ |
| 51 ± 13 | 2 | BADIER | 72 | HBC | —0 | Lower mass |
| 58 ± 13 | 2 | BADIER | 72 | HBC | —0 | Higher mass |
| 103 ± 38 | 3 | CRENNELL | 70B | DBC | —0 | 3.6, 3.9 GeV/c |
| —24 | | CRENNELL | 70B | DBC | —0 | 3.6, 3.9 GeV/c |
| 48 ± 36 | 4 | CRENNELL | 70B | DBC | —0 | 3.6, 3.9 GeV/c |
| —19 | | ALITTI | 69 | HBC | — | $\Lambda, \Sigma\bar{K}$ |
| 55 ± 40 | | BADIER | 65 | HBC | 0 | $\Lambda\bar{K}^0$ |
| 12 ± 4 | | SMITH | 65B | HBC | —0 | $\Lambda\bar{K}$ |
| 30 ± 7 | | HALSTEINSLID63 | FBC | | —0 | K^- freon 3.5 GeV/c |
| <80 | | | | | | |



$\Xi(1820)$ DECAY MODES

| Mode | Fraction (Γ_i/Γ) |
|--|--------------------------------|
| $\Gamma_1 \Lambda\bar{K}$ | large |
| $\Gamma_2 \Sigma\bar{K}$ | small |
| $\Gamma_3 \Xi\pi$ | small |
| $\Gamma_4 \Xi(1530)\pi$ | small |
| $\Gamma_5 \Xi\pi\pi(\text{not } \Xi(1530)\pi)$ | |

$\Xi(1820)$ BRANCHING RATIOS

The dominant modes seem to be $\Lambda\bar{K}$ and (perhaps) $\Xi(1530)\pi$, but the branching fractions are very poorly determined.

$\Gamma(\Lambda\bar{K})/\Gamma_{\text{total}}$

| VALUE |
|---|
| 0.25 ± 0.05 OUR AVERAGE |

0.24 ± 0.05

0.30 ± 0.15

| DOCUMENT ID | TECN | CHG | COMMENT |
|-------------|------|-----|---------|
|-------------|------|-----|---------|

| | | | |
|-----------|-----|------|---------------------|
| ANISOVICH | 12A | DPWA | Multichannel |
| ALITTI | 69 | HBC | $K^- p$ 3.9–5 GeV/c |

$\Gamma(\Xi\pi)/\Gamma_{\text{total}}$

| VALUE |
|-----------------------------------|
| 0.10 ± 0.10 |

| DOCUMENT ID | TECN | CHG | COMMENT |
|-------------|------|-----|---------------------|
| ALITTI | 69 | HBC | $K^- p$ 3.9–5 GeV/c |

$\Gamma(\Xi\pi)/\Gamma(\Lambda\bar{K})$

| VALUE | CL% |
|-----------------------------------|-----|
| <0.36 | 95 |
| 0.20 ± 0.20 | |

| DOCUMENT ID | TECN | CHG | COMMENT |
|-------------|------|-----|-------------------|
| GAY | 76C | HBC | $K^- p$ 4.2 GeV/c |
| BADIER | 65 | HBC | $K^- p$ 3 GeV/c |

$\Gamma(\Xi\pi)/\Gamma(\Xi(1530)\pi)$

| VALUE |
|---------------------------------------|
| $1.5^{+0.6}_{-0.4}$ |

| DOCUMENT ID | TECN | CHG | COMMENT |
|-------------|------|-----|--------------------|
| APSELL | 70 | HBC | $K^- p$ 2.87 GeV/c |

$\Gamma(\Sigma\bar{K})/\Gamma_{\text{total}}$

| VALUE |
|-----------------------------------|
| 0.30 ± 0.15 |

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.02

| DOCUMENT ID | TECN | CHG | COMMENT |
|-------------|------|-----|---------------------|
| ALITTI | 69 | HBC | $K^- p$ 3.9–5 GeV/c |

| | | | |
|-------|----|------|---------------|
| TRIPP | 67 | RVUE | Use SMITH 65C |
|-------|----|------|---------------|

$\Gamma(\Sigma\bar{K})/\Gamma(\Lambda\bar{K})$

| VALUE |
|-----------------------------------|
| 0.24 ± 0.10 |

| DOCUMENT ID | TECN | CHG | COMMENT |
|-------------|------|-----|-------------------|
| GAY | 76C | HBC | $K^- p$ 4.2 GeV/c |

$\Gamma(\Xi(1530)\pi)/\Gamma_{\text{total}}$

| VALUE |
|-----------------------------------|
| 0.30 ± 0.15 |

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen

| DOCUMENT ID | TECN | CHG | COMMENT |
|-------------|------|-----|---------------------|
| ALITTI | 69 | HBC | $K^- p$ 3.9–5 GeV/c |

| | | | |
|-------|-----|------|------------------|
| ASTON | 85B | LASS | $K^- p$ 11 GeV/c |
|-------|-----|------|------------------|

| | | | | |
|----------|----------------------|----|-----|-------------------|
| not seen | ⁵ HASSALL | 81 | HBC | $K^- p$ 6.5 GeV/c |
| <0.25 | ⁶ DAUBER | 69 | HBC | $K^- p$ 2.7 GeV/c |

 $\Gamma(\Xi(1530)\pi)/\Gamma(\Lambda\bar{K})$

| VALUE | DOCUMENT ID | TECN | CHG | COMMENT |
|------------------------------|-------------|------|-----|-------------------------------------|
| 0.38±0.27 OUR AVERAGE | | | | Error includes scale factor of 2.3. |
| 1.0 ± 0.3 | GAY | 76C | HBC | — $K^- p$ 4.2 GeV/c |
| 0.26±0.13 | SMITH | 65C | HBC | —0 $K^- p$ 2.45–2.7 GeV/c |

 $\Gamma(\Xi\pi\pi(\text{not } \Xi(1530)\pi))/\Gamma(\Lambda\bar{K})$

| VALUE | DOCUMENT ID | TECN | CHG | COMMENT |
|---|---------------------|------|------|---------------------------|
| 0.30±0.20 | BIAGI | 87 | SPEC | — Ξ^- Be 116 GeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| <0.14 | ⁷ BADIER | 65 | HBC | 0 1 st. dev. limit |
| >0.1 | SMITH | 65C | HBC | —0 $K^- p$ 2.45–2.7 GeV/c |

 $\Gamma(\Xi\pi\pi(\text{not } \Xi(1530)\pi))/\Gamma(\Xi(1530)\pi)$

| VALUE | DOCUMENT ID | TECN | CHG | COMMENT |
|---|---------------------|------|-----|----------------------|
| consistent with zero | GAY | 76C | HBC | — $K^- p$ 4.2 GeV/c |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.3±0.5 | ⁸ APSELL | 70 | HBC | 0 $K^- p$ 2.87 GeV/c |

 $\Xi(1820)$ FOOTNOTES

¹ BIAGI 87 also sees weak signals in the in the $\Xi^-\pi^+\pi^-$ channel at 1782.6 ± 1.4 MeV ($\Gamma = 6.0 \pm 1.5$ MeV) and 1831.9 ± 2.8 MeV ($\Gamma = 9.6 \pm 9.9$ MeV).

² BADIER 72 adds all channels and divides the peak into lower and higher mass regions. The data can also be fitted with a single Breit-Wigner of mass 1800 MeV and width 150 MeV.

³ From a fit to inclusive $\Xi\pi$, $\Xi\pi\pi$, and ΛK^- spectra.

⁴ From a fit to inclusive $\Xi\pi$ and $\Xi\pi\pi$ spectra only.

⁵ Including $\Xi\pi\pi$.

⁶ DAUBER 69 uses in part the same data as SMITH 65C.

⁷ For the decay mode $\Xi^-\pi^+\pi^0$ only. This limit includes $\Xi(1530)\pi$.

⁸ Or less. Upper limit for the 3-body decay.

 $\Xi(1820)$ REFERENCES

| | | | | |
|-----------|-----|----------------|------------------------------|--------------------------|
| ABLIKIM | 20C | PRL 124 032002 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ANISOVICH | 12A | EPJ A48 15 | A.V. Anisovich <i>et al.</i> | (BONN, PNPI) |
| ADAMOVICH | 99B | EPJ C11 271 | M.I. Adamovich <i>et al.</i> | (CERN WA89 Collab.) |
| BIAGI | 87 | ZPHY C34 15 | S.F. Biagi <i>et al.</i> | (BRIS, CERN, GEVA+) |
| BIAGI | 87C | ZPHY C34 175 | S.F. Biagi <i>et al.</i> | (BRIS, CERN, GEVA+) JP |
| ASTON | 85B | PR D32 2270 | D. Aston <i>et al.</i> | (SLAC, CARL, CNRC, CINC) |
| JENKINS | 83 | PRL 51 951 | C.M. Jenkins <i>et al.</i> | (FSU, BRAN, LBL+) |
| BIAGI | 81 | ZPHY C9 305 | S.F. Biagi <i>et al.</i> | (BRIS, CAVE, GEVA+) |
| HASSALL | 81 | NP B189 397 | J.K. Hassall <i>et al.</i> | (CAVE, MSU) |
| TEODORO | 78 | PL 77B 451 | D. Teodoro <i>et al.</i> | (AMST, CERN, NIJM+) JP |
| BRIEFEL | 77 | PR D16 2706 | E. Briefel <i>et al.</i> | (BRAN, UMD, SYRA+) |
| Also | | PRL 23 884 | S.P. Apsell <i>et al.</i> | (BRAN, UMD, SYRA+) |
| GAY | 76C | PL 62B 477 | J.B. Gay <i>et al.</i> | (AMST, CERN, NIJM) IJ |
| DIBIANCA | 75 | NP B98 137 | F.A. Dibianca, R.J. Endorf | (CMU) |
| BADIER | 72 | NP B37 429 | J. Badier <i>et al.</i> | (EPOL) |
| APSELL | 70 | PRL 24 777 | S.P. Apsell <i>et al.</i> | (BRAN, UMD, SYRA+) I |
| CRENNELL | 70B | PR D1 847 | D.J. Crennell <i>et al.</i> | (BNL) |

| | | | | |
|--------------|-----|------------------|-------------------------------|-----------------------|
| ALITTI | 69 | PRL 22 79 | J. Alitti <i>et al.</i> | (BNL, SYRA) I |
| DAUBER | 69 | PR 179 1262 | P.M. Dauber <i>et al.</i> | (LRL) |
| TRIPP | 67 | NP B3 10 | R.D. Tripp <i>et al.</i> | (LRL, SLAC, CERN+) |
| BADIER | 65 | PL 16 171 | J. Badier <i>et al.</i> | (EPOL, SACL, AMST) I |
| SMITH | 65B | Athens Conf. 251 | G.A. Smith, J.S. Lindsey | (LRL) |
| SMITH | 65C | PRL 14 25 | G.A. Smith <i>et al.</i> | (LRL) IJP |
| HALSTEINSLID | 63 | Siena Conf. 1 73 | A. Halsteinslid <i>et al.</i> | (BERG, CERN, EPOL+) I |

OTHER RELATED PAPERS

| | | | | |
|---------|----|------------------|--------------------------------|------------------------|
| TEODORO | 78 | PL 77B 451 | D. Teodoro <i>et al.</i> | (AMST, CERN, NIJM+) JP |
| BRIEFEL | 75 | PR D12 1859 | E. Briefel <i>et al.</i> | (BRAN, UMD, SYRA+) |
| SCHMIDT | 73 | Purdue Conf. 363 | P.E. Schmidt | (BRAN) |
| MERRILL | 68 | PR 167 1202 | D.W. Merrill, J. Button-Shafer | (LRL) |
| SMITH | 64 | PRL 13 61 | G.A. Smith <i>et al.</i> | (LRL) IJP |