

# $T_{b\bar{b}1}(10650)^+$

$I^G(J^{PC}) = 1^+(1^{+-})$   
 $I, G, C$  need confirmation.

was  $Z_b(10650)$ ,  $X(10650)^\pm$

Properties incompatible with a  $q\bar{q}$  structure (exotic state). See the review on non- $q\bar{q}$  states.

Observed by BONDAR 12 in  $\Upsilon(5S)$  decays to  $\Upsilon(nS)\pi^+\pi^-$  ( $n = 1, 2, 3$ ) and  $h_b(mP)\pi^+\pi^-$  ( $m = 1, 2$ ).  $J^P = 1^+$  is favored from angular analyses.

## $T_{b\bar{b}1}(10650)^+$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>10652.2±1.5</b>	<sup>1</sup> BONDAR	12	BELL $e^+e^- \rightarrow$ hadrons
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>			
10656.7±5.0 <sup>+1.1</sup> <sub>-3.1</sub>	<sup>2</sup> GARMASH	15	BELL $e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$
10650.7±1.5 <sup>+0.5</sup> <sub>-0.2</sub>	<sup>2</sup> GARMASH	15	BELL $e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$
10651.2±1.0 <sup>+0.4</sup> <sub>-0.3</sub>	<sup>2</sup> GARMASH	15	BELL $e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$
10657 ± 6 ± 3	<sup>3</sup> BONDAR	12	BELL $e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$
10651 ± 2 ± 3	<sup>3</sup> BONDAR	12	BELL $e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$
10652 ± 1 ± 2	<sup>3</sup> BONDAR	12	BELL $e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$
10654 ± 3 ± 1	<sup>3</sup> BONDAR	12	BELL $e^+e^- \rightarrow h_b(1P)\pi^+\pi^-$
10651 ± 2 ± 3	<sup>3</sup> BONDAR	12	BELL $e^+e^- \rightarrow h_b(2P)\pi^+\pi^-$

<sup>1</sup> Average of the BONDAR 12 measurements in separate channels.

<sup>2</sup> Correlated with the corresponding result from BONDAR 12.

<sup>3</sup> Superseded by the average measurement of BONDAR 12.

## $T_{b\bar{b}1}(10650)^+$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>11.5± 2.2</b>	<sup>1</sup> BONDAR	12	BELL $e^+e^- \rightarrow$ hadrons
<b>• • •</b> We do not use the following data for averages, fits, limits, etc. <b>• • •</b>			
12.1 <sup>+11.3 + 2.7</sup> <sub>-4.8 - 0.6</sub>	<sup>2</sup> GARMASH	15	BELL $e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$
14.2± 3.7 <sup>+ 0.9</sup> <sub>- 0.4</sub>	<sup>2</sup> GARMASH	15	BELL $e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$
9.3± 2.2 <sup>+ 0.3</sup> <sub>- 0.5</sub>	<sup>2</sup> GARMASH	15	BELL $e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$
16.3± 9.8 <sup>+ 6.0</sup> <sub>- 2.0</sub>	<sup>3</sup> BONDAR	12	BELL $e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$
13.3± 3.3 <sup>+ 4.0</sup> <sub>- 3.0</sub>	<sup>3</sup> BONDAR	12	BELL $e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$
8.4± 2.0± 2.0	<sup>3</sup> BONDAR	12	BELL $e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$
20.9 <sup>+ 5.4 + 2.1</sup> <sub>- 4.7 - 5.7</sub>	<sup>3</sup> BONDAR	12	BELL $e^+e^- \rightarrow h_b(1P)\pi^+\pi^-$
19 ± 7 ± 11	<sup>3</sup> BONDAR	12	BELL $e^+e^- \rightarrow h_b(2P)\pi^+\pi^-$

<sup>1</sup> Average of the BONDAR 12 measurements in separate channels.<sup>2</sup> Correlated with the corresponding result from BONDAR 12.<sup>3</sup> Superseded by the average measurement of BONDAR 12.

## $T_{b\bar{b}1}(10650)^+$ DECAY MODES

$T_{b\bar{b}1}(10650)^-$  decay modes are charge conjugates of the modes below.

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 \quad \gamma(1S)\pi^+$	( $1.7^{+0.8}_{-0.6}$ ) $\times 10^{-3}$
$\Gamma_2 \quad \gamma(2S)\pi^+$	( $1.4^{+0.6}_{-0.4}$ ) %
$\Gamma_3 \quad \gamma(3S)\pi^+$	( $1.6^{+0.7}_{-0.5}$ ) %
$\Gamma_4 \quad h_b(1P)\pi^+$	( $8.4^{+2.9}_{-2.4}$ ) %
$\Gamma_5 \quad h_b(2P)\pi^+$	( $15 \pm 4$ ) %
$\Gamma_6 \quad B^+\bar{B}^0$	not seen
$\Gamma_7 \quad B^+\bar{B}^{*0} + B^{*+}\bar{B}^0$	not seen
$\Gamma_8 \quad B^{*+}\bar{B}^{*0}$	( $74^{+4}_{-6}$ ) %

## $T_{b\bar{b}1}(10650)^+$ BRANCHING RATIOS

$\Gamma(\gamma(1S)\pi^+)/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma$
<u>VALUE (units <math>10^{-3}</math>)</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
<b><math>1.7^{+0.7+0.3}_{-0.6-0.2}</math></b>	<sup>1</sup> GARMASH    16    BELL $e^+e^- \rightarrow \pi^- B^{*+}\bar{B}^{*0}$
• • • We do not use the following data for averages, fits, limits, etc. • • •	
seen	GARMASH    15    BELL $e^+e^- \rightarrow \gamma(1S)\pi^+\pi^-$
seen	BONDAR    12    BELL $e^+e^- \rightarrow \gamma(1S)\pi^+\pi^-$

<sup>1</sup> Assuming the  $T_{b\bar{b}1}(10650)$  decay width is saturated by the channels  $\pi^+ \gamma(1S, 2S, 3S)$ ,  $\pi^+ h_b(1P, 2P)$ , and  $B^{*+}\bar{B}^{*0}$ , and using the results from BONDAR 12 and MIZUK 16.

$\Gamma(\gamma(2S)\pi^+)/\Gamma_{\text{total}}$	$\Gamma_2/\Gamma$
<u>VALUE (units <math>10^{-2}</math>)</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
<b><math>1.39^{+0.48+0.34}_{-0.38-0.23}</math></b>	<sup>1</sup> GARMASH    16 $e^+e^- \rightarrow \pi^- B^{*+}\bar{B}^{*0}$
• • • We do not use the following data for averages, fits, limits, etc. • • •	
seen	GARMASH    15    BELL $e^+e^- \rightarrow \gamma(2S)\pi^+\pi^-$
seen	BONDAR    12    BELL $e^+e^- \rightarrow \gamma(2S)\pi^+\pi^-$

<sup>1</sup> Assuming the  $T_{b\bar{b}1}(10650)$  decay width is saturated by the channels  $\pi^+ \gamma(1S, 2S, 3S)$ ,  $\pi^+ h_b(1P, 2P)$ , and  $B^{*+}\bar{B}^{*0}$ , and using the results from BONDAR 12 and MIZUK 16.

### $\Gamma(\Upsilon(3S)\pi^+)/\Gamma_{\text{total}}$

$\Gamma_3/\Gamma$

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b><math>1.63^{+0.53+0.39}_{-0.42-0.28}</math></b>	<sup>1</sup> GARMASH	16	BELL $e^+ e^- \rightarrow \pi^- B^* + \bar{B}^* 0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	GARMASH	15	BELL $e^+ e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$
seen	BONDAR	12	BELL $e^+ e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$

<sup>1</sup> Assuming the  $T_{b\bar{b}1}(10650)$  decay width is saturated by the channels  $\pi^+ \Upsilon(1S, 2S, 3S)$ ,  $\pi^+ h_b(1P, 2P)$ , and  $B^* + \bar{B}^* 0$ , and using the results from BONDAR 12 and MIZUK 16.

### $\Gamma(h_b(1P)\pi^+)/\Gamma_{\text{total}}$

$\Gamma_4/\Gamma$

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b><math>8.41^{+2.43+1.49}_{-2.12-1.06}</math></b>	<sup>1</sup> GARMASH	16	BELL $e^+ e^- \rightarrow \pi^- B^* + \bar{B}^* 0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	<sup>2</sup> MIZUK	16	BELL $e^+ e^- \rightarrow h_b(1P)\pi^+\pi^-$
seen	<sup>3</sup> BONDAR	12	BELL $e^+ e^- \rightarrow h_b(1P)\pi^+\pi^-$

<sup>1</sup> Assuming the  $T_{b\bar{b}1}(10650)$  decay width is saturated by the channels  $\pi^+ \Upsilon(1S, 2S, 3S)$ ,  $\pi^+ h_b(1P, 2P)$ , and  $B^* + \bar{B}^* 0$ , and using the results from BONDAR 12 and MIZUK 16.

<sup>2</sup> Using  $e^+ e^-$  energies near the  $\Upsilon(11020)$ .

<sup>3</sup> Using  $e^+ e^-$  energies near the  $\Upsilon(10860)$ .

### $\Gamma(h_b(2P)\pi^+)/\Gamma_{\text{total}}$

$\Gamma_5/\Gamma$

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b><math>14.7^{+3.2+2.8}_{-2.8-2.3}</math></b>	<sup>1</sup> GARMASH	16	BELL $e^+ e^- \rightarrow \pi^- B^* + \bar{B}^* 0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

possibly seen	<sup>2</sup> MIZUK	16	BELL $e^+ e^- \rightarrow h_b(2P)\pi^+\pi^-$
seen	<sup>3</sup> BONDAR	12	BELL $e^+ e^- \rightarrow h_b(2P)\pi^+\pi^-$

<sup>1</sup> Assuming the  $T_{b\bar{b}1}(10650)$  decay width is saturated by the channels  $\pi^+ \Upsilon(1S, 2S, 3S)$ ,  $\pi^+ h_b(1P, 2P)$ , and  $B^* + \bar{B}^* 0$ , and using the results from BONDAR 12 and MIZUK 16.

<sup>2</sup> Using  $e^+ e^-$  energies near the  $\Upsilon(11020)$ .

<sup>3</sup> Using  $e^+ e^-$  energies near the  $\Upsilon(10860)$ .

### $\Gamma(B^+\bar{B}^0)/\Gamma_{\text{total}}$

$\Gamma_6/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>not seen</b>	GARMASH	16	BELL $e^+ e^- \rightarrow \pi^- B^+ \bar{B}^0$

### $[\Gamma(B^+\bar{B}^0) + \Gamma(B^*\bar{B}^0)]/\Gamma_{\text{total}}$

$\Gamma_7/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>not seen</b>	GARMASH	16	BELL $e^+ e^- \rightarrow \pi^- B^+ \bar{B}^0, \pi^- \bar{B}^0 B^*$

$\Gamma(B^{*+}\bar{B}^{*0})/\Gamma_{\text{total}}$  $\Gamma_8/\Gamma$ 

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>73.7<sup>+3.4+2.7</sup><sub>-4.4-3.5</sub></b>	161	<sup>1</sup> GARMASH	16	BELL $e^+ e^- \rightarrow \pi^- B^{*+} \bar{B}^{*0}$

<sup>1</sup> Assuming the  $T_{b\bar{b}1}(10650)$  decay width is saturated by the channels  $\pi^+ \gamma(1S, 2S, 3S)$ ,  $\pi^+ h_b(1P, 2P)$ , and  $B^{*+}\bar{B}^{*0}$ , and using the results from BONDAR 12 and MIZUK 16. Using the mass and width of the  $T_{b\bar{b}1}(10650)$  from BONDAR 12.

$$\begin{aligned} & \Gamma(B^{*+}\bar{B}^{*0}) / [\Gamma(\gamma(1S)\pi^+) + \Gamma(\gamma(2S)\pi^+) + \Gamma(\gamma(3S)\pi^+) + \\ & \Gamma(h_b(1P)\pi^+) + \Gamma(h_b(2P)\pi^+)] \quad \Gamma_8 / (\Gamma_1 + \Gamma_2 + \Gamma_3 + \Gamma_4 + \Gamma_5) \end{aligned}$$

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.80^{+0.69+0.54}_{-0.40-0.36}$	161	<sup>1</sup> GARMASH	16	BELL $e^+ e^- \rightarrow \pi^- B^{*+} \bar{B}^{*0}$
----------------------------------	-----	----------------------	----	--

<sup>1</sup> Combined with the results of BONDAR 12 and MIZUK 16. Not independent from  $T_{b\bar{b}1}(10650)$  branching fractions to  $\pi^+ \gamma(1S, 2S, 3S)$ ,  $\pi^+ h_b(1P, 2P)$ , and  $B^{*+}\bar{B}^{*0}$ .

### $T_{b\bar{b}1}(10650)^+$ REFERENCES

GARMASH	16	PRL 116 212001	A. Garmash <i>et al.</i>	(BELLE Collab.)
MIZUK	16	PRL 117 142001	R. Mizuk <i>et al.</i>	(BELLE Collab.)
GARMASH	15	PR D91 072003	A. Garmash <i>et al.</i>	(BELLE Collab.)
BONDAR	12	PRL 108 122001	A. Bondar <i>et al.</i>	(BELLE Collab.)