

$\chi_{c2}(1P)$ $I^G(J^{PC}) = 0^+(2^{++})$

See the Review on “Branching Ratios of $\psi(2S)$, $\chi_{c0,1,2}$ and $\eta_c(1S)$ ”
before the $\chi_{c0}(1P)$ Listings.

 $\chi_{c2}(1P)$ MASS

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|--------------------------|-----------|---|
| 3556.17 ± 0.07 OUR AVERAGE | | | | |
| 3557.3 ± 1.7 ± 0.7 | 611 | ¹ AAIJ | 17BB LHCb | $p\bar{p} \rightarrow b\bar{b}X \rightarrow 2(K^+K^-)X$ |
| 3556.10 ± 0.06 ± 0.11 | 4.0k | ² AAIJ | 17BI LHCb | $\chi_{c2} \rightarrow J/\psi\mu^+\mu^-$ |
| 3555.3 ± 0.6 ± 2.2 | 2.5k | UEHARA | 08 BELL | $\gamma\gamma \rightarrow \text{hadrons}$ |
| 3555.70 ± 0.59 ± 0.39 | | ABLIKIM | 05G BES2 | $\psi(2S) \rightarrow \gamma\chi_{c2}$ |
| 3556.173 ± 0.123 ± 0.020 | | ANDREOTTI | 05A E835 | $p\bar{p} \rightarrow e^+e^-\gamma$ |
| 3559.9 ± 2.9 | | EISENSTEIN | 01 CLE2 | $e^+e^- \rightarrow e^+e^-\chi_{c2}$ |
| 3556.4 ± 0.7 | | BAI | 99B BES | $\psi(2S) \rightarrow \gamma X$ |
| 3556.22 ± 0.131 ± 0.020 | 585 | ³ ARMSTRONG | 92 E760 | $\bar{p}p \rightarrow e^+e^-\gamma$ |
| 3556.9 ± 0.4 ± 0.5 | 50 | BAGLIN | 86B SPEC | $\bar{p}p \rightarrow e^+e^-X$ |
| 3557.8 ± 0.2 ± 4 | | ⁴ GAISER | 86 CBAL | $\psi(2S) \rightarrow \gamma X$ |
| 3553.4 ± 2.2 | 66 | ⁵ LEMOIGNE | 82 GOLI | $185\pi^-Be \rightarrow \gamma\mu^+\mu^-A$ |
| 3555.9 ± 0.7 | | ⁶ OREGLIA | 82 CBAL | $e^+e^- \rightarrow J/\psi 2\gamma$ |
| 3557 ± 1.5 | 69 | ⁷ HIMEL | 80 MRK2 | $e^+e^- \rightarrow J/\psi 2\gamma$ |
| 3551 ± 11 | 15 | BRANDELIK | 79B DASP | $e^+e^- \rightarrow J/\psi 2\gamma$ |
| 3553 ± 4 | | ⁷ BARTEL | 78B CNTR | $e^+e^- \rightarrow J/\psi 2\gamma$ |
| 3553 ± 4 ± 4 | | ^{7,8} TANENBAUM | 78 MRK1 | e^+e^- |
| 3563 ± 7 | 360 | ⁷ BIDDICK | 77 CNTR | $e^+e^- \rightarrow \gamma X$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 3555.4 ± 1.3 | 53 | UEHARA | 13 BELL | $\gamma\gamma \rightarrow K_S^0 K_S^0$ |
| 3543 ± 10 | 4 | WHITAKER | 76 MRK1 | $e^+e^- \rightarrow J/\psi 2\gamma$ |

¹ From a fit of the $\phi\phi$ invariant mass with the width of $\chi_{c2}(1P)$ fixed to the PDG 16 value.

² AAIJ 17BI reports also $m(\chi_{c2}) - m(\chi_{c1}) = 45.39 \pm 0.07 \pm 0.03$ MeV.

³ Recalculated by ANDREOTTI 05A, using the value of $\psi(2S)$ mass from AULCHENKO 03.

⁴ Using mass of $\psi(2S) = 3686.0$ MeV.

⁵ $J/\psi(1S)$ mass constrained to 3097 MeV.

⁶ Assuming $\psi(2S)$ mass = 3686 MeV and $J/\psi(1S)$ mass = 3097 MeV.

⁷ Mass value shifted by us by amount appropriate for $\psi(2S)$ mass = 3686 MeV and $J/\psi(1S)$ mass = 3097 MeV.

⁸ From a simultaneous fit to radiative and hadronic decay channels.

$\chi_{c2}(1P)$ WIDTH

| <i>VALUE (MeV)</i> | <i>EVTS</i> | <i>DOCUMENT ID</i> | <i>TECN</i> | <i>COMMENT</i> |
|-------------------------------|-------------|-------------------------------------|-------------|--|
| 1.98 ±0.09 OUR FIT | | Error includes scale factor of 1.1. | | |
| 2.00 ±0.11 OUR AVERAGE | | | | |
| 2.10 ±0.20 ±0.02 | 4.0k | AAIJ | 17BI LHCb | $\chi_{c2} \rightarrow J/\psi \mu^+ \mu^-$ |
| 1.915 ±0.188 ±0.013 | | ANDREOTTI | 05A E835 | $p\bar{p} \rightarrow e^+ e^- \gamma$ |
| 1.96 ±0.17 ±0.07 | 585 | ¹ ARMSTRONG | 92 E760 | $\bar{p}p \rightarrow e^+ e^- \gamma$ |
| 2.6 ±1.4 -1.0 | 50 | BAGLIN | 86B SPEC | $\bar{p}p \rightarrow e^+ e^- X$ |
| 2.8 ±2.1 -2.0 | | ² GAISER | 86 CBAL | $\psi(2S) \rightarrow \gamma X$ |

¹ Recalculated by ANDREOTTI 05A.

² Errors correspond to 90% confidence level; authors give only width range.

$\chi_{c2}(1P)$ DECAY MODES

| Mode | Fraction (Γ_i/Γ) | Scale factor/ Confidence level |
|---|-----------------------------------|-----------------------------------|
| Hadronic decays | | |
| $\Gamma_1 2(\pi^+ \pi^-)$ | (1.00 ±0.13) % | S=1.4 |
| $\Gamma_2 \rho\rho$ | | |
| $\Gamma_3 \pi^+ \pi^- \pi^0 \pi^0$ | (1.86 ±0.24) % | |
| $\Gamma_4 \rho^+ \pi^- \pi^0 + \text{c.c.}$ | (2.22 ±0.35) % | |
| $\Gamma_5 4\pi^0$ | (1.13 ±0.15) × 10 ⁻³ | |
| $\Gamma_6 K^+ K^- \pi^0 \pi^0$ | (2.1 ±0.4) × 10 ⁻³ | |
| $\Gamma_7 K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.}$ | (1.41 ±0.20) % | |
| $\Gamma_8 \rho^- K^+ \bar{K}^0 + \text{c.c.}$ | (4.2 ±1.3) × 10 ⁻³ | |
| $\Gamma_9 K^*(892)^0 K^- \pi^+ \rightarrow$ $K^- \pi^+ K^0 \pi^0 + \text{c.c.}$ | (3.0 ±0.8) × 10 ⁻³ | |
| $\Gamma_{10} K^*(892)^0 \bar{K}^0 \pi^0 \rightarrow$ $K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.}$ | (3.9 ±0.9) × 10 ⁻³ | |
| $\Gamma_{11} K^*(892)^- K^+ \pi^0 \rightarrow$ $K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.}$ | (3.8 ±0.8) × 10 ⁻³ | |
| $\Gamma_{12} K^*(892)^+ \bar{K}^0 \pi^- \rightarrow$ $K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.}$ | (3.0 ±0.8) × 10 ⁻³ | |
| $\Gamma_{13} K^+ K^- \eta \pi^0$ | (1.3 ±0.4) × 10 ⁻³ | |
| $\Gamma_{14} K^+ K^- \pi^+ \pi^-$ | (8.3 ±1.1) × 10 ⁻³ | S=1.2 |
| $\Gamma_{15} K^+ K^- \pi^+ \pi^- \pi^0$ | (1.17 ±0.13) % | |
| $\Gamma_{16} K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$ | (7.3 ±0.8) × 10 ⁻³ | |
| $\Gamma_{17} K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.}$ | (2.1 ±1.0) × 10 ⁻³ | |
| $\Gamma_{18} K^*(892)^0 \bar{K}^*(892)^0$ | (2.2 ±0.9) × 10 ⁻³ | S=2.3 |
| $\Gamma_{19} 3(\pi^+ \pi^-)$ | (1.53 ±0.19) % | S=3.8 |
| $\Gamma_{20} \phi\phi$ | (1.23 ±0.07) × 10 ⁻³ | S=1.9 |
| $\Gamma_{21} \phi\phi\eta$ | (5.4 ±0.7) × 10 ⁻⁴ | |
| $\Gamma_{22} \omega\omega$ | (8.6 ±1.0) × 10 ⁻⁴ | |
| $\Gamma_{23} \omega K^+ K^-$ | (7.3 ±0.9) × 10 ⁻⁴ | |

| | | | |
|---------------|--|------------------------------------|--------|
| Γ_{24} | $\omega\phi$ | $(9.7 \pm 2.8) \times 10^{-6}$ | |
| Γ_{25} | $\pi\pi$ | $(2.27 \pm 0.10) \times 10^{-3}$ | |
| Γ_{26} | $\rho^0\pi^+\pi^-$ | $(3.6 \pm 1.5) \times 10^{-3}$ | |
| Γ_{27} | $\pi^+\pi^-\pi^0$ (non-resonant) | $(2.0 \pm 0.4) \times 10^{-5}$ | |
| Γ_{28} | $\rho(770)^{\pm}\pi^{\mp}$ | $(6 \pm 4) \times 10^{-6}$ | |
| Γ_{29} | $\pi^+\pi^-\eta$ | $(4.9 \pm 1.3) \times 10^{-4}$ | |
| Γ_{30} | $\pi^+\pi^-\eta'$ | $(5.1 \pm 1.9) \times 10^{-4}$ | |
| Γ_{31} | $\eta\eta$ | $(5.5 \pm 0.5) \times 10^{-4}$ | |
| Γ_{32} | K^+K^- | $(1.02 \pm 0.15) \times 10^{-3}$ | S=2.3 |
| Γ_{33} | $K_S^0K_S^0$ | $(5.3 \pm 0.4) \times 10^{-4}$ | |
| Γ_{34} | $K^*(892)^{\pm}K^{\mp}$ | $(1.46 \pm 0.21) \times 10^{-4}$ | |
| Γ_{35} | $K^*(892)^0\bar{K}^0 + \text{c.c.}$ | $(1.27 \pm 0.27) \times 10^{-4}$ | |
| Γ_{36} | $K_2^*(1430)^{\pm}K^{\mp}$ | $(1.51 \pm 0.13) \times 10^{-3}$ | |
| Γ_{37} | $K_2^*(1430)^0\bar{K}^0 + \text{c.c.}$ | $(1.27 \pm 0.17) \times 10^{-3}$ | |
| Γ_{38} | $K_3^*(1780)^{\pm}K^{\mp}$ | $(5.3 \pm 0.8) \times 10^{-4}$ | |
| Γ_{39} | $K_3^*(1780)^0\bar{K}^0 + \text{c.c.}$ | $(5.7 \pm 2.1) \times 10^{-4}$ | |
| Γ_{40} | $a_2(1320)^0\pi^0$ | $(1.31 \pm 0.35) \times 10^{-3}$ | |
| Γ_{41} | $a_2(1320)^{\pm}\pi^{\mp}$ | $(1.8 \pm 0.6) \times 10^{-3}$ | |
| Γ_{42} | $\bar{K}^0K^+\pi^- + \text{c.c.}$ | $(1.30 \pm 0.19) \times 10^{-3}$ | |
| Γ_{43} | $K^+K^-\pi^0$ | $(3.1 \pm 0.8) \times 10^{-4}$ | |
| Γ_{44} | $K^+K^-\eta$ | $< 3.3 \times 10^{-4}$ | CL=90% |
| Γ_{45} | $K^+K^-\eta'(958)$ | $(1.94 \pm 0.34) \times 10^{-4}$ | |
| Γ_{46} | $\eta\eta'$ | $(2.2 \pm 0.5) \times 10^{-5}$ | |
| Γ_{47} | $\eta'\eta'$ | $(4.6 \pm 0.6) \times 10^{-5}$ | |
| Γ_{48} | $\pi^+\pi^-K_S^0K_S^0$ | $(2.2 \pm 0.5) \times 10^{-3}$ | |
| Γ_{49} | $K^+K^-K_S^0K_S^0$ | $< 4 \times 10^{-4}$ | CL=90% |
| Γ_{50} | $K_S^0K_S^0K_S^0K_S^0$ | $(1.15 \pm 0.18) \times 10^{-4}$ | |
| Γ_{51} | $K^+K^-K^+K^-$ | $(1.67 \pm 0.22) \times 10^{-3}$ | S=1.1 |
| Γ_{52} | $K^+K^-\phi$ | $(1.45 \pm 0.30) \times 10^{-3}$ | |
| Γ_{53} | $\bar{K}^0K^+\pi^-\phi + \text{c.c.}$ | $(4.8 \pm 0.7) \times 10^{-3}$ | |
| Γ_{54} | $K^+K^-\pi^0\phi$ | $(2.7 \pm 0.5) \times 10^{-3}$ | |
| Γ_{55} | $\phi\pi^+\pi^-\pi^0$ | $(9.3 \pm 1.2) \times 10^{-4}$ | |
| Γ_{56} | $p\bar{p}$ | $(7.3 \pm 0.4) \times 10^{-5}$ | S=1.1 |
| Γ_{57} | $p\bar{p}\pi^0$ | $(4.7 \pm 0.4) \times 10^{-4}$ | |
| Γ_{58} | $p\bar{p}\eta$ | $(1.77 \pm 0.25) \times 10^{-4}$ | |
| Γ_{59} | $p\bar{p}\omega$ | $(3.7 \pm 0.4) \times 10^{-4}$ | |
| Γ_{60} | $p\bar{p}\phi$ | $(2.8 \pm 0.9) \times 10^{-5}$ | |
| Γ_{61} | $p\bar{p}\pi^+\pi^-$ | $(1.32 \pm 0.34) \times 10^{-3}$ | |
| Γ_{62} | $p\bar{p}\pi^0\pi^0$ | $(8.0 \pm 2.4) \times 10^{-4}$ | |
| Γ_{63} | $p\bar{p}K^+K^-$ (non-resonant) | $(1.94 \pm 0.33) \times 10^{-4}$ | |
| Γ_{64} | $p\bar{p}K_S^0K_S^0$ | $< 7.9 \times 10^{-4}$ | CL=90% |
| Γ_{65} | $p\bar{n}\pi^-$ | $(8.7 \pm 1.0) \times 10^{-4}$ | |
| Γ_{66} | $\bar{p}n\pi^+$ | $(9.1 \pm 0.8) \times 10^{-4}$ | |

| | | | |
|---------------|--|----------------------------------|--------|
| Γ_{67} | $p\bar{n}\pi^-\pi^0$ | $(2.21 \pm 0.18) \times 10^{-3}$ | |
| Γ_{68} | $\bar{p}n\pi^+\pi^0$ | $(2.15 \pm 0.19) \times 10^{-3}$ | |
| Γ_{69} | $\Lambda\bar{\Lambda}$ | $(1.86 \pm 0.16) \times 10^{-4}$ | |
| Γ_{70} | $\Lambda\bar{\Lambda}\pi^+\pi^-$ | $(1.28 \pm 0.16) \times 10^{-3}$ | |
| Γ_{71} | $\Lambda\bar{\Lambda}\pi^+\pi^-$ (non-resonant) | $(6.7 \pm 1.5) \times 10^{-4}$ | |
| Γ_{72} | $\Sigma(1385)^+\bar{\Lambda}\pi^- + \text{c.c.}$ | $< 4 \times 10^{-4}$ | CL=90% |
| Γ_{73} | $\Sigma(1385)^-\bar{\Lambda}\pi^+ + \text{c.c.}$ | $< 6 \times 10^{-4}$ | CL=90% |
| Γ_{74} | $\Lambda\bar{\Lambda}\eta$ | $(1.07 \pm 0.26) \times 10^{-4}$ | |
| Γ_{75} | $K^+\bar{p}\Lambda + \text{c.c.}$ | $(7.9 \pm 0.6) \times 10^{-4}$ | |
| Γ_{76} | $nK_S^0\bar{\Lambda} + \text{c.c.}$ | $(3.64 \pm 0.29) \times 10^{-4}$ | |
| Γ_{77} | $K^*(892)^+\bar{p}\Lambda + \text{c.c.}$ | $(8.3 \pm 1.2) \times 10^{-4}$ | |
| Γ_{78} | $K^+\bar{p}\Lambda(1520) + \text{c.c.}$ | $(2.9 \pm 0.7) \times 10^{-4}$ | |
| Γ_{79} | $\Lambda(1520)\bar{\Lambda}(1520)$ | $(4.7 \pm 1.5) \times 10^{-4}$ | |
| Γ_{80} | $\Sigma^0\bar{\Sigma}^0$ | $(3.7 \pm 0.6) \times 10^{-5}$ | |
| Γ_{81} | $\Sigma^+\bar{p}K_S^0 + \text{c.c.}$ | $(8.4 \pm 1.0) \times 10^{-5}$ | |
| Γ_{82} | $\Sigma^0\bar{p}K^+ + \text{c.c.}$ | $(9.3 \pm 0.8) \times 10^{-5}$ | |
| Γ_{83} | $\Sigma^+\bar{\Sigma}^-$ | $(3.4 \pm 0.7) \times 10^{-5}$ | |
| Γ_{84} | $\Sigma^-\bar{\Sigma}^+$ | $(4.5 \pm 1.8) \times 10^{-5}$ | |
| Γ_{85} | $\Sigma(1385)^+\bar{\Sigma}(1385)^-$ | $< 1.6 \times 10^{-4}$ | CL=90% |
| Γ_{86} | $\Sigma(1385)^-\bar{\Sigma}(1385)^+$ | $< 8 \times 10^{-5}$ | CL=90% |
| Γ_{87} | $K^-\Lambda\Xi^+ + \text{c.c.}$ | $(1.80 \pm 0.32) \times 10^{-4}$ | |
| Γ_{88} | $\Xi^0\bar{\Xi}^0$ | $(1.86 \pm 0.22) \times 10^{-4}$ | |
| Γ_{89} | $\Xi^-\bar{\Xi}^+$ | $(1.46 \pm 0.12) \times 10^{-4}$ | |
| Γ_{90} | $\Omega^-\bar{\Omega}^+$ | $(4.52 \pm 0.30) \times 10^{-5}$ | |
| Γ_{91} | $J/\psi(1S)\pi^+\pi^-\pi^0$ | $< 1.5 \%$ | CL=90% |
| Γ_{92} | $\pi^0\eta_c$ | $< 3.2 \times 10^{-3}$ | CL=90% |
| Γ_{93} | $\eta_c(1S)\pi^+\pi^-$ | $< 5.4 \times 10^{-3}$ | CL=90% |

Radiative decays

| | | | |
|----------------|------------------------|----------------------------------|--------|
| Γ_{94} | $\gamma J/\psi(1S)$ | $(19.5 \pm 0.8) \%$ | S=1.5 |
| Γ_{95} | $\gamma\rho^0$ | $< 1.9 \times 10^{-5}$ | CL=90% |
| Γ_{96} | $\gamma\omega$ | $< 6 \times 10^{-6}$ | CL=90% |
| Γ_{97} | $\gamma\phi$ | $< 8 \times 10^{-6}$ | CL=90% |
| Γ_{98} | $\gamma\gamma$ | $(2.92 \pm 0.12) \times 10^{-4}$ | S=1.3 |
| Γ_{99} | $e^+e^-J/\psi(1S)$ | $(2.20 \pm 0.15) \times 10^{-3}$ | |
| Γ_{100} | $\mu^+\mu^-J/\psi(1S)$ | $(2.07 \pm 0.34) \times 10^{-4}$ | |

CONSTRAINED FIT INFORMATION

A multiparticle fit to $\chi_{c1}(1P)$, $\chi_{c0}(1P)$, $\chi_{c2}(1P)$, and $\psi(2S)$ with 4 total widths, a partial width, 25 combinations of partial widths obtained from integrated cross section, and 86 branching ratios uses 253 measurements to determine 49 parameters. The overall fit has a $\chi^2 = 389.6$ for 204 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$.

| | | | | | | | | | | | |
|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|--|
| x_{14} | 7 | | | | | | | | | | |
| x_{17} | 2 | 26 | | | | | | | | | |
| x_{18} | 2 | 2 | 0 | | | | | | | | |
| x_{20} | 4 | 4 | 1 | 1 | | | | | | | |
| x_{25} | 4 | 4 | 1 | 2 | 5 | | | | | | |
| x_{26} | 26 | 2 | 1 | 1 | 1 | 1 | | | | | |
| x_{31} | 2 | 2 | 0 | 1 | 3 | 16 | 1 | | | | |
| x_{32} | 3 | 3 | 1 | 1 | 2 | 7 | 1 | 3 | | | |
| x_{33} | 3 | 3 | 1 | 1 | 3 | 16 | 1 | 9 | 4 | | |
| x_{42} | 1 | 1 | 0 | 1 | 2 | 8 | 0 | 5 | 2 | 5 | |
| x_{51} | 3 | 3 | 1 | 1 | 3 | 9 | 1 | 5 | 3 | 5 | |
| x_{56} | 6 | 6 | 1 | 2 | 4 | 12 | 2 | 6 | 4 | 7 | |
| x_{69} | 2 | 2 | 0 | 1 | 3 | 15 | 0 | 8 | 3 | 8 | |
| x_{94} | 12 | 12 | 3 | 3 | 9 | 25 | 4 | 13 | 8 | 15 | |
| x_{98} | -10 | -10 | -3 | -1 | -3 | 19 | -3 | 12 | 1 | 9 | |
| Γ | -18 | -18 | -5 | -5 | -12 | -26 | -6 | -13 | -10 | -16 | |
| | x_1 | x_{14} | x_{17} | x_{18} | x_{20} | x_{25} | x_{26} | x_{31} | x_{32} | x_{33} | |
| | x_{51} | 3 | | | | | | | | | |
| | x_{56} | 4 | 5 | | | | | | | | |
| | x_{69} | 4 | 4 | 6 | | | | | | | |
| | x_{94} | 7 | 10 | -12 | 12 | | | | | | |
| | x_{98} | 6 | 2 | 27 | 11 | 11 | | | | | |
| | Γ | -8 | -12 | -42 | -13 | -48 | -47 | | | | |
| | | x_{42} | x_{51} | x_{56} | x_{69} | x_{94} | x_{98} | | | | |

$\chi_{c2}(1P)$ PARTIAL WIDTHS

$\chi_{c2}(1P) \Gamma(i)\Gamma(\gamma J/\psi(1S))/\Gamma(\text{total})$

$$\Gamma(p\bar{p}) \times \Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}} \quad \Gamma_{56}\Gamma_{94}/\Gamma$$

| VALUE (eV) | DOCUMENT ID | TECN | COMMENT |
|------------|-------------|------|---------|
|------------|-------------|------|---------|

28.0±1.3 OUR FIT Error includes scale factor of 1.1.

27.5±1.5 OUR AVERAGE

| | | |
|--------------|----------------------------------|---------------------------------------|
| 27.0±1.5±1.1 | ¹ ANDREOTTI 05A E835 | $p\bar{p} \rightarrow e^+ e^- \gamma$ |
| 27.7±1.5±2.0 | ^{1,2} ARMSTRONG 92 E760 | $\bar{p}p \rightarrow e^+ e^- \gamma$ |
| 36 ± 8 | ¹ BAGLIN 86B SPEC | $\bar{p}p \rightarrow e^+ e^- X$ |

¹ Calculated by us using $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.0593 \pm 0.0010$.

² Recalculated by ANDREOTTI 05A.

$$\Gamma(\gamma\gamma) \times \Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}} \quad \Gamma_{98}\Gamma_{94}/\Gamma$$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------|------|-------------|------|---------|
|------------|------|-------------|------|---------|

113 ± 5 OUR FIT Error includes scale factor of 1.3.

123 ± 6 OUR AVERAGE

| | | | | |
|-----------------|------|-----------------------------|-----|--|
| 124.1± 2.5± 5.9 | 4960 | ¹ SEINO | 23 | BELL $e^+ e^- \rightarrow e^+ e^- \chi_{c2}$ |
| 111 ± 12 ± 9 | 147 | ² DOBBS | 06 | CLE3 $e^+ e^- \rightarrow e^+ e^- \chi_{c2}$ |
| 139 ± 55 ± 21 | | ^{2,3} ACCIARRI | 99E | L3 $e^+ e^- \rightarrow e^+ e^- \chi_{c2}$ |
| 242 ± 65 ± 51 | | ^{2,4} ACKER...K... | 98 | OPAL $e^+ e^- \rightarrow e^+ e^- \chi_{c2}$ |
| 150 ± 42 ± 36 | | ^{2,5} DOMINICK | 94 | CLE2 $e^+ e^- \rightarrow e^+ e^- \chi_{c2}$ |
| 470 ± 240 ± 120 | | ^{2,6} BAUER | 93 | TPC $e^+ e^- \rightarrow e^+ e^- \chi_{c2}$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|--------------|-----|--------------------|-----|--|
| 114 ± 11 ± 9 | 136 | ^{2,7} ABE | 02T | BELL $e^+ e^- \rightarrow e^+ e^- \chi_{c2}$ |
|--------------|-----|--------------------|-----|--|

¹ Calculated from the measured $\Gamma_{\gamma\gamma} \times B(\chi_{c2}(1S) \rightarrow \gamma J/\psi(1S)) \times B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 14.8 \pm 0.3 \pm 0.7$ eV, using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 11.93 \pm 0.05\%$.

² Calculated by us using $B(J/\psi \rightarrow \ell^+ \ell^-) = 0.1187 \pm 0.0008$.

³ The value for $\Gamma(\chi_{c2} \rightarrow \gamma\gamma)$ reported in ACCIARRI 99E is derived using $B(\chi_{c2} \rightarrow \gamma J/\psi(1S)) \times B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.0162 \pm 0.0014$.

⁴ The value for $\Gamma(\chi_{c2} \rightarrow \gamma\gamma)$ reported in ACKERSTAFF,K 98 is derived using $B(\chi_{c2} \rightarrow \gamma J/\psi(1S)) = 0.135 \pm 0.011$ and $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1203 \pm 0.0038$.

⁵ The value for $\Gamma(\chi_{c2} \rightarrow \gamma\gamma)$ reported in DOMINICK 94 is derived using $B(\chi_{c2} \rightarrow \gamma J/\psi(1S)) = 0.135 \pm 0.011$, $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.0627 \pm 0.0020$, and $B(J/\psi(1S) \rightarrow \mu^+ \mu^-) = 0.0597 \pm 0.0025$.

⁶ The value for $\Gamma(\chi_{c2} \rightarrow \gamma\gamma)$ reported in BAUER 93 is derived using $B(\chi_{c2} \rightarrow \gamma J/\psi(1S)) = 0.135 \pm 0.011$, $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.0627 \pm 0.0020$, and $B(J/\psi(1S) \rightarrow \mu^+ \mu^-) = 0.0597 \pm 0.0025$.

⁷ All systematic errors added in quadrature. Superseded by SEINO 23.

 $\chi_{c2}(1P) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$$\Gamma(2(\pi^+\pi^-)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_1\Gamma_{98}/\Gamma$$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------|------|-------------|------|---------|
|------------|------|-------------|------|---------|

5.8 ±0.7 OUR FIT Error includes scale factor of 1.4.

5.2 ±0.7 OUR AVERAGE

| | | | | |
|-----------------|----------|---------------|------|--|
| 5.01±0.44±0.55 | 1597±138 | UEHARA 08 | BELL | $\gamma\gamma \rightarrow \chi_{c2} \rightarrow 2(\pi^+\pi^-)$ |
| 6.4 ± 1.8 ± 0.8 | | EISENSTEIN 01 | CLE2 | $e^+ e^- \rightarrow e^+ e^- \chi_{c2}$ |

$\Gamma(\rho\rho) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_2\Gamma_{98}/\Gamma$

| VALUE (eV) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-----|------|-------------|------|--|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| <7.8 | 90 | <598 | UEHARA | 08 | BELL $\gamma\gamma \rightarrow \chi_{c2} \rightarrow 2(\pi^+ \pi^-)$ |

$\Gamma(K^+ K^- \pi^+ \pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{14}\Gamma_{98}/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------------|----------|-------------------------------------|------|---|
| 4.8 ± 0.6 OUR FIT | | Error includes scale factor of 1.2. | | |
| 4.42 ± 0.42 ± 0.53 | 780 ± 74 | UEHARA | 08 | BELL $\gamma\gamma \rightarrow \chi_{c2} \rightarrow K^+ K^- \pi^+ \pi^-$ |

$\Gamma(K^+ K^- \pi^+ \pi^- \pi^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{15}\Gamma_{98}/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------|------|-----------------|------|--|
| 6.5 ± 0.9 ± 1.5 | 1250 | DEL-AMO-SA..11M | BABR | $\gamma\gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ |

$\Gamma(K^*(892)^0 \bar{K}^*(892)^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{18}\Gamma_{98}/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|----------|-------------------------------------|------|---|
| 1.3 ± 0.5 OUR FIT | | Error includes scale factor of 2.3. | | |
| 0.8 ± 0.17 ± 0.27 | 151 ± 30 | UEHARA | 08 | BELL $\gamma\gamma \rightarrow \chi_{c2} \rightarrow K^+ K^- \pi^+ \pi^-$ |

$\Gamma(\phi\phi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{20}\Gamma_{98}/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------------|-------------------------------------|------|--|
| 0.71 ± 0.05 OUR FIT | | Error includes scale factor of 1.5. | | |
| 0.62 ± 0.07 ± 0.05 | 89 ± 11 | ¹ LIU | 12B | BELL $\gamma\gamma \rightarrow 2(K^+ K^-)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.58 ± 0.18 ± 0.16 | 26.5 ± 8.1 | UEHARA | 08 | BELL $\gamma\gamma \rightarrow \chi_{c2} \rightarrow 2(K^+ K^-)$ |

¹ Supersedes UEHARA 08. Using $B(\phi \rightarrow K^+ K^-) = (48.9 \pm 0.5)\%$.

$\Gamma(\omega\omega) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{22}\Gamma_{98}/\Gamma$

| VALUE (eV) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|------------------|------|--|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| <0.64 | 90 | ¹ LIU | 12B | BELL $\gamma\gamma \rightarrow 2(\pi^+ \pi^- \pi^0)$ |
| ¹ Using $B(\omega \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7)\%$. | | | | |

$\Gamma(\omega\phi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{24}\Gamma_{98}/\Gamma$

| VALUE (eV) | CL% | DOCUMENT ID | TECN | COMMENT |
|--|-----|------------------|------|---|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| <0.04 | 90 | ¹ LIU | 12B | BELL $\gamma\gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ |
| ¹ Using $B(\phi \rightarrow K^+ K^-) = (48.9 \pm 0.5)\%$ and $B(\omega \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7)\%$. | | | | |

$\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{25}\Gamma_{98}/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------------|------|-------------------------------------|------|---------|
| 1.31 ± 0.08 OUR FIT | | Error includes scale factor of 1.2. | | |
| 1.18 ± 0.25 OUR AVERAGE | | | | |

| | | | | |
|--------------------|---------|-----------------------|----|---|
| 1.44 ± 0.54 ± 0.47 | 34 ± 13 | ¹ UEHARA | 09 | BELL $10.6 e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$ |
| 1.14 ± 0.21 ± 0.17 | 54 ± 10 | ² NAKAZAWA | 05 | BELL $10.6 e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$ |

¹ We multiplied the measurement by 3 to convert from $\pi^0 \pi^0$ to $\pi\pi$. Interference with the continuum included.

² We have multiplied $\pi^+ \pi^-$ measurement by 3/2 to obtain $\pi\pi$.

| $\Gamma(\rho^0 \pi^+ \pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ | $\Gamma_{26}\Gamma_{98}/\Gamma$ | | | |
|--|---------------------------------|--------------------|-------------|--|
| <u>VALUE (eV)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
| 2.1±0.9 OUR FIT | | | | |
| 3.2±1.9±0.5 | 986 ± 578 | UEHARA | 08 | BELL $\gamma\gamma \rightarrow \chi_{c2} \rightarrow 2(\pi^+ \pi^-)$ |

| $\Gamma(\eta\eta) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ | $\Gamma_{31}\Gamma_{98}/\Gamma$ | | | |
|--|---------------------------------|--------------------|-------------|--|
| <u>VALUE (eV)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
| 0.53±0.22±0.09 | 8 | 1 UEHARA | 10A | BELL $10.6 e^+ e^- \rightarrow e^+ e^- \eta\eta$ |

¹ Interference with the continuum not included.

| $\Gamma(K^+ K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ | $\Gamma_{32}\Gamma_{98}/\Gamma$ | | | |
|---|-------------------------------------|--------------------|-------------|--|
| <u>VALUE (eV)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
| 0.59±0.09 OUR FIT | Error includes scale factor of 2.1. | | | |
| 0.44±0.11±0.07 | 33 ± 8 | NAKAZAWA 05 | BELL | $10.6 e^+ e^- \rightarrow e^+ e^- K^+ K^-$ |

| $\Gamma(K_S^0 K_S^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ | $\Gamma_{33}\Gamma_{98}/\Gamma$ | | | |
|---|-------------------------------------|--------------------|-------------|--|
| <u>VALUE (eV)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
| 0.307±0.026 OUR FIT | Error includes scale factor of 1.1. | | | |
| 0.27 ±0.07 ±0.03 | 53 | 1 UEHARA | 13 | BELL $\gamma\gamma \rightarrow K_S^0 K_S^0$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.31 ± 0.05 ± 0.03 | 38 ± 7 | CHEN | 07B | BELL $e^+ e^- \rightarrow e^+ e^- \chi_{c2}$ |

¹ Supersedes CHEN 07B.

| $\Gamma(\bar{K}^0 K^+ \pi^- + \text{c.c.}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ | $\Gamma_{42}\Gamma_{98}/\Gamma$ | | | |
|---|---------------------------------|------------------------|--|----------------|
| <u>VALUE (eV)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
| 0.75±0.11 OUR FIT | | | | |
| 1.20±0.33±0.13 | 126 | 1 DEL-AMO-SA..11M BABR | $\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$ | |

¹ We have multiplied $\bar{K}K\pi$ by 2/3 to obtain $\bar{K}^0 K^+ \pi^- + \text{c.c.}$

| $\Gamma(K^+ K^- K^+ K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ | $\Gamma_{51}\Gamma_{98}/\Gamma$ | | | |
|---|-------------------------------------|--------------------|-------------|--|
| <u>VALUE (eV)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
| 0.97±0.13 OUR FIT | Error includes scale factor of 1.1. | | | |
| 1.10±0.21±0.15 | 126 ± 24 | UEHARA | 08 | BELL $\gamma\gamma \rightarrow \chi_{c2} \rightarrow 2(K^+ K^-)$ |

| $\Gamma(\eta_c(1S)\pi^+\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ | $\Gamma_{93}\Gamma_{98}/\Gamma$ | | | |
|--|---------------------------------|--------------------|-------------|--|
| <u>VALUE (eV)</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
| <15.7 | 90 | LEES | 12AE BABR | $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \eta_c$ |

$\chi_{c2}(1P)$ BRANCHING RATIOS

— HADRONIC DECAYS —

| $\Gamma(\rho^0 \pi^+ \pi^-)/\Gamma(2(\pi^+ \pi^-))$ | Γ_{26}/Γ_1 | | |
|---|------------------------|-------------|---|
| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
| 0.36±0.15 OUR FIT | | | |
| 0.31±0.17 | TANENBAUM 78 | MRK1 | $\psi(2S) \rightarrow \gamma \chi_{c2}$ |

$\Gamma(\pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_3/Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-------|-------------|----------|--|
| 1.86±0.23±0.05 | 903.5 | 1 HE | 08B CLEO | $e^+e^- \rightarrow \gamma h^+h^-h^0h^0$ |

¹ HE 08B reports $1.87 \pm 0.07 \pm 0.22 \pm 0.13$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\rho^+\pi^-\pi^0+c.c.)/\Gamma_{\text{total}}$ Γ_4/Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|--------|-------------|----------|--|
| 2.22±0.34±0.05 | 1031.9 | 1,2 HE | 08B CLEO | $e^+e^- \rightarrow \gamma h^+h^-h^0h^0$ |

¹ HE 08B reports $2.23 \pm 0.11 \pm 0.32 \pm 0.16$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \rho^+\pi^-\pi^0 + c.c.)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Calculated by us. We have added the values from HE 08B for $\rho^+\pi^-\pi^0$ and $\rho^-\pi^+\pi^0$ decays assuming uncorrelated statistical and fully correlated systematic uncertainties.

 $\Gamma(4\pi^0)/\Gamma_{\text{total}}$ Γ_5/Γ

| VALUE (units 10^{-3}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|----------|---|
| 1.13±0.15±0.03 | 1164 | 1 ABLIKIM | 11A BES3 | $e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c2}$ |

¹ ABLIKIM 11A reports $(1.21 \pm 0.05 \pm 0.16) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow 4\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(K^+K^-\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_6/Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|------|-------------|----------|--|
| 0.21±0.04±0.01 | 76.9 | 1 HE | 08B CLEO | $e^+e^- \rightarrow \gamma h^+h^-h^0h^0$ |

¹ HE 08B reports $0.21 \pm 0.03 \pm 0.03 \pm 0.01$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+K^-\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(K^+\pi^-\bar{K}^0\pi^0+c.c.)/\Gamma_{\text{total}}$ Γ_7/Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-------|-------------|----------|--|
| 1.41±0.20±0.03 | 211.6 | 1 HE | 08B CLEO | $e^+e^- \rightarrow \gamma h^+h^-h^0h^0$ |

¹ HE 08B reports $1.41 \pm 0.11 \pm 0.16 \pm 0.10$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+\pi^-\bar{K}^0\pi^0 + c.c.)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\rho^- K^+ \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_8/Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|------|-------------|----------|--|
| 0.42±0.13±0.01 | 62.9 | 1 HE | 08B CLEO | $e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$ |

¹ HE 08B reports $0.42 \pm 0.11 \pm 0.06 \pm 0.03$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \rho^- K^+ \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))] \text{ assuming } \text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(K^*(892)^0 K^- \pi^+ \rightarrow K^- \pi^+ K^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_9/Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|------|-------------|----------|--|
| 0.30±0.08±0.01 | 38.7 | 1 HE | 08B CLEO | $e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$ |

¹ HE 08B reports $0.30 \pm 0.07 \pm 0.04 \pm 0.02$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^0 K^- \pi^+ \rightarrow K^- \pi^+ K^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))] \text{ assuming } \text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(K^*(892)^0 \bar{K}^0 \pi^0 \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{10}/Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|------|-------------|----------|--|
| 0.39±0.09±0.01 | 63.0 | 1 HE | 08B CLEO | $e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$ |

¹ HE 08B reports $0.39 \pm 0.07 \pm 0.05 \pm 0.03$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^0 \bar{K}^0 \pi^0 \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))] \text{ assuming } \text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(K^*(892)^- K^+ \pi^0 \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{11}/Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|------|-------------|----------|--|
| 0.38±0.08±0.01 | 51.1 | 1 HE | 08B CLEO | $e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$ |

¹ HE 08B reports $0.38 \pm 0.07 \pm 0.04 \pm 0.03$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^- K^+ \pi^0 \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))] \text{ assuming } \text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(K^*(892)^+ \bar{K}^0 \pi^- \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{12}/Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|------|-------------|----------|--|
| 0.30±0.08±0.01 | 39.3 | 1 HE | 08B CLEO | $e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$ |

¹ HE 08B reports $0.30 \pm 0.07 \pm 0.04 \pm 0.02$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^+ \bar{K}^0 \pi^- \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))] \text{ assuming } \text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+ K^- \eta\pi^0)/\Gamma_{\text{total}}$ Γ_{13}/Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|----------|--|
| 0.130±0.045±0.003 | 22.9 | 1 HE | 08B CLEO | $e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$ |

¹ HE 08B reports $0.13 \pm 0.04 \pm 0.02 \pm 0.01$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- \eta\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_{15}/Γ

| VALUE (units 10^{-3}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|----------|--|
| 11.69±0.13±1.31 | 11k | 1 ABLIKIM | 13B BES3 | $e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c2}$ |

¹ Using 1.06×10^8 $\psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c2}\gamma) = (8.72 \pm 0.34)\%$.

 $\Gamma(K_S^0 K^\pm \pi^\mp \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{16}/Γ

| VALUE (units 10^{-3}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|----------|--|
| 7.30±0.11±0.75 | 4.5k | 1 ABLIKIM | 13B BES3 | $e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c2}$ |

¹ Using 1.06×10^8 $\psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c2}\gamma) = (8.72 \pm 0.34)\%$.

 $\Gamma(K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.})/\Gamma(K^+ K^- \pi^+ \pi^-)$ Γ_{17}/Γ_{14}

| VALUE | | DOCUMENT ID | TECN | COMMENT |
|--------------------------|--|--------------|------|--|
| 0.25±0.13 OUR FIT | | | | |
| 0.25±0.13 | | TANENBAUM 78 | MRK1 | $\psi(2S) \rightarrow \gamma\chi_{c2}$ |

 $\Gamma(3(\pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{19}/Γ

| VALUE (units 10^{-3}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------------|-------------------------------------|----------------|----------|--|
| 15.3±1.9 OUR AVERAGE | Error includes scale factor of 3.8. | | | |
| 15.9±0.4±0.4 | 112K | 1 ABLIKIM | 22Q BES3 | $\psi(2S) \rightarrow \gamma 3(\pi^+ \pi^-)$ |
| 8.6±0.9±1.6 | | 2 BAI | 99B BES | $\psi(2S) \rightarrow \gamma\chi_{c2}$ |
| 8.7±5.9±0.4 | | 2 TANENBAUM 78 | MRK1 | $\psi(2S) \rightarrow \gamma\chi_{c2}$ |

¹ ABLIKIM 22Q reports $(1.565 \pm 0.005 \pm 0.048) \times 10^{-2}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow 3(\pi^+ \pi^-))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Rescaled by us using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (8.3 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (32.6 \pm 0.5)\%$. Multiplied by a factor of 2 to convert from $K_S^0 K^+ \pi^-$ to $K^0 K^+ \pi^-$ decay.

 $\Gamma(\phi\phi)/\Gamma_{\text{total}}$ Γ_{20}/Γ

| VALUE (units 10^{-3}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------------|-------------------------------------|-------------|----------|---------------------------------------|
| 1.23 ±0.07 OUR FIT | Error includes scale factor of 1.9. | | | |
| 1.267±0.028±0.033 | 4247 | 1 ABLIKIM | 23N BES3 | $\psi(2S) \rightarrow \gamma$ hadrons |

¹ Measured using $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ and $B(\phi \rightarrow K^+ K^-) = (49.2 \pm 0.5) \times 10^{-2}$ from PDG 22.

$\Gamma(\phi\phi\eta)/\Gamma_{\text{total}}$ Γ_{21}/Γ

| <i>VALUE</i> (units 10^{-4}) | <i>EVTS</i> | <i>DOCUMENT ID</i> | <i>TECN</i> | <i>COMMENT</i> |
|---------------------------------|-------------|--------------------|-------------|---|
| 5.4±0.7±0.1 | 143.6 | 1 ABLIKIM | 20B BES3 | $\psi(2S) \rightarrow \gamma\phi\phi\eta$ |

¹ ABLIKIM 20B reports $(5.33 \pm 0.52 \pm 0.39) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \phi\phi\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\omega\omega)/\Gamma_{\text{total}}$ Γ_{22}/Γ

| <i>VALUE</i> (units 10^{-3}) | <i>EVTS</i> | <i>DOCUMENT ID</i> | <i>TECN</i> | <i>COMMENT</i> |
|---------------------------------|-------------|--------------------|-------------|----------------|
| 0.86±0.10 OUR AVERAGE | | | | |

| | | | | |
|--------------------------|----------------|-----------|----------|--|
| $0.83 \pm 0.10 \pm 0.02$ | 762 | 1 ABLIKIM | 11K BES3 | $\psi(2S) \rightarrow \gamma$ hadrons |
| $1.76 \pm 0.58 \pm 0.04$ | 27.7 ± 7.4 | 2 ABLIKIM | 05N BES2 | $\psi(2S) \rightarrow \gamma\chi_{c2} \rightarrow \gamma 6\pi$ |

¹ ABLIKIM 11K reports $(8.9 \pm 0.3 \pm 1.1) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \omega\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ABLIKIM 05N reports $[\Gamma(\chi_{c2}(1P) \rightarrow \omega\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] = (0.165 \pm 0.044 \pm 0.032) \times 10^{-3}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\omega K^+ K^-)/\Gamma_{\text{total}}$ Γ_{23}/Γ

| <i>VALUE</i> (units 10^{-3}) | <i>EVTS</i> | <i>DOCUMENT ID</i> | <i>TECN</i> | <i>COMMENT</i> |
|---------------------------------|-------------|--------------------|-------------|--|
| 0.73±0.04±0.08 | 512 | 1 ABLIKIM | 13B BES3 | $e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c2}$ |

¹ Using 1.06×10^8 $\psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c2}\gamma) = (8.72 \pm 0.34)\%$.

 $\Gamma(\omega\phi)/\Gamma_{\text{total}}$ Γ_{24}/Γ

| <i>VALUE</i> (units 10^{-6}) | <i>CL%</i> | <i>EVTS</i> | <i>DOCUMENT ID</i> | <i>TECN</i> | <i>COMMENT</i> |
|---------------------------------|------------|-------------|--------------------|---------------------------------------|----------------|
| 9.7±2.8±0.2 | 33 | 1 ABLIKIM | 19J BES3 | $\psi(2S) \rightarrow \gamma$ hadrons | |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-----|----|-------------------------|----------|---------------------------------------|
| <19 | 90 | 2, ³ ABLIKIM | 11K BES3 | $\psi(2S) \rightarrow \gamma$ hadrons |
|-----|----|-------------------------|----------|---------------------------------------|

¹ ABLIKIM 19J reports $[\Gamma(\chi_{c2}(1P) \rightarrow \omega\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] = (0.91 \pm 0.23 \pm 0.12) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ABLIKIM 11K reports $< 2 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \omega\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.36 \times 10^{-2}$.

³ Superseded by ABLIKIM 19J.

$\Gamma(\pi^+\pi^-\pi^0\text{(non-resonant)})/\Gamma_{\text{total}}$ Γ_{27}/Γ

| <u>VALUE</u> (units 10^{-5}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|----------------------|-------------|--|
| $2.0 \pm 0.4 \pm 0.1$ | 64 | ¹ ABLIKIM | 17AG BES3 | $\psi(2S) \rightarrow \gamma\pi^+\pi^-\pi^0$ |

¹ ABLIKIM 17AG reports $(2.1 \pm 0.4 \pm 0.2) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+\pi^-\pi^0\text{(non-resonant)})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\rho(770)\pm\pi^\mp)/\Gamma_{\text{total}}$ Γ_{28}/Γ

| <u>VALUE</u> (units 10^{-5}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|-------------|----------------------|-------------|--|
| $0.62 \pm 0.39 \pm 0.02$ | 15 | ¹ ABLIKIM | 17AG BES3 | $\psi(2S) \rightarrow \gamma\pi^+\pi^-\pi^0$ |

¹ ABLIKIM 17AG reports $(0.64 \pm 0.39 \pm 0.07) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \rho(770)\pm\pi^\mp)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\pi^+\pi^-\eta)/\Gamma_{\text{total}}$ Γ_{29}/Γ

| <u>VALUE</u> (units 10^{-3}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|------------|--------------------|-------------|--|
| $0.49 \pm 0.13 \pm 0.01$ | | ¹ ATHAR | 07 | CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|------|----|----------------------|----------|--|
| <1.5 | 90 | ² ABLIKIM | 06R BES2 | $\psi(2S) \rightarrow \gamma\chi_{c2}$ |
|------|----|----------------------|----------|--|

¹ ATHAR 07 reports $(0.49 \pm 0.12 \pm 0.06) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+\pi^-\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ABLIKIM 06R reports $< 1.7 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+\pi^-\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.1 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.36 \times 10^{-2}$.

 $\Gamma(\pi^+\pi^-\eta')/\Gamma_{\text{total}}$ Γ_{30}/Γ

| <u>VALUE</u> (units 10^{-3}) | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|--------------------|-------------|--|
| $0.51 \pm 0.19 \pm 0.01$ | ¹ ATHAR | 07 | CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$ |

¹ ATHAR 07 reports $(0.51 \pm 0.18 \pm 0.06) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+\pi^-\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(K^*(892)\pm K^\mp)/\Gamma_{\text{total}}$ Γ_{34}/Γ

| <u>VALUE</u> (units 10^{-4}) | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|----------------------|-------------|---|
| $1.46 \pm 0.21 \pm 0.04$ | ¹ ABLIKIM | 17AG BES3 | $\psi(2S) \rightarrow \gamma K\bar{K}\pi$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|--------------------------|----------------------|-----------|---|
| $1.75 \pm 0.27 \pm 0.04$ | ² ABLIKIM | 17AG BES3 | $\psi(2S) \rightarrow \gamma K^+ K^- \pi^0$ |
| $1.36 \pm 0.27 \pm 0.03$ | ³ ABLIKIM | 17AG BES3 | $\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp$ |

¹ ABLIKIM 17AG reports $(1.5 \pm 0.1 \pm 0.2) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^{\pm} K^{\mp})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ABLIKIM 17AG reports $(1.8 \pm 0.2 \pm 0.2) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^{\pm} K^{\mp})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ ABLIKIM 17AG reports $(1.4 \pm 0.2 \pm 0.2) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^{\pm} K^{\mp})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | Γ_{35}/Γ |
|--|----------------------|-----------|---|
| $1.27 \pm 0.27 \pm 0.03$ | ¹ ABLIKIM | 17AG BES3 | $\psi(2S) \rightarrow \gamma K_S^0 K^{\pm} \pi^{\mp}$ |

¹ ABLIKIM 17AG reports $(1.3 \pm 0.2 \pm 0.2) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K_2^*(1430)^{\pm} K^{\mp})/\Gamma_{\text{total}}$

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | Γ_{36}/Γ |
|--|----------------------|-----------|---|
| $15.1 \pm 1.2 \pm 0.4$ | ¹ ABLIKIM | 17AG BES3 | $\psi(2S) \rightarrow \gamma K \bar{K} \pi$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|------------------------|----------------------|-----------|---|
| $17.7 \pm 1.6 \pm 0.4$ | ² ABLIKIM | 17AG BES3 | $\psi(2S) \rightarrow \gamma K^+ K^- \pi^0$ |
| $13.2 \pm 1.5 \pm 0.3$ | ³ ABLIKIM | 17AG BES3 | $\psi(2S) \rightarrow \gamma K_S^0 K^{\pm} \pi^{\mp}$ |

¹ ABLIKIM 17AG reports $(15.5 \pm 0.6 \pm 1.2) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K_2^*(1430)^{\pm} K^{\mp})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ABLIKIM 17AG reports $(18.2 \pm 0.8 \pm 1.6) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K_2^*(1430)^{\pm} K^{\mp})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ ABLIKIM 17AG reports $(13.6 \pm 0.8 \pm 1.4) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K_2^*(1430)^{\pm} K^{\mp})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K_2^*(1430)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{37}/Γ

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|----------------------|-----------|---|
| 12.7±1.7±0.3 | ¹ ABLIKIM | 17AG BES3 | $\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp$ |

¹ ABLIKIM 17AG reports $(13.0 \pm 1.0 \pm 1.5) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K_2^*(1430)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))] \text{ assuming } \mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(K_3^*(1780)^\pm K^\mp)/\Gamma_{\text{total}}$ Γ_{38}/Γ

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|---|----------------------|-----------|---|
| 5.3±0.8±0.1 | ¹ ABLIKIM | 17AG BES3 | $\psi(2S) \rightarrow \gamma K \bar{K} \pi$ |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | |
| 5.2±1.0±0.1 | ² ABLIKIM | 17AG BES3 | $\psi(2S) \rightarrow \gamma K^+ K^- \pi^0$ |
| 5.7±1.8±0.1 | ³ ABLIKIM | 17AG BES3 | $\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp$ |

¹ ABLIKIM 17AG reports $(5.4 \pm 0.5 \pm 0.7) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K_3^*(1780)^\pm K^\mp)/\Gamma_{\text{total}}] \times [\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))] \text{ assuming } \mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ABLIKIM 17AG reports $(5.3 \pm 0.5 \pm 0.9) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K_3^*(1780)^\pm K^\mp)/\Gamma_{\text{total}}] \times [\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))] \text{ assuming } \mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ ABLIKIM 17AG reports $(5.9 \pm 1.1 \pm 1.5) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K_3^*(1780)^\pm K^\mp)/\Gamma_{\text{total}}] \times [\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))] \text{ assuming } \mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(K_3^*(1780)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{39}/Γ

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|----------------------|-----------|---|
| 5.7±2.1±0.1 | ¹ ABLIKIM | 17AG BES3 | $\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp$ |

¹ ABLIKIM 17AG reports $(5.9 \pm 1.6 \pm 1.5) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K_3^*(1780)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))] \text{ assuming } \mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(a_2(1320)^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{40}/Γ

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|----------------------|-----------|---|
| 13.1±3.5±0.3 | ¹ ABLIKIM | 17AG BES3 | $\psi(2S) \rightarrow \gamma K^+ K^- \pi^0$ |

¹ ABLIKIM 17AG reports $(13.5 \pm 1.6 \pm 3.2) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow a_2(1320)^0 \pi^0) / \Gamma_{\text{total}}] \times [\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(a_2(1320)^{\pm} \pi^{\mp})/\Gamma_{\text{total}}$ Γ_{41}/Γ

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|--|----------------------|-----------|---|
| $17.9 \pm 6.2 \pm 0.4$ | ¹ ABLIKIM | 17AG BES3 | $\psi(2S) \rightarrow \gamma K_S^0 K^{\pm} \pi^{\mp}$ |

¹ ABLIKIM 17AG reports $(18.4 \pm 3.3 \pm 5.5) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow a_2(1320)^{\pm} \pi^{\mp}) / \Gamma_{\text{total}}] \times [\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{43}/Γ

| VALUE (units 10^{-3}) | DOCUMENT ID | TECN | COMMENT |
|--|--------------------|------|---|
| $0.31 \pm 0.08 \pm 0.01$ | ¹ ATHAR | 07 | $\psi(2S) \rightarrow \gamma h^+ h^- h^0$ |

¹ ATHAR 07 reports $(0.31 \pm 0.07 \pm 0.04) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- \pi^0) / \Gamma_{\text{total}}] \times [\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+ K^- \eta)/\Gamma_{\text{total}}$ Γ_{44}/Γ

| VALUE (units 10^{-3}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|--------------------|------|---|
| <0.33 | 90 | ¹ ATHAR | 07 | $\psi(2S) \rightarrow \gamma h^+ h^- h^0$ |

¹ ATHAR 07 reports $< 0.33 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- \eta) / \Gamma_{\text{total}}] \times [\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.36 \times 10^{-2}$.

$\Gamma(K^+ K^- \eta'(958))/\Gamma_{\text{total}}$ Γ_{45}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------------------|------|----------------------|----------|--|
| 1.94 ± 0.34 | 107 | ¹ ABLIKIM | 14J BES3 | $\psi(2S) \rightarrow \gamma K^+ K^- \eta'(958)$ |

¹ Derived using $\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c2}) = (8.72 \pm 0.34)\%$. Uncertainty includes both statistical and systematic contributions combined in quadrature.

$\Gamma(\eta \eta')/\Gamma_{\text{total}}$ Γ_{46}/Γ

| VALUE (units 10^{-5}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-----|----------------------|-------------|--|---------|
| $2.2 \pm 0.5 \pm 0.1$ | 20 | ¹ ABLIKIM | 17AI BES3 | $\psi(2S) \rightarrow \gamma \eta' \eta$ | |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | | |
|------|----|---------------|--------------------|----|------|--|
| < 6 | 90 | 3.3 ± 8.0 | ² ASNER | 09 | CLEO | $\psi(2S) \rightarrow \gamma \eta \eta'$ |
| < 23 | 90 | | ³ ADAMS | 07 | CLEO | $\psi(2S) \rightarrow \gamma \chi_{c2}$ |

¹ ABLIKIM 17AI reports $(2.27 \pm 0.43 \pm 0.25) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \eta\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ASNER 09 reports $< 0.6 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \eta\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.36 \times 10^{-2}$.

³ Superseded by ASNER 09. ADAMS 07 reports $< 2.3 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \eta\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 0.0933 \pm 0.0014 \pm 0.0061$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.36 \times 10^{-2}$.

$\Gamma(\eta'\eta')/\Gamma_{\text{total}}$

Γ_{47}/Γ

| VALUE (units 10^{-5}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|------|----------------------|------|---|
| 4.6±0.6±0.1 | | 60 | ¹ ABLIKIM | 17AI | $\psi(2S) \rightarrow \gamma\eta'\eta'$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | | |
|-----|----|------------|--------------------|----|------|---|
| <10 | 90 | 12 ± 7 | ² ASNER | 09 | CLEO | $\psi(2S) \rightarrow \gamma\eta'\eta'$ |
| <31 | 90 | | ³ ADAMS | 07 | CLEO | $\psi(2S) \rightarrow \gamma\chi_{c2}$ |

¹ ABLIKIM 17AI reports $(4.76 \pm 0.56 \pm 0.38) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \eta'\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ASNER 09 reports $< 1.0 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \eta'\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.36 \times 10^{-2}$.

³ Superseded by ASNER 09. ADAMS 07 reports $< 3.1 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \eta'\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 0.0933 \pm 0.0014 \pm 0.0061$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.36 \times 10^{-2}$.

$\Gamma(\pi^+\pi^-K_S^0K_S^0)/\Gamma_{\text{total}}$

Γ_{48}/Γ

| VALUE (units 10^{-3}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|----------------------|----------|--|
| 2.2±0.5±0.1 | 57 ± 11 | ¹ ABLIKIM | 050 BES2 | $\psi(2S) \rightarrow \gamma\chi_{c2}$ |

¹ ABLIKIM 050 reports $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+\pi^-K_S^0K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] = (0.207 \pm 0.039 \pm 0.033) \times 10^{-3}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+K^-K_S^0K_S^0)/\Gamma_{\text{total}}$

Γ_{49}/Γ

| VALUE (units 10^{-4}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|---------------|----------------------|----------|--------------------------------------|
| <4 | 90 | 2.3 ± 2.2 | ¹ ABLIKIM | 050 BES2 | $e^+e^- \rightarrow \chi_{c2}\gamma$ |

¹ ABLIKIM 050 reports $[\Gamma(\chi_{c2}(1P) \rightarrow K^+K^-K_S^0K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] < 3.5 \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.36 \times 10^{-2}$.

| $\Gamma(K_S^0 K_S^0 K_S^0 K_S^0)/\Gamma_{\text{total}}$ | Γ_{50}/Γ | | | |
|---|----------------------|----------------------|-------------|--------------------------------------|
| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
| 1.15±0.18±0.03 | 68 | ¹ ABLIKIM | 19AA BES3 | $\psi(2S) \rightarrow \gamma 4K_S^0$ |

¹ Using $B(K_S^0 \rightarrow \pi^+ \pi^-) = (69.20 \pm 0.05)\%$. ABLIKIM 19AA reports $[\Gamma(\chi_{c2}(1P) \rightarrow K_S^0 K_S^0 K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))] = (10.8 \pm 1.5 \pm 0.8) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value..

| $\Gamma(K^+ K^- \phi)/\Gamma_{\text{total}}$ | Γ_{52}/Γ | | | |
|--|----------------------|----------------------|-------------|---|
| <u>VALUE (units 10^{-3})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
| 1.45±0.30±0.04 | 52 | ¹ ABLIKIM | 06T BES2 | $\psi(2S) \rightarrow \gamma 2K^+ 2K^-$ |

¹ ABLIKIM 06T reports $(1.67 \pm 0.26 \pm 0.24) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- \phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.1 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

| $\Gamma(\bar{K}^0 K^+ \pi^- \phi + \text{c.c.})/\Gamma_{\text{total}}$ | Γ_{53}/Γ | | |
|--|----------------------|-------------|---|
| <u>VALUE (units 10^{-3})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
| 4.83±0.32±0.66 | ABLIKIM | 15M BES3 | $\psi(2S) \rightarrow \gamma \chi_{c2}$ |

| $\Gamma(K^+ K^- \pi^0 \phi)/\Gamma_{\text{total}}$ | Γ_{54}/Γ | | |
|--|----------------------|-------------|---|
| <u>VALUE (units 10^{-3})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
| 2.74±0.16±0.44 | ABLIKIM | 15M BES3 | $\psi(2S) \rightarrow \gamma \chi_{c2}$ |

| $\Gamma(\phi \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$ | Γ_{55}/Γ | | | |
|--|----------------------|----------------------|-------------|---|
| <u>VALUE (units 10^{-3})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
| 0.93±0.06±0.10 | 408 | ¹ ABLIKIM | 13B BES3 | $e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c2}$ |

¹ Using $1.06 \times 10^8 \psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c2}\gamma) = (8.72 \pm 0.34)\%$.

| $\Gamma(p\bar{p}\pi^0)/\Gamma_{\text{total}}$ | Γ_{57}/Γ | | |
|---|----------------------|-------------|----------------|
| <u>VALUE (units 10^{-3})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
| 0.47±0.04 OUR AVERAGE | | | |

| | | | | |
|--|---------------------|----|------|---|
| 0.48±0.04±0.01 | ¹ ONYISI | 10 | CLE3 | $\psi(2S) \rightarrow \gamma p\bar{p}X$ |
| 0.44±0.09±0.01 | ² ATHAR | 07 | CLEO | $\psi(2S) \rightarrow \gamma h^+ h^- h^0$ |
| ¹ ONYISI 10 reports $(4.83 \pm 0.25 \pm 0.35 \pm 0.31) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. | | | | |
| ² ATHAR 07 reports $(0.44 \pm 0.08 \pm 0.05) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. | | | | |

$\Gamma(p\bar{p}\eta)/\Gamma_{\text{total}}$ Γ_{58}/Γ

| VALUE (units 10^{-3}) | DOCUMENT ID | TECN | COMMENT |
|--|------------------------|------|---|
| 0.177±0.025 OUR AVERAGE | | | |
| 0.175±0.027±0.004 | ¹ ONYISI 10 | CLE3 | $\psi(2S) \rightarrow \gamma p\bar{p}X$ |
| 0.189±0.071±0.005 | ² ATHAR 07 | CLEO | $\psi(2S) \rightarrow \gamma h^+ h^- h^0$ |
| ¹ ONYISI 10 reports $(1.76 \pm 0.23 \pm 0.14 \pm 0.11) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. ² ATHAR 07 reports $(0.19 \pm 0.07 \pm 0.02) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. | | | |

 $\Gamma(p\bar{p}\omega)/\Gamma_{\text{total}}$ Γ_{59}/Γ

| VALUE (units 10^{-3}) | DOCUMENT ID | TECN | COMMENT |
|--|-------------|------|---|
| 0.37±0.04±0.01 | | | |
| ¹ ONYISI 10 | 10 | CLE3 | $\psi(2S) \rightarrow \gamma p\bar{p}X$ |
| ¹ ONYISI 10 reports $(3.68 \pm 0.35 \pm 0.26 \pm 0.24) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. | | | |

 $\Gamma(p\bar{p}\phi)/\Gamma_{\text{total}}$ Γ_{60}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------------|--------------------------|------|---|
| 2.8±0.9±0.1 | 24 ± 7 | ¹ ABLIKIM 11F | BES3 | $\psi(2S) \rightarrow \gamma p\bar{p}K^+ K^-$ |
| ¹ ABLIKIM 11F reports $(3.04 \pm 0.85 \pm 0.43) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. | | | | |

 $\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{61}/Γ

| VALUE (units 10^{-3}) | DOCUMENT ID | TECN | COMMENT |
|---|---------------------------|------|--|
| 1.32±0.34 OUR EVALUATION Treating systematic error as correlated. | | | |
| 1.3 ± 0.4 OUR AVERAGE Error includes scale factor of 1.3. | | | |
| 1.17±0.19±0.30 | ¹ BAI 99B | BES | $\psi(2S) \rightarrow \gamma\chi_{c2}$ |
| 2.64±1.03±0.14 | ¹ TANENBAUM 78 | MRK1 | $\psi(2S) \rightarrow \gamma\chi_{c2}$ |
| ¹ Rescaled by us using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (8.3 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$. Multiplied by a factor of 2 to convert from $K_S^0 K^+ \pi^-$ to $K^0 K^+ \pi^-$ decay. | | | |

$\Gamma(p\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_{62}/Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|----------|--|
| 0.080±0.024±0.002 | 29.2 | 1 HE | 08B CLEO | $e^+e^- \rightarrow \gamma h^+h^-h^0h^0$ |

¹ HE 08B reports $0.08 \pm 0.02 \pm 0.01 \pm 0.01$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(p\bar{p}K^+K^-(\text{non-resonant}))/\Gamma_{\text{total}}$ Γ_{63}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|--------------|-------------|----------|--|
| 1.94±0.32±0.05 | 131 ± 12 | 1 ABLIKIM | 11F BES3 | $\psi(2S) \rightarrow \gamma p\bar{p}K^+K^-$ |

¹ ABLIKIM 11F reports $(2.08 \pm 0.19 \pm 0.30) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}K^+K^-(\text{non-resonant}))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(p\bar{p}K_S^0K_S^0)/\Gamma_{\text{total}}$ Γ_{64}/Γ

| VALUE (units 10^{-4}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|----------|--|
| <7.9 | 90 | 1 ABLIKIM | 06D BES2 | $\psi(2S) \rightarrow \chi_{c2}\gamma$ |

¹ Using $B(\psi(2S) \rightarrow \chi_{c2}\gamma) = (9.3 \pm 0.6)\%$.

 $\Gamma(p\bar{n}\pi^-)/\Gamma_{\text{total}}$ Γ_{65}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------|------|-------------|------|---------|
| 8.7±1.0 OUR AVERAGE | | | | |

| | | | | |
|--------------|------|-----------|----------|---|
| 8.5±1.0±0.2 | 3309 | 1 ABLIKIM | 12J BES3 | $\psi(2S) \rightarrow \gamma p\bar{n}\pi^-$ |
| 10.4±3.5±0.3 | | 2 ABLIKIM | 06I BES2 | $\psi(2S) \rightarrow \gamma p\pi^-X$ |

¹ ABLIKIM 12J reports $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{n}\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] = (0.80 \pm 0.02 \pm 0.09) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ABLIKIM 06I reports $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{n}\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] = (0.97 \pm 0.20 \pm 0.26) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\bar{p}n\pi^+)/\Gamma_{\text{total}}$ Γ_{66}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|----------|--|
| 9.1±0.8±0.2 | 3732 | 1 ABLIKIM | 12J BES3 | $\psi(2S) \rightarrow \gamma\bar{p}n\pi^+$ |

¹ ABLIKIM 12J reports $[\Gamma(\chi_{c2}(1P) \rightarrow \bar{p}n\pi^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] = (0.85 \pm 0.02 \pm 0.07) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(p\bar{n}\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{67}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|----------|--|
| 22.1±1.7±0.5 | 2128 | 1 ABLIKIM | 12J BES3 | $\psi(2S) \rightarrow \gamma p\bar{n}\pi^-\pi^0$ |

¹ ABLIKIM 12J reports $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{n}\pi^-\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] = (2.07 \pm 0.06 \pm 0.15) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\bar{p}n\pi^+\pi^0)/\Gamma_{\text{total}}$ Γ_{68}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|----------|---|
| 21.5±1.8±0.5 | 2352 | 1 ABLIKIM | 12J BES3 | $\psi(2S) \rightarrow \gamma\bar{p}n\pi^+\pi^0$ |

¹ ABLIKIM 12J reports $[\Gamma(\chi_{c2}(1P) \rightarrow \bar{p}n\pi^+\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] = (2.01 \pm 0.06 \pm 0.16) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{70}/Γ

| VALUE (units 10^{-5}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-----------|-------------|---|---------|
| 128±15±3 | 371 | 1 ABLIKIM | 12I BES3 | $\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}\pi^+\pi^-$ | |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|------|----|-----------|----------|--|
| <350 | 90 | 2 ABLIKIM | 06D BES2 | $\psi(2S) \rightarrow \chi_{c2}\gamma$ |
|------|----|-----------|----------|--|

¹ ABLIKIM 12I reports $(137.0 \pm 7.6 \pm 15.7) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using $B(\psi(2S) \rightarrow \chi_{c2}\gamma) = (9.3 \pm 0.6)\%$.

 $\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^- \text{ (non-resonant)})/\Gamma_{\text{total}}$ Γ_{71}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|----------|---|
| 67±15±2 | 36 | 1 ABLIKIM | 12I BES3 | $\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}\pi^+\pi^-$ |

¹ ABLIKIM 12I reports $(71.8 \pm 14.5 \pm 8.2) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Lambda\bar{\Lambda}\pi^+\pi^- \text{ (non-resonant)})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\Sigma(1385)^+\bar{\Lambda}\pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{72}/Γ

| VALUE (units 10^{-5}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|----------|---|
| <40 | 90 | 1 ABLIKIM | 12I BES3 | $\psi(2S) \rightarrow \gamma\Sigma(1385)^+\bar{\Lambda}\pi^-$ |

¹ ABLIKIM 12I reports $< 42 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma(1385)^+\bar{\Lambda}\pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.36 \times 10^{-2}$.

$\Gamma(\Sigma(1385)^-\bar{\Lambda}\pi^++\text{c.c.})/\Gamma_{\text{total}}$ Γ_{73}/Γ

| <u>VALUE</u> (units 10^{-5}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|------------|----------------------|-------------|--|
| <60 | 90 | ¹ ABLIKIM | 12I | $BES3 \quad \psi(2S) \rightarrow \gamma\Sigma(1385)^-\bar{\Lambda}\pi^+$ |
| ¹ ABLIKIM 12I reports $< 61 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma(1385)^-\bar{\Lambda}\pi^+ + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.36 \times 10^{-2}$. | | | | |

 $\Gamma(K^+\bar{p}\Lambda+\text{c.c.})/\Gamma_{\text{total}}$ Γ_{75}/Γ

| <u>VALUE</u> (units 10^{-4}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|------------------------|-------------|---|
| 7.9 ± 0.6 OUR AVERAGE | | | | |
| 7.8 \pm 0.5 \pm 0.2 | 5k | ^{1,2} ABLIKIM | 13D | $BES3 \quad \psi(2S) \rightarrow \gamma\Lambda\bar{p}K^+$ |
| 8.5 \pm 1.6 \pm 0.2 | | ³ ATHAR | 07 | $CLEO \quad \psi(2S) \rightarrow \gamma h^+h^-h^0$ |
| ¹ ABLIKIM 13D reports $(8.4 \pm 0.3 \pm 0.6) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. | | | | |
| ² Using $B(\Lambda \rightarrow p\pi^-) = 63.9\%$. | | | | |
| ³ ATHAR 07 reports $(8.5 \pm 1.4 \pm 1.0) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. | | | | |

 $\Gamma(nK_S^0\bar{\Lambda}+\text{c.c.})/\Gamma_{\text{total}}$ Γ_{76}/Γ

| <u>VALUE</u> (units 10^{-4}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|-------------|----------------------|-------------|--|
| $3.64 \pm 0.27 \pm 0.09$ | 879 | ¹ ABLIKIM | 21AV | $BES3 \quad \psi(2S) \rightarrow \gamma nK_S^0\bar{\Lambda} + \text{c.c.}$ |
| ¹ ABLIKIM 21AV reports $(3.58 \pm 0.16 \pm 0.23) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow nK_S^0\bar{\Lambda} + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 0.0952 \pm 0.0020$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Also uses $B(\bar{\Lambda} \rightarrow \bar{p}\pi^+) = (63.9 \pm 0.5)\%$ and $B(K_S^0 \rightarrow \pi^+\pi^-) = (69.20 \pm 0.05)\%$. | | | | |

 $\Gamma(K^*(892)^+\bar{p}\Lambda+\text{c.c.})/\Gamma_{\text{total}}$ Γ_{77}/Γ

| <u>VALUE</u> (units 10^{-4}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|----------------------|-------------|---|
| $8.3 \pm 1.2 \pm 0.2$ | 476 | ¹ ABLIKIM | 19AU | $BES3 \quad \psi(2S) \rightarrow \gamma K^{*+}\bar{p}\Lambda$ |
| ¹ ABLIKIM 19AU reports $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] = (7.8 \pm 0.9 \pm 0.6) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. | | | | |

$\Gamma(K^+\bar{p}\Lambda(1520)+\text{c.c.})/\Gamma_{\text{total}}$ Γ_{78}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|----------------------|----------|---|
| 2.9±0.7±0.1 | 79 ± 13 | ¹ ABLIKIM | 11F BES3 | $\psi(2S) \rightarrow \gamma p \bar{p} K^+ K^-$ |

¹ ABLIKIM 11F reports $(3.06 \pm 0.50 \pm 0.54) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ \bar{p}\Lambda(1520) + \text{c.c.})/\Gamma_{\text{total}}] \times [\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))] \approx (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\Lambda(1520)\bar{\Lambda}(1520))/\Gamma_{\text{total}}$ Γ_{79}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------------|----------------------|----------|---|
| 4.7±1.5±0.1 | 29 ± 7 | ¹ ABLIKIM | 11F BES3 | $\psi(2S) \rightarrow \gamma p \bar{p} K^+ K^-$ |

¹ ABLIKIM 11F reports $(5.05 \pm 1.29 \pm 0.93) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Lambda(1520)\bar{\Lambda}(1520))/\Gamma_{\text{total}}] \times [\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))] \approx (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}$ Γ_{80}/Γ

| VALUE (units 10^{-5}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|------|----------------------|----------|---|
| 3.7±0.6±0.1 | | 91 | ¹ ABLIKIM | 18V BES3 | $\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|----|----|----------------------|-------------------|---|
| <6 | 90 | ² ABLIKIM | 13H BES3 | $\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$ |
| <7 | 90 | 7.5 ± 3.4 | ³ NAIK | 08 CLEO $\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$ |

¹ ABLIKIM 18V reports $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))] = (0.35 \pm 0.05 \pm 0.02) \times 10^{-5}$ which we divide by our best value $\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ABLIKIM 13H reports $< 0.65 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))] \approx (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.36 \times 10^{-2}$.

³ NAIK 08 reports $< 0.75 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))] \approx (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.36 \times 10^{-2}$.

 $\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$ Γ_{83}/Γ

| VALUE (units 10^{-5}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|------|----------------------|----------|---|
| 3.4±0.7±0.1 | | 55 | ¹ ABLIKIM | 18V BES3 | $\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|----|----|----------------------|-------------------|---|
| <8 | 90 | ² ABLIKIM | 13H BES3 | $\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$ |
| <7 | 90 | 4.0 ± 3.5 | ³ NAIK | 08 CLEO $\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$ |

¹ ABLIKIM 18V reports $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))] = (0.32 \pm 0.06 \pm 0.03) \times 10^{-5}$ which we divide by our best value $\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ABLIKIM 13H reports $< 0.88 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.36 \times 10^{-2}$.

³ NAIK 08 reports $< 0.67 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.36 \times 10^{-2}$.

$\Gamma(\Sigma^- \bar{\Sigma}^+)/\Gamma_{\text{total}}$

Γ_{84}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|----------------------|------|--|
| 4.5±1.8±0.1 | 131 | ¹ ABLIKIM | 20I | BES3 $\psi(2S) \rightarrow \gamma \Sigma^- \bar{\Sigma}^+$ |

¹ ABLIKIM 20I reports $(4.4 \pm 1.7 \pm 0.5) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^- \bar{\Sigma}^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\Sigma(1385)^+ \bar{\Sigma}(1385)^-)/\Gamma_{\text{total}}$

Γ_{85}/Γ

| VALUE (units 10^{-5}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|----------------------|------|--|
| <16 | 90 | ¹ ABLIKIM | 12I | BES3 $\psi(2S) \rightarrow \gamma \Sigma(1385)^+ \bar{\Sigma}(1385)^-$ |

¹ ABLIKIM 12I reports $< 17 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma(1385)^+ \bar{\Sigma}(1385)^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.36 \times 10^{-2}$.

$\Gamma(\Sigma(1385)^- \bar{\Sigma}(1385)^+)/\Gamma_{\text{total}}$

Γ_{86}/Γ

| VALUE (units 10^{-5}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|----------------------|------|--|
| <8 | 90 | ¹ ABLIKIM | 12I | BES3 $\psi(2S) \rightarrow \gamma \Sigma(1385)^- \bar{\Sigma}(1385)^+$ |

¹ ABLIKIM 12I reports $< 8.5 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma(1385)^- \bar{\Sigma}(1385)^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.36 \times 10^{-2}$.

$\Gamma(K^- \Lambda \bar{\Xi}^+ + \text{c.c.})/\Gamma_{\text{total}}$

Γ_{87}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|----------------------|------|--|
| 1.80±0.32±0.04 | 51 | ¹ ABLIKIM | 15I | BES3 $\psi(2S) \rightarrow \gamma K^- \Lambda \bar{\Xi}^+ + \text{c.c.}$ |

¹ ABLIKIM 15I reports $[\Gamma(\chi_{c2}(1P) \rightarrow K^- \Lambda \bar{\Xi}^+ + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))] = (1.68 \pm 0.26 \pm 0.15) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\Xi^0 \bar{\Xi}^0)/\Gamma_{\text{total}}$

Γ_{88}/Γ

| VALUE (units 10^{-4}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|----------------------|-------------|--|---------|
| 1.86±0.22±0.05 | 804 | ¹ ABLIKIM | 220 | BES3 $\psi(2S) \rightarrow \gamma \Xi^0 \bar{\Xi}^0$ | |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|------|----|---|-------------------|----|--|
| <1.1 | 90 | 3 | ² NAIK | 08 | CLEO $\psi(2S) \rightarrow \gamma \Xi^0 \bar{\Xi}^0$ |
|------|----|---|-------------------|----|--|

¹ ABLIKIM 220 reports $(1.83 \pm 0.15 \pm 0.16) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Xi^0 \bar{\Xi}^0) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² NAIK 08 reports $< 1.06 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Xi^0 \bar{\Xi}^0) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.36 \times 10^{-2}$.

$\Gamma(\Xi^- \bar{\Xi}^+)/\Gamma_{\text{total}}$

Γ_{89}/Γ

| VALUE (units 10^{-4}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|-----|------|-------------|------|---------|
| $1.46 \pm 0.12 \pm 0.12$ OUR AVERAGE | | | | | |

| | | | | | |
|--------------------------|------------|----------------------|-----|------|---|
| $1.47 \pm 0.12 \pm 0.04$ | 1691 | ¹ ABLIKIM | 220 | BES3 | $\psi(2S) \rightarrow \gamma \Xi^- \bar{\Xi}^+$ |
| $1.45 \pm 0.32 \pm 0.04$ | 29 ± 5 | ² NAIK | 08 | CLEO | $\psi(2S) \rightarrow \gamma \Xi^+ \bar{\Xi}^-$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|---------|----|----------------------|-----|------|---|
| < 3.7 | 90 | ³ ABLIKIM | 06D | BES2 | $\psi(2S) \rightarrow \chi_{c2} \gamma$ |
|---------|----|----------------------|-----|------|---|

¹ ABLIKIM 220 reports $(1.44 \pm 0.06 \pm 0.11) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Xi^- \bar{\Xi}^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² NAIK 08 reports $(1.45 \pm 0.30 \pm 0.15) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Xi^- \bar{\Xi}^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Using $B(\psi(2S) \rightarrow \chi_{c2} \gamma) = (9.3 \pm 0.6)\%$.

$\Gamma(\Omega^- \bar{\Omega}^+)/\Gamma_{\text{total}}$

Γ_{90}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|-------------|------|---------|
| $4.52 \pm 0.24 \pm 0.18$ | 1038 | ABLIKIM | 23T | BES3 |

$\Gamma(J/\psi(1S)\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$

Γ_{91}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------|-----|-------------|------|---------|
| < 0.015 | 90 | BARATE | 81 | SPEC |

$\Gamma(\pi^0 \eta_c)/\Gamma_{\text{total}}$

Γ_{92}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|------------------------|-----|----------------------|------|---------|
| $< 3.2 \times 10^{-3}$ | 90 | ¹ ABLIKIM | 15N | BES3 |

¹ Using $B(\eta_c \rightarrow K_S^0 K^\pm \pi^\mp) \times B(K_S^0 \rightarrow \pi^+ \pi^-) \times B(\pi^0 \rightarrow \gamma \gamma) = (1.66 \pm 0.11) \times 10^{-2}$.

$\Gamma(\eta_c(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$

Γ_{93}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|--|-----|-------------|------|---------|
| $< 0.54 \times 10^{-2}$ | 90 | 1,2 ABLIKIM | 13B | BES3 |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|------------------------|----|-------------|-----|------|
| $< 1.2 \times 10^{-2}$ | 90 | 1,3 ABLIKIM | 13B | BES3 |
|------------------------|----|-------------|-----|------|

¹ Using $1.06 \times 10^8 \psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c2} \gamma) = (8.72 \pm 0.34)\%$.

² From the $\eta_c \rightarrow K_S^0 K^\pm \pi^\mp$ decays.

³ From the $\eta_c \rightarrow K^+ K^- \pi^0$ decays.

$\Gamma(\eta_c(1S)\pi^+\pi^-)/\Gamma(K^0K^+\pi^- + \text{c.c.})$ Γ_{93}/Γ_{42}

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|------------|--------------------|-------------|---|
| <16.4 | 90 | 1 LEES | 12AE BABR | $e^+e^- \rightarrow e^+e^-\pi^+\pi^-\eta_c$ |

¹ We divided the reported limit by 2 to take into account the $K_L^0 K^+ \pi^-$ mode.

RADIATIVE DECAYS

 $\Gamma(\gamma\rho^0)/\Gamma_{\text{total}}$ Γ_{95}/Γ

| <u>VALUE (units 10^{-6})</u> | <u>CL%</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|-------------|--------------------|-------------|---|
| <19 | 90 | 13 ± 11 | 1 ABLIKIM | 11E BES3 | $\psi(2S) \rightarrow \gamma\gamma\rho^0$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-----|----|----------------|-----------|----------|---|
| <40 | 90 | 17.2 ± 6.8 | 2 BENNETT | 08A CLEO | $\psi(2S) \rightarrow \gamma\gamma\rho^0$ |
|-----|----|----------------|-----------|----------|---|

¹ ABLIKIM 11E reports $< 20.8 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \gamma\rho^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.36 \times 10^{-2}$.

² BENNETT 08A reports $< 50 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \gamma\rho^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.1 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.36 \times 10^{-2}$.

 $\Gamma(\gamma\omega)/\Gamma_{\text{total}}$ Γ_{96}/Γ

| <u>VALUE (units 10^{-6})</u> | <u>CL%</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|-------------|--------------------|-------------|---|
| <6 | 90 | 1 ± 6 | 1 ABLIKIM | 11E BES3 | $\psi(2S) \rightarrow \gamma\gamma\omega$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|----|----|---------------|-----------|----------|---|
| <6 | 90 | 0.0 ± 1.8 | 2 BENNETT | 08A CLEO | $\psi(2S) \rightarrow \gamma\gamma\omega$ |
|----|----|---------------|-----------|----------|---|

¹ ABLIKIM 11E reports $< 6.1 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \gamma\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.36 \times 10^{-2}$.

² BENNETT 08A reports $< 7.0 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \gamma\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.1 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.36 \times 10^{-2}$.

 $\Gamma(\gamma\phi)/\Gamma_{\text{total}}$ Γ_{97}/Γ

| <u>VALUE (units 10^{-6})</u> | <u>CL%</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|-------------|--------------------|-------------|---|
| < 8 | 90 | 5 ± 5 | 1 ABLIKIM | 11E BES3 | $\psi(2S) \rightarrow \gamma\gamma\phi$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-----|----|---------------|-----------|----------|---|
| <11 | 90 | 1.3 ± 2.5 | 2 BENNETT | 08A CLEO | $\psi(2S) \rightarrow \gamma\gamma\phi$ |
|-----|----|---------------|-----------|----------|---|

¹ ABLIKIM 11E reports $< 8.1 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \gamma\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.36 \times 10^{-2}$.

² BENNETT 08A reports $< 13 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \gamma\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.1 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.36 \times 10^{-2}$.

$\Gamma(e^+ e^- J/\psi(1S))/\Gamma_{\text{total}}$ Γ_{99}/Γ

| VALUE (units 10^{-3}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|------------------------|------|---|
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | |
| $2.41 \pm 0.15 \pm 0.06$ | 1.3k | ^{1,2} ABLIKIM | 17I | BES3 $\psi(2S) \rightarrow \gamma e^+ e^- J/\psi$ |
| ¹ ABLIKIM 17I reports $(2.48 \pm 0.08 \pm 0.16) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow e^+ e^- J/\psi(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. | | | | |

² Not independent from other measurements reported by ABLIKIM 17I

 $\Gamma(e^+ e^- J/\psi(1S))/\Gamma(\gamma J/\psi(1S))$ Γ_{99}/Γ_{94}

| VALUE (units 10^{-3}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|----------------------|------|---|
| $11.3 \pm 0.4 \pm 0.5$ | 1.3k | ¹ ABLIKIM | 17I | BES3 $\psi(2S) \rightarrow e^+ e^- \gamma J/\psi$ |
| ¹ Uses $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) \times B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) = (199.6 \pm 0.8 \pm 7.0) \times 10^{-4}$ from ABLIKIM 17N and accounts for common systematic errors. | | | | |

 $\Gamma(\mu^+ \mu^- J/\psi(1S))/\Gamma(e^+ e^- J/\psi(1S))$ Γ_{100}/Γ_{99}

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|-------------|------|---|
| $9.40 \pm 0.79 \pm 1.15$ | 219 | ABLIKIM | 19Z | BES3 $\psi(2S) \rightarrow \gamma \chi_c \rightarrow \gamma (\mu^+ \mu^- J/\psi)$ |

 $\Gamma(\gamma\gamma)/\Gamma(\gamma J/\psi(1S))$ Γ_{98}/Γ_{94}

| VALUE (units 10^{-3}) | DOCUMENT ID | TECN | COMMENT |
|---|-------------------------------------|------|--|
| 1.50 ± 0.08 OUR FIT | Error includes scale factor of 1.5. | | |
| 0.99 ± 0.18 | ¹ AMBROGIANI 00B | E835 | $\bar{p}p \rightarrow \chi_{c2} \rightarrow \gamma\gamma, \gamma J/\psi$ |

¹ Calculated by us using $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.0593 \pm 0.0010$.

 $\Gamma(\gamma\gamma)/\Gamma_{\text{total}} \times \Gamma(p\bar{p})/\Gamma_{\text{total}}$ $\Gamma_{98}/\Gamma \times \Gamma_{56}/\Gamma$

| VALUE (units 10^{-8}) | DOCUMENT ID | TECN | COMMENT |
|---|-------------------------------------|------|---------|
| 2.12 ± 0.15 OUR FIT | Error includes scale factor of 1.2. | | |
| 1.7 ± 0.4 OUR AVERAGE | | | |

| | | | |
|-----------------|--------------|------|--|
| 1.60 ± 0.42 | ARMSTRONG 93 | E760 | $\bar{p}p \rightarrow \gamma\gamma X$ |
| 9.9 ± 4.5 | BAGLIN | 87B | SPEC $\bar{p}p \rightarrow \gamma\gamma X$ |

$\chi_{c2}(1P)$ CROSS-PARTICLE BRANCHING RATIOS

 $\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)$ $\Gamma_{14}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}$

| VALUE (units 10^{-3}) | DOCUMENT ID | TECN | COMMENT |
|---|-------------------------------------|------|---------|
| 2.24 ± 0.30 OUR FIT | Error includes scale factor of 1.2. | | |
| 2.5 ± 0.9 OUR AVERAGE | Error includes scale factor of 2.3. | | |

| | | | |
|--------------------------|---------------------------|------|---|
| 1.90 $\pm 0.14 \pm 0.44$ | BAI | 99B | BES $\psi(2S) \rightarrow \gamma \chi_{c2}$ |
| 3.8 ± 0.67 | ¹ TANENBAUM 78 | MRK1 | $\psi(2S) \rightarrow \gamma \chi_{c2}$ |

¹ The reported value is derived using $B(\psi(2S) \rightarrow \pi^+ \pi^- J/\psi) \times B(J/\psi \rightarrow \ell^+ \ell^-) = (4.6 \pm 0.7)\%$. Calculated by us using $B(J/\psi \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$.

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^0 \bar{K}^*(892)^0) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}}}{\Gamma_{\text{total}} \times \Gamma_{18} / \Gamma \times \Gamma_{181}^{\psi(2S)} / \Gamma^{\psi(2S)}}$$

| <u>VALUE (units 10^{-4})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------------------------------|-------------|---|
| 2.1 ± 0.9 OUR FIT | Error includes scale factor of 2.3. | | |
| 3.11 ± 0.36 ± 0.48 | ABLIKIM | 04H BES2 | $\psi(2S) \rightarrow \gamma \chi_{c2}$ |

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}{\Gamma_{\text{total}} \times \Gamma_{56} / \Gamma \times \Gamma_{181}^{\psi(2S)} / \Gamma_{12}^{\psi(2S)}}$$

| <u>VALUE (units 10^{-5})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------------------------------|-------------|---|
| 1.96 ± 0.10 OUR FIT | Error includes scale factor of 1.1. | | |
| 1.4 ± 1.1 | ¹ BAI | 98I BES | $\psi(2S) \rightarrow \gamma \chi_{c2} \rightarrow \gamma p\bar{p}$ |

¹ Calculated by us. The value for $B(\chi_{c2} \rightarrow p\bar{p})$ reported in BAI 98I is derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (7.8 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}}}{\Gamma_{\text{total}} \times \Gamma_{56} / \Gamma \times \Gamma_{181}^{\psi(2S)} / \Gamma^{\psi(2S)}}$$

| <u>VALUE (units 10^{-6})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------------------------------|--------------------|-------------|--|
| 6.79 ± 0.34 OUR FIT | Error includes scale factor of 1.1. | | | |
| 7.1 ± 0.5 OUR AVERAGE | Error includes scale factor of 1.2. | | | |
| 7.3 ± 0.4 ± 0.3 | 405 | ABLIKIM | 13V BES3 | $\psi(2S) \rightarrow \gamma p\bar{p}$ |

| | | | | |
|-----------------|----------------|-------------------|---------|---|
| 7.2 ± 0.7 ± 0.4 | 121 ± 12 | ¹ NAIK | 08 CLEO | $\psi(2S) \rightarrow \gamma p\bar{p}$ |
| 4.4 ± 1.6 ± 0.6 | 14.3 ± 5.2 | BAI | 04F BES | $\psi(2S) \rightarrow \gamma \chi_{c2}(1P) \rightarrow \gamma p\bar{p}$ |

¹ Calculated by us. NAIK 08 reports $B(\chi_{c2} \rightarrow p\bar{p}) = (7.7 \pm 0.8 \pm 0.4 \pm 0.5) \times 10^{-5}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$.

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \Lambda\bar{\Lambda}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}}}{\Gamma_{\text{total}} \times \Gamma_{69} / \Gamma \times \Gamma_{181}^{\psi(2S)} / \Gamma^{\psi(2S)}}$$

| <u>VALUE (units 10^{-6})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|----------------|
| 17.4 ± 1.4 OUR FIT | | | | |
| 17.3 ± 1.5 OUR AVERAGE | | | | |

| | | | | |
|------------------|-----|-------------------|----------|--|
| 18.2 ± 0.8 ± 1.7 | 670 | ABLIKIM | 21L BES3 | $\psi(2S) \rightarrow \gamma p\pi^-\bar{p}\pi^+$ |
| 15.9 ± 2.1 ± 1.0 | 71 | ¹ NAIK | 08 CLEO | $\psi(2S) \rightarrow \gamma \Lambda\bar{\Lambda}$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|------------------|-----|------------------------|----------|--|
| 18.2 ± 1.4 ± 0.9 | 207 | ^{2,3} ABLIKIM | 13H BES3 | $\psi(2S) \rightarrow \gamma \Lambda\bar{\Lambda}$ |
|------------------|-----|------------------------|----------|--|

¹ Calculated by us. NAIK 08 reports $B(\chi_{c2} \rightarrow \Lambda\bar{\Lambda}) = (17.0 \pm 2.2 \pm 1.1 \pm 1.1) \times 10^{-5}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$.

² Superseded by ABLIKIM 21L

³ Calculated by us. ABLIKIM 13H reports $B(\chi_{c2} \rightarrow \Lambda\bar{\Lambda}) = (20.8 \pm 1.6 \pm 2.3) \times 10^{-5}$ from a measurement of $B(\chi_{c2} \rightarrow \Lambda\bar{\Lambda}) \times B(\psi(2S) \rightarrow \gamma \chi_{c2})$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (8.74 \pm 0.35)\%$.

$$\Gamma(\chi_{c2}(1P) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)$$

$$\Gamma_{69}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}$$

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
| 5.0±0.4 OUR FIT | | | | |

7.1^{+3.1}_{-2.9}±1.3 $8.3^{+3.7}_{-3.4}$ ¹ BAI 03E BES $\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}$

¹ BAI 03E reports [$B(\chi_{c2} \rightarrow \Lambda\bar{\Lambda}) B(\psi(2S) \rightarrow \gamma\chi_{c2}) / B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-)$] \times $[B^2(\Lambda \rightarrow \pi^- p) / B(J/\psi \rightarrow p\bar{p})] = (1.33^{+0.59}_{-0.55} \pm 0.25)\%$. We calculate from this measurement the presented value using $B(\Lambda \rightarrow \pi^- p) = (63.9 \pm 0.5)\%$ and $B(J/\psi \rightarrow p\bar{p}) = (2.17 \pm 0.07) \times 10^{-3}$.

$$\Gamma(\chi_{c2}(1P) \rightarrow \Lambda\bar{\Lambda}\eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma_{\text{total}}$$

$$\Gamma_{74}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}$$

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|-----------|--|
| 1.00±0.20±0.14 | 32 | ABLIKIM | 22AO BES3 | $\psi(2S) \rightarrow \gamma p\pi^-\bar{p}\pi^+\gamma\gamma$ |

$$\Gamma(\chi_{c2}(1P) \rightarrow \pi\pi)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma_{\text{total}}$$

$$\Gamma_{25}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}$$

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
| 2.12±0.08 OUR FIT | | | | |

2.17±0.09 OUR AVERAGE

| | | | | |
|--------------------------|------|----------------------|----------|---|
| $2.19 \pm 0.05 \pm 0.15$ | 4.5k | ¹ ABLIKIM | 10A BES3 | $e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c2}$ |
| $2.23 \pm 0.06 \pm 0.10$ | 2.5k | ² ASNER | 09 CLEO | $\psi(2S) \rightarrow \gamma\pi^+\pi^-$ |
| $1.90 \pm 0.08 \pm 0.20$ | 0.8k | ³ ASNER | 09 CLEO | $\psi(2S) \rightarrow \gamma\pi^0\pi^0$ |

¹ Calculated by us. ABLIKIM 10A reports $B(\chi_{c2} \rightarrow \pi^0\pi^0) = (0.88 \pm 0.02 \pm 0.06 \pm 0.04) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (8.3 \pm 0.4)\%$. We have multiplied the $\pi^0\pi^0$ measurement by 3 to obtain $\pi\pi$.

² Calculated by us. ASNER 09 reports $B(\chi_{c2} \rightarrow \pi^+\pi^-) = (1.59 \pm 0.04 \pm 0.07 \pm 0.10) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$. We have multiplied the $\pi^+\pi^-$ measurement by 3/2 to obtain $\pi\pi$.

³ Calculated by us. ASNER 09 reports $B(\chi_{c2} \rightarrow \pi^0\pi^0) = (0.68 \pm 0.03 \pm 0.07 \pm 0.04) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$. We have multiplied the $\pi^0\pi^0$ measurement by 3 to obtain $\pi\pi$.

$$\Gamma(\chi_{c2}(1P) \rightarrow \pi\pi)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)$$

$$\Gamma_{25}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}$$

| VALUE (units 10^{-3}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------|------|-------------|------|---------|
| 0.611±0.024 OUR FIT | | | | |

0.54 ±0.06 OUR AVERAGE

| | | | | |
|--------------------------|--------------|------------------|---------|---|
| $0.66 \pm 0.18 \pm 0.37$ | 21 ± 6 | ¹ BAI | 03C BES | $\psi(2S) \rightarrow \gamma\pi^0\pi^0$ |
| $0.54 \pm 0.05 \pm 0.04$ | 185 ± 16 | ² BAI | 98I BES | $\psi(2S) \rightarrow \gamma\pi^+\pi^-$ |

¹ We have multiplied $\pi^0\pi^0$ measurement by 3 to obtain $\pi\pi$.

² Calculated by us. The value for $B(\chi_{c2} \rightarrow \pi^+\pi^-)$ reported by BAI 98I is derived using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (7.8 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D]. We have multiplied $\pi^+\pi^-$ measurement by 3/2 to obtain $\pi\pi$.

$$\Gamma(\chi_{c2}(1P) \rightarrow \eta\eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma_{\text{total}}$$

$$\Gamma_{31}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

| <u>VALUE (units 10^{-4})</u> | <u>CL%</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|-------------|----------------------|-------------|---|
| 0.52±0.04 OUR FIT | | | | | |
| 0.52±0.04 OUR AVERAGE | | | | | |
| 0.54±0.03±0.04 | | 386 | ¹ ABLIKIM | 10A | BES3 $e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c2}$ |
| 0.47±0.05±0.05 | | 156 | ASNER | 09 | CLEO $\psi(2S) \rightarrow \gamma\eta\eta$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| < 0.44 | | 90 | ² ADAMS | 07 | CLEO $\psi(2S) \rightarrow \gamma\chi_{c2}$ |
| < 3 | | 90 | BAI | 03C | BES $\psi(2S) \rightarrow \gamma\eta\eta \rightarrow 5\gamma$ |
| 0.62±0.31±0.19 | | | LEE | 85 | CBAL $\psi(2S) \rightarrow \text{photons}$ |

¹ Calculated by us. ABLIKIM 10A reports $B(\chi_{c2} \rightarrow \eta\eta) = (0.65 \pm 0.04 \pm 0.05 \pm 0.03) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (8.3 \pm 0.4)\%$.

² Superseded by ASNER 09.

$$\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma_{\text{total}}$$

$$\Gamma_{32}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|--------------------|--------------------|-------------|---------------------------------------|
| 9.5±1.4 OUR FIT Error includes scale factor of 2.4. | | | | |
| 10.5±0.3±0.6 | | | | |
| 1.6k | ¹ ASNER | 09 | CLEO | $\psi(2S) \rightarrow \gamma K^+ K^-$ |
| ¹ Calculated by us. ASNER 09 reports $B(\chi_{c2} \rightarrow K^+ K^-) = (1.13 \pm 0.03 \pm 0.06 \pm 0.07) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$. | | | | |

$$\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)$$

$$\Gamma_{32}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}$$

| <u>VALUE (units 10^{-3})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------------|--------------------|-------------|---------------------------------------|
| 0.27 ±0.04 OUR FIT Error includes scale factor of 2.4. | | | | |
| 0.190±0.034±0.019 | | | | |
| 115 ± 13 | ¹ BAI | 98I | BES | $\psi(2S) \rightarrow \gamma K^+ K^-$ |
| ¹ Calculated by us. The value for $B(\chi_{c2} \rightarrow K^+ K^-)$ reported by BAI 98I is derived using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (7.8 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D]. | | | | |

$$\Gamma(\chi_{c2}(1P) \rightarrow K_S^0 K_S^0)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma_{\text{total}}$$

$$\Gamma_{33}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|-------------|--------------------|-------------|--|
| 5.0 ±0.4 OUR FIT | | | | |
| 5.0 ±0.4 OUR AVERAGE | | | | |
| 4.9 ± 0.3 ± 0.3 | 373 ± 20 | ¹ ASNER | 09 | CLEO $\psi(2S) \rightarrow \gamma K_S^0 K_S^0$ |
| 5.72±0.76±0.63 | 65 | ABLIKIM | 05O | BES2 $\psi(2S) \rightarrow \gamma K_S^0 K_S^0$ |
| ¹ Calculated by us. ASNER 09 reports $B(\chi_{c2} \rightarrow K_S^0 K_S^0) = (0.53 \pm 0.03 \pm 0.03 \pm 0.03) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$. | | | | |

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow K_S^0 K_S^0)/\Gamma_{\text{total}}}{J/\psi(1S)\pi^+\pi^-} \times \frac{\Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}{\Gamma_{33}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

| VALUE (units 10^{-5}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|------|---------|
|--------------------------|-------------|------|---------|

14.3±1.1 OUR FIT**14.7±4.1±3.3** ${}^1 \text{BAI}$ 99B BES $\psi(2S) \rightarrow \gamma K_S^0 K_S^0$

¹ Calculated by us. The value of $B(\chi_{c2} \rightarrow K_S^0 K_S^0)$ reported by BAI 99B was derived using $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (7.8 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}}{J/\psi(1S)\pi^+\pi^-} \times \frac{\Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma_{\text{total}}}{\Gamma_{42}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

1.22±0.17 OUR FIT**1.15±0.18 OUR AVERAGE**

| | | | | |
|--------------------------|----|----------------------|-----|--|
| $1.21 \pm 0.19 \pm 0.09$ | 37 | ¹ ATHAR | 07 | CLEO $\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp$ |
| $0.97 \pm 0.32 \pm 0.13$ | 28 | ² ABLIKIM | 06R | BES2 $\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp$ |

¹ Calculated by us. ATHAR 07 reports $B(\chi_{c2} \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.}) = (1.3 \pm 0.2 \pm 0.1 \pm 0.1) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$.

² Calculated by us. ABLIKIM 06R reports $B(\chi_{c2} \rightarrow K_S^0 K^\pm \pi^\mp) = (0.6 \pm 0.2 \pm 0.1) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (8.1 \pm 0.6)\%$. We have multiplied by 2 to obtain $\bar{K}^0 K^+ \pi^- + \text{c.c.}$ from $K_S^0 K^\pm \pi^\mp$.

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow 2(\pi^+\pi^-))/\Gamma_{\text{total}}}{J/\psi(1S)\pi^+\pi^-} \times \frac{\Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}{\Gamma_1/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

| VALUE (units 10^{-3}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|------|---------|
|--------------------------|-------------|------|---------|

2.7±0.4 OUR FIT Error includes scale factor of 1.4.**3.1±1.0 OUR AVERAGE** Error includes scale factor of 2.5.

| | | | | |
|-----------------------|------------------------|--------------|--------------|--|
| $2.3 \pm 0.1 \pm 0.5$ | ¹ BAI | 99B | BES | $\psi(2S) \rightarrow \gamma\chi_{c2}$ |
| 4.3 ± 0.6 | ² TANENBAUM | 78 | MRK1 | $\psi(2S) \rightarrow \gamma\chi_{c2}$ |

¹ Calculated by us. The value for $B(\chi_{c2} \rightarrow 2\pi^+ 2\pi^-)$ reported in BAI 99B is derived using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (7.8 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

² The value for $B(\psi(2S) \rightarrow \gamma\chi_{c2}) \times B(\chi_{c2} \rightarrow 2\pi^+ \pi^-)$ reported in TANENBAUM 78 is derived using $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times B(J/\psi(1S)\ell^+\ell^-) = (4.6 \pm 0.7)\%$. Calculated by us using $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1181 \pm 0.0020$.

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- K^+ K^-)/\Gamma_{\text{total}}}{J/\psi(1S)\pi^+\pi^-} \times \frac{\Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma_{\text{total}}}{\Gamma_{51}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

1.56±0.21 OUR FIT Error includes scale factor of 1.1.**1.76±0.16±0.24** ${}^1 \text{ABLIKIM}$ 06T BES2 $\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

¹ Calculated by us. The value of $B(\chi_{c2} \rightarrow 2K^+ 2K^-)$ reported by ABLIKIM 06T was derived using $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.1 \pm 0.4)\%$.

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- K^+ K^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))}{\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)} \frac{\Gamma_{51}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

| <u>VALUE (units 10^{-4})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|----------------|
|---|--------------------|-------------|----------------|

4.5±0.6 OUR FIT Error includes scale factor of 1.1.**3.6±0.6±0.6** ¹ BAI 99B BES $\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

¹ Calculated by us. The value of $B(\chi_{c2} \rightarrow 2K^+ 2K^-)$ reported by BAI 99B was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (7.8 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \phi\phi)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma_{\text{total}}}{\Gamma_{20}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|----------------|
|---|-------------|--------------------|-------------|----------------|

1.15±0.07 OUR FIT Error includes scale factor of 1.8.**0.98±0.13 OUR AVERAGE** Error includes scale factor of 1.3.

| | | | | |
|--------------------------|-----|----------------------|----------|---|
| $0.94 \pm 0.03 \pm 0.10$ | 849 | ¹ ABLIKIM | 11K BES3 | $\psi(2S) \rightarrow \gamma$ hadrons |
| $1.38 \pm 0.24 \pm 0.23$ | 41 | ² ABLIKIM | 06T BES2 | $\psi(2S) \rightarrow \gamma 2K^+ 2K^-$ |

¹ Calculated by us. The value of $B(\chi_{c2} \rightarrow \phi\phi)$ reported by ABLIKIM 11K was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35)\%$.

² Calculated by us. The value of $B(\chi_{c2} \rightarrow \phi\phi)$ reported by ABLIKIM 06T was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.1 \pm 0.4)\%$.

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \phi\phi)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)}{\Gamma_{20}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

| <u>VALUE (units 10^{-4})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|----------------|
|---|--------------------|-------------|----------------|

3.31±0.21 OUR FIT Error includes scale factor of 1.7.**4.8 ± 1.3 ± 1.3** ¹ BAI 99B BES $\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

¹ Calculated by us. The value of $B(\chi_{c2} \rightarrow \phi\phi)$ reported by BAI 99B was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (7.8 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^+ \bar{p} K_S^0 + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma_{\text{total}}}{\Gamma_{81}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

| <u>VALUE (units 10^{-6})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|----------------------|-------------|--|
| $7.85 \pm 0.77 \pm 0.44$ | 129 | ¹ ABLIKIM | 19BB BES3 | $\psi(2S) \rightarrow \gamma \Sigma^+ \bar{p} K_S^0 + \text{c.c.}$ |

¹ Calculated by us. ABLIKIM 19BB reports $B(\chi_{c2} \rightarrow \Sigma^+ \bar{p} K_S^0 + \text{c.c.}) = (8.25 \pm 0.83 \pm 0.49) \times 10^{-5}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (9.52 \pm 0.20)\%$ and other branching fractions from PDG 18.

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^0 \bar{p} K^+ + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma_{\text{total}}}{\Gamma_{82}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|----------------------|-------------|--|
| $0.87 \pm 0.06 \pm 0.04$ | 271 | ¹ ABLIKIM | 20AE BES3 | $\psi(2S) \rightarrow \gamma \Sigma^0 \bar{p} K^+ + \text{c.c.}$ |

¹ Calculated by us. ABLIKIM 20AE reports $B(\chi_{c2} \rightarrow \Sigma^0 \bar{p} K^+ + \text{c.c.}) = (0.91 \pm 0.06 \pm 0.05) \times 10^{-4}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (9.52 \pm 0.20)\%$ and other branching fractions from PDG 20.

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma_{\text{total}}}{\Gamma_{94}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma^{\psi(2S)}}$$

| <u>VALUE (units 10^{-2})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|---|-------------|--|
| 1.82 ±0.07 OUR FIT | | Error includes scale factor of 1.9. | | |
| 1.69 ±0.16 OUR AVERAGE | | Error includes scale factor of 3.4. See the ideogram below. | | |
| 1.996±0.008±0.070 | 81k | ¹ ABLIKIM | 17N BES3 | $\psi(2S) \rightarrow \gamma\gamma J/\psi$ |
| 1.793±0.008±0.163 | 1.0M | ABLIKIM | 17U BES3 | $e^+ e^- \rightarrow \gamma X$ |
| 1.62 ±0.04 ±0.12 | 5.8k | BAI | 04I BES2 | $\psi(2S) \rightarrow J/\psi\gamma\gamma$ |
| 0.99 ±0.10 ±0.08 | | GAISER | 86 CBAL | $\psi(2S) \rightarrow \gamma X$ |
| 1.47 ±0.17 | | ² OREGLIA | 82 CBAL | $\psi(2S) \rightarrow \gamma\chi_{c2}$ |
| 1.8 ±0.5 | | ³ BRANDELIK | 79B DASP | $\psi(2S) \rightarrow \gamma\chi_{c2}$ |
| 1.2 ±0.2 | | ³ BARTEL | 78B CNTR | $\psi(2S) \rightarrow \gamma\chi_{c2}$ |
| 2.2 ±1.2 | | ⁴ BIDDICK | 77 CNTR | $e^+ e^- \rightarrow \gamma X$ |
| 1.2 ±0.7 | | ² WHITAKER | 76 MRK1 | $e^+ e^-$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 1.874±0.007±0.102 | 76k | ⁵ ABLIKIM | 120 BES3 | $\psi(2S) \rightarrow \gamma\chi_{c2}$ |
| 1.95 ±0.02 ±0.07 | 12.4k | ⁶ MENDEZ | 08 CLEO | $\psi(2S) \rightarrow \gamma\chi_{c2}$ |
| 1.85 ±0.04 ±0.07 | 1.9k | ⁷ ADAM | 05A CLEO | Repl. by MENDEZ 08 |

¹ Uses $B(J/\psi \rightarrow e^+ e^-) = (5.971 \pm 0.032)\%$ and $B(J/\psi \rightarrow \mu^+ \mu^-) = (5.961 \pm 0.033)\%$.

² Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$.


³ Recalculated by us using $B(J/\psi(1S) \rightarrow \mu^+ \mu^-) = 0.0588 \pm 0.0010$.

⁴ Assumes isotropic gamma distribution.

⁵ Superseded by ABLIKIM 17N.

⁶ Not independent from other measurements of MENDEZ 08.

⁷ Not independent from other values reported by ADAM 05A.



$$\Gamma(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}} (\text{units } 10^{-2})$$

$$\Gamma(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \quad \frac{\Gamma_{94}/\Gamma}{\Gamma_{181}^{\psi(2S)} / \Gamma_{12}^{\psi(2S)}}$$

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------------------------------|-------------|------|---------|
| 5.26±0.19 OUR FIT | Error includes scale factor of 1.8. | | | |

5.53±0.17 OUR AVERAGE

| | | | | |
|--------------------------|-------|----------------------|-----|--|
| $5.56 \pm 0.05 \pm 0.16$ | 12.4k | MENDEZ | 08 | CLEO $\psi(2S) \rightarrow \gamma \chi_{c2}$ |
| 6.0 ± 2.8 | 1.3k | ¹ ABLIKIM | 04B | BES $\psi(2S) \rightarrow J/\psi X$ |
| 3.9 ± 1.2 | | ² HIMEL | 80 | MRK2 $\psi(2S) \rightarrow \gamma \chi_{c2}$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|--------------------------|------|-------------------|-----|-------------------------|
| $5.52 \pm 0.13 \pm 0.13$ | 1.9k | ³ ADAM | 05A | CLEO Repl. by MENDEZ 08 |
|--------------------------|------|-------------------|-----|-------------------------|

¹ From a fit to the J/ψ recoil mass spectra.

² The value for $B(\psi(2S) \rightarrow \gamma \chi_{c2}) \times B(\chi_{c2} \rightarrow \gamma J/\psi(1S))$ reported in HIMEL 80 is derived using $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (33 \pm 3)\%$ and $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.138 \pm 0.018$. Calculated by us using $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = (0.1181 \pm 0.0020)$.

³ Not independent from other values reported by ADAM 05A.

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}}{\Gamma_{98}/\Gamma} \times \frac{\Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma_{\text{total}}}{\Gamma_{181}^{\psi(2S)}/\Gamma^{\psi(2S)}}$$

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------------------------------|-----------|--|
| 2.74±0.10 OUR FIT | | Error includes scale factor of 1.3. | | |
| 2.82±0.10 OUR AVERAGE | | | | |
| 2.83±0.08±0.06 | 5k | ¹ ABLIKIM | 17AE BES3 | $\psi(2S) \rightarrow \gamma\chi_{c2} \rightarrow 3\gamma$ |
| 2.68±0.28±0.15 | 0.3k | ECKLUND | 08A CLEO | $\psi(2S) \rightarrow \gamma\chi_{c2} \rightarrow 3\gamma$ |
| 7.0 ± 2.1 ± 2.0 | | LEE | 85 CBAL | $\psi(2S) \rightarrow \gamma\chi_{c2}$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 2.81±0.17±0.15 | 1.1k | ² ABLIKIM | 12A BES3 | $\psi(2S) \rightarrow \gamma\chi_{c2} \rightarrow 3\gamma$ |

¹ ABLIKIM 17AE measures the ratio of two-photon partial widths for the helicity $\lambda = 0$ and helicity $\lambda = 2$ components to be $f_{0/2} = \frac{\Gamma_{\gamma\gamma}^{\lambda=0}}{\Gamma_{\gamma\gamma}^{\lambda=2}} = 0.000 \pm 0.006 \pm 0.012$.

² ABLIKIM 12A measures the ratio of two-photon partial widths for the helicity $\lambda = 0$ and helicity $\lambda = 2$ components to be $f_{0/2} = \frac{\Gamma_{\gamma\gamma}^{\lambda=0}}{\Gamma_{\gamma\gamma}^{\lambda=2}} = 0.00 \pm 0.02 \pm 0.02$. Superseded by ABLIKIM 17AE.

| VALUE | EVTS | DOCUMENT ID | TECN | $\Gamma_{98}/\Gamma_{98}^{\chi_{c0}(1P)}$ |
|---|------|------------------------|-----------|--|
| 0.292±0.028 OUR AVERAGE | | | | |
| 0.295±0.014±0.028 | 8k | ¹ ABLIKIM | 17AE BES3 | $\psi(2S) \rightarrow \gamma\chi_{cJ} \rightarrow 3\gamma$ |
| 0.278±0.050±0.036 | 0.5k | ¹ ECKLUND | 08A CLEO | $\psi(2S) \rightarrow \gamma\chi_{cJ} \rightarrow 3\gamma$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.271±0.029±0.030 | 1.9k | ^{1,2} ABLIKIM | 12A BES3 | $\psi(2S) \rightarrow \gamma\chi_{cJ} \rightarrow 3\gamma$ |
| ¹ Not independent from the values of $\Gamma(\chi_{c0}, \chi_{c2})$ and $B(\psi(2S) \rightarrow \chi_{c0}, \chi_{c2})$. | | | | |
| ² Superseded by ABLIKIM 17AE. | | | | |

MULTIPOLE AMPLITUDES IN $\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)$ RADIATIVE DECAY

$a_2 = M2/\sqrt{E1^2 + M2^2 + E3^2}$ Magnetic quadrupole fractional transition amplitude


| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-------|-------------------------|----------|---|
| -11.0± 1.0 OUR AVERAGE | | | | |
| -12.0± 1.3±0.4 | 89k | ¹ ABLIKIM | 17N BES3 | $\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$ |
| -9.3± 1.6±0.3 | 19.8k | ² ARTUSO | 09 CLEO | $\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$ |
| -9.3 ^{+3.9} _{-4.1} ±0.6 | 5.9k | ³ AMBROGIANI | 02 E835 | $p\bar{p} \rightarrow \chi_{c2} \rightarrow J/\psi\gamma$ |
| -14 ± 6 | 1.9k | ³ ARMSTRONG | 93E E760 | $p\bar{p} \rightarrow \chi_{c2} \rightarrow J/\psi\gamma$ |
| -33.3 ^{+11.6} _{-29.2} | 441 | ³ OREGLIA | 82 CBAL | $\psi(2S) \rightarrow \chi_{c1}\gamma \rightarrow J/\psi\gamma\gamma$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| -7.9± 1.9±0.3 | 19.8k | ⁴ ARTUSO | 09 CLEO | $\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$ |
| ¹ Correlated with a_3 , b_2 , and b_3 with correlation coefficients $\rho_{a_2 a_3} = 0.733$, $\rho_{a_2 b_2} = -0.605$, and $\rho_{a_2 b_3} = -0.095$. | | | | |
| ² From a fit with floating $M2$ amplitudes a_2 and b_2 , and fixed $E3$ amplitudes $a_3=b_3=0$. | | | | |
| ³ Assuming $a_3=0$. | | | | |
| ⁴ From a fit with floating $M2$ and $E3$ amplitudes a_2 , b_2 , and a_3 , and b_3 . | | | | |

$a_3 = E3/\sqrt{E1^2 + M2^2 + E3^2}$ Electric octupole fractional transition amplitude

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|---|----------------------|----------|---|
| -0.3±1.0 OUR AVERAGE | Error includes scale factor of 1.3. See the ideogram below. | | | |
| -1.3±0.9±0.4 | 89k | ¹ ABLIKIM | 17N BES3 | $\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$ |
| 1.7±1.4±0.3 | 19.8k | ² ARTUSO | 09 CLEO | $\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$ |
| 2.0 ^{+5.5} _{-4.4} ±0.9 | 5908 | AMBROGIANI | 02 E835 | $p\bar{p} \rightarrow \chi_{c2} \rightarrow J/\psi\gamma$ |
| 0 ⁺⁶ ₋₅ | 1904 | ARMSTRONG | 93E E760 | $p\bar{p} \rightarrow \chi_{c2} \rightarrow J/\psi\gamma$ |

¹ Correlated with a_2 , b_2 , and b_3 with correlation coefficients $\rho_{a_2 a_3} = 0.733$, $\rho_{a_3 b_2} = -0.422$, and $\rho_{a_3 b_3} = -0.024$.

² From a fit with floating $M2$ and $E3$ amplitudes a_2 , b_2 , and a_3 , and b_3 .



$$a_3 = E3/\sqrt{E1^2 + M2^2 + E3^2} \text{ Electric octupole fractional transition amplitude (units } 10^{-2})$$

MULTIPOLE AMPLITUDES IN $\psi(2S) \rightarrow \gamma\chi_{c2}(1P)$ RADIATIVE DECAY

$b_2 = M2/\sqrt{E1^2 + M2^2 + E3^2}$ Magnetic quadrupole fractional transition amplitude

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------|---|----------------------|----------|--|
| 1.9±0.9 OUR AVERAGE | Error includes scale factor of 1.4. See the ideogram below. | | | |
| 1.7±0.8±0.2 | 89k | ¹ ABLIKIM | 17N BES3 | $\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$ |
| 4.6±1.0±1.3 | 13.8k | ² ABLIKIM | 11I BES3 | $\psi(2S) \rightarrow \gamma\pi^+\pi^-, \gamma K^+K^-$ |
| 0.2±1.5±0.4 | 19.8k | ³ ARTUSO | 09 CLEO | $\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$ |

| | | | | | |
|----------------------|-----|----------------------|-----|------|--|
| $-5.1^{+5.4}_{-3.6}$ | 721 | ² ABLIKIM | 04I | BES2 | $\psi(2S) \rightarrow \gamma\pi^+\pi^-, \gamma K^+K^-$ |
| $13.2^{+9.8}_{-7.5}$ | 441 | ⁴ OREGLIA | 82 | CBAL | $\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •


| | | | | | |
|-----------------------|-------|---------------------|----|------|---|
| $1.0 \pm 1.3 \pm 0.3$ | 19.8k | ⁴ ARTUSO | 09 | CLEO | $\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$ |
|-----------------------|-------|---------------------|----|------|---|

¹ Correlated with a_2 , a_3 , and b_3 with correlation coefficients $\rho_{a_2} b_2 = -0.605$, $\rho_{a_3} b_2 = -0.422$, and $\rho_{b_2} b_3 = 0.384$.

² From a fit with floating $M2$ and $E3$ amplitudes b_2 and b_3 .

³ From a fit with floating $M2$ and $E3$ amplitudes a_2 , b_2 , and a_3 , and b_3 .

⁴ From a fit with floating $M2$ amplitudes a_2 and b_2 , and fixed $E3$ amplitudes $a_3=b_3=0$.



$$b_2 = M2/\sqrt{E1^2 + M2^2 + E3^2} \quad \text{Magnetic quadrupole fractional transition amplitude (units } 10^{-2})$$

$b_3 = E3/\sqrt{E1^2 + M2^2 + E3^2}$ Electric octupole fractional transition amplitude

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------------------|-------|----------------------|------|---|
| -1.0±0.6 OUR AVERAGE | | | | |
| -1.4±0.7±0.4 | 89k | ¹ ABLIKIM | 17N | BES3 $\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$ |
| 1.5±0.8±1.8 | 13.8k | ² ABLIKIM | 11I | BES3 $\psi(2S) \rightarrow \gamma\pi^+\pi^-, \gamma K^+K^-$ |
| -0.8±1.2±0.2 | 19.8k | ARTUSO | 09 | CLEO $\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$ |
| -2.7 ^{+4.3} _{-2.9} | 721 | ² ABLIKIM | 04I | BES2 $\psi(2S) \rightarrow \gamma\pi^+\pi^-, \gamma K^+K^-$ |

¹ Correlated with a_2 , a_3 , and b_2 with correlation coefficients $\rho_{a_2} b_3 = -0.095$, $\rho_{a_3} b_3 = -0.024$, and $\rho_{b_2} b_3 = 0.384$.

² From a fit with floating $M2$ and $E3$ amplitudes b_2 and b_3 .

MULTIPOLE AMPLITUDE RATIOS IN RADIATIVE DECAYS

$\psi(2S) \rightarrow \gamma\chi_{c2}(1P)$ and $\chi_{c2} \rightarrow \gamma J/\psi(1S)$

b_2/a_2 Magnetic quadrupole transition amplitude ratio

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------|---------------------|------|--|
| -11^{+14}_{-15} | 19.8k | ¹ ARTUSO | 09 | CLEO $\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$ |

¹ Statistical and systematic errors combined. From a fit with floating $M2$ amplitudes a_2 and b_2 , and fixed $E3$ amplitudes $a_3=b_3=0$. Not independent of values for $a_2(\chi_{c2}(1P))$ and $b_2(\chi_{c2}(1P))$ from ARTUSO 09.

$\chi_{c2}(1P)$ REFERENCES

| | | | | |
|---------------|------|------------------|----------------------------------|------------------|
| ABLIKIM | 23N | JHEP 2305 069 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 23T | PR D107 092004 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| SEINO | 23 | JHEP 2301 160 | Y. Seino <i>et al.</i> | (BELLE Collab.) |
| ABLIKIM | 22AO | PR D106 072004 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 22O | JHEP 2206 074 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 22Q | PR D106 032014 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| PDG | 22 | PTEP 2022 083C01 | R.L. Workman <i>et al.</i> | (PDG Collab.) |
| ABLIKIM | 21AV | JHEP 2111 217 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 21L | PR D103 112004 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 20AE | PR D102 092006 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 20B | PR D101 012012 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 20I | PR D101 092002 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| PDG | 20 | PTEP 2020 083C01 | P.A. Zyla <i>et al.</i> | (PDG Collab.) |
| ABLIKIM | 19AA | PR D99 052008 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 19AU | PR D100 052010 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 19BB | PR D100 092006 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 19J | PR D99 012015 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 19Z | PR D99 051101 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 18V | PR D97 052011 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| PDG | 18 | PR D98 030001 | M. Tanabashi <i>et al.</i> | (PDG Collab.) |
| AAIJ | 17BB | EPJ C77 609 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 17BI | PRL 119 221801 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| ABLIKIM | 17AE | PR D96 092007 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 17AG | PR D96 111102 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 17AI | PR D96 112006 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 17I | PRL 118 221802 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 17N | PR D95 072004 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 17U | PR D96 032001 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| PDG | 16 | CP C40 100001 | C. Patrignani <i>et al.</i> | (PDG Collab.) |
| ABLIKIM | 15I | PR D91 092006 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 15M | PR D91 112008 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 15N | PR D91 112018 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 14J | PR D89 074030 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 13B | PR D87 012002 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 13D | PR D87 012007 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 13H | PR D87 032007 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 13V | PR D88 112001 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| UEHARA | 13 | PTEP 2013 123C01 | S. Uehara <i>et al.</i> | (BELLE Collab.) |
| ABLIKIM | 12A | PR D85 112008 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 12I | PR D86 052004 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 12J | PR D86 052011 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 12O | PRL 109 172002 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| LEES | 12AE | PR D86 092005 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| LIU | 12B | PRL 108 232001 | Z.Q. Liu <i>et al.</i> | (BELLE Collab.) |
| ABLIKIM | 11A | PR D83 012006 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 11E | PR D83 112005 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 11F | PR D83 112009 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 11I | PR D84 092006 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 11K | PRL 107 092001 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| DEL-AMO-SA... | 11M | PR D84 012004 | P. del Amo Sanchez <i>et al.</i> | (BABAR Collab.) |
| ABLIKIM | 10A | PR D81 052005 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ONYISI | 10 | PR D82 011103 | P.U.E. Onyisi <i>et al.</i> | (CLEO Collab.) |
| UEHARA | 10A | PR D82 114031 | S. Uehara <i>et al.</i> | (BELLE Collab.) |

| | | | | |
|---------------|-----|----------------|-------------------------------|---------------------------------|
| ARTUSO | 09 | PR D80 112003 | M. Artuso <i>et al.</i> | (CLEO Collab.) |
| ASNER | 09 | PR D79 072007 | D.M. Asner <i>et al.</i> | (CLEO Collab.) |
| UEHARA | 09 | PR D79 052009 | S. Uehara <i>et al.</i> | (BELLE Collab.) |
| BENNETT | 08A | PRL 101 151801 | J.V. Bennett <i>et al.</i> | (CLEO Collab.) |
| ECKLUND | 08A | PR D78 091501 | K.M. Ecklund <i>et al.</i> | (CLEO Collab.) |
| HE | 08B | PR D78 092004 | Q. He <i>et al.</i> | (CLEO Collab.) |
| MENDEZ | 08 | PR D78 011102 | H. Mendez <i>et al.</i> | (CLEO Collab.) |
| NAIK | 08 | PR D78 031101 | P. Naik <i>et al.</i> | (CLEO Collab.) |
| UEHARA | 08 | EPJ C53 1 | S. Uehara <i>et al.</i> | (BELLE Collab.) |
| ADAMS | 07 | PR D75 071101 | G.S. Adams <i>et al.</i> | (CLEO Collab.) |
| ATHAR | 07 | PR D75 032002 | S.B. Athar <i>et al.</i> | (CLEO Collab.) |
| CHEN | 07B | PL B651 15 | W.T. Chen <i>et al.</i> | (BELLE Collab.) |
| ABLIKIM | 06D | PR D73 052006 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 06I | PR D74 012004 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 06R | PR D74 072001 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 06T | PL B642 197 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| DOBBS | 06 | PR D73 071101 | S. Dobbs <i>et al.</i> | (CLEO Collab.) |
| ABLIKIM | 05G | PR D71 092002 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 05N | PL B630 7 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 05O | PL B630 21 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ADAM | 05A | PRL 94 232002 | N.E. Adam <i>et al.</i> | (CLEO Collab.) |
| ANDREOTTI | 05A | NP B717 34 | M. Andreotti <i>et al.</i> | (FNAL E835 Collab.) |
| NAKAZAWA | 05 | PL B615 39 | H. Nakazawa <i>et al.</i> | (BELLE Collab.) |
| ABLIKIM | 04B | PR D70 012003 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 04H | PR D70 092003 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 04I | PR D70 092004 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| BAI | 04F | PR D69 092001 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| BAI | 04I | PR D70 012006 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| AULCHENKO | 03 | PL B573 63 | V.M. Aulchenko <i>et al.</i> | (KEDR Collab.) |
| BAI | 03C | PR D67 032004 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| BAI | 03E | PR D67 112001 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| ABE | 02T | PL B540 33 | K. Abe <i>et al.</i> | (BELLE Collab.) |
| AMBROGIANI | 02 | PR D65 052002 | M. Ambrogiani <i>et al.</i> | (FNAL E835 Collab.) |
| EISENSTEIN | 01 | PRL 87 061801 | B.I. Eisenstein <i>et al.</i> | (CLEO Collab.) |
| AMBROGIANI | 00B | PR D62 052002 | M. Ambrogiani <i>et al.</i> | (FNAL E835 Collab.) |
| ACCIARRI | 99E | PL B453 73 | M. Acciari <i>et al.</i> | (L3 Collab.) |
| BAI | 99B | PR D60 072001 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| ACKER...,K... | 98 | PL B439 197 | K. Ackerstaff <i>et al.</i> | (OPAL Collab.) |
| BAI | 98D | PR D58 092006 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| BAI | 98I | PRL 81 3091 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| DOMINICK | 94 | PR D50 4265 | J. Dominick <i>et al.</i> | (CLEO Collab.) |
| ARMSTRONG | 93 | PRL 70 2988 | T.A. Armstrong <i>et al.</i> | (FNAL E760 Collab.) |
| ARMSTRONG | 93E | PR D48 3037 | T.A. Armstrong <i>et al.</i> | (FNAL-E760 Collab.) |
| BAUER | 93 | PL B302 345 | D.A. Bauer <i>et al.</i> | (TPC Collab.) |
| ARMSTRONG | 92 | NP B373 35 | T.A. Armstrong <i>et al.</i> | (FNAL, FERR, GENO+) |
| Also | | PRL 68 1468 | T.A. Armstrong <i>et al.</i> | (FNAL, FERR, GENO+) |
| BAGLIN | 87B | PL B187 191 | C. Baglin <i>et al.</i> | (R704 Collab.) |
| BAGLIN | 86B | PL B172 455 | C. Baglin | (LAPP, CERN, GENO, LYON, OSLO+) |
| GAISER | 86 | PR D34 711 | J. Gaiser <i>et al.</i> | (Crystal Ball Collab.) |
| LEE | 85 | SLAC 282 | R.A. Lee | (SLAC) |
| LEMOIGNE | 82 | PL 113B 509 | Y. Lemoigne <i>et al.</i> | (SACL, LOIC, SHMP+) |
| OREGLIA | 82 | PR D25 2259 | M.J. Oreglia <i>et al.</i> | (SLAC, CIT, HARV+) |
| Also | | Private Comm. | M.J. Oreglia | (EFI) |
| BARATE | 81 | PR D24 2994 | R. Barate <i>et al.</i> | (SACL, LOIC, SHMP, CERN+) |
| HIMEL | 80 | PRL 44 920 | T. Himel <i>et al.</i> | (LBL, SLAC) |
| Also | | Private Comm. | G. Trilling | (LBL, UCB) |
| BRANDELIK | 79B | NP B160 426 | R. Brandelik <i>et al.</i> | (DASP Collab.) |
| BARTEL | 78B | PL 79B 492 | W. Bartel <i>et al.</i> | (DESY, HEIDP) |
| TANENBAUM | 78 | PR D17 1731 | W.M. Tanenbaum <i>et al.</i> | (SLAC, LBL) |
| Also | | Private Comm. | G. Trilling | (LBL, UCB) |
| BIDDICK | 77 | PRL 38 1324 | C.J. Biddick <i>et al.</i> | (UCSD, UMD, PAVI+) |
| WHITAKER | 76 | PRL 37 1596 | J.S. Whitaker <i>et al.</i> | (SLAC, LBL) |