

$f_0(1710)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

See the review on "Spectroscopy of Light Meson Resonances."

$f_0(1710)$ T-MATRIX POLE \sqrt{s}

Note that $\Gamma = -2 \text{Im}(\sqrt{s})$.

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
(1680–1820) – i (50–180) OUR ESTIMATE			
$(1769 \pm 8) - i (78 \pm 6)$	¹ RODAS	22	RVUE $J/\psi(1S) \rightarrow \gamma (\pi\pi, K\bar{K})$
$(1700 \pm 18) - i (127 \pm 12)$	SARANTSEV	21	RVUE $J/\psi(1S) \rightarrow \gamma (\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
$(1803 \pm 3.5^{+45.5}_{-10.4}) - i (145 \pm 2.5^{+16.3}_{-9.6})$	² ALBRECHT	20	RVUE $0.9 \bar{p}p \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta\eta, \pi^0 K^+ K^-$
$(1732 \pm 15) - i (160^{+25}_{-10})$	³ ANISOVICH	03	RVUE $\pi\pi, K\bar{K}, \eta\eta, \eta\eta', \pi\pi\pi\pi$
$(1698 \pm 18) - i (60 \pm 13)$	BARBERIS	00E	OMEG 450 $pp \rightarrow p_f \eta\eta p_S$
$(1770 \pm 12) - i (110 \pm 20)$	⁴ ANISOVICH	99B	SPEC $0.6\text{--}1.2 p\bar{p} \rightarrow \eta\eta\pi^0$
$(1727 \pm 12 \pm 11) - i (63 \pm 8 \pm 9)$	BARBERIS	99D	OMEG 450 $pp \rightarrow K^+ K^-, \pi^+ \pi^-$
$(1750 \pm 30) - i (125 \pm 70)$	ANISOVICH	98B	RVUE Compilation

¹ T-matrix pole from coupled channel K-matrix fit to data on $J/\psi \rightarrow \gamma \pi^0 \pi^0$ (ABLIKIM 15AE) and $J/\psi \rightarrow \gamma K_S^0 K_S^0$ (ABLIKIM 18AA).

² T-matrix pole, 5 poles, 5 channels, including scattering data from HYAMS 75 ($\pi\pi$), LONGACRE 86 ($K\bar{K}$), BINON 83 ($\eta\eta$), and BINON 84C ($\eta\eta'$).

³ Solution I.

⁴ Not seen by AMSLER 02.

$f_0(1710)$ Breit-Wigner MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1733^{+8}_{-7} OUR AVERAGE Error includes scale factor of 1.5. See the ideogram below.				
$1757 \pm 24 \pm 9$		LEES	21A	BABR $\eta_c(1S) \rightarrow \eta' K^+ K^-$
$1759 \pm 6^{+14}_{-25}$	5.5k	¹ ABLIKIM	13N	BES3 $e^+ e^- \rightarrow J/\psi \rightarrow \gamma\eta\eta$
$1750^{+6}_{-7}^{+29}_{-18}$		² UEHARA	13	BELL $\gamma\gamma \rightarrow K_S^0 K_S^0$
$1701 \pm 5^{+9}_{-2}$	4k	³ CHEKANOV	08	ZEUS $ep \rightarrow K_S^0 K_S^0 X$
$1765^{+4}_{-3} \pm 13$		ABLIKIM	06V	BES2 $e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$
1738 ± 30		ABLIKIM	04E	BES2 $J/\psi \rightarrow \omega K^+ K^-$
$1740 \pm 4^{+10}_{-25}$		BAI	03G	BES $J/\psi \rightarrow \gamma K\bar{K}$
1740^{+30}_{-25}		BAI	00A	BES $J/\psi \rightarrow \gamma (\pi^+ \pi^- \pi^+ \pi^-)$
1710 ± 25		⁴ FRENCH	99	300 $pp \rightarrow p_f (K^+ K^-) p_S$

• • • We do not use the following data for averages, fits, limits, etc. • • •


1744 ± 7 ± 5	381	5,6	DOBBS	15		$J/\psi \rightarrow \gamma \pi^+ \pi^-$
1705 ± 11 ± 5	237	5,6	DOBBS	15		$\psi(2S) \rightarrow \gamma \pi^+ \pi^-$
1706 ± 4 ± 5	1.0k	5,6	DOBBS	15		$J/\psi \rightarrow \gamma K^+ K^-$
1690 ± 8 ± 3	349	5,6	DOBBS	15		$\psi(2S) \rightarrow \gamma K^+ K^-$
1750 ± 13			AMSLER	06	CBAR	$1.64 \bar{p} p \rightarrow K^+ K^- \pi^0$
1747 ± 5	80k	7	UMAN	06	E835	$5.2 \bar{p} p \rightarrow \eta \eta \pi^0$
1776 ± 15			VLADIMIRSK...	06	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
1670 ± 20			BINON	05	GAMS	$33 \pi^- p \rightarrow \eta \eta n$
1682 ± 16			TIKHOMIROV	03	SPEC	$40.0 \pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
1670 ± 26	3.6k	8	NICHITIU	02	OBLX	$0 \bar{p} p \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
1730 ± 15			BARBERIS	99	OMEG	$450 p p \rightarrow p_S p_f K^+ K^-$
1750 ± 20			BARBERIS	99B	OMEG	$450 p p \rightarrow p_S p_f \pi^+ \pi^-$
1720 ± 39			BAI	98H	BES	$J/\psi \rightarrow \gamma \pi^0 \pi^0$
1775 ± 1.5	57	9	BARKOV	98		$\pi^- p \rightarrow K_S^0 K_S^0 n$
1690 ± 11		10	ABREU	96C	DLPH	$Z^0 \rightarrow K^+ K^- + X$
1696 ± 5 ⁺⁹ ₋₃₄		11	BAI	96C	BES	$J/\psi \rightarrow \gamma K^+ K^-$
1781 ± 8 ⁺¹⁰ ₋₃₁			BAI	96C	BES	$J/\psi \rightarrow \gamma K^+ K^-$
1768 ± 14			BALOSHIN	95	SPEC	$40 \pi^- C \rightarrow K_S^0 K_S^0 X$
1750 ± 15		12	BUGG	95	MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$
1620 ± 16		11	BUGG	95	MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$
1748 ± 10		13	ARMSTRONG	93C	E760	$\bar{p} p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$
~ 1750			BREAKSTONE	93	SFM	$p p \rightarrow p p \pi^+ \pi^- \pi^+ \pi^-$
1744 ± 15		14	ALDE	92D	GAM2	$38 \pi^- p \rightarrow \eta \eta n$
1713 ± 10		15	ARMSTRONG	89D	OMEG	$300 p p \rightarrow p p K^+ K^-$
1706 ± 10		15	ARMSTRONG	89D	OMEG	$300 p p \rightarrow p p K_S^0 K_S^0$
1707 ± 10		13	AUGUSTIN	88	DM2	$J/\psi \rightarrow \gamma K^+ K^-, K_S^0 K_S^0$
1700 ± 15		11	BOLONKIN	88	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
1720 ± 60			BOLONKIN	88	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
1638 ± 10		16	FALVARD	88	DM2	$J/\psi \rightarrow \phi K^+ K^-, K_S^0 K_S^0$
1690 ± 4		17	FALVARD	88	DM2	$J/\psi \rightarrow \phi K^+ K^-, K_S^0 K_S^0$
1698 ± 15		13	AUGUSTIN	87	DM2	$J/\psi \rightarrow \gamma \pi^+ \pi^-$
1720 ± 10 ± 10		11	BALTRUSAIT..	87	MRK3	$J/\psi \rightarrow \gamma K^+ K^-$
1755 ± 8		18	ALDE	86C	GAM2	$38 \pi^- p \rightarrow n 2\eta$
1730 ⁺² ₋₁₀		19	LONGACRE	86	RVUE	$22 \pi^- p \rightarrow n 2K_S^0$
1742 ± 15		13	WILLIAMS	84	MPSF	$200 \pi^- N \rightarrow 2K_S^0 X$
1670 ± 50			BLOOM	83	CBAL	$J/\psi \rightarrow \gamma 2\eta$
1650 ± 50			BURKE	82	MRK2	$J/\psi \rightarrow \gamma 2\rho$
1640 ± 50		20,21	EDWARDS	82D	CBAL	$J/\psi \rightarrow \gamma 2\eta$
1730 ± 10 ± 20		22	ETKIN	82C	MPS	$23 \pi^- p \rightarrow n 2K_S^0$

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

² Spin 0 favored over spin 2.

³ In the SU(3) based model with a specific interference pattern of the $f_2(1270)$, $a_2^0(1320)$, and $f_2'(1525)$ mesons incoherently added to the $f_0(1710)$ and non-resonant background.

- 4 $J^P = 0^+$, supersedes ARMSTRONG 89D.
- 5 Using CLEO-c data but not authored by the CLEO Collaboration.
- 6 From a fit to a Breit-Wigner line shape with fixed $\Gamma = 135$ MeV.
- 7 Systematic errors not estimated.
- 8 Decaying to $f_0(1370)\pi\pi$.
- 9 No J^{PC} determination.
- 10 No J^{PC} determination, width not determined.
- 11 $J^P = 2^+$.
- 12 From a fit to the 0^+ partial wave.
- 13 No J^{PC} determination.
- 14 ALDE 92D combines all the GAMS-2000 data.
- 15 $J^P = 2^+$, superseded by FRENCH 99.
- 16 From an analysis ignoring interference with $f'_2(1525)$.
- 17 From an analysis including interference with $f'_2(1525)$.
- 18 Superseded by ALDE 92D.
- 19 Uses MRK3 data. From a partial-wave analysis of data using a K-matrix formalism with 5 poles, but assuming spin 2. Fit with constrained inelasticity.
- 20 $J^P = 2^+$ preferred.
- 21 From fit neglecting nearby $f'_2(1525)$. Replaced by BLOOM 83.
- 22 Superseded by LONGACRE 86.



$f_0(1710)$ Breit-Wigner WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
150 \pm $\frac{12}{10}$	OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.		
175 \pm 23 \pm 4		LEES	21A BABR	$\eta_c(1S) \rightarrow \eta' K^+ K^-$
172 \pm 10 $\frac{+32}{-16}$	5.5k	¹ ABLIKIM	13N BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \eta \eta$
139 $\frac{+11}{-12}$ $\frac{+96}{-50}$		² UEHARA	13 BELL	$\gamma \gamma \rightarrow K_S^0 K_S^0$
100 \pm 24 $\frac{+7}{-22}$	4k	³ CHEKANOV	08 ZEUS	$e p \rightarrow K_S^0 K_S^0 X$
145 \pm 8 \pm 69		ABLIKIM	06V BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$
125 \pm 20		ABLIKIM	04E BES2	$J/\psi \rightarrow \omega K^+ K^-$
166 $\frac{+5}{-8}$ $\frac{+15}{-10}$		BAI	03G BES	$J/\psi \rightarrow \gamma K \bar{K}$
120 $\frac{+50}{-40}$		BAI	00A BES	$J/\psi \rightarrow \gamma (\pi^+ \pi^- \pi^+ \pi^-)$
105 \pm 34		⁴ FRENCH	99	300 $p p \rightarrow p_f (K^+ K^-) p_s$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
148 $\frac{+40}{-30}$		AMSLER	06 CBAR	1.64 $\bar{p} p \rightarrow K^+ K^- \pi^0$
188 \pm 13	80k	⁵ UMAN	06 E835	5.2 $\bar{p} p \rightarrow \eta \eta \pi^0$
250 \pm 30		VLADIMIRSK...	06 SPEC	40 $\pi^- p \rightarrow K_S^0 K_S^0 n$
260 \pm 50		BINON	05 GAMS	33 $\pi^- p \rightarrow \eta \eta n$
102 \pm 26		TIKHOMIROV	03 SPEC	40.0 $\pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
267 \pm 44	3651	⁶ NICHITIU	02 OBLX	0 $\bar{p} p \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
100 \pm 25		BARBERIS	99 OMEG	450 $p p \rightarrow p_s p_f K^+ K^-$
160 \pm 30		BARBERIS	99B OMEG	450 $p p \rightarrow p_s p_f \pi^+ \pi^-$
30 \pm 7	57	⁷ BARKOV	98	$\pi^- p \rightarrow K_S^0 K_S^0 n$
103 \pm 18 $\frac{+30}{-11}$		⁸ BAI	96C BES	$J/\psi \rightarrow \gamma K^+ K^-$
85 \pm 24 $\frac{+22}{-19}$		BAI	96C BES	$J/\psi \rightarrow \gamma K^+ K^-$
56 \pm 19		BALOSHIN	95 SPEC	40 $\pi^- C \rightarrow K_S^0 K_S^0 X$
160 \pm 40		⁹ BUGG	95 MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$
160 $\frac{+60}{-20}$		⁸ BUGG	95 MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$
264 \pm 25		¹⁰ ARMSTRONG	93C E760	$\bar{p} p \rightarrow \pi^0 \eta \eta \rightarrow 6 \gamma$
200 to 300		BREAKSTONE	93 SFM	$p p \rightarrow p p \pi^+ \pi^- \pi^+ \pi^-$
< 80 90% CL		¹¹ ALDE	92D GAM2	38 $\pi^- p \rightarrow \eta \eta N^*$
181 \pm 30		¹² ARMSTRONG	89D OMEG	300 $p p \rightarrow p p K^+ K^-$
104 \pm 30		¹² ARMSTRONG	89D OMEG	300 $p p \rightarrow p p K_S^0 K_S^0$
166.4 \pm 33.2		¹⁰ AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K^+ K^-, K_S^0 K_S^0$
30 \pm 20		⁸ BOLONKIN	88 SPEC	40 $\pi^- p \rightarrow K_S^0 K_S^0 n$
350 \pm 150		BOLONKIN	88 SPEC	40 $\pi^- p \rightarrow K_S^0 K_S^0 n$
148 \pm 17		¹³ FALVARD	88 DM2	$J/\psi \rightarrow \phi K^+ K^-, K_S^0 K_S^0$

184 ± 6	¹⁴ FALVARD	88	DM2	$J/\psi \rightarrow \phi K^+ K^-, K_S^0 K_S^0$
136 ± 28	¹⁰ AUGUSTIN	87	DM2	$J/\psi \rightarrow \gamma \pi^+ \pi^-$
130 ± 20	⁸ BALTRUSAIT..	87	MRK3	$J/\psi \rightarrow \gamma K^+ K^-$
122 + 74 - 15	¹⁵ LONGACRE	86	RVUE	$22 \pi^- p \rightarrow n 2 K_S^0$
57 ± 38	¹⁶ WILLIAMS	84	MPSF	$200 \pi^- N \rightarrow 2 K_S^0 X$
160 ± 80	BLOOM	83	CBAL	$J/\psi \rightarrow \gamma 2\eta$
200 ± 100	BURKE	82	MRK2	$J/\psi \rightarrow \gamma 2\rho$
220 + 100 - 70	^{17,18} EDWARDS	82D	CBAL	$J/\psi \rightarrow \gamma 2\eta$
200 + 156 - 9	¹⁹ ETKIN	82B	MPS	$23 \pi^- p \rightarrow n 2 K_S^0$

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

² Spin 0 favored over spin 2.

³ In the SU(3) based model with a specific interference pattern of the $f_2(1270)$, $a_2^0(1320)$, and $f_2'(1525)$ mesons incoherently added to the $f_0(1710)$ and non-resonant background.

⁴ $J^P = 0^+$, supersedes ARMSTRONG 89D.

⁵ Systematic errors not estimated.

⁶ Decaying to $f_0(1370) \pi \pi$.

⁷ No J^{PC} determination.

⁸ $J^P = 2^+$.

⁹ From a fit to the 0^+ partial wave.

¹⁰ No J^{PC} determination.

¹¹ ALDE 92D combines all the GAMS-2000 data.

¹² $J^P = 2^+$, (0^+ excluded).

¹³ From an analysis ignoring interference with $f_2'(1525)$.

¹⁴ From an analysis including interference with $f_2'(1525)$.

¹⁵ Uses MRK3 data. From a partial-wave analysis of data using a K-matrix formalism with 5 poles, but assuming spin 2. Fit with constrained inelasticity.


¹⁶ No J^{PC} determination.

¹⁷ $J^P = 2^+$ preferred.

¹⁸ From fit neglecting nearby $f_2'(1525)$. Replaced by BLOOM 83.

¹⁹ From an amplitude analysis of the $K_S^0 K_S^0$ system, superseded by LONGACRE 86.

WEIGHTED AVERAGE
150+12-10 (Error scaled by 1.3)



$f_0(1710)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K\bar{K}$	seen
Γ_2 $\eta\eta$	seen
Γ_3 $\eta\eta'$	not seen
Γ_4 $\pi\pi$	seen
Γ_5 $\gamma\gamma$	seen
Γ_6 $\omega\omega$	seen

$f_0(1710)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_1\Gamma_5/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
$12^{+3}_{-2} + \frac{227}{8}$		UEHARA	13	BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<480	95	ALBRECHT	90G	ARG	$\gamma\gamma \rightarrow K^+ K^-$
<110	95	¹ BEHREND	89C	CELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$
<280	95	¹ ALTHOFF	85B	TASS	$\gamma\gamma \rightarrow K\bar{K}\pi$

¹ Assuming helicity 2.

$\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_4\Gamma_5/\Gamma$
VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT	
<0.82	95	¹ BARATE	00E	ALEP	$\gamma\gamma \rightarrow \pi^+\pi^-$

¹ Assuming spin 0.

$f_0(1710)$ BRANCHING RATIOS

$\Gamma(K\bar{K})/\Gamma_{\text{total}}$					Γ_1/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
seen	1004	¹ DOBBS	15		$J/\psi \rightarrow \gamma K^+ K^-$
seen	349	¹ DOBBS	15		$\psi(2S) \rightarrow \gamma K^+ K^-$
0.36 ± 0.12		ALBALADEJO	08	RVUE	
$0.38^{+0.09}_{-0.19}$		² LONGACRE	86	MPS	$22 \pi^- p \rightarrow n 2K_S^0$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² From a partial-wave analysis of data using a K-matrix formalism with 5 poles, but assuming spin 2. Fit with constrained inelasticity.

$\Gamma(\eta\eta)/\Gamma_{\text{total}}$					Γ_2/Γ
VALUE		DOCUMENT ID	TECN		
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
0.22 ± 0.12		ALBALADEJO	08	RVUE	
$0.18^{+0.03}_{-0.13}$		¹ LONGACRE	86	RVUE	

¹ From a partial-wave analysis of data using a K-matrix formalism with 5 poles, but assuming spin 2. Fit with constrained inelasticity.

$\Gamma(\eta\eta)/\Gamma(K\bar{K})$					Γ_2/Γ_1
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
0.48 ± 0.15		BARBERIS	00E		$450 p p \rightarrow p_f \eta \eta p_S$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
$0.46^{+0.70}_{-0.38}$		¹ ANISOVICH	02D	SPEC	Combined fit
<0.02	90	² PROKOSHKIN	91	GA24	$300 \pi^- p \rightarrow \pi^- p \eta \eta$

¹ From a combined K-matrix analysis of Crystal Barrel ($0. p\bar{p} \rightarrow \pi^0 \pi^0 \pi^0, \pi^0 \eta \eta, \pi^0 \pi^0 \eta$), GAMS ($\pi p \rightarrow \pi^0 \pi^0 n, \eta \eta n, \eta \eta' n$), and BNL ($\pi p \rightarrow K\bar{K} n$) data.

² Combining results of GAM4 with those of ARMSTRONG 89D.

$\Gamma(\pi\pi)/\Gamma_{\text{total}}$					Γ_4/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
seen	381	¹ DOBBS	15		$J/\psi \rightarrow \gamma \pi^+ \pi^-$
seen	237	¹ DOBBS	15		$\psi(2S) \rightarrow \gamma \pi^+ \pi^-$
not seen		AMSLER	02	CBAR	$0.9 \bar{p} p \rightarrow \pi^0 \eta \eta, \pi^0 \pi^0 \pi^0$
$0.039^{+0.002}_{-0.024}$		² LONGACRE	86	RVUE	

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² From a partial-wave analysis of data using a K-matrix formalism with 5 poles, but assuming spin 2. Fit with constrained inelasticity.

$\Gamma(\pi\pi)/\Gamma(K\bar{K})$ Γ_4/Γ_1

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
0.23±0.05	OUR AVERAGE	Error includes scale factor of 1.2.		
0.64±0.27	±0.18	LEES	18A BABR	$\Upsilon(1S) \rightarrow \gamma\pi^+\pi^-$, γK^+K^-
0.41 ^{+0.11} _{-0.17}		ABLIKIM	06V BES2	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
0.2 ±0.024±0.036		BARBERIS	99D OMEG 450	$pp \rightarrow K^+K^-, \pi^+\pi^-$
0.39±0.14		ARMSTRONG	91 OMEG 300	$pp \rightarrow pp\pi\pi, ppK\bar{K}$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.32±0.14		ALBALADEJO	08 RVUE	
< 0.11	95	¹ ABLIKIM	04E BES2	$J/\psi \rightarrow \omega K^+K^-$
5.8 ^{+9.1} _{-5.5}		² ANISOVICH	02D SPEC	Combined fit

¹ Using data from ABLIKIM 04A.

² From a combined K-matrix analysis of Crystal Barrel (0. $p\bar{p} \rightarrow \pi^0\pi^0\pi^0, \pi^0\eta\eta, \pi^0\pi^0\eta$), GAMS ($\pi p \rightarrow \pi^0\pi^0 n, \eta\eta n, \eta\eta' n$), and BNL ($\pi p \rightarrow K\bar{K}n$) data.

$\Gamma(\eta\eta')/\Gamma(\pi\pi)$ Γ_3/Γ_4

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.87 × 10⁻³	90	¹ ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma\eta\eta'$

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta\eta'$ P-wave.

$\Gamma(\omega\omega)/\Gamma_{total}$ Γ_6/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	180	ABLIKIM	06H BES	$J/\psi \rightarrow \gamma\omega\omega$

$f_0(1710)$ REFERENCES

ABLIKIM	22AS	PR D106 072012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
Also		PR D107 079901 (errat.)	M. Ablikim <i>et al.</i>	(BESIII Collab.)
RODAS	22	EPJ C82 80	A. Rodas <i>et al.</i>	(JPAC Collab.)
LEES	21A	PR D104 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
SARANTSEV	21	PL B816 136227	A.V. Sarantsev <i>et al.</i>	(BONN, PNPI)
ALBRECHT	20	EPJ C80 453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)
ABLIKIM	18AA	PR D98 072003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	18A	PR D97 112006	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	15AE	PR D92 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)
ABLIKIM	13N	PR D87 092009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
UEHARA	13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)
ALBALADEJO	08	PRL 101 252002	M. Albaladejo, J.A. Oller	
CHEKANOV	08	PRL 101 112003	S. Chekanov <i>et al.</i>	(ZEUS Collab.)
ABLIKIM	06H	PR D73 112007	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06V	PL B642 441	M. Ablikim <i>et al.</i>	(BES Collab.)
AMSLER	06	PL B639 165	C. AMSler <i>et al.</i>	(Crystal Barrel Collab.)
UMAN	06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)
VLADIMIRSK...	06	PAN 69 493	V.V. Vladimirovsky <i>et al.</i>	(ITEP, Moscow)
		Translated from YAF 69 515.		
BINON	05	PAN 68 960	F. Binon <i>et al.</i>	
		Translated from YAF 68 998.		
ABLIKIM	04A	PL B598 149	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	04E	PL B603 138	M. Ablikim <i>et al.</i>	(BES Collab.)
ANISOVICH	03	EPJ A16 229	V.V. Anisovich <i>et al.</i>	
BAI	03G	PR D68 052003	J.Z. Bai <i>et al.</i>	(BES Collab.)
TIKHOMIROV	03	PAN 66 828	G.D. Tikhomirov <i>et al.</i>	
		Translated from YAF 66 860.		

AMSLER	02	EPJ C23 29	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
ANISOVICH	02D	PAN 65 1545	V.V. Anisovich <i>et al.</i>	
		Translated from YAF 65 1583.		
NICHITIU	02	PL B545 261	F. Nichitiu <i>et al.</i>	(OBELIX Collab.)
BAI	00A	PL B472 207	J.Z. Bai <i>et al.</i>	(BES Collab.)
BARATE	00E	PL B472 189	R. Barate <i>et al.</i>	(ALEPH Collab.)
BARBERIS	00E	PL B479 59	D. Barberis <i>et al.</i>	(WA 102 Collab.)
ANISOVICH	99B	PL B449 154	A.V. Anisovich <i>et al.</i>	
BARBERIS	99	PL B453 305	D. Barberis <i>et al.</i>	(Omega Expt.)
BARBERIS	99B	PL B453 316	D. Barberis <i>et al.</i>	(Omega Expt.)
BARBERIS	99D	PL B462 462	D. Barberis <i>et al.</i>	(Omega Expt.)
FRENCH	99	PL B460 213	B. French <i>et al.</i>	(WA76 Collab.)
ANISOVICH	98B	SPU 41 419	V.V. Anisovich <i>et al.</i>	
		Translated from UFN 168 481.		
BAI	98H	PRL 81 1179	J.Z. Bai <i>et al.</i>	(BES Collab.)
BARKOV	98	JETPL 68 764	B.P. Barkov <i>et al.</i>	
ABREU	96C	PL B379 309	P. Abreu <i>et al.</i>	(DELPHI Collab.)
BAI	96C	PRL 77 3959	J.Z. Bai <i>et al.</i>	(BES Collab.)
BALOSHIN	95	PAN 58 46	O.N. Baloshin <i>et al.</i>	(ITEP)
		Translated from YAF 58 50.		
BUGG	95	PL B353 378	D.V. Bugg <i>et al.</i>	(LOQM, PNPI, WASH)
ARMSTRONG	93C	PL B307 394	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
BREAKSTONE	93	ZPHY C58 251	A.M. Breakstone <i>et al.</i>	(IOWA, CERN, DORT+)
ALDE	92D	PL B284 457	D.M. Alde <i>et al.</i>	(GAM2 Collab.)
Also		SJNP 54 451	D.M. Alde <i>et al.</i>	(GAM2 Collab.)
		Translated from YAF 54 745.		
ARMSTRONG	91	ZPHY C51 351	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)
PROKOSHKIN	91	SPD 36 155	Y.D. Prokoshkin	(GAM2 and GAM4 Collab.)
		Translated from DANS 316 900.		
ALBRECHT	90G	ZPHY C48 183	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ARMSTRONG	89D	PL B227 186	T.A. Armstrong, M. Benayoun	(ATHU, BARI, BIRM+)
BEHREND	89C	ZPHY C43 91	H.J. Behrend <i>et al.</i>	(CELLO Collab.)
AUGUSTIN	88	PRL 60 2238	J.E. Augustin <i>et al.</i>	(DM2 Collab.)
BOLONKIN	88	NP B309 426	B.V. Bolonkin <i>et al.</i>	(ITEP, SERP)
FALVARD	88	PR D38 2706	A. Falvard <i>et al.</i>	(CLER, FRAS, LALO+)
AUGUSTIN	87	ZPHY C36 369	J.E. Augustin <i>et al.</i>	(LALO, CLER, FRAS+)
BALTRUSAITIS...	87	PR D35 2077	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
ALDE	86C	PL B182 105	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP)
LONGACRE	86	PL B177 223	R.S. Longacre <i>et al.</i>	(BNL, BRAN, CUNY+)
ALTHOFF	85B	ZPHY C29 189	M. Althoff <i>et al.</i>	(TASSO Collab.)
BINON	84C	NC 80A 363	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)
WILLIAMS	84	PR D30 877	E.G.H. Williams <i>et al.</i>	(VAND, NDAM, TUFTS+)
BINON	83	NC 78A 313	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)
BLOOM	83	ARNS 33 143	E.D. Bloom, C. Peck	(SLAC, CIT)
BURKE	82	PRL 49 632	D.L. Burke <i>et al.</i>	(LBL, SLAC)
EDWARDS	82D	PRL 48 458	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
ETKIN	82B	PR D25 1786	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)
ETKIN	82C	PR D25 2446	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)
HYAMS	75	NP B100 205	B.D. Hyams <i>et al.</i>	(CERN, MPIM)