

$f_2(2150)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

OMITTED FROM SUMMARY TABLE

This entry was previously called T_0 .

$f_2(2150)$ MASS

$f_2(2150)$ MASS, COMBINED MODES (MeV)

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

2157 ± 12 OUR AVERAGE Includes data from the datablock that follows this one.

• • • We do not use the following data for averages, fits, limits, etc. • • •

2170 ± 6	80k	¹ UMAN	06 E835	5.2 $\bar{p}p \rightarrow \eta\eta\pi^0$
----------	-----	-------------------	---------	--

¹Statistical error only.

$\eta\eta$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

The data in this block is included in the average printed for a previous datablock.

2157 ± 12 OUR AVERAGE

2151 ± 16	BARBERIS	00E	450 $pp \rightarrow p_f\eta\eta p_s$
2175 ± 20	PROKOSHKIN	95D GAM4	300 $\pi^- N \rightarrow \pi^- N 2\eta$, 450 $pp \rightarrow pp 2\eta$
2130 ± 35	SINGOVSKI	94 GAM4	450 $pp \rightarrow pp 2\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2140 ± 30	² ABELE	99B CBAR	1.94 $\bar{p}p \rightarrow \pi^0\eta\eta$
2104 ± 20	³ ARMSTRONG	93C E760	$\bar{p}p \rightarrow \pi^0\eta\eta \rightarrow 6\gamma$

²Spin not determined.

³No J^{PC} determination.

$\eta\pi\pi$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

2135 ± 20 ± 45	⁴ ADOMEIT	96 CBAR	0	1.94 $\bar{p}p \rightarrow \eta 3\pi^0$
----------------	----------------------	---------	---	---

⁴ANISOVICH 00E recommends to withdraw ADOMEIT 96 that assumed a single $J^P = 2^+$ resonance.

$\bar{p}p \rightarrow \pi\pi$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 2090	⁵ OAKDEN	94 RVUE	0.36–1.55 $\bar{p}p \rightarrow \pi\pi$
~ 2120	⁶ OAKDEN	94 RVUE	0.36–1.55 $\bar{p}p \rightarrow \pi\pi$
~ 2170	⁷ MARTIN	80B RVUE	
~ 2150	⁷ MARTIN	80C RVUE	
~ 2150	⁸ DULUDE	78B OSPK	1–2 $\bar{p}p \rightarrow \pi^0\pi^0$

⁵ OAKDEN 94 makes an amplitude analysis of LEAR data on $\bar{p}p \rightarrow \pi\pi$ using a method based on Barrelet zeros. This is solution A. The amplitude analysis of HASAN 94 includes earlier data as well, and assume that the data can be parametrized in terms of towers of nearly degenerate resonances on the leading Regge trajectory. See also KLOET 96 and MARTIN 97 who make related analyses.

⁶ From solution B of amplitude analysis of data on $\bar{p}p \rightarrow \pi\pi$.

⁷ $I(J^P) = 0(2^+)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^-\pi^+$ and $\pi^0\pi^0$.

⁸ $I^G(J^P) = 0^+(2^+)$ from partial-wave amplitude analysis.

S-CHANNEL $\bar{p}p$, $\bar{N}N$ or $\bar{K}K$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2139^{+8}_{-9}	⁹ EVANGELIS...	97	SPEC	0.6-2.4 $\bar{p}p \rightarrow K_S^0 K_S^0$
~ 2190	⁹ CUTTS	78B	CNTR	0.97-3 $\bar{p}p \rightarrow \bar{N}N$
2155 ± 15	^{9,10} COUPLAND	77	CNTR 0	0.7-2.4 $\bar{p}p \rightarrow \bar{p}p$
2193 ± 2	^{9,11} ALSPECTOR	73	CNTR	$\bar{p}p$ S channel

⁹ Isospins 0 and 1 not separated.

¹⁰ From a fit to the total elastic cross section.

¹¹ Referred to as T or T region by ALSPECTOR 73.

$K\bar{K}$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
2200 ± 13	VLADIMIRSK...06	SPEC	40 $\pi^- p \rightarrow K_S^0 K_S^0 n$
2150 ± 20	ABLIKIM	04E	BES2 $J/\psi \rightarrow \omega K^+ K^-$
2130 ± 35	BARBERIS	99	OMEG 450 $p\bar{p} \rightarrow p_S p_f K^+ K^-$

$f_2(2150)$ WIDTH


$f_2(2150)$ WIDTH, COMBINED MODES (MeV)

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
152 ± 30 OUR AVERAGE	Includes data from the datablock that follows this one. Error includes scale factor of 1.4. See the ideogram below.			

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

182 ± 11 80k ¹² UMAN 06 E835 5.2 $\bar{p}p \rightarrow \eta\eta\pi^0$

¹² Statistical error only.



$\eta\eta$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

The data in this block is included in the average printed for a previous datablock.

152 ± 30 OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below.


280 ± 70	BARBERIS	00E	450 $pp \rightarrow p_f \eta \eta p_s$
150 ± 35	PROKOSHKIN	95D	GAM4 300 $\pi^- N \rightarrow \pi^- N 2\eta$, 450 $pp \rightarrow pp 2\eta$
130 ± 30	SINGOVSKI	94	GAM4 450 $pp \rightarrow pp 2\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

310 ± 50	¹³ ABELE	99B	CBAR 1.94 $\bar{p}p \rightarrow \pi^0 \eta \eta$
203 ± 10	¹⁴ ARMSTRONG	93C	E760 $\bar{p}p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$

¹³ Spin not determined.

¹⁴ No J^{PC} determination.



$\eta\pi\pi$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
250±25±45	¹⁵ ADOMEIT 96	CBAR	0	1.94 $\bar{p}p \rightarrow \eta 3\pi^0$
	¹⁵ ANISOVICH 00E			recommends to withdraw ADOMEIT 96 that assumed a single $J^P = 2^+$ resonance.

$\bar{p}p \rightarrow \pi\pi$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
250 OUR ESTIMATE				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
~ 70	¹⁶ OAKDEN 94	RVUE	0.36–1.55	$\bar{p}p \rightarrow \pi\pi$
~ 250	¹⁷ MARTIN 80B	RVUE		
~ 250	¹⁷ MARTIN 80C	RVUE		
~ 250	¹⁸ DULUDE 78B	OSPK	1–2	$\bar{p}p \rightarrow \pi^0\pi^0$
	¹⁶			See however KLOET 96 who fit $\pi^+\pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.
	¹⁷			$I(J^P) = 0(2^+)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^-\pi^+$ and $\pi^0\pi^0$.
	¹⁸			$I^G(J^P) = 0^+(2^+)$ from partial-wave amplitude analysis.

S-CHANNEL $\bar{p}p$, $\bar{N}N$ or $\bar{K}K$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
56 ⁺³¹ ₋₁₆	¹⁹ EVANGELIS... 97	SPEC		0.6–2.4 $\bar{p}p \rightarrow K_S^0 K_S^0$
135±75	^{20,21} COUPLAND 77	CNTR	0	0.7–2.4 $\bar{p}p \rightarrow \bar{p}p$
98±8	²¹ ALSPECTOR 73	CNTR		$\bar{p}p$ S channel

- ¹⁹ Isospin 0 and 2 not separated.
- ²⁰ From a fit to the total elastic cross section.
- ²¹ Isospins 0 and 1 not separated.

$K\bar{K}$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
91 ± 62	VLADIMIRSK...06	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
150 ± 30	ABLIKIM	04E BES2	$J/\psi \rightarrow \omega K^+ K^-$
270 ± 50	BARBERIS	99 OMEG	$450 pp \rightarrow p_S p_f K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$f_2(2150)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad \pi\pi$	
$\Gamma_2 \quad \eta\eta$	seen
$\Gamma_3 \quad K\bar{K}$	seen
$\Gamma_4 \quad f_2(1270)\eta$	seen
$\Gamma_5 \quad a_2(1320)\pi$	seen
$\Gamma_6 \quad p\bar{p}$	seen

$f_2(2150)$ BRANCHING RATIOS

$\Gamma(K\bar{K})/\Gamma(\eta\eta)$					Γ_3/Γ_2
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
1.28 ± 0.23		BARBERIS	00E	$450 pp \rightarrow p_f \eta \eta p_S$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.1	95	²² PROKOSHKIN 95D	GAM4	$300 \pi^- N \rightarrow \pi^- N 2\eta,$ $450 pp \rightarrow p p 2\eta$
--------	----	------------------------------	------	--

²² Using data from ARMSTRONG 89D.

$\Gamma(\pi\pi)/\Gamma(\eta\eta)$					Γ_1/Γ_2
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.33	95	²³ PROKOSHKIN 95D	GAM4	$300 \pi^- N \rightarrow \pi^- N 2\eta,$ $450 pp \rightarrow p p 2\eta$
---------	----	------------------------------	------	--

²³ Derived from a $\pi^0\pi^0/\eta\eta$ limit.

$\Gamma(f_2(1270)\eta)/\Gamma(a_2(1320)\pi)$					Γ_4/Γ_5
VALUE		DOCUMENT ID	TECN	COMMENT	

0.79 ± 0.11		²⁴ ADOMEIT	96	CBAR	$1.94 \bar{p}p \rightarrow \eta 3\pi^0$
-----------------------------------	--	-----------------------	----	------	---

²⁴ Using $B(a_2(1320) \rightarrow \eta\pi) = 0.145$

$\Gamma(p\bar{p})/\Gamma_{total}$					Γ_6/Γ
VALUE	EVT%	DOCUMENT ID	TECN	COMMENT	

seen	73	ALEXANDER	10	CLEO	$\psi(2S) \rightarrow \gamma p\bar{p}$
-------------	----	-----------	----	------	--

$f_2(2150)$ REFERENCES

ALEXANDER	10	PR D82 092002	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
UMAN	06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)
VLADIMIRSK...	06	PAN 69 493	V.V. Vladimisky <i>et al.</i>	(ITEP, Moscow)
ABLIKIM	04E	Translated from YAF 69 515. PL B603 138	M. Ablikim <i>et al.</i>	(BES Collab.)
ANISOVICH	00E	PL B477 19	A.V. Anisovich <i>et al.</i>	
BARBERIS	00E	PL B479 59	D. Barberis <i>et al.</i>	(WA 102 Collab.)
ABELE	99B	EPJ C8 67	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
BARBERIS	99	PL B453 305	D. Barberis <i>et al.</i>	(Omega Expt.)
EVANGELIS...	97	PR D56 3803	C. Evangelista <i>et al.</i>	(LEAR Collab.)
MARTIN	97	PR C56 1114	B.R. Martin, G.C. Oades	(LOUC, AARH)
ADOMEIT	96	ZPHY C71 227	J. Adomeit <i>et al.</i>	(Crystal Barrel Collab.)
KLOET	96	PR D53 6120	W.M. Kloet, F. Myhrer	(RUTG, NORD)
PROKOSHKIN	95D	PD 40 495	Y.D. Prokoshkin	(SERP) IGJPC
HASAN	94	Translated from DANS 344 469. PL B334 215	A. Hasan, D.V. Bugg	(LOQM)
OAKDEN	94	NP A574 731	M.N. Oakden, M.R. Pennington	(DURH)
SINGOVSKI	94	NC A107 1911	A.V. Singovsky	(SERP)
ARMSTRONG	93C	PL B307 394	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
ARMSTRONG	89D	PL B227 186	T.A. Armstrong, M. Benayoun	(ATHU, BARI, BIRM+)
MARTIN	80B	NP B176 355	B.R. Martin, D. Morgan	(LOUC, RHEL) JP
MARTIN	80C	NP B169 216	A.D. Martin, M.R. Pennington	(DURH) JP
CUTTS	78B	PR D17 16	D. Cutts <i>et al.</i>	(STON, WISC)
DULUDE	78B	PL 79B 335	R.S. Dulude <i>et al.</i>	(BROW, MIT, BARI) JP
COUPLAND	77	PL 71B 460	M. Coupland <i>et al.</i>	(LOQM, RHEL)
ALSPECTOR	73	PRL 30 511	J. Alspector <i>et al.</i>	(RUTG, UPNJ)
