

$\Lambda(1690) \ 3/2^-$ $I(J^P) = 0(\frac{3}{2}^-)$ Status: * * * *

The measurements of the mass, width, and elasticity published before 1974 are now obsolete and have been omitted. They were last listed in our 1982 edition Physics Letters **111B** 1 (1982).

$\Lambda(1690)$ POLE POSITION

REAL PART

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------------	--------------------	-------------	----------------

1680 to 1700 (≈ 1690) OUR ESTIMATE

1683 \pm 3	SARANTSEV	19	DPWA $\bar{K}N$ multichannel
1697 $^{+6}_{-6}$	¹ KAMANO	15	DPWA $\bar{K}N$ multichannel

• • • We do not use the following data for averages, fits, limits, etc. • • •

1689	ZHANG	13A	DPWA $\bar{K}N$ multichannel
------	-------	-----	------------------------------

¹From the preferred solution A in KAMANO 15.

–2×IMAGINARY PART

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------------	--------------------	-------------	----------------

60 to 80 (≈ 70) OUR ESTIMATE

72 \pm 5	SARANTSEV	19	DPWA $\bar{K}N$ multichannel
65 \pm 14	¹ KAMANO	15	DPWA $\bar{K}N$ multichannel

• • • We do not use the following data for averages, fits, limits, etc. • • •

53	ZHANG	13A	DPWA $\bar{K}N$ multichannel
----	-------	-----	------------------------------

¹From the preferred solution A in KAMANO 15.

$\Lambda(1690)$ POLE RESIDUES

The normalized residue is the residue divided by $\Gamma_{pole}/2$.

Normalized residue in $N\bar{K} \rightarrow \Lambda(1690) \rightarrow N\bar{K}$

<u>MODULUS</u>	<u>PHASE ($^\circ$)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
----------------	------------------------------------	--------------------	-------------	----------------

0.24 \pm0.05	–28 \pm 5	SARANTSEV	19	DPWA $\bar{K}N$ multichannel
----------------------------------	-------------------------------	-----------	----	------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.251	3	¹ KAMANO	15	DPWA Multichannel
-------	---	---------------------	----	-------------------

¹From the preferred solution A in KAMANO 15.

Normalized residue in $N\bar{K} \rightarrow \Lambda(1690) \rightarrow \Sigma\pi$

<u>MODULUS</u>	<u>PHASE ($^\circ$)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
----------------	------------------------------------	--------------------	-------------	----------------

0.35 \pm0.07	175 \pm 6	SARANTSEV	19	DPWA $\bar{K}N$ multichannel
----------------------------------	-------------------------------	-----------	----	------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.315	–173	¹ KAMANO	15	DPWA $\bar{K}N$ multichannel
-------	------	---------------------	----	------------------------------

¹From the preferred solution A in KAMANO 15.

Normalized residue in $N\bar{K} \rightarrow \Lambda(1690) \rightarrow \Lambda\eta$

<u>MODULUS</u>	<u>PHASE ($^{\circ}$)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.05 ± 0.02	88 ± 8	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel
0.00567	81	¹ KAMANO 15	DPWA	Multichannel

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹From the preferred solution A in KAMANO 15.

Normalized residue in $N\bar{K} \rightarrow \Lambda(1690) \rightarrow \Lambda\sigma$

<u>MODULUS</u>	<u>PHASE ($^{\circ}$)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.08 ± 0.02	-10 ± 6	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel

Normalized residue in $N\bar{K} \rightarrow \Lambda(1690) \rightarrow \Sigma(1385)\pi$, S-wave

<u>MODULUS</u>	<u>PHASE ($^{\circ}$)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.11 ± 0.06	170 ± 70	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel
0.134	168	¹ KAMANO 15	DPWA	$\bar{K}N$ multichannel

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹From the preferred solution A in KAMANO 15.

Normalized residue in $N\bar{K} \rightarrow \Lambda(1690) \rightarrow \Sigma(1385)\pi$, D-wave

<u>MODULUS</u>	<u>PHASE ($^{\circ}$)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.06 ± 0.04	164 ± 15	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel
0.319	-22	¹ KAMANO 15	DPWA	$\bar{K}N$ multichannel

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹From the preferred solution A in KAMANO 15.

Normalized residue in $N\bar{K} \rightarrow \Lambda(1690) \rightarrow N\bar{K}^*(892)$, S-wave

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.05 ± 0.04	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel

Normalized residue in $N\bar{K} \rightarrow \Lambda(1690) \rightarrow N\bar{K}^*(892)$, D-wave

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.18 ± 0.05 @ -110 ± 45	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel

$\Lambda(1690)$ MASS

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1685 to 1695 (≈ 1690) OUR ESTIMATE			
1689 ± 3	SARANTSEV 19	DPWA	$\bar{K}N$ multichannel
1691 ± 3	ZHANG 13A	DPWA	$\bar{K}N$ multichannel
1695.7 ± 2.6	KOISO 85	DPWA	$K^- p \rightarrow \Sigma\pi$
1690 ± 5	GOPAL 80	DPWA	$\bar{K}N \rightarrow \bar{K}N$
1692 ± 5	ALSTON-... 78	DPWA	$\bar{K}N \rightarrow \bar{K}N$
1690 ± 3	HEPP 76B	DPWA	$K^- N \rightarrow \Sigma\pi$
1689 ± 1	KANE 74	DPWA	$K^- p \rightarrow \Sigma\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1690 ± 5	GOPAL 77	DPWA	$\bar{K}N$ multichannel
1687 or 1689	¹ MARTIN 77	DPWA	$\bar{K}N$ multichannel
1692 ± 4	CARROLL 76	DPWA	Isospin-0 total σ

¹The two MARTIN 77 values are from a T-matrix pole and from a Breit-Wigner fit. Another D_{03} Λ at 1966 MeV is also suggested by MARTIN 77, but is very uncertain.

$\Lambda(1690)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
60 to 80 (≈ 70) OUR ESTIMATE			
75 \pm 5	SARANTSEV	19	DPWA $\overline{K}N$ multichannel
54 \pm 5	ZHANG	13A	DPWA $\overline{K}N$ multichannel
67.2 \pm 5.6	KOISO	85	DPWA $K^- p \rightarrow \Sigma \pi$
61 \pm 5	GOPAL	80	DPWA $\overline{K}N \rightarrow \overline{K}N$
64 \pm 10	ALSTON-...	78	DPWA $\overline{K}N \rightarrow \overline{K}N$
82 \pm 8	HEPP	76B	DPWA $K^- N \rightarrow \Sigma \pi$
60 \pm 4	KANE	74	DPWA $K^- p \rightarrow \Sigma \pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
60 \pm 5	GOPAL	77	DPWA $\overline{K}N$ multichannel
62 or 62	¹ MARTIN	77	DPWA $\overline{K}N$ multichannel
38	CARROLL	76	DPWA Isospin-0 total σ

¹The two MARTIN 77 values are from a T-matrix pole and from a Breit-Wigner fit. Another D_{03} Λ at 1966 MeV is also suggested by MARTIN 77, but is very uncertain.

$\Lambda(1690)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\overline{K}$	20–30 %
Γ_2 $\Sigma \pi$	20–40 %
Γ_3 $\Lambda \sigma$	(5.0 \pm 2.0) %
Γ_4 $\Lambda \pi \pi$	\sim 25 %
Γ_5 $\Sigma \pi \pi$	\sim 20 %
Γ_6 $\Lambda \eta$	
Γ_7 $\Sigma(1385)\pi$, S-wave	(9 \pm 5) %
Γ_8 $\Sigma(1385)\pi$, D-wave	(3.0 \pm 2.0) %
Γ_9 $N\overline{K}^*(892)$, S=1/2, D-wave	
Γ_{10} $N\overline{K}^*(892)$, S=3/2, S-wave	
Γ_{11} $N\overline{K}^*(892)$, S=3/2, D-wave	

$\Lambda(1690)$ BRANCHING RATIOS

$\Gamma(N\overline{K})/\Gamma_{\text{total}}$				Γ_1/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
0.20 to 0.28 OUR ESTIMATE				
0.23 \pm 0.05	SARANTSEV	19	DPWA $\overline{K}N$ multichannel	
0.25 \pm 0.04	ZHANG	13A	DPWA $\overline{K}N$ multichannel	
0.23 \pm 0.03	GOPAL	80	DPWA $\overline{K}N \rightarrow \overline{K}N$	
0.22 \pm 0.03	ALSTON-...	78	DPWA $\overline{K}N \rightarrow \overline{K}N$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.239	¹ KAMANO	15	DPWA	$\bar{K}N$ multichannel
0.24 ± 0.03	GOPAL	77	DPWA	See GOPAL 80
0.28 or 0.26	² MARTIN	77	DPWA	$\bar{K}N$ multichannel

¹ From the preferred solution A in KAMANO 15.

² The two MARTIN 77 values are from a T-matrix pole and from a Breit-Wigner fit. Another $D_{03} \Lambda$ at 1966 MeV is also suggested by MARTIN 77, but is very uncertain.

$\Gamma(\Sigma\pi)/\Gamma_{\text{total}}$ Γ_2/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.50 ± 0.10	SARANTSEV	19	DPWA $\bar{K}N$ multichannel

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.387	¹ KAMANO	15	DPWA	$\bar{K}N$ multichannel
-------	---------------------	----	------	-------------------------

¹ From the preferred solution A in KAMANO 15.

$\Gamma(\Lambda\eta)/\Gamma_{\text{total}}$ Γ_6/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
~ 0.01	SARANTSEV	19	DPWA $\bar{K}N$ multichannel

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	¹ KAMANO	15	DPWA	Multichannel
----------	---------------------	----	------	--------------

¹ From the preferred solution A in KAMANO 15.

$\Gamma(\Lambda\sigma)/\Gamma_{\text{total}}$ Γ_3/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.05 ± 0.02	SARANTSEV	19	DPWA $\bar{K}N$ multichannel

$\Gamma(\Sigma(1385)\pi, S\text{-wave})/\Gamma_{\text{total}}$ Γ_7/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.09 ± 0.05	SARANTSEV	19	DPWA $\bar{K}N$ multichannel

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.062	¹ KAMANO	15	DPWA	$\bar{K}N$ multichannel
-------	---------------------	----	------	-------------------------

¹ From the preferred solution A in KAMANO 15.

$\Gamma(\Sigma(1385)\pi, D\text{-wave})/\Gamma_{\text{total}}$ Γ_8/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.03 ± 0.02	SARANTSEV	19	DPWA $\bar{K}N$ multichannel

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.308	¹ KAMANO	15	DPWA	$\bar{K}N$ multichannel
-------	---------------------	----	------	-------------------------

¹ From the preferred solution A in KAMANO 15.

$\Gamma(N\bar{K}^*(892), S=1/2, D\text{-wave})/\Gamma_{\text{total}}$ Γ_9/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
not seen	SARANTSEV	19	DPWA $\bar{K}N$ multichannel
not seen	¹ KAMANO	15	DPWA $\bar{K}N$ multichannel

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	SARANTSEV	19	DPWA	$\bar{K}N$ multichannel
not seen	¹ KAMANO	15	DPWA	$\bar{K}N$ multichannel

¹ From the preferred solution A in KAMANO 15.

$\Gamma(N\bar{K}^*(892), S=3/2, S\text{-wave})/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.003 KAMANO 15 DPWA Multichannel

$\Gamma(N\bar{K}^*(892), S=3/2, D\text{-wave})/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen ¹KAMANO 15 DPWA Multichannel

¹ From the preferred solution A in KAMANO 15.

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Lambda(1690) \rightarrow \Sigma\pi$ $(\Gamma_1\Gamma_2)^{1/2}/\Gamma$

VALUE DOCUMENT ID TECN COMMENT

-0.27±0.03 ZHANG 13A DPWA Multichannel

-0.34±0.02 KOISO 85 DPWA $K^-p \rightarrow \Sigma\pi$

-0.25±0.03 GOPAL 77 DPWA $\bar{K}N$ multichannel

-0.29±0.03 HEPP 76B DPWA $K^-N \rightarrow \Sigma\pi$

-0.28±0.03 LONDON 75 HLBC $K^-p \rightarrow \Sigma^0\pi^0$

-0.28±0.02 KANE 74 DPWA $K^-p \rightarrow \Sigma\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

-0.30 or -0.28 ¹MARTIN 77 DPWA $\bar{K}N$ multichannel

¹ The two MARTIN 77 values are from a T-matrix pole and from a Breit-Wigner fit. Another $D_{03} \Lambda$ at 1966 MeV is also suggested by MARTIN 77, but is very uncertain.

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Lambda(1690) \rightarrow \Lambda\pi\pi$ $(\Gamma_1\Gamma_4)^{1/2}/\Gamma$

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.25±0.02 ¹BARTLEY 68 HDBC $K^-p \rightarrow \Lambda\pi\pi$

¹ BARTLEY 68 uses only cross-section data. The enhancement is not seen by PREVOST 71.

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Lambda(1690) \rightarrow \Sigma\pi\pi$ $(\Gamma_1\Gamma_5)^{1/2}/\Gamma$

VALUE DOCUMENT ID TECN COMMENT

0.21 ARMENTEROS68C HDBC $K^-N \rightarrow \Sigma\pi\pi$

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Lambda(1690) \rightarrow \Lambda\eta$ $(\Gamma_1\Gamma_6)^{1/2}/\Gamma$

VALUE DOCUMENT ID TECN COMMENT

0.00±0.03 BAXTER 73 DPWA $K^-p \rightarrow$ neutrals

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Lambda(1690) \rightarrow \Sigma(1385)\pi, S\text{-wave}$ $(\Gamma_1\Gamma_7)^{1/2}/\Gamma$

VALUE DOCUMENT ID TECN COMMENT

-0.28±0.06 ZHANG 13A DPWA Multichannel

+0.27±0.04 PREVOST 74 DPWA $K^-N \rightarrow \Sigma(1385)\pi$

$\Lambda(1690)$ REFERENCES

SARANTSEV	19	EPJ A55 180	A.V. Sarantsev <i>et al.</i>	(BONN, PNPI)
KAMANO	15	PR C92 025205	H. Kamano <i>et al.</i>	(ANL, OSAK)
ZHANG	13A	PR C88 035205	H. Zhang <i>et al.</i>	(KSU)
KOISO	85	NP A433 619	H. Koiso <i>et al.</i>	(TOKY, MASA)
PDG	82	PL 111B 1	M. Roos <i>et al.</i>	(HELS, CIT, CERN)
GOPAL	80	Toronto Conf. 159	G.P. Gopal	(RHEL) IJP
ALSTON-...	78	PR D18 182	M. Alston-Garnjost <i>et al.</i>	(LBL, MTHO+) IJP
Also		PRL 38 1007	M. Alston-Garnjost <i>et al.</i>	(LBL, MTHO+) IJP
GOPAL	77	NP B119 362	G.P. Gopal <i>et al.</i>	(LOIC, RHEL) IJP
MARTIN	77	NP B127 349	B.R. Martin, M.K. Pidcock, R.G. Moorhouse	(LOUC+) IJP
Also		NP B126 266	B.R. Martin, M.K. Pidcock	(LOUC)
Also		NP B126 285	B.R. Martin, M.K. Pidcock	(LOUC) IJP
CARROLL	76	PRL 37 806	A.S. Carroll <i>et al.</i>	(BNL) I
HEPP	76B	PL 65B 487	V. Hepp <i>et al.</i>	(CERN, HEIDH, MPIM) IJP
LONDON	75	NP B85 289	G.W. London <i>et al.</i>	(BNL, CERN, EPOL+)
KANE	74	LBL-2452	D.F. Kane	(LBL) IJP
PREVOST	74	NP B69 246	J. Prevost <i>et al.</i>	(SACL, CERN, HEID)
BAXTER	73	NP B67 125	D.F. Baxter <i>et al.</i>	(OXF) IJP
PREVOST	71	Amsterdam Conf.	J. Prevost	(CERN, HEID, SACL)
ARMENTEROS	68C	NP B8 216	R. Armenteros <i>et al.</i>	(CERN, HEID, SACL) I
BARTLEY	68	PRL 21 1111	J.H. Bartley <i>et al.</i>	(TUFTS, FSU, BRAN) I
