$I(J^P) = 0(\frac{5}{2})$ Status: ****

For results published before 1973 (they are now obsolete), see our 1982 edition Physics Letters **111B** 1 (1982).

The best evidence for this resonance is in the $\Sigma \pi$ channel.

A(1830) POLE POSITION

REAL PAF	T						
VALUE (MeV)		DOCUMENT ID		TECN	COMMENT		
1800 to 1860 (≈ 1830) OUR ESTIMATE							
$1819.5\pm$ 3.0	1	SARANTSEV	19	DPWA	$\overline{K}N$ multichannel		
$\begin{array}{rrr} 1899 & +35 \\ -37 \end{array}$		1 KAMANO	15	DPWA	Multichannel		
• • • We do	not use the following	data for averages	, fits,	limits, e	tc. ● ● ●		
1766 $^{+37}_{-34}$		² KAMANO	15	DPWA	Multichannel		
1809		ZHANG	13A	DPWA	Multichannel		
¹ The pref	erred solution A in K	AMANO 15 repoi	rts tw	o poles.	This entry is from the		

preferred solution A.

 $^2\,{\rm From}$ the preferred solution A in KAMANO 15. Not seen in solution B.

-2×IMAGINARY PART VALUE (MeV) DOCUMENT ID TECN COMMENT 50 to 80 (\approx 65) OUR ESTIMATE SARANTSEV DPWA $\overline{K}N$ multichannel 62± 5 19 80^{+100}_{-34} ¹ KAMANO 15 DPWA Multichannel • • • We do not use the following data for averages, fits, limits, etc. • • • 212^{+}_{-} $^{94}_{62}$ ² KAMANO 15 DPWA Multichannel ZHANG 109 13A DPWA Multichannel ¹The preferred solution A in KAMANO 15 reports two poles. This entry is from the preferred solution A.

² From the preferred solution A in KAMANO 15. Not seen in solution B.

A(1830) POLE RESIDUES

The normalized residue is the residue divided by $\Gamma_{pole}/2$.

Normalized residue in $N\overline{K} \rightarrow \Lambda(1830) \rightarrow N\overline{K}$

MODULUS	PHASE (°)	DOCUMENT ID		TECN	COMMENT
0.055 ±0.010	20 ± 14	SARANTSEV	19	DPWA	K N multichannel
• • • We do not	t use the following d	ata for averages,	fits, l	imits, eto	2. ● ● ●
0.00502	-80	¹ KAMANO	15	DPWA	Multichannel
1	с I I .: А :				

¹ From the preferred solution A in KAMANO 15.

Normalized residue in $N\overline{K} \rightarrow \Lambda(1830) \rightarrow \Sigma \pi$ MODULUS PHASE (°) DOCUMENT ID COMMENT TECN SARANTSEV 19 DPWA $\overline{K}N$ multichannel 0.15 ± 0.03 180 \pm 10 • • • We do not use the following data for averages, fits, limits, etc. • • • ¹ KAMANO 179 15 DPWA Multichannel 0.00581 ¹From the preferred solution A in KAMANO 15. Normalized residue in $N\overline{K} \rightarrow \Lambda(1830) \rightarrow \Lambda \eta$ PHASE (°) MODULUS DOCUMENT ID TECN COMMENT • • • We do not use the following data for averages, fits, limits, etc. • • • ¹ KAMANO 0.00941 -6515 DPWA Multichannel ¹From the preferred solution A in KAMANO 15. Normalized residue in $N\overline{K} \rightarrow \Lambda(1830) \rightarrow \Xi K$ MODULUS PHASE (°) DOCUMENT ID TECN COMMENT $0.010 \pm 0.005 65 \pm 20$ DPWA $\overline{K}N$ multichannel SARANTSEV 19 • • • We do not use the following data for averages, fits, limits, etc. • • • ¹ KAMANO 0.0477 94 15 **DPWA** Multichannel ¹From the preferred solution A in KAMANO 15. Normalized residue in $N\overline{K} \rightarrow \Lambda(1830) \rightarrow \Sigma(1385)\pi$, D-wave TECN COMMENT MODULUS PHASE (°) DOCUMENT ID SARANTSEV 19 DPWA $\overline{K}N$ multichannel 0.10 ±0.04 10 ± 25 • • • We do not use the following data for averages, fits, limits, etc. • • • 113 ¹ KAMANO 15 DPWA Multichannel 0.0237 ¹ From the preferred solution A in KAMANO 15. Normalized residue in $N\overline{K} \rightarrow \Lambda(1830) \rightarrow \Sigma(1385)\pi$, G-wave MODULUS PHASE (°) DOCUMENT ID TECN COMMENT SARANTSEV 19 DPWA $\overline{K}N$ multichannel ±0.02 0.03 • • We do not use the following data for averages, fits, limits, etc. • • • 1 kamano 15 **DPWA** Multichannel 0.000726 127 ¹ From the preferred solution A in KAMANO 15. Normalized residue in $N\overline{K} \rightarrow \Lambda(1830) \rightarrow N\overline{K}^*(892)$, S=1/2, D-wave PHASE (°) DOCUMENT ID TECN COMMENT MODULUS • • • We do not use the following data for averages, fits, limits, etc. • • • 0.0278 -177¹ KAMANO 15 DPWA Multichannel ¹From the preferred solution A in KAMANO 15. Normalized residue in $N\overline{K} \rightarrow \Lambda(1830) \rightarrow N\overline{K}^*(892)$, S=3/2, D-wave MODULUS PHASE (°) DOCUMENT ID TECN COMMENT • • • We do not use the following data for averages, fits, limits, etc. • • • ¹ KAMANO 3 0.0255 15 DPWA Multichannel ¹ From the preferred solution A in KAMANO 15.

Citation: R.L. Workman et al. (Particle Data Group), Prog. Theor. Exp. Phys. 2022, 083C01 (2022) and 2023 update

Normalized	residue in $N\overline{K} \rightarrow PHASE(^{\circ})$	$\Lambda(1830) \rightarrow N\overline{K}^*$	(892), S=3/2 , G-wave				
• • • We do r	not use the following	data for averages, fits,	limits, etc. ● ● ●				
0.00773	-17	¹ KAMANO 15	DPWA Multichannel				
1 From the μ	1 From the preferred solution A in KAMANO 15.						
Normalized	residue in $N\overline{K} \rightarrow$	Λ (1830) $\rightarrow \Lambda \omega$, S	S=1/2 , <i>D</i> -wave				
MODULUS	PHASE (°)	DOCUMENT ID	TECN COMMENT				
0.04±0.03		SARANTSEV 19	DPWA $\overline{K}N$ multichannel				
Normalized residue in $N\overline{K} ightarrow \Lambda(1830) ightarrow \Lambda\omega$, $S=3/2$, D-wave							
MODULUS	PHASE (°)	DOCUMENT ID	TECN COMMENT				
0.05 ± 0.03	-110 ± 35	SARANTSEV 19	DPWA $\overline{K}N$ multichannel				

Л(1830) MASS

VALUE (MeV)	DOCUMENT ID		TECN	COMMENT			
1820 to 1830 (≈ 1825) OUR ESTIMATE							
$1821\pm$ 3	SARANTSEV	19	DPWA	K N multichannel			
$1820\pm$ 4	ZHANG	13A	DPWA	Multichannel			
1831 ± 10	GOPAL	80	DPWA	$\overline{K}N \rightarrow \overline{K}N$			
1825 ± 10	GOPAL	77	DPWA	K N multichannel			
$1825\pm$ 1	KANE	74	DPWA	$K^- p \rightarrow \Sigma \pi$			
ullet $ullet$ $ullet$ We do not use the following data for averages, fits, limits, etc. $ullet$ $ullet$							
1817 or 1818	¹ MARTIN	77	DPWA	K N multichannel			
1 The two MARTIN 77 values are	e from a T-matri×	pole	and from	n a Breit-Wigner fit.			

Л(1830) WIDTH

VALUE (MeV)	DOCUMENT ID		TECN	COMMENT
60 to 120 (\approx 90) OUR ESTIMATE				
64± 7	SARANTSEV	19	DPWA	K N multichannel
114 ± 10	ZHANG	13A	DPWA	Multichannel
100 ± 10	GOPAL	80	DPWA	$\overline{K}N \rightarrow \overline{K}N$
94±10	GOPAL	77	DPWA	K N multichannel
$119\pm$ 3	KANE	74	DPWA	$K^- p \rightarrow \Sigma \pi$
\bullet \bullet \bullet We do not use the following a	lata for averages	, fits,	limits, e	tc. ● ● ●
56 or 56	^L MARTIN	77	DPWA	K N multichannel
1 The two MARTIN 77 values are	from a T-matrix	pole	and from	n a Breit-Wigner fit.

	Mode	Fraction (Γ_i/Γ)	Scale factor
Г1	NK	0.04 to 0.08	
Γ2	$\Sigma \pi$	35-75 %	
Γ ₃	ΞK		
Γ ₄	$\Sigma(1385)\pi$	>15 %	
Γ ₅	$\Sigma(1385)\pi$, <i>D</i> -wave	(40 ±15) %	3.2
Γ ₆	$\Sigma(1385)\pi$, <i>G</i> -wave		
Γ ₇	$\Lambda\eta$		
Γ ₈	<i>N</i> K *(892), <i>S</i> =1/2, <i>D</i> -wave		
Γ9	<i>NK</i> [*] (892), <i>S</i> =3/2, <i>D</i> -wave		
Γ_{10}	$N\overline{K}^{*}(892)$, $S=3/2$, G -wave		

A(1830) DECAY MODES

A(1830) BRANCHING RATIOS

See "Sign conventions for resonance couplings" in the Note on ${\it A}$ and ${\it \Sigma}$ Resonances.

$\Gamma(N\overline{K})/\Gamma_{\text{total}}$				Γ_1/Γ
VALUE	DOCUMENT ID		TECN	COMMENT
0.04 to 0.08 OUR ESTIMATE				
0.055 ± 0.010	SARANTSEV	19	DPWA	$\overline{K}N$ multichannel
0.041 ± 0.005	ZHANG	13A	DPWA	Multichannel
0.08 ± 0.03	GOPAL	80	DPWA	$\overline{K}N \rightarrow \overline{K}N$
0.02 ± 0.02	ALSTON	78	DPWA	$\overline{K}N \rightarrow \overline{K}N$
ullet $ullet$ $ullet$ We do not use the following	data for averages	s, fits,	limits, e	etc. • • •
0.006	1 KAMANO	15	DPWA	Multichannel
0.04 ± 0.03	GOPAL	77	DPWA	See GOPAL 80
0.04 or 0.04	² MARTIN	77	DPWA	K N multichannel
¹ From the preferred solution A ² The two MARTIN 77 values ar	in KAMANO 15. re from a T-matriz	x pole	and fro	m a Breit-Wigner fit.
$\Gamma(\Sigma\pi)/\Gamma_{\text{total}}$				Γ ₂ /Γ
$\Gamma(\Sigma \pi)/\Gamma_{\text{total}}$	DOCUMENT ID	·	TECN	Г2/Г соммент
$\frac{\Gamma(\Sigma\pi)/\Gamma_{\text{total}}}{0.42 \pm 0.08}$	<u>DOCUMENT ID</u> SARANTSEV	19	<u>tecn</u> DPWA	Γ2/Γ <u>COMMENT</u> <u>K</u> N multichannel
$\frac{\Gamma(\Sigma \pi) / \Gamma_{\text{total}}}{0.42 \pm 0.08}$ • • • We do not use the following	DOCUMENT ID SARANTSEV data for averages	19 s, fits,	<u>TECN</u> DPWA limits, e	$\frac{COMMENT}{\overline{K}N \text{ multichannel}}$
$\frac{\Gamma(\Sigma \pi) / \Gamma_{\text{total}}}{0.42 \pm 0.08}$ ••• We do not use the following 0.017	DOCUMENT ID SARANTSEV data for averages ¹ KAMANO	19 s, fits, 15	<u>TECN</u> DPWA limits, e DPWA	Γ_2/Γ $\frac{COMMENT}{\overline{K}N \text{ multichannel}}$ etc. • • •
$\Gamma(\Sigma \pi)/\Gamma_{\text{total}}$ <u>VALUE</u> 0.42 ±0.08 ••• We do not use the following 0.017 ¹ From the preferred solution A is	<u>DOCUMENT ID</u> SARANTSEV data for averages ¹ KAMANO in KAMANO 15.	19 s, fits, 15	<u>TECN</u> DPWA limits, e DPWA	Γ_2/Γ $\frac{COMMENT}{\overline{K}N \text{ multichannel}}$ etc. • • •
$\frac{\Gamma(\Sigma \pi) / \Gamma_{\text{total}}}{\frac{VALUE}{0.42 \pm 0.08}}$ • • • We do not use the following 0.017 ¹ From the preferred solution A is $\Gamma(\Xi K) / \Gamma_{\text{total}}$	<u>DOCUMENT ID</u> SARANTSEV data for averages ¹ KAMANO in KAMANO 15.	19 s, fits, 15	<u>TECN</u> DPWA limits, e DPWA	$ Γ2/Γ \overline{K} N \text{ multichannel} etc. • • • Multichannel Γ3/Γ $
$\Gamma(\Sigma \pi)/\Gamma_{\text{total}}$ <u>VALUE</u> 0.42 ±0.08 • • • We do not use the following 0.017 ¹ From the preferred solution A is $\Gamma(\Xi K)/\Gamma_{\text{total}}$ <u>VALUE</u>	<u>DOCUMENT ID</u> SARANTSEV data for averages ¹ KAMANO in KAMANO 15. <u>DOCUMENT ID</u>	19 s, fits, 15	<u>TECN</u> DPWA limits, e DPWA	Γ_2/Γ <u>COMMENT</u> $\overline{K} N \text{ multichannel}$ etc. • • • Multichannel Γ_3/Γ <u>COMMENT</u>
$\Gamma(\Sigma \pi)/\Gamma_{\text{total}}$ <u>VALUE</u> 0.42 ±0.08 ••• We do not use the following 0.017 ¹ From the preferred solution A is $\Gamma(\Xi K)/\Gamma_{\text{total}}$ <u>VALUE</u> ••• We do not use the following	<u>DOCUMENT ID</u> SARANTSEV data for averages ¹ KAMANO in KAMANO 15. <u>DOCUMENT ID</u> data for averages	19 s, fits, 15 s, fits,	<u>TECN</u> DPWA limits, e DPWA <u>TECN</u> limits, e	Γ_2/Γ $\frac{COMMENT}{\overline{K}N \text{ multichannel}}$ etc. • • • Multichannel Γ_3/Γ $COMMENT$ etc. • • •
$\Gamma(\Sigma \pi)/\Gamma_{\text{total}}$ \xrightarrow{VALUE} 0.42 ±0.08 ••• We do not use the following 0.017 ¹ From the preferred solution A is $\Gamma(\Xi K)/\Gamma_{\text{total}}$ \xrightarrow{VALUE} ••• We do not use the following 0.562	DOCUMENT ID SARANTSEV data for averages ¹ KAMANO in KAMANO 15. <u>DOCUMENT ID</u> data for averages ¹ KAMANO	19 5, fits, 15 s, fits, 15	<u>TECN</u> DPWA limits, e DPWA <u>TECN</u> limits, e DPWA	$ Γ2/Γ \overline{K} N \text{ multichannel} etc. • • • Multichannel Γ3/Γ \\ \underline{COMMENT} etc. • • • Multichannel $

¹ From the preferred solution A in KAMANO 15.

$\Gamma(\Sigma(1385)\pi$, <i>D</i> -wave $)/\Gamma_{ m tet}$	otal				Г ₅ /Г
VALUE	DOCUMENT ID		TECN	COMMENT	
0.40 \pm 0.15 OUR AVERAGE	Error includes scale f	actor	of 3.2.	_	
0.20 ± 0.08	SARANTSEV	19	DPWA	KN multichannel	
0.52 ± 0.06	ZHANG	13A	DPWA	Multichannel	
• • We do not use the follow	wing data for averages	s, fits,	limits, e	tc. ● ● ●	
).134	¹ KAMANO	15	DPWA	Multichannel	
1 From the preferred solution	n A in KAMANO 15.				
$(\Sigma(1385)\pi, G-wave)/\Gamma_{tot}$	otal				Г ₆ /Г
ALUE	DOCUMENT ID		TECN	COMMENT	
0.020 ± 0.015	SARANTSEV	19	DPWA	K N multichannel	
$(\Lambda n)/\Gamma_{total}$					Γ ₇ /Γ
ALUE	DOCUMENT ID		TECN	COMMENT	• /
• • We do not use the follow	wing data for averages	s, fits,	limits, e	tc. ● ● ●	
).024	¹ KAMANO	15	DPWA	Multichannel	
1 From the preferred solution	Δ in KAMANO 15				
Trom the preferred solution	I A III NAMANO 13.				
-(<i>NT</i> *(892), <i>S</i> =1/2, <i>D</i> -w	/ave)/Γ _{total}				Γ ₈ /Γ
ALUE	DOCUMENT ID		TECN	COMMENT	-
• • We do not use the follow	wing data for averages	s, fits,	limits, e	tc. ● ● ●	
).134	¹ KAMANO	15	DPWA	Multichannel	
1 From the preferred solution	A in KAMANO 15				
	N				
~(<i>NK</i> *(892), <i>S</i> =3/2, <i>D</i> -w	/ave)/F _{total}				٦/و٦
ALUE	DOCUMENT ID		TECN	COMMENT	
• • We do not use the follow	wing data for averages	s, fits,	limits, e	tc. ● ● ●	
0.115	¹ KAMANO	15	DPWA	Multichannel	
¹ From the preferred solutior	n A in KAMANO 15.				
$(N\overline{K}*(902)) \subset 2/2 \subset$					Г., /Г
(IN A (092), 3=3/2, G-W			TECH	COMMENT	10/1
ALUE	<u>DOCUMENT ID</u>	<i></i>	<u>TECN</u>	<u>COMMENT</u>	
$\bullet \bullet \bullet$ vve ao not use the follow	ving data for averages	s, tits,	iimits, e	τς. ● ● ●	
.009	¹ KAMANO	15	DPWA	Multichannel	
¹ From the preferred solutior	n A in KAMANO 15.				
$(\Gamma,\Gamma_{c})^{\frac{1}{2}}/\Gamma_{c}$, in $N\overline{K} \rightarrow C$	Λ(1830) → Σπ			([1]	<u>\</u> %/г
\' i' f) / ' total ''' / ' · · · · /Al UF	DOCUMENT ID		TECN	COMMENT	<i>''</i>
-0.13+0.01	ZHANG	134	DPWA	Multichannel	
-0.17 ± 0.03	GOPAL	77	DPWA	$\overline{K}N$ multichannel	
-0.15 ± 0.01	KANE	74	DPWA	$K^- p \rightarrow \Sigma \pi$	
• • We do not use the follow	wing data for averages	s, fits,	limits, e	tc. ● ● ●	
-0.17 or -0.17	¹ MARTIN	77		$\overline{K}N$ multichannel	
	······································	••			··.

¹ The two MARTIN 77 values are from a T-matrix pole and from a Breit-Wigner fit.

Citation: R.L. Workman et al. (Particle Data Group), Prog.Theor.Exp.Phys. 2022, 083C01 (2022) and 2023 update

$(\Gamma_i \Gamma_f)^{\frac{1}{2}} / \Gamma_{\text{total}} \text{ in } N\overline{K} \to \Lambda(1830) \to \Sigma(1385)\pi$					(Γ ₁ Γ ₄) ^½ /Γ
VALUE	DOCUMENT ID		TECN	<u>COMMENT</u>	
0.20 to 0.50 OUR ESTIMAT	Έ				
$+0.141\!\pm\!0.014$	¹ CAMERON	78	DPWA	$K^- p \rightarrow$	$\Sigma(1385)\pi$
$+0.13 \pm 0.03$	PREVOST	74	DPWA	$K^-N \rightarrow$	$\Sigma(1385)\pi$
-					

¹ The CAMERON 78 upper limit on *G*-wave decay is 0.03. The published sign has been changed to be in accord with the baryon-first convention.

$(\Gamma_i \Gamma_f)^{\frac{1}{2}} / \Gamma_{\text{total}} \text{ in } N\overline{K} \to \Lambda(1830) \to \Lambda \eta$							
VALUE	DOCUMENT ID		TECN				
-0.044 ± 0.020	RADER	73	MPWA				

A(1830) REFERENCES

 $(\Gamma_1\Gamma_7)^{\frac{1}{2}}/\Gamma$