

$\phi(1020)$ $I^G(J^{PC}) = 0^-(1^{--})$ **$\phi(1020)$ MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1019.461 ± 0.016 OUR AVERAGE				
1019.463 ± 0.061	2.3M	¹ KOZYREV	18	CMD3 $e^+ e^- \rightarrow K^+ K^-, K_S^0 K_L^0$
1019.462 $\pm 0.042 \pm 0.056$	28k	² LEES	14H	BABR $e^+ e^- \rightarrow K_S^0 K_L^0 \gamma$
1019.51 $\pm 0.02 \pm 0.05$		³ LEES	13Q	BABR $e^+ e^- \rightarrow K^+ K^- \gamma$
1019.30 $\pm 0.02 \pm 0.10$	105k	AKHMETSHIN 06	CMD2	$0.98-1.06 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
1019.52 $\pm 0.05 \pm 0.05$	17.4k	AKHMETSHIN 05	CMD2	$0.60-1.38 e^+ e^- \rightarrow \eta \gamma$
1019.483 $\pm 0.011 \pm 0.025$	272k	⁴ AKHMETSHIN 04	CMD2	$e^+ e^- \rightarrow K_L^0 K_S^0$
1019.42 ± 0.05	1900k	⁵ ACHASOV	01E	SND $e^+ e^- \rightarrow K^+ K^-, K_S K_L, \pi^+ \pi^- \pi^0$
1019.40 $\pm 0.04 \pm 0.05$	23k	AKHMETSHIN 01B	CMD2	$e^+ e^- \rightarrow \eta \gamma$
1019.36 ± 0.12		⁶ ACHASOV	00B	SND $e^+ e^- \rightarrow \eta \gamma$
1019.38 $\pm 0.07 \pm 0.08$	2200	⁷ AKHMETSHIN 99F	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \geq 2\gamma$
1019.51 $\pm 0.07 \pm 0.10$	11169	AKHMETSHIN 98	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
1019.5 ± 0.4		BARBERIS	98	OMEG $450 pp \rightarrow pp 2K^+ 2K^-$
1019.42 ± 0.06	55600	AKHMETSHIN 95	CMD2	$e^+ e^- \rightarrow \text{hadrons}$
1019.7 ± 0.3	2012	DAVENPORT	86	MPSF $400 \mu\text{A} \rightarrow 4KX$
1019.7 $\pm 0.1 \pm 0.1$	5079	ALBRECHT	85D	ARG $10 e^+ e^- \rightarrow K^+ K^- X$
1019.3 ± 0.1	1500	ARENTON	82	AEMS $11.8 \text{ polar. } pp \rightarrow KK$
1019.67 ± 0.17	25080	⁸ PELLINEN	82	RVUE
1019.52 ± 0.13	3681	BUKIN	78C	OLYA $e^+ e^- \rightarrow \text{hadrons}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1018.4 $\pm 0.5 \pm 0.1$		⁹ ALBRECHT	20	CBAR $0.9 \bar{p}p \rightarrow K^+ K^- \pi^0$
1019.21 $\pm 0.04 \pm 0.03$		¹⁰ HOID	20	RVUE $e^+ e^- \rightarrow \pi^0 \gamma$
1019.54 $\pm 0.10 \pm 0.51$		¹¹ AAIJ	19H	LHCb $pp \rightarrow D^\pm X$
1019.20 $\pm 0.02 \pm 0.01$		¹² HOFERICHT...	19	RVUE $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
1019.469 ± 0.061	1.7M	KOZYREV	18	CMD3 $e^+ e^- \rightarrow K^+ K^-$
1019.457 ± 0.061	610k	KOZYREV	16	CMD3 $e^+ e^- \rightarrow K_S^0 K_L^0$
1019.48 ± 0.01		LEES	13F	BABR $D^+ \rightarrow K^+ K^- \pi^+$
1019.441 $\pm 0.008 \pm 0.080$	542k	¹³ AKHMETSHIN 08	CMD2	$1.02 e^+ e^- \rightarrow K^+ K^-$
1019.63 ± 0.07	12540	¹⁴ AUBERT,B	05J	BABR $D^0 \rightarrow \bar{K}^0 K^+ K^-$
1019.8 ± 0.7		ARMSTRONG	86	OMEG $85 \pi^+ / pp \rightarrow \pi^+ / p4Kp$
1020.1 ± 0.11	5526	¹⁴ ATKINSON	86	OMEG $20-70 \gamma p$
1019.7 ± 1.0		BEBEK	86	CLEO $e^+ e^- \rightarrow \Upsilon(4S)$
1019.411 ± 0.008	642k	¹⁵ DIJKSTRA	86	SPEC $100-200 \pi^\pm, \bar{p}, p, K^\pm, \text{on Be}$

1020.9	± 0.2		¹⁴ FRAME	86	OMEG	13	$K^+ p \rightarrow \phi K^+ p$
1021.0	± 0.2		¹⁴ ARMSTRONG	83B	OMEG	18.5	$K^- p \rightarrow K^- K^+ \Lambda$
1020.0	± 0.5		¹⁴ ARMSTRONG	83B	OMEG	18.5	$K^- p \rightarrow K^- K^+ \Lambda$
1019.7	± 0.3		¹⁴ BARATE	83	GOLI	190	$\pi^- Be \rightarrow 2\mu X$
1019.8	± 0.2	± 0.5	766	IVANOV	81	OLYA	$1-1.4 e^+ e^- \rightarrow K^+ K^-$
1019.4	± 0.5		337	COOPER	78B	HBC	$0.7-0.8 \bar{p}p \rightarrow K_S^0 K_L^0 \pi^+ \pi^-$
1020	± 1		383	¹⁴ BALDI	77	CNTR	$10 \pi^- p \rightarrow \pi^- \phi p$
1018.9	± 0.6		800	COHEN	77	ASPK	$6 \pi^\pm N \rightarrow K^+ K^- N$
1019.7	± 0.5		454	KALBFLEISCH	76	HBC	$2.18 K^- p \rightarrow \Lambda K\bar{K}$
1019.4	± 0.8		984	BESCH	74	CNTR	$2 \gamma p \rightarrow p K^+ K^-$
1020.3	± 0.4		100	BALLAM	73	HBC	$2.8-9.3 \gamma p$
1019.4	± 0.7			BINNIE	73B	CNTR	$\pi^- p \rightarrow \phi n$
1019.6	± 0.5		120	¹⁶ AGUILAR-...	72B	HBC	$3.9, 4.6 K^- p \rightarrow \Lambda K^+ K^-$
1019.9	± 0.5		100	¹⁶ AGUILAR-...	72B	HBC	$3.9, 4.6 K^- p \rightarrow K^- p K^+ K^-$
1020.4	± 0.5		131	COLLEY	72	HBC	$10 K^+ p \rightarrow K^+ p \phi$
1019.9	± 0.3		410	STOTTLE...	71	HBC	$2.9 K^- p \rightarrow \Sigma/\Lambda K\bar{K}$

¹ Average of KOZYREV 16 and KOZYREV 18 values taking into account the correlated uncertainties. Supersedes individual KOZYREV 16 and KOZYREV 18 results.

² Using a vector meson dominance model with contribution from $\phi(1020)$ and higher mass excitations of $\rho(770)$, $\omega(782)$, and $\phi(1020)$.

³ Using a phenomenological model based on KUHN 90 with a sum of Breit-Wigner resonances for $\rho(770)$, $\omega(782)$, $\phi(1020)$ and their higher mass excitations.

⁴ Update of AKHMETSHIN 99D

⁵ From the combined fit assuming that the total $\phi(1020)$ production cross section is saturated by those of $K^+ K^-$, $K_S K_L$, $\pi^+ \pi^- \pi^0$, and $\eta \gamma$ decays modes and using ACHASOV 00B for the $\eta \gamma$ decay mode.

⁶ Using a total width of 4.43 ± 0.05 MeV. Systematic uncertainty included.

⁷ Using a total width of 4.43 ± 0.05 MeV.

⁸ PELLINEN 82 review includes AKERLOF 77, DAUM 81, BALDI 77, AYRES 74, DEGROOT 74.

⁹ Width fixed at 4.2 MeV.

¹⁰ The values were extracted from a dispersively improved Breit-Wigner parameterization and do not include vacuum polarization. Inclusion of vacuum polarization gives 1019.457 ± 0.020 MeV.

¹¹ From the $D^\pm \rightarrow K^\pm K^+ K^-$ Dalitz plot fit with the Triple-M amplitude in the multi-meson model of AOUDE 18.

¹² The values were extracted from a dispersively improved Breit-Wigner parameterization and do not include vacuum polarization.

¹³ Strongly correlated with AKHMETSHIN 04.

¹⁴ Systematic errors not evaluated.

¹⁵ Weighted and scaled average of 12 measurements of DIJKSTRA 86.

¹⁶ Mass errors enlarged by us to Γ/\sqrt{N} ; see the note with the $K^*(892)$ mass.

$\phi(1020)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
4.249±0.013 OUR AVERAGE				Error includes scale factor of 1.1.
4.245±0.013	2.3M	¹ KOZYREV	18	CMD3 $e^+ e^- \rightarrow K^+ K^-$, $K_S^0 K_L^0$
4.205±0.103±0.067	28k	² LEES	14H	BABR $e^+ e^- \rightarrow K_S^0 K_L^0 \gamma$
4.29 ± 0.04 ± 0.07		³ LEES	13Q	BABR $e^+ e^- \rightarrow K^+ K^- \gamma$
4.30 ± 0.06 ± 0.17	105k	AKHMETSHIN 06	CMD2	0.98–1.06 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
4.280±0.033±0.025	272k	⁴ AKHMETSHIN 04	CMD2	$e^+ e^- \rightarrow K_L^0 K_S^0$
4.21 ± 0.04	1900k	⁵ ACHASOV	01E	SND $e^+ e^- \rightarrow K^+ K^-$, $K_S K_L, \pi^+ \pi^- \pi^0$
4.44 ± 0.09	55600	AKHMETSHIN 95	CMD2	$e^+ e^- \rightarrow$ hadrons
4.5 ± 0.7	1500	ARENTON	82	AEMS 11.8 polar. $p p \rightarrow K K$
4.2 ± 0.6	766	⁶ IVANOV	81	OLYA 1–1.4 $e^+ e^- \rightarrow K^+ K^-$
4.3 ± 0.6		⁶ CORDIER	80	DM1 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
4.36 ± 0.29	3681	⁶ BUKIN	78C	OLYA $e^+ e^- \rightarrow$ hadrons
4.4 ± 0.6	984	⁶ BESCH	74	CNTR $2 \gamma p \rightarrow p K^+ K^-$
4.67 ± 0.72	681	⁶ BALAKIN	71	OSPK $e^+ e^- \rightarrow$ hadrons
4.09 ± 0.29		BIZOT	70	OSPK $e^+ e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4.07 ± 0.13 ± 0.01		⁷ HOID	20	RVUE $e^+ e^- \rightarrow \pi^0 \gamma$
4.23 ± 0.04 ± 0.02		⁸ HOFERICHT...	19	RVUE $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
4.249±0.015	1.7M	KOZYREV	18	CMD3 $e^+ e^- \rightarrow K^+ K^-$
4.240±0.017	610k	KOZYREV	16	CMD3 $e^+ e^- \rightarrow K_S^0 K_L^0$
4.37 ± 0.02		LEES	13F	BABR $D^+ \rightarrow K^+ K^- \pi^+$
4.24 ± 0.02 ± 0.03	542k	⁹ AKHMETSHIN 08	CMD2	1.02 $e^+ e^- \rightarrow K^+ K^-$
4.28 ± 0.13	12540	¹⁰ AUBERT,B	05J	BABR $D^0 \rightarrow \bar{K}^0 K^+ K^-$
4.45 ± 0.06	271k	DIJKSTRA	86	SPEC 100 π^- Be
3.6 ± 0.8	337	⁶ COOPER	78B	HBC 0.7–0.8 $\bar{p} p \rightarrow K_S^0 K_L^0 \pi^+ \pi^-$
4.5 ± 0.50	1300	^{6,10} AKERLOF	77	SPEC 400 $p\bar{A} \rightarrow K^+ K^- X$
4.5 ± 0.8	500	^{6,10} AYRES	74	ASPK 3–6 $\pi^- p \rightarrow K^+ K^- n, K^- p \rightarrow K^+ K^- \Lambda/\Sigma^0$
3.81 ± 0.37		COSME	74B	OSPK $e^+ e^- \rightarrow K_L^0 K_S^0$
3.8 ± 0.7	454	⁶ BORENSTEIN	72	HBC 2.18 $K^- p \rightarrow K \bar{K} n$

¹ Average of KOZYREV 16 and KOZYREV 18 values taking into account the correlated uncertainties. Supersedes individual KOZYREV 16 and KOZYREV 18 results.

² Using a vector meson dominance model with contribution from $\phi(1020)$ and higher mass excitations of $\rho(770)$, $\omega(782)$, and $\phi(1020)$.

³ Using a phenomenological model based on KUHN 90 with a sum of Breit-Wigner resonances for $\rho(770)$, $\omega(782)$, $\phi(1020)$ and their higher mass excitations.

⁴ Update of AKHMETSHIN 99D

⁵ From the combined fit assuming that the total $\phi(1020)$ production cross section is saturated by those of $K^+ K^-$, $K_S K_L$, $\pi^+ \pi^- \pi^0$, and $\eta \gamma$ decays modes and using ACHASOV 00B for the $\eta \gamma$ decay mode.

⁶ Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.

⁷ The values were extracted from a dispersively improved Breit-Wigner parameterization.⁸ The values were extracted from a dispersively improved Breit-Wigner parameterization and do not include vacuum polarization.⁹ Strongly correlated with AKHMETSHIN 04.¹⁰ Systematic errors not evaluated.

$\phi(1020)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
$\Gamma_1 K^+ K^-$	(49.1 \pm 0.5) %	S=1.3
$\Gamma_2 K_L^0 K_S^0$	(33.9 \pm 0.4) %	S=1.2
$\Gamma_3 \rho\pi + \pi^+\pi^-\pi^0$	(15.4 \pm 0.4) %	S=1.2
$\Gamma_4 \pi^+\pi^-\pi^0$		
$\Gamma_5 \eta\gamma$	(1.301 \pm 0.024) %	S=1.2
$\Gamma_6 \pi^0\gamma$	(1.32 \pm 0.05) $\times 10^{-3}$	
$\Gamma_7 \ell^+\ell^-$	—	
$\Gamma_8 e^+e^-$	(2.979 \pm 0.033) $\times 10^{-4}$	S=1.2
$\Gamma_9 \mu^+\mu^-$	(2.85 \pm 0.22) $\times 10^{-4}$	S=1.2
$\Gamma_{10} \eta e^+e^-$	(1.08 \pm 0.04) $\times 10^{-4}$	
$\Gamma_{11} \pi^+\pi^-$	(7.3 \pm 1.3) $\times 10^{-5}$	
$\Gamma_{12} \omega\pi^0$	(4.7 \pm 0.5) $\times 10^{-5}$	
$\Gamma_{13} \omega\gamma$	< 5 %	CL=84%
$\Gamma_{14} \rho\gamma$	< 1.2 $\times 10^{-5}$	CL=90%
$\Gamma_{15} \pi^+\pi^-\gamma$	(4.1 \pm 1.3) $\times 10^{-5}$	
$\Gamma_{16} f_0(980)\gamma$	(3.22 \pm 0.19) $\times 10^{-4}$	S=1.1
$\Gamma_{17} \pi^0\pi^0\gamma$	(1.12 \pm 0.06) $\times 10^{-4}$	
$\Gamma_{18} \pi^+\pi^-\pi^+\pi^-$	(3.9 \pm 2.8) $\times 10^{-6}$	
$\Gamma_{19} \pi^+\pi^+\pi^-\pi^-\pi^0$	< 4.6 $\times 10^{-6}$	CL=90%
$\Gamma_{20} \pi^0e^+e^-$	(1.33 \pm 0.07) $\times 10^{-5}$	
$\Gamma_{21} \pi^0\eta\gamma$	(7.27 \pm 0.30) $\times 10^{-5}$	S=1.5
$\Gamma_{22} a_0(980)\gamma$	(7.6 \pm 0.6) $\times 10^{-5}$	
$\Gamma_{23} K^0\bar{K}^0\gamma$	< 1.9 $\times 10^{-8}$	CL=90%
$\Gamma_{24} \eta'(958)\gamma$	(6.21 \pm 0.20) $\times 10^{-5}$	
$\Gamma_{25} \eta\pi^0\pi^0\gamma$	< 2 $\times 10^{-5}$	CL=90%
$\Gamma_{26} \mu^+\mu^-\gamma$	(1.4 \pm 0.5) $\times 10^{-5}$	
$\Gamma_{27} \rho\gamma\gamma$	< 1.2 $\times 10^{-4}$	CL=90%
$\Gamma_{28} \eta\pi^+\pi^-$	< 1.8 $\times 10^{-5}$	CL=90%
$\Gamma_{29} \eta\mu^+\mu^-$	< 9.4 $\times 10^{-6}$	CL=90%
$\Gamma_{30} \eta U \rightarrow \eta e^+e^-$	< 1 $\times 10^{-6}$	CL=90%
Γ_{31} invisible	< 1.7 $\times 10^{-4}$	CL=90%
Lepton Family number (<i>LF</i>) violating modes		
$\Gamma_{32} e^\pm\mu^\mp$	<i>LF</i> < 2 $\times 10^{-6}$	CL=90%

CONSTRAINED FIT INFORMATION

An overall fit to 30 branching ratios uses 80 measurements and one constraint to determine 14 parameters. The overall fit has a $\chi^2 = 61.8$ for 67 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-72									
x_3	-61 -12									
x_5	-19 17 2									
x_6	-11 11 1 8									
x_8	47 -50 -8 -35 -22									
x_9	-5 4 1 3 2 -9									
x_{11}	-3 3 0 2 1 -6 1									
x_{12}	-4 4 1 3 2 -8 1 0									
x_{16}	0 0 0 0 0 0 0 0 0									
x_{17}	-9 9 1 18 4 -18 2 1 1 0									
x_{18}	-1 1 0 1 0 -2 0 0 0 0									
x_{22}	0 0 0 0 0 0 0 0 0 0									
x_{24}	-6 5 1 32 2 -11 1 1 1 0									
	x_1	x_2	x_3	x_5	x_6	x_8	x_9	x_{11}	x_{12}	x_{16}
x_{18}	0									
x_{22}	0 0									
x_{24}	6 0 0									
	x_{17}	x_{18}	x_{22}							

$\phi(1020)$ PARTIAL WIDTHS

$\Gamma(\eta\gamma)$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT	Γ_5
-------------	-------------	------	---------	------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$58.9 \pm 0.5 \pm 2.4$ ACHASOV 00 SND $e^+ e^- \rightarrow \eta\gamma$

$\Gamma(\pi^0\gamma)$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT	Γ_6
-------------	-------------	------	---------	------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$5.40 \pm 0.16^{+0.43}_{-0.40}$ ACHASOV 00 SND $e^+ e^- \rightarrow \pi^0\gamma$

$\Gamma(\ell^+ \ell^-)$ Γ_7

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
$1.320 \pm 0.017 \pm 0.015$	¹ AMBROSINO 05	KLOE	$1.02 e^+ e^- \rightarrow \mu^+ \mu^-$
¹ Weighted average of Γ_{ee} and $\sqrt{\Gamma_{ee}\Gamma_{\mu\mu}}$ from AMBROSINO 05 assuming lepton universality.			

$\Gamma(e^+ e^-)$ Γ_8

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
1.27 ± 0.04 OUR EVALUATION			
1.251 ± 0.021 OUR AVERAGE Error includes scale factor of 1.1.			
$1.235 \pm 0.006 \pm 0.022$	¹ AKHMETSHIN 11	CMD2	$1.02 e^+ e^- \rightarrow \phi$
$1.32 \pm 0.05 \pm 0.03$	² AMBROSINO 05	KLOE	$1.02 e^+ e^- \rightarrow e^+ e^-$
1.28 ± 0.05	AKHMETSHIN 95	CMD2	$1.02 e^+ e^- \rightarrow \phi$
¹ Combined analysis of the CMD-2 data on $\phi \rightarrow K^+ K^-$, $K_S^0 K_L^0$, $\pi^+ \pi^- \pi^0$, $\eta \gamma$ assuming that the sum of their branching fractions is 0.99741 ± 0.00007 .			
² From forward-backward asymmetry and using $\Gamma_{\text{total}} = 4.26 \pm 0.05$ MeV from the 2004 edition of this Review.			

$$(\Gamma(e^+ e^-) \times \Gamma(\mu^+ \mu^-))^{\frac{1}{2}} \quad (\Gamma_8 \Gamma_9)^{\frac{1}{2}}$$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
$1.320 \pm 0.018 \pm 0.017$	AMBROSINO 05	KLOE	$1.02 e^+ e^- \rightarrow \mu^+ \mu^-$

$$\phi(1020) \Gamma(i) \Gamma(e^+ e^-) / \Gamma(\text{total})$$

$\Gamma(K^+ K^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_1 \Gamma_8 / \Gamma$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
$0.6340 \pm 0.0070 \pm 0.0039$	¹ LEES	13Q	BABR	$e^+ e^- \rightarrow K^+ K^- \gamma$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$0.669 \pm 0.001 \pm 0.023$ 1.7M	KOZYREV	18	CMD3	$e^+ e^- \rightarrow K^+ K^-$
¹ Using a phenomenological model based on KUHN 90 with a sum of Breit-Wigner resonances for $\rho(770)$, $\omega(782)$, $\phi(1020)$ and their higher mass excitations. The first error combines statistical and systematic uncertainties. The second one is due to the parametrization of the charged kaon form factor and mass calibration.				

$\Gamma(K_L^0 K_S^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_2 \Gamma_8 / \Gamma$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
$0.4200 \pm 0.0033 \pm 0.0123$	28k	¹ LEES	14H	BABR $e^+ e^- \rightarrow K_S^0 K_L^0 \gamma$

¹ Using a vector meson dominance model with contribution from $\phi(1020)$ and higher mass excitations of $\rho(770)$, $\omega(782)$, and $\phi(1020)$.

$[\Gamma(\rho\pi) + \Gamma(\pi^+ \pi^- \pi^0)] \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_3 \Gamma_8 / \Gamma$


VALUE (eV)	DOCUMENT ID	TECN	COMMENT
$184.1 \pm 2.1 \pm 8.0$	1 LEES	21B	BABR $10.5 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \gamma$
¹ From the cross section for $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$ with contributions from $\rho(770)$, $\omega(782)$, $\phi(1020)$, $\omega(1420)$, and $\omega(1650)$.			

$\phi(1020) \Gamma(i)\Gamma(e^+e^-)/\Gamma^2(\text{total})$

$$\Gamma(K^+K^-)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_1/\Gamma \times \Gamma_8/\Gamma$$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
14.64 ± 0.27 OUR FIT		Error includes scale factor of 1.4.		
14.6 ± 0.5 OUR AVERAGE		Error includes scale factor of 1.8. See the ideogram below.		
15.789 ± 0.541	1.7M	KOZYREV 18	CMD3	$e^+e^- \rightarrow K^+K^-$
14.27 ± 0.05 ± 0.31	542k	AKHMETSHIN 08	CMD2	$1.02 e^+e^- \rightarrow K^+K^-$
13.93 ± 0.14 ± 0.99	1000k	¹ ACHASOV 01E	SND	$e^+e^- \rightarrow K^+K^-$, $K_S K_L, \pi^+\pi^-\pi^0$

¹ From the combined fit assuming that the total $\phi(1020)$ production cross section is saturated by those of K^+K^- , $K_S K_L$, $\pi^+\pi^-\pi^0$, and $\eta\gamma$ decays modes and using ACHASOV 00B for the $\eta\gamma$ decay mode.



$$\Gamma(K_L^0 K_S^0)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$$

$$\Gamma_2/\Gamma \times \Gamma_8/\Gamma$$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
10.11 ± 0.12 OUR FIT				
10.07 ± 0.13 OUR AVERAGE				
10.078 ± 0.223	610k	¹ KOZYREV 16	CMD3	$e^+e^- \rightarrow K_S^0 K_L^0$
10.01 ± 0.04 ± 0.17	272k	² AKHMETSHIN 04	CMD2	$e^+e^- \rightarrow K_L^0 K_S^0$
10.27 ± 0.07 ± 0.34	500k	³ ACHASOV 01E	SND	$e^+e^- \rightarrow K^+K^-$, $K_S K_L, \pi^+\pi^-\pi^0$

¹ KOZYREV 16 also reports $\Gamma(e^+ e^-) B(\phi \rightarrow K_S^0 K_L^0) = (0.428 \pm 0.001 \pm 0.009)$ keV.

² Update of AKHMETSHIN 99D

³ From the combined fit assuming that the total $\phi(1020)$ production cross section is saturated by those of $K^+ K^-$, $K_S K_L$, $\pi^+ \pi^- \pi^0$, and $\eta \gamma$ decays modes and using ACHASOV 00B for the $\eta \gamma$ decay mode.

[$\Gamma(\rho\pi) + \Gamma(\pi^+ \pi^- \pi^0)$] / Γ_{total}	$\Gamma_3/\Gamma \times \Gamma_8/\Gamma$
---	--

[$\Gamma(\rho\pi) + \Gamma(\pi^+ \pi^- \pi^0)$] / Γ_{total}	$\Gamma_3/\Gamma \times \Gamma_8/\Gamma$			
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
4.58 ± 0.11 OUR FIT		Error includes scale factor of 1.1.		
4.51 ± 0.14 OUR AVERAGE				
4.51 ± 0.16 ± 0.11	105k	AKHMETSHIN 06	CMD2	$0.98-1.06 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
4.665 ± 0.042 ± 0.261	400k	¹ ACHASOV 01E	SND	$e^+ e^- \rightarrow K^+ K^-$, $K_S K_L, \pi^+ \pi^- \pi^0$
4.35 ± 0.27 ± 0.08	11169	² AKHMETSHIN 98	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4.38 ± 0.12		BENAYOUN 10	RVUE	$0.4-1.05 e^+ e^-$
4.30 ± 0.08 ± 0.21		³ AUBERT,B 04N	BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \gamma$

¹ From the combined fit assuming that the total $\phi(1020)$ production cross section is saturated by those of $K^+ K^-$, $K_S K_L$, $\pi^+ \pi^- \pi^0$, and $\eta \gamma$ decays modes and using ACHASOV 00B for the $\eta \gamma$ decay mode.

² Recalculated by us from the cross section in the peak.

³ Superseeded by LEES 21B.

[$\Gamma(\eta\gamma) / \Gamma_{\text{total}} \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$]	$\Gamma_5/\Gamma \times \Gamma_8/\Gamma$
---	--

[$\Gamma(\eta\gamma) / \Gamma_{\text{total}} \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$]	$\Gamma_5/\Gamma \times \Gamma_8/\Gamma$			
VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
3.88 ± 0.07 OUR FIT		Error includes scale factor of 1.2.		
3.93 ± 0.09 OUR AVERAGE		Error includes scale factor of 1.3. See the ideogram below.		
4.050 ± 0.067 ± 0.118	33k	¹ ACHASOV 07B	SND	$0.6-1.38 e^+ e^- \rightarrow \eta \gamma$
4.093 ^{+0.040} _{-0.043} ± 0.247	17.4k	² AKHMETSHIN 05	CMD2	$0.60-1.38 e^+ e^- \rightarrow \eta \gamma$
3.850 ± 0.041 ± 0.159	23k	^{3,4} AKHMETSHIN 01B	CMD2	$e^+ e^- \rightarrow \eta \gamma$
4.00 ± 0.04 ± 0.11		⁵ ACHASOV 00	SND	$e^+ e^- \rightarrow \eta \gamma$
3.53 ± 0.08 ± 0.17	2200	^{6,7} AKHMETSHIN 99F	CMD2	$e^+ e^- \rightarrow \eta \gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4.19 ± 0.06		⁸ BENAYOUN 10	RVUE	$0.4-1.05 e^+ e^-$

¹ From a combined fit of $\sigma(e^+ e^- \rightarrow \eta \gamma)$ with $\eta \rightarrow 3\pi^0$ and $\eta \rightarrow \pi^+ \pi^- \pi^0$, and fixing $B(\eta \rightarrow 3\pi^0) / B(\eta \rightarrow \pi^+ \pi^- \pi^0) = 1.44 \pm 0.04$. Recalculated by us from the cross section at the peak. Supersedes ACHASOV 00D and ACHASOV 06A.

² From the $\eta \rightarrow 2\gamma$ decay and using $B(\eta \rightarrow \gamma\gamma) = 39.43 \pm 0.26\%$.

³ From the $\eta \rightarrow 3\pi^0$ decay and using $B(\eta \rightarrow 3\pi^0) = (32.24 \pm 0.29) \times 10^{-2}$.


⁴ The combined fit from 600 to 1380 MeV taking into account $\rho(770)$, $\omega(782)$, $\phi(1020)$, and $\rho(1450)$ (mass and width fixed at 1450 MeV and 310 MeV respectively).

⁵ From the $\eta \rightarrow 2\gamma$ decay and using $B(\eta \rightarrow 2\gamma) = (39.21 \pm 0.34) \times 10^{-2}$.

⁶ Recalculated by the authors from the cross section in the peak.

⁷ From the $\eta \rightarrow \pi^+ \pi^- \pi^0$ decay and using $B(\eta \rightarrow \pi^+ \pi^- \pi^0) = (23.1 \pm 0.5) \times 10^{-2}$.

⁸ A simultaneous fit of $e^+ e^- \rightarrow \pi^+ \pi^-$, $\pi^+ \pi^- \pi^0$, $\pi^0 \gamma$, $\eta \gamma$ data.



$$\Gamma(\eta\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$$

$$\Gamma_5/\Gamma \times \Gamma_8/\Gamma$$

$$\Gamma(\pi^0\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$$

$$\Gamma_6/\Gamma \times \Gamma_8/\Gamma$$

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

3.94±0.16 OUR FIT

3.95±0.17 OUR AVERAGE

4.04±0.09±0.19	1	ACHASOV	16A	SND	0.60–1.38 $e^+e^- \rightarrow \pi^0\gamma$
3.75±0.11±0.29	18k	AKHMETSHIN	05	CMD2	0.60–1.38 $e^+e^- \rightarrow \pi^0\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.29±0.11	2	BENAYOUN	10	RVUE	$0.4–1.05 e^+e^-$
-----------	---	----------	----	------	-------------------

3.67±0.10 ^{+0.27} _{-0.25}	3	ACHASOV	00	SND	$e^+e^- \rightarrow \pi^0\gamma$
---	---	---------	----	-----	----------------------------------

¹ From the VMD model with the interfering $\rho(770)$, $\omega(782)$, $\phi(1020)$ resonances, and an additional resonance describing the total contribution of the $\rho(1450)$ and $\omega(1420)$ states. Supersedes ACHASOV 00.

² A simultaneous fit of $e^+e^- \rightarrow \pi^+\pi^-$, $\pi^+\pi^-\pi^0$, $\pi^0\gamma$, $\eta\gamma$ data.

³ From the $\pi^0 \rightarrow 2\gamma$ decay and using $B(\pi^0 \rightarrow 2\gamma) = (98.798 \pm 0.032) \times 10^{-2}$.

$$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$$

$$\Gamma_9/\Gamma \times \Gamma_8/\Gamma$$

VALUE (units 10^{-8})		DOCUMENT ID	TECN	COMMENT
--------------------------	--	-------------	------	---------


8.5 ±0.6 OUR FIT Error includes scale factor of 1.2.

8.8 ±0.9 OUR AVERAGE Error includes scale factor of 1.5. See the ideogram below.

8.36±0.59±0.37	ACHASOV	01G	SND	$e^+e^- \rightarrow \mu^+\mu^-$	
9.9 ±1.4 ±0.9	1	ACHASOV	99C	SND	$e^+e^- \rightarrow \mu^+\mu^-$
14.4 ±3.0	2	VASSERMAN	81	OLYA	$e^+e^- \rightarrow \mu^+\mu^-$
8.6 ±5.9	2	AUGUSTIN	73	OSPK	$e^+e^- \rightarrow \mu^+\mu^-$

¹ Recalculated by the authors from the cross section in the peak.

² Recalculated by us from the cross section in the peak.



$$\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$$

VALUE (units 10^{-8})

2.2 ± 0.4 OUR FIT

2.2 ± 0.4 OUR AVERAGE

$2.1 \pm 0.3 \pm 0.3$

$1.95^{+1.15}_{-0.87}$

$6.01^{+3.19}_{-2.51}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.31 ± 0.99

DOCUMENT ID

TECN

COMMENT

¹ ACHASOV 00C SND $e^+e^- \rightarrow \pi^+\pi^-$

² GOLUBEV 86 ND $e^+e^- \rightarrow \pi^+\pi^-$

² VASSERMAN 81 OLYA $e^+e^- \rightarrow \pi^+\pi^-$

$$\Gamma_{11}/\Gamma \times \Gamma_8/\Gamma$$

$$\Gamma(\omega\pi^0)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$$

$$\Gamma_{12}/\Gamma \times \Gamma_8/\Gamma$$

VALUE (units 10^{-8})

1.40±0.15 OUR FIT

1.37±0.17±0.01

DOCUMENT ID

TECN

COMMENT

¹ Recalculated by the authors from the cross section in the peak.

² Recalculated by us from the cross section in the peak.

³ A simultaneous fit to $e^+e^- \rightarrow \pi^+\pi^-, \pi^+\pi^-\pi^0, \pi^0\gamma, \eta\gamma, K\bar{K}$, and $\tau^- \rightarrow \pi^-\pi^0\nu_\tau$ data.

¹ Recalculated by the authors from the cross section at the peak.

² AMBROSINO 08G reports $[\Gamma(\phi(1020) \rightarrow \omega\pi^0)/\Gamma_{\text{total}} \times \Gamma(\phi(1020) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = (1.22 \pm 0.13 \pm 0.08) \times 10^{-8}$ which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\pi^0\pi^0\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{17}/\Gamma \times \Gamma_8/\Gamma$

VALUE (units 10^{-8})	DOCUMENT ID	TECN	COMMENT
3.34 ± 0.17 OUR FIT			

$3.33^{+0.04+0.19}_{-0.09-0.20}$ ¹ AMBROSINO 07 KLOE $e^+e^- \rightarrow \pi^0\pi^0\gamma$

¹ Calculated by the authors from the cross section at the peak.

$\Gamma(\pi^+\pi^-\pi^+\pi^-)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{18}/\Gamma \times \Gamma_8/\Gamma$

VALUE (units 10^{-9})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$1.2^{+0.8}_{-0.7}$ OUR FIT

$1.17 \pm 0.52 \pm 0.64$ 3285 ¹ AKHMETSHIN 00E CMD2 $e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$

¹ Recalculated by the authors from the cross section in the peak.

$\phi(1020)$ BRANCHING RATIOS

$\Gamma(K^+K^-)/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.491 ± 0.005 OUR FIT Error includes scale factor of 1.3.

0.493 ± 0.010 OUR AVERAGE

0.492 ± 0.012	2913	AKHMETSHIN 95	CMD2	$e^+e^- \rightarrow K^+K^-$
0.44 ± 0.05	321	KALBFLEISCH 76	HBC	$2.18 K^- p \rightarrow \Lambda K^+ K^-$
0.49 ± 0.06	270	DEGROOT 74	HBC	$4.2 K^- p \rightarrow \Lambda \phi$
0.540 ± 0.034	565	BALAKIN 71	OSPK	$e^+e^- \rightarrow K^+K^-$
0.48 ± 0.04	252	LINDSEY	HBC	$2.1-2.7 K^- p \rightarrow \Lambda K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.493 \pm 0.003 \pm 0.007$ ¹ AKHMETSHIN 11 CMD2 $1.02 e^+e^- \rightarrow K^+K^-$

0.476 ± 0.017 1000k ² ACHASOV 01E SND $e^+e^- \rightarrow K^+K^-, K_S K_L, \pi^+\pi^-\pi^0$

¹ Combined analysis of the CMD-2 data on $\phi \rightarrow K^+K^-$, $K_S^0 K_L^0$, $\pi^+\pi^-\pi^0$, $\eta\gamma$ assuming that the sum of their branching fractions is 0.99741 ± 0.00007 .

² Using $B(\phi \rightarrow e^+e^-) = (2.93 \pm 0.14) \times 10^{-4}$.

$\Gamma(K_L^0 K_S^0)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.339 ± 0.004 OUR FIT Error includes scale factor of 1.2.

0.331 ± 0.009 OUR AVERAGE

0.335 ± 0.010	40644	AKHMETSHIN 95	CMD2	$e^+e^- \rightarrow K_L^0 K_S^0$
0.326 ± 0.035		DOLINSKY 91	ND	$e^+e^- \rightarrow K_L^0 K_S^0$
0.310 ± 0.024		DRUZHININ 84	ND	$e^+e^- \rightarrow K_L^0 K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.336 \pm 0.002 \pm 0.006$	1 AKHMETSHIN 11	CMD2	$1.02 e^+ e^- \rightarrow K_S^0 K_L^0$
0.351 ± 0.013	500k	2 ACHASOV 01E	$e^+ e^- \rightarrow K^+ K^-, K_S^0 K_L^0, \pi^+ \pi^- \pi^0$
0.27 ± 0.03	133	KALBFLEISCH 76	$HBC \quad 2.18 K^- p \rightarrow \Lambda K_L^0 K_S^0$
0.257 ± 0.030	95	3 BALAKIN 71	$OSPK \quad e^+ e^- \rightarrow K_L^0 K_S^0$
0.40 ± 0.04	167	LINDSEY	$HBC \quad 2.1-2.7 K^- p \rightarrow \Lambda K_L^0 K_S^0$

1 Combined analysis of the CMD-2 data on $\phi \rightarrow K^+ K^-, K_S^0 K_L^0, \pi^+ \pi^- \pi^0, \eta \gamma$ assuming that the sum of their branching fractions is 0.99741 ± 0.00007 .

2 Using $B(\phi \rightarrow e^+ e^-) = (2.93 \pm 0.14) \times 10^{-4}$.

3 Balakin error increased by Paul.

$\Gamma(K_L^0 K_S^0)/\Gamma(K^+ K^-)$

Γ_2/Γ_1

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.690 ± 0.014 OUR FIT		Error includes scale factor of 1.3.		
0.740 ± 0.031 OUR AVERAGE				

0.70 ± 0.06 2732 BUKIN 78C OLYA $e^+ e^- \rightarrow K_L^0 K_S^0$

0.82 ± 0.08 LOSTY 78 HBC $4.2 K^- p \rightarrow \phi$ hyperon

0.71 ± 0.05 LAVEN 77 HBC $10 K^- p \rightarrow K^+ K^- \Lambda$

0.71 ± 0.08 LYONS 77 HBC $3-4 K^- p \rightarrow \Lambda \phi$

0.89 ± 0.10 144 AGUILAR... 72B HBC $3.9, 4.6 K^- p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.638 ± 0.022 2.3M 1 KOZYREV 18 CMD3 $e^+ e^- \rightarrow K_L^0 K_S^0$

0.68 ± 0.03 2 AKHMETSHIN 95 CMD2 $e^+ e^- \rightarrow K_L^0 K_S^0, K^+ K^-$

1 The prediction taking into account phase-space difference, radiative corrections, isospin breaking, and the Sommerfeld-Gamow-Sakharov factor gives 0.630.

2 Theoretical analysis of BRAMON 00 taking into account phase-space difference, electromagnetic radiative corrections, as well as isospin breaking, predicts 0.62. FLOREZ-BAEZ 08 predicts 0.63 considering also structure-dependent radiative corrections. FIS-CHBACH 02 calculates additional corrections caused by the close threshold and predicts 0.68. See also BENAYOUN 01 and DUBYNSKIY 07. BENAYOUN 12 obtains 0.71 ± 0.01 in the HLS model.

$\Gamma(K_L^0 K_S^0)/\Gamma(K\bar{K})$

$\Gamma_2/(\Gamma_1 + \Gamma_2)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.408 ± 0.005 OUR FIT		Error includes scale factor of 1.3.		
0.45 ± 0.04 OUR AVERAGE				

0.44 ± 0.07 1 LONDON 66 HBC $2.24 K^- p \rightarrow \Lambda K\bar{K}$

0.48 ± 0.07 52 BADIER 65B HBC $3 K^- p$

0.40 ± 0.10 34 SCHLEIN 63 HBC $1.95 K^- p \rightarrow \Lambda K\bar{K}$

1 This is probably not affected by their controversial background subtraction; the value is from their numbers of $K_1 K_2$ vs $K^+ K^-$ events.

$[\Gamma(\rho\pi) + \Gamma(\pi^+\pi^-\pi^0)]/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.154±0.004 OUR FIT				Error includes scale factor of 1.2.
0.151±0.009 OUR AVERAGE				Error includes scale factor of 1.7.
0.161±0.008	11761	AKHMETSHIN 95	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.143±0.007		DOLINSKY 91	ND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.155±0.002±0.005		¹ AKHMETSHIN 11	CMD2	$1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.159±0.008	400k	² ACHASOV 01E	SND	$e^+e^- \rightarrow K^+K^-$, $K_S^0 K_L^0, \pi^+\pi^-\pi^0$
0.145±0.009±0.003	11169	³ AKHMETSHIN 98	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.139±0.007		⁴ PARROUR 76B	OSPK	e^+e^-

¹ Combined analysis of the CMD-2 data on $\phi \rightarrow K^+K^-$, $K_S^0 K_L^0$, $\pi^+\pi^-\pi^0$, $\eta\gamma$ assuming that the sum of their branching fractions is 0.99741 ± 0.00007 .

² Using $B(\phi \rightarrow e^+e^-) = (2.93 \pm 0.14) \times 10^{-4}$.

³ Using $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$.

⁴ Using $\Gamma(\phi) = 4.1$ MeV. If interference between the $\rho\pi$ and 3π modes is neglected, the fraction of the $\rho\pi$ is more than 80% at the 90% confidence level.

 $[\Gamma(\rho\pi) + \Gamma(\pi^+\pi^-\pi^0)]/\Gamma(K^+K^-)$ Γ_3/Γ_1

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.313±0.010 OUR FIT				Error includes scale factor of 1.2.
0.28 ±0.09	34	AGUILAR-...	72B	HBC 3.9,4.6 $K^- p$

 $[\Gamma(\rho\pi) + \Gamma(\pi^+\pi^-\pi^0)]/\Gamma(K\bar{K})$ $\Gamma_3/(\Gamma_1+\Gamma_2)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.185±0.005 OUR FIT				Error includes scale factor of 1.2.
0.24 ±0.04 OUR AVERAGE				
0.237±0.039		CERRADA 77B	HBC	$4.2 K^- p \rightarrow \Lambda 3\pi$
0.30 ±0.15		LONDON 66	HBC	$2.24 K^- p \rightarrow \Lambda \pi^+ \pi^- \pi^0$

 $[\Gamma(\rho\pi) + \Gamma(\pi^+\pi^-\pi^0)]/\Gamma(K_L^0 K_S^0)$ Γ_3/Γ_2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.453±0.012 OUR FIT				Error includes scale factor of 1.1.
0.51 ±0.05 OUR AVERAGE				
0.56 ±0.07	3681	BUKIN 78C	OLYA	$e^+e^- \rightarrow K_L^0 K_S^0, \pi^+\pi^-\pi^0$
0.47 ±0.06	516	COSME 74	OSPK	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$

 $\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$\simeq 0.0087$	1.98M	^{1,2} ALOISIO 03	KLOE	$1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^0$	
<0.0006	90	³ ACHASOV 02	SND	$1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^0$	
<0.23	90	³ CORDIER 80	DM1	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$	
<0.20	90	³ PARROUR 76B	OSPK	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$	

¹ From a fit without limitations on charged and neutral ρ masses and widths.

² Adding the direct and $\omega\pi$ contributions and considering the interference between the $\rho\pi$ and $\pi^+\pi^-\pi^0$.

³ Neglecting the interference between the $\rho\pi$ and $\pi^+\pi^-\pi^0$.

$\Gamma(\eta\gamma)/\Gamma_{\text{total}}$ Γ_5/Γ

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.301 ± 0.024 OUR FIT		Error includes scale factor of 1.2.		
1.26 ± 0.04 OUR AVERAGE				
1.246 ± 0.025 ± 0.057	10k	¹ ACHASOV 98F	SND	$e^+ e^- \rightarrow 7\gamma$
1.18 ± 0.11	279	² AKHMETSHIN 95	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- 3\gamma$
1.30 ± 0.06		³ DRUZHININ 84	ND	$e^+ e^- \rightarrow 3\gamma$
1.4 ± 0.2		⁴ DRUZHININ 84	ND	$e^+ e^- \rightarrow 6\gamma$
0.88 ± 0.20	290	KURDADZE 83C	OLYA	$e^+ e^- \rightarrow 3\gamma$
1.35 ± 0.29		ANDREWS 77	CNTR	6.7–10 γ Cu
1.5 ± 0.4	54	³ COSME 76	OSPK	$e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.38 ± 0.02 ± 0.02		⁵ AKHMETSHIN 11	CMD2	$1.02 e^+ e^- \rightarrow \eta\gamma$
1.36 ± 0.05 ± 0.01	33k	⁶ ACHASOV 07B	SND	$0.6–1.38 e^+ e^- \rightarrow \eta\gamma$
1.373 ± 0.014 ± 0.085	17.4k	^{7,8} AKHMETSHIN 05	CMD2	$0.60–1.38 e^+ e^- \rightarrow \eta\gamma$
1.287 ± 0.013 ± 0.063		^{9,10} AKHMETSHIN 01B	CMD2	$e^+ e^- \rightarrow \eta\gamma$
1.338 ± 0.012 ± 0.052		¹¹ ACHASOV 00	SND	$e^+ e^- \rightarrow \eta\gamma$
1.18 ± 0.03 ± 0.06	2200	¹² AKHMETSHIN 99F	CMD2	$e^+ e^- \rightarrow \eta\gamma$
1.21 ± 0.07		¹³ BENAYOUN 96	RVUE	$0.54–1.04 e^+ e^- \rightarrow \eta\gamma$

¹ Using $B(\phi \rightarrow e^+ e^-) = (2.99 \pm 0.08) \times 10^{-4}$ and $B(\eta \rightarrow 3\pi^0) = (32.2 \pm 0.4) \times 10^{-2}$.

² From $\pi^+ \pi^- \pi^0$ decay mode of η .

³ From 2γ decay mode of η .

⁴ From $3\pi^0$ decay mode of η .

⁵ Combined analysis of the CMD-2 data on $\phi \rightarrow K^+ K^-, K_S^0 K_L^0, \pi^+ \pi^- \pi^0, \eta\gamma$ assuming that the sum of their branching fractions is 0.99741 ± 0.00007 .

⁶ ACHASOV 07B reports $[\Gamma(\phi(1020) \rightarrow \eta\gamma)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow e^+ e^-)] = (4.050 \pm 0.067 \pm 0.118) \times 10^{-6}$ which we divide by our best value $B(\phi(1020) \rightarrow e^+ e^-) = (2.979 \pm 0.033) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Supersedes ACHASOV 00D and ACHASOV 06A.

⁷ Using $B(\phi \rightarrow e^+ e^-) = (2.98 \pm 0.04) \times 10^{-4}$ and $B(\eta \rightarrow \gamma\gamma) = 39.43 \pm 0.26\%$.

⁸ Not independent of the corresponding $\Gamma(e^+ e^-) \times \Gamma(\eta\gamma)/\Gamma_{\text{total}}^2$.

⁹ Using $B(\phi \rightarrow e^+ e^-) = (2.99 \pm 0.08) \times 10^{-4}$ and $B(\eta \rightarrow 3\pi^0) = (32.24 \pm 0.29) \times 10^{-2}$.

¹⁰ The combined fit from 600 to 1380 MeV taking into account $\rho(770)$, $\omega(782)$, $\phi(1020)$, and $\rho(1450)$ (mass and width fixed at 1450 MeV and 310 MeV respectively).

¹¹ From the $\eta \rightarrow 2\gamma$ decay and using $B(\phi \rightarrow e^+ e^-) = (2.99 \pm 0.08) \times 10^{-4}$.

¹² From $\pi^+ \pi^- \pi^0$ decay mode of η and using $B(\phi \rightarrow e^+ e^-) = (2.99 \pm 0.08) \times 10^{-4}$.

¹³ Reanalysis of DRUZHININ 84, DOLINSKY 89, and DOLINSKY 91 taking into account a triangle anomaly contribution.

$\Gamma(\pi^0\gamma)/\Gamma_{\text{total}}$ Γ_6/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.32 ± 0.05 OUR FIT				
1.31 ± 0.13 OUR AVERAGE				
1.30 ± 0.13		DRUZHININ 84	ND	$e^+ e^- \rightarrow 3\gamma$
1.4 ± 0.5	32	COSME 76	OSPK	$e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.367 ± 0.072	¹ ACHASOV	16A	SND	$0.60\text{--}1.38$	$e^+ e^- \rightarrow \pi^0 \gamma$
$1.258 \pm 0.037 \pm 0.077$	18k	^{2,3} AKHMETSHIN	05	CMD2	$0.60\text{--}1.38$ $e^+ e^- \rightarrow \pi^0 \gamma$
$1.226 \pm 0.036^{+0.096}_{-0.089}$		⁴ ACHASOV	00	SND	$e^+ e^- \rightarrow \pi^0 \gamma$
1.26 ± 0.17		⁵ BENAYOUN	96	RVUE	$0.54\text{--}1.04$ $e^+ e^- \rightarrow \pi^0 \gamma$

¹ Using $B(\phi \rightarrow e^+ e^-)$ from PDG 15. Supersedes ACHASOV 00.

² Using $B(\phi \rightarrow e^+ e^-) = (2.98 \pm 0.04) \times 10^{-4}$.

³ Not independent of the corresponding $\Gamma(e^+ e^-) \times \Gamma(\pi^0 \gamma)/\Gamma_{\text{total}}^2$.

⁴ From the $\pi^0 \rightarrow 2\gamma$ decay and using $B(\phi \rightarrow e^+ e^-) = (2.99 \pm 0.08) \times 10^{-4}$.

⁵ Reanalysis of DRUZHININ 84, DOLINSKY 89, and DOLINSKY 91 taking into account a triangle anomaly contribution.

$\Gamma(\eta\gamma)/\Gamma(\pi^0\gamma)$

Γ_5/Γ_6

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$10.9 \pm 0.3^{+0.7}_{-0.8}$	ACHASOV	00	SND $e^+ e^- \rightarrow \eta\gamma, \pi^0\gamma$

$\Gamma(e^+ e^-)/\Gamma_{\text{total}}$

Γ_8/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.979 ± 0.033 OUR FIT	Error includes scale factor of 1.2.			
2.98 ± 0.07 OUR AVERAGE	Error includes scale factor of 1.1.			
2.93 ± 0.14	1900k	¹ ACHASOV	01E	SND $e^+ e^- \rightarrow K^+ K^-, K_S K_L, \pi^+ \pi^- \pi^0$
2.88 ± 0.09	55600	AKHMETSHIN	95	CMD2 $e^+ e^- \rightarrow \text{hadrons}$
3.00 ± 0.21	3681	BUKIN	78C	OLYA $e^+ e^- \rightarrow \text{hadrons}$
3.10 ± 0.14		² PARROUR	76	OSPK $e^+ e^-$
3.3 ± 0.3		COSME	74	OSPK $e^+ e^- \rightarrow \text{hadrons}$
2.81 ± 0.25	681	BALAKIN	71	OSPK $e^+ e^- \rightarrow \text{hadrons}$
3.50 ± 0.27		CHATELUS	71	OSPK $e^+ e^-$

¹ From the combined fit assuming that the total $\phi(1020)$ production cross section is saturated by those of $K^+ K^-$, $K_S K_L$, $\pi^+ \pi^- \pi^0$, and $\eta\gamma$ decays modes and using ACHASOV 00B for the $\eta\gamma$ decay mode.

² Using total width 4.2 MeV. They detect 3π mode and observe significant interference with ω tail. This is accounted for in the result quoted above.

$\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$

Γ_9/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
2.85 ± 0.22 OUR FIT	Error includes scale factor of 1.2.		
2.5 ± 0.4 OUR AVERAGE			
2.69 ± 0.46	¹ HAYES	71	CNTR $8.3, 9.8 \gamma C \rightarrow \mu^+ \mu^- X$
2.17 ± 0.60	¹ EARLES	70	CNTR $6.0 \gamma C \rightarrow \mu^+ \mu^- X$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2.87 $\pm 0.20 \pm 0.14$	² ACHASOV	01G	SND $e^+ e^- \rightarrow \mu^+ \mu^-$
3.30 $\pm 0.45 \pm 0.32$	³ ACHASOV	99C	SND $e^+ e^- \rightarrow \mu^+ \mu^-$
4.83 ± 1.02	⁴ VASSERMAN	81	OLYA $e^+ e^- \rightarrow \mu^+ \mu^-$
2.87 ± 1.98	⁴ AUGUSTIN	73	OSPK $e^+ e^- \rightarrow \mu^+ \mu^-$

¹ Neglecting interference between resonance and continuum.

² Using $B(\phi \rightarrow e^+ e^-) = (2.91 \pm 0.07) \times 10^{-4}$.

³ Using $B(\phi \rightarrow e^+ e^-) = (2.99 \pm 0.08) \times 10^{-4}$.

⁴ Recalculated by us using $B(\phi \rightarrow e^+ e^-) = (2.99 \pm 0.08) \times 10^{-4}$.

$\Gamma(\eta e^+ e^-)/\Gamma_{\text{total}}$

Γ_{10}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.08 ± 0.04 OUR AVERAGE				
1.075 ± 0.007 ± 0.038	30k	¹ BABUSCI 15	KLOE	$1.02 e^+ e^- \rightarrow \eta e^+ e^-$
1.19 ± 0.19 ± 0.12	213	² ACHASOV 01B	SND	$e^+ e^- \rightarrow \eta e^+ e^-$
1.14 ± 0.10 ± 0.06	355	³ AKHMETSHIN 01	CMD2	$e^+ e^- \rightarrow \eta e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.13 ± 0.14 ± 0.07	183	⁴ AKHMETSHIN 01	CMD2	$e^+ e^- \rightarrow \eta e^+ e^-$
1.21 ± 0.14 ± 0.09	130	⁵ AKHMETSHIN 01	CMD2	$e^+ e^- \rightarrow \eta e^+ e^-$
1.04 ± 0.20 ± 0.08	42	⁶ AKHMETSHIN 01	CMD2	$e^+ e^- \rightarrow \eta e^+ e^-$
1.3 +0.8 -0.6	7	GOLUBEV 85	ND	$e^+ e^- \rightarrow \eta e^+ e^-$

¹ Using $B(\eta \rightarrow 3\pi^0) = (32.57 \pm 0.23)\%$ from PDG 12.

² Using $B(\eta \rightarrow \gamma\gamma) = (39.25 \pm 0.32)\%$, $B(\phi \rightarrow \eta\gamma) = (1.26 \pm 0.06)\%$, and $B(\phi \rightarrow e^+ e^-) = (3.00 \pm 0.06) \times 10^{-4}$.

³ The average of the branching ratios separately obtained from the $\eta \rightarrow \gamma\gamma$, $3\pi^0$, $\pi^+\pi^-\pi^0$ decays.

⁴ From $\eta \rightarrow \gamma\gamma$ decays and using $B(\eta \rightarrow \gamma\gamma) = (39.33 \pm 0.25) \times 10^{-2}$, $B(\eta \rightarrow \pi^+\pi^-\gamma) = (4.75 \pm 11) \times 10^{-2}$, and $B(\phi \rightarrow \eta\gamma) = (1.297 \pm 0.033) \times 10^{-2}$.

⁵ From $\eta \rightarrow 3\pi^0$ decays and using $B(\pi^0 \rightarrow \gamma\gamma) = (98.798 \pm 0.033) \times 10^{-2}$, $B(\eta \rightarrow 3\pi^0) = (32.24 \pm 0.29) \times 10^{-2}$, $B(\eta \rightarrow \pi^+\pi^-\gamma) = (4.75 \pm 0.11) \times 10^{-2}$, and $B(\phi \rightarrow \eta\gamma) = (1.297 \pm 0.033) \times 10^{-2}$.

⁶ From $\eta \rightarrow \pi^+\pi^-\pi^0$ decays and using $B(\pi^0 \rightarrow \gamma\gamma) = (98.798 \pm 0.033) \times 10^{-2}$, $B(\pi^0 \rightarrow e^+e^-\gamma) = (1.198 \pm 0.032) \times 10^{-2}$, $B(\eta \rightarrow \pi^+\pi^-\pi^0) = (23.0 \pm 0.4) \times 10^{-2}$, $B(\phi \rightarrow \pi^+\pi^-\pi^0) = (15.5 \pm 0.6) \times 10^{-2}$, and $B(\phi \rightarrow \eta\gamma) = (1.297 \pm 0.033) \times 10^{-2}$.

$\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$

Γ_{11}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.71 ± 0.11 ± 0.09		¹ ACHASOV 00C	SND	$e^+ e^- \rightarrow \pi^+\pi^-$
0.65 +0.38 -0.29		¹ GOLUBEV 86	ND	$e^+ e^- \rightarrow \pi^+\pi^-$
2.01 +1.07 -0.84		¹ VASSERMAN 81	OLYA	$e^+ e^- \rightarrow \pi^+\pi^-$
<6.6	95	BUKIN 78B	OLYA	$e^+ e^- \rightarrow \pi^+\pi^-$
<2.7	95	ALVENSLEB... 72	CNTR	$6.7 \gamma C \rightarrow C\pi^+\pi^-$

¹ Using $B(\phi \rightarrow e^+ e^-) = (2.99 \pm 0.08) \times 10^{-4}$.

$\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$

Γ_{12}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
4.7 ± 0.5 OUR FIT			
5.2 +1.3 -1.1	^{1,2} AULCHENKO 00A	SND	$e^+ e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.4 ± 0.6	³ AMBROSINO 08G KLOE $e^+ e^- \rightarrow \pi^+ \pi^- 2\pi^0, 2\pi^0 \gamma$
~ 5.4	⁴ ACHASOV 00E SND $e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
$5.5^{+1.6}_{-1.4} \pm 0.3$	^{2,5} AULCHENKO 00A SND $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \pi^0$
$4.8^{+1.9}_{-1.7} \pm 0.8$	⁴ ACHASOV 99 SND $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \pi^0$

¹ Using the 1996 and 1998 data.

² $(2.3 \pm 0.3)\%$ correction for other decay modes of the $\omega(782)$ applied.

³ Not independent of the corresponding $\Gamma(\omega\pi^0) \times \Gamma(e^+ e^-) / \Gamma^2(\text{total})$.

⁴ Using the 1996 data.

⁵ Using the 1998 data.

$\Gamma(\omega\gamma)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{13}/Γ
<0.05	84	LINDSEY	66	HBC	$2.1-2.7 K^- p \rightarrow \Lambda \pi^+ \pi^- \text{ neutrals}$

$\Gamma(\rho\gamma)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{14}/Γ
< 0.12	90	¹ AKHMETSHIN 99B	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \gamma$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 7	90	AKHMETSHIN 97C	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \gamma$
-----	----	----------------	------	--

<200	84	LINDSEY	66	HBC	$2.1-2.7 K^- p \rightarrow \Lambda \pi^+ \pi^- \text{ neutrals}$
------	----	---------	----	-----	--

¹ Supersedes AKHMETSHIN 97C.

$\Gamma(\pi^+ \pi^- \gamma)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{15}/Γ
$0.41 \pm 0.12 \pm 0.04$	30175	¹ AKHMETSHIN 99B	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \gamma$		

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.3	90	² AKHMETSHIN 97C	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \gamma$
<600	90	KALBFLEISCH 75	HBC	$2.18 K^- p \rightarrow \Lambda \pi^+ \pi^- \gamma$
< 70	90	COSME	74	OSPK $e^+ e^- \rightarrow \pi^+ \pi^- \gamma$
<400	90	LINDSEY	65	HBC $2.1-2.7 K^- p \rightarrow \Lambda \pi^+ \pi^- \text{ neutrals}$

¹ For $E_\gamma > 20$ MeV and assuming that $B(\phi(1020) \rightarrow f_0(980)\gamma)$ is negligible. Supersedes AKHMETSHIN 97C.

² For $E_\gamma > 20$ MeV and assuming that $B(\phi(1020) \rightarrow f_0(980)\gamma)$ is negligible.

$\Gamma(f_0(980)\gamma)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{16}/Γ
--------------------------	-----	------	-------------	------	---------	----------------------

$3.22 \pm 0.19 \text{ OUR FIT}$ Error includes scale factor of 1.1.

$3.21 \pm 0.19 \text{ OUR AVERAGE}$

$3.21^{+0.03}_{-0.09} \pm 0.18$	¹ AMBROSINO 07 KLOE $e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
$2.90 \pm 0.21 \pm 1.54$	² AKHMETSHIN 99C CMD2 $e^+ e^- \rightarrow \pi^+ \pi^- \gamma, \pi^0 \pi^0 \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.47 ± 0.21	2438	³ ALOISIO	02D	KLOE	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
3.5 ± 0.3 $^{+1.3}_{-0.5}$	419	^{4,5} ACHASOV	00H	SND	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
$1.93 \pm 0.46 \pm 0.50$	27188	⁶ AKHMETSHIN 99B	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \gamma$	
$3.05 \pm 0.25 \pm 0.72$	268	⁷ AKHMETSHIN 99C	CMD2	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$	
1.5 ± 0.5	268	⁸ AKHMETSHIN 99C	CMD2	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$	
$3.42 \pm 0.30 \pm 0.36$	164	⁴ ACHASOV	98I	SND	$e^+ e^- \rightarrow 5\gamma$
< 1	90	⁹ AKHMETSHIN 97C	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \gamma$	
< 7	90	¹⁰ AKHMETSHIN 97C	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \gamma$	
< 20	90	DRUZHININ	87	ND	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$

¹ Obtained by the authors taking into account the $\pi^+ \pi^-$ decay mode. Includes a component due to $\pi\pi$ production via the $f_0(500)$ meson. Supersedes ALOISIO 02D.

² From the combined fit of the photon spectra in the reactions $e^+ e^- \rightarrow \pi^+ \pi^- \gamma$, $\pi^0 \pi^0 \gamma$.

³ From the negative interference with the $f_0(500)$ meson of AITALA 01B using the ACHASOV 89 parameterization for the $f_0(980)$, a Breit-Wigner for the $f_0(500)$, and ACHASOV 01F for the $\rho\pi$ contribution. Superseded by AMBROSINO 07.

⁴ Assuming that the $\pi^0 \pi^0 \gamma$ final state is completely determined by the $f_0 \gamma$ mechanism, neglecting the decay $B(\phi \rightarrow K\bar{K}\gamma)$ and using $B(f_0 \rightarrow \pi^+ \pi^-) = 2B(f_0 \rightarrow \pi^0 \pi^0)$.

⁵ Using the value $B(\phi \rightarrow \eta\gamma) = (1.338 \pm 0.053) \times 10^{-2}$.

⁶ For $E_\gamma > 20$ MeV. Supersedes AKHMETSHIN 97C.

⁷ Neglecting other intermediate mechanisms ($\rho\pi$, $\sigma\gamma$).

⁸ A narrow pole fit taking into account $f_0(980)$ and $f_0(1200)$ intermediate mechanisms.

⁹ For destructive interference with the Bremsstrahlung process

¹⁰ For constructive interference with the Bremsstrahlung process

$\Gamma(f_0(980)\gamma)/\Gamma(\eta\gamma)$

Γ_{16}/Γ_5

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.48 ± 0.15 OUR FIT		Error includes scale factor of 1.1.		

2.6 ± 0.2 $^{+0.8}_{-0.3}$	419	¹ ACHASOV	00H	SND	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
--	-----	----------------------	-----	-----	--

¹ Assuming that the $\pi^0 \pi^0 \gamma$ final state is completely determined by the $f_0 \gamma$ mechanism, neglecting the decay $B(\phi \rightarrow K\bar{K}\gamma)$ and using $B(f_0 \rightarrow \pi^+ \pi^-) = 2B(f_0 \rightarrow \pi^0 \pi^0)$.

$\Gamma(\pi^0 \pi^0 \gamma)/\Gamma_{\text{total}}$

Γ_{17}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.07 ± 0.06 OUR AVERAGE					

1.07 $^{+0.01}_{-0.03}$ $^{+0.06}_{-0.06}$		¹ AMBROSINO 07	KLOE	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
--	--	---------------------------	------	--

$1.08 \pm 0.17 \pm 0.09$	268	AKHMETSHIN 99C	CMD2	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
--------------------------	-----	----------------	------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.09 \pm 0.03 \pm 0.05$	2438	ALOISIO	02D	KLOE	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
--------------------------	------	---------	-----	------	--

$1.158 \pm 0.093 \pm 0.052$	419	^{2,3} ACHASOV	00H	SND	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
-----------------------------	-----	------------------------	-----	-----	--

< 10	90	DRUZHININ	87	ND	$e^+ e^- \rightarrow 5\gamma$
--------	----	-----------	----	----	-------------------------------

¹ Supersedes ALOISIO 02D.

² Using the value $B(\phi \rightarrow \eta\gamma) = (1.338 \pm 0.053) \times 10^{-2}$.

³ Supersedes ACHASOV 98I. Excluding $\omega\pi^0$.

$\Gamma(\pi^0\pi^0\gamma)/\Gamma(\eta\gamma)$ Γ_{17}/Γ_5

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.86 ± 0.04 OUR FIT				
0.865 ± 0.070 ± 0.017	419	¹ ACHASOV	00H SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.90 ± 0.08 ± 0.07	164	ACHASOV	98I SND	$e^+e^- \rightarrow 5\gamma$
1 Supersedes ACHASOV 98I. Excluding $\omega\pi^0$.				

 $\Gamma(\pi^+\pi^-\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{18}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
6.5 ± 2.7 ± 1.6	6.8k	¹ AKHMETSHIN 17	CMD3	$e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
3.93 ± 1.74 ± 2.14	3.3k	AKHMETSHIN 00E	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$	
< 870	90	CORDIER	79	WIRE	$e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$

¹ Using the cross section at the ϕ meson peak $\sigma(\phi) = 4172 \pm 42$ nb, the nonresonant cross section $\sigma(0) = 1.263 \pm 0.027$ nb and $\text{Re}(Z) = 0.146 \pm 0.030$, $\text{Im}(Z) = -0.002 \pm 0.024$ for the complex amplitude of the $\phi \rightarrow \pi^+\pi^-\pi^+\pi^-$ transition.

 $\Gamma(\pi^+\pi^+\pi^-\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{19}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 4.6	90	AKHMETSHIN 00E	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-\pi^0$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 150	95	BARKOV	88	CMD	$e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-\pi^0$

 $\Gamma(\pi^0e^+e^-)/\Gamma_{\text{total}}$ Γ_{20}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.33 ± 0.07 OUR AVERAGE					

1.35 ± 0.05 ± 0.05	9.5k	¹ ANASTASI	16B KLOE	$e^+e^- \rightarrow \pi^0e^+e^-$
1.01 ± 0.28 ± 0.29	52	² ACHASOV	02D SND	$e^+e^- \rightarrow \pi^0e^+e^-$
1.22 ± 0.34 ± 0.21	46	³ AKHMETSHIN 01C	CMD2	$e^+e^- \rightarrow \pi^0e^+e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 12	90	DOLINSKY	88 ND	$e^+e^- \rightarrow \pi^0e^+e^-$

¹ Using $B(\pi^0 \rightarrow \gamma\gamma)$ from the 2014 Edition of this Review (PDG 14).

² Using various branching ratios from the 2000 Edition of this Review (PDG 00).

³ Using $B(\pi^0 \rightarrow \gamma\gamma) = 0.98798 \pm 0.00032$, $B(\phi \rightarrow \eta\gamma) = (1.297 \pm 0.033) \times 10^{-2}$, and $B(\eta \rightarrow \pi^+\pi^-\gamma) = (4.75 \pm 0.11) \times 10^{-2}$.

 $\Gamma(\pi^0\eta\gamma)/\Gamma_{\text{total}}$ Γ_{21}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
7.27 ± 0.30 OUR AVERAGE					Error includes scale factor of 1.5. See the ideogram below.
7.06 ± 0.22	16.9k	¹ AMBROSINO	09F KLOE	1.02 $e^+e^- \rightarrow \eta\pi^0\gamma$	
8.51 ± 0.51 ± 0.57	607	² ALOISIO	02C KLOE	$e^+e^- \rightarrow \eta\pi^0\gamma$	
7.96 ± 0.60 ± 0.40	197	³ ALOISIO	02C KLOE	$e^+e^- \rightarrow \eta\pi^0\gamma$	
8.8 ± 1.4 ± 0.9	36	⁴ ACHASOV	00F SND	$e^+e^- \rightarrow \eta\pi^0\gamma$	
9.0 ± 2.4 ± 1.0	80	AKHMETSHIN 99C	CMD2	$e^+e^- \rightarrow \eta\pi^0\gamma$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

$7.01 \pm 0.10 \pm 0.20$	13.3k	2,5 AMBROSINO 09F	KLOE	$1.02 e^+ e^- \rightarrow \eta \pi^0 \gamma$
$7.12 \pm 0.13 \pm 0.22$	3.6k	3,6 AMBROSINO 09F	KLOE	$1.02 e^+ e^- \rightarrow \eta \pi^0 \gamma$
$8.3 \pm 2.3 \pm 1.2$	20	ACHASOV 98B	SND	$e^+ e^- \rightarrow 5\gamma$
<250	90	DOLINSKY	ND	$e^+ e^- \rightarrow \pi^0 \eta \gamma$

¹ Combined results of $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow \pi^+ \pi^- \pi^0$ decay modes measurements.

² From the decay mode $\eta \rightarrow \gamma\gamma$.


³ From the decay mode $\eta \rightarrow \pi^+ \pi^- \pi^0$.

⁴ Supersedes ACHASOV 98B.

⁵ Using $B(\phi \rightarrow \eta\gamma) = (1.304 \pm 0.025)\%$, $B(\eta \rightarrow 3\pi^0) = (32.56 \pm 0.23)\%$, and $B(\eta \rightarrow \gamma\gamma) = (39.31 \pm 0.20)\%$.

⁶ Using $B(\phi \rightarrow \eta\gamma) = (1.304 \pm 0.025)\%$, $B(\eta \rightarrow 3\pi^0) = (32.56 \pm 0.23)\%$, and $B(\eta \rightarrow \pi^+ \pi^- \pi^0) = (22.73 \pm 0.28)\%$.

WEIGHTED AVERAGE
7.27±0.30 (Error scaled by 1.5)



$\Gamma(a_0(980)\gamma)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
7.6±0.6 OUR FIT					
7.6±0.6 OUR AVERAGE					
7.4±0.7			¹ ALOISIO 02C	KLOE	$e^+ e^- \rightarrow \eta \pi^0 \gamma$
8.8±1.7		36	² ACHASOV 00F	SND	$e^+ e^- \rightarrow \eta \pi^0 \gamma$
11 ± 2			³ GOKALP 02	RVUE	$e^+ e^- \rightarrow \eta \pi^0 \gamma$
<500		90	DOLINSKY	ND	$e^+ e^- \rightarrow \pi^0 \eta \gamma$

¹ Using $M_{a_0(980)}=984.8$ MeV and assuming $a_0(980)\gamma$ dominance.

² Assuming $a_0(980)\gamma$ dominance in the $\eta \pi^0 \gamma$ final state.

³ Using data of ACHASOV 00F.

$\Gamma(f_0(980)\gamma)/\Gamma(a_0(980)\gamma)$ Γ_{16}/Γ_{22}

VALUE	DOCUMENT ID	TECN	COMMENT
6.1±0.6	¹ ALOISIO 02C	KLOE	$e^+ e^- \rightarrow \eta\pi^0\gamma$

¹ Using results of ALOISIO 02D and assuming that $f_0(980)$ decays into $\pi\pi$ only and $a_0(980)$ into $\eta\pi$ only.

 $\Gamma(K^0\bar{K}^0\gamma)/\Gamma_{\text{total}}$ Γ_{23}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.9 \times 10^{-8}$	90	AMBROSINO 09C	KLOE	$e^+ e^- \rightarrow K_S^0 \bar{K}_S^0 \gamma$

 $\Gamma(\eta'(958)\gamma)/\Gamma_{\text{total}}$ Γ_{24}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
6.21±0.20 OUR FIT					
6.21±0.29 OUR AVERAGE					

$6.21 \pm 0.27 \pm 0.11$	3407	¹ AMBROSINO 07A	KLOE	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- 7\gamma$
$6.7 \begin{array}{l} +2.8 \\ -2.4 \end{array} \pm 0.8$	12	² AULCHENKO 03B	SND	$e^+ e^- \rightarrow \eta' \gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$6.7 \begin{array}{l} +5.0 \\ -4.2 \end{array} \pm 1.5$	7	AULCHENKO 03B	SND	$e^+ e^- \rightarrow 7\gamma$
$6.10 \pm 0.61 \pm 0.43$	120	³ ALOISIO 02E	KLOE	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- 3\gamma$
$8.2 \begin{array}{l} +2.1 \\ -1.9 \end{array} \pm 1.1$	21	⁴ AKHMETSHIN 00B	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- 3\gamma$
$4.9 \begin{array}{l} +2.2 \\ -1.8 \end{array} \pm 0.6$	9	⁵ AKHMETSHIN 00F	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^+ \pi^- \geq 2\gamma$
6.4 ± 1.6	30	⁶ AKHMETSHIN 00F	CMD2	$e^+ e^- \rightarrow \eta'(958)\gamma$
$6.7 \begin{array}{l} +3.4 \\ -2.9 \end{array} \pm 1.0$	5	⁷ AULCHENKO 99	SND	$e^+ e^- \rightarrow \pi^+ \pi^- 3\gamma$
<11	90	AULCHENKO 98	SND	$e^+ e^- \rightarrow 7\gamma$
$12 \begin{array}{l} +7 \\ -5 \end{array} \pm 2$	6	⁴ AKHMETSHIN 97B	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- 3\gamma$
<41	90	DRUZHININ 87	ND	$e^+ e^- \rightarrow \gamma\eta\pi^+\pi^-$

¹ AMBROSINO 07A reports $[\Gamma(\phi(1020) \rightarrow \eta'(958)\gamma)/\Gamma_{\text{total}}] / [\mathcal{B}(\phi(1020) \rightarrow \eta\gamma)] = (4.77 \pm 0.09 \pm 0.19) \times 10^{-3}$ which we multiply by our best value $\mathcal{B}(\phi(1020) \rightarrow \eta\gamma) = (1.301 \pm 0.024) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Averaging AULCHENKO 03B with AULCHENKO 99.

³ Using $\mathcal{B}(\phi \rightarrow \eta\gamma) = (1.297 \pm 0.033)\%$.

⁴ Using the value $\mathcal{B}(\phi \rightarrow \eta\gamma) = (1.26 \pm 0.06) \times 10^{-2}$.

⁵ Using $\mathcal{B}(\phi \rightarrow K_L^0 K_S^0) = (33.8 \pm 0.6)\%$.

⁶ Averaging AKHMETSHIN 00B with AKHMETSHIN 00F.

⁷ Using the value $\mathcal{B}(\eta' \rightarrow \eta\pi^+\pi^-) = (43.7 \pm 1.5) \times 10^{-2}$ and $\mathcal{B}(\eta \rightarrow \gamma\gamma) = (39.25 \pm 0.31) \times 10^{-2}$.

$\Gamma(\eta'(958)\gamma)/\Gamma(K_L^0 K_S^0)$ Γ_{24}/Γ_2

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.83 ± 0.06 OUR FIT				
$1.46^{+0.64}_{-0.54} \pm 0.18$	9	¹ AKHMETSHIN 00F	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^+ \pi^- \geq 2\gamma$

¹ Using various branching ratios of K_S^0 , K_L^0 , η , η' from the 2000 edition (The European Physical Journal **C15** 1 (2000)) of this Review.

 $\Gamma(\eta'(958)\gamma)/\Gamma(\eta\gamma)$ Γ_{24}/Γ_5

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
4.77 ± 0.15 OUR FIT				
4.78 ± 0.20 OUR AVERAGE				
$4.77 \pm 0.09 \pm 0.19$	3407	AMBROSINO 07A	KLOE	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- 7\gamma$
$4.70 \pm 0.47 \pm 0.31$	120	¹ ALOISIO 02E	KLOE	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- 3\gamma$
$6.5^{+1.7}_{-1.5} \pm 0.8$	21	AKHMETSHIN 00B	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- 3\gamma$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$9.5^{+5.2}_{-4.0} \pm 1.4$	6	² AKHMETSHIN 97B	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- 3\gamma$

¹ From the decay mode $\eta' \rightarrow \eta \pi^+ \pi^-$, $\eta \rightarrow \gamma \gamma$.

² Superseded by AKHMETSHIN 00B.

 $\Gamma(\eta\pi^0\pi^0\gamma)/\Gamma_{\text{total}}$ Γ_{25}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2	90	AULCHENKO 98	SND	$e^+ e^- \rightarrow 7\gamma$

 $\Gamma(\mu^+\mu^-\gamma)/\Gamma_{\text{total}}$ Γ_{26}/Γ

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.43 \pm 0.45 \pm 0.14$	27188	¹ AKHMETSHIN 99B	CMD2	$e^+ e^- \rightarrow \mu^+ \mu^- \gamma$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				

2.3 ± 1.0 824 ± 33 ² AKHMETSHIN 97C CMD2 $e^+ e^- \rightarrow \mu^+ \mu^- \gamma$

¹ For $E_\gamma > 20$ MeV. Supersedes AKHMETSHIN 97C.

² For $E_\gamma > 20$ MeV.

 $\Gamma(\rho\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{27}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.2	90	AULCHENKO 08	CMD2	$\phi \rightarrow \pi^+ \pi^- \gamma \gamma$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				

<5 90 AKHMETSHIN 98 CMD2 $e^+ e^- \rightarrow \pi^+ \pi^- \gamma \gamma$

 $\Gamma(\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{28}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 1.8	90	AKHMETSHIN 00E	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^+ \pi^- \pi^0$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				

< 6.1 90 AULCHENKO 08 CMD2 $\phi \rightarrow \eta\pi^+\pi^-$

<30 90 AKHMETSHIN 98 CMD2 $e^+ e^- \rightarrow \pi^+ \pi^- \gamma \gamma$

$\Gamma(\eta\mu^+\mu^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{29}/Γ
<9.4	90	AKHMETSHIN 01	CMD2	$e^+e^- \rightarrow \eta e^+e^-$	

$\Gamma(\eta U \rightarrow \eta e^+e^-)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{30}/Γ
$<1 \times 10^{-6}$	90	¹ BABUSCI	13B	KLOE	$1.02 e^+e^- \rightarrow \eta e^+e^-$

¹ For a narrow vector U with mass between 5 and 470 MeV, from the combined analysis of $\eta \rightarrow \pi^+\pi^-\pi^0$ and $\eta \rightarrow \pi^0\pi^0\pi^0$ from ARCHILLI 12. Measured 90% CL limits as a function of m_U range from 2.2×10^{-8} to 10^{-6} .

$\Gamma(\text{invisible})/\Gamma(K^+K^-)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{31}/Γ_1
$<3.4 \times 10^{-4}$	90	ABLIKIM	18S	BES3	$J/\psi \rightarrow \phi\eta \rightarrow \phi\pi^+\pi^-\pi^0$

———— Lepton Family number (LF) violating modes ——

$\Gamma(e^\pm\mu^\mp)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{32}/Γ
$<2 \times 10^{-6}$	90	ACHASOV	10A	SND	$e^+e^- \rightarrow e^\pm\mu^\mp$

$\pi^+\pi^-\pi^0 / \rho\pi$ AMPLITUDE RATIO a_1 IN DECAY OF $\phi \rightarrow \pi^+\pi^-\pi^0$

NIECKNIG 12 describes final-state interactions between the three pions in a dispersive framework using data on the $\pi\pi$ P -wave scattering phase shift.

VALUE (units 10^{-2})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
9.1 ± 1.2 OUR AVERAGE					

$10.1 \pm 4.4 \pm 1.7$	80k	¹ AKHMETSHIN 06	CMD2	$1.017-1.021 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
$9.0 \pm 1.1 \pm 0.6$	1.98M	^{2,3} ALOISIO	03	KLOE $1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$-6 < a_1 < 6$	500k	³ ACHASOV	02	SND $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
$-16 < a_1 < 11$	90	9.8k	^{1,4} AKHMETSHIN 98	CMD2 $e^+e^- \rightarrow \pi^+\pi^-\gamma\gamma$

¹ Dalitz plot analysis taking into account interference between the contact and $\rho\pi$ amplitudes.

² From a fit without limitations on charged and neutral ρ masses and widths.

³ Recalculated by us to match the notations of AKHMETSHIN 98.

⁴ Assuming zero phase for the contact term.

PARAMETER β IN $\phi \rightarrow Pe^+e^-$ DECAYS

In the one-pole approximation the electromagnetic transition form factor for $\phi \rightarrow Pe^+e^-$ ($P = \pi, \eta$) is given as a function of the e^+e^- invariant mass squared, q^2 , by the expression:

$$|F(q^2)|^2 = (1 - q^2/\Lambda^2)^{-2},$$

where vector meson dominance predicts parameter $\Lambda \approx 0.770$ GeV ($\Lambda^{-2} \approx 1.687$ GeV $^{-2}$). The slope of this form factor, $\beta = dF/dq^2(q^2=0)$, equals Λ^{-2} in this approximation.

The measurements below obtain β in the one-pole approximation.

PARAMETER β IN $\phi \rightarrow \pi^0 e^+e^-$ DECAY

VALUE (GeV $^{-2}$)	EVTS	DOCUMENT ID	TECN	COMMENT
2.02±0.11	9.5k	¹ ANASTASI	16B KLOE	$1.02 e^+e^- \rightarrow \pi^0 e^+e^-$

¹ The error combines statistical and systematic uncertainties.

PARAMETER β IN $\phi \rightarrow \eta e^+e^-$ DECAY

VALUE (GeV $^{-2}$)	EVTS	DOCUMENT ID	TECN	COMMENT
1.29±0.13 OUR AVERAGE				
1.28 ± 0.10 ^{+0.09} _{-0.08}	30k	BABUSCI	15 KLOE	$1.02 e^+e^- \rightarrow \eta e^+e^-$
3.8 ± 1.8	213	¹ ACHASOV	01B SND	$1.02 e^+e^- \rightarrow \eta e^+e^-$

¹ The uncertainty is statistical only. The systematic one is negligible, in comparison.

φ(1020) REFERENCES

LEES	21B	PR D104 112003	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ALBRECHT	20	EPJ C80 453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)
HOID	20	EPJ C80 988	B.-L. Hoid, M. Hoferichter, B. Kubis	(BONN, BERN)
AAIJ	19H	JHEP 1904 063	R. Aaij <i>et al.</i>	(LHCb Collab.)
HOFERICHT...	19	JHEP 1908 137	M. Hoferichter, B.-L. Hoid, B. Kubis	(WASH, BONN)
ABLIKIM	18S	PR D98 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AOUDE	18	PR D98 056021	R.T. Aoude <i>et al.</i>	
KOZYREV	18	PL B779 64	E.A. Kozyrev <i>et al.</i>	(CMD-3 Collab.)
AKHMETSHIN	17	PL B768 345	R.R. Akhmetshin <i>et al.</i>	(CMD-3 Collab.)
ACHASOV	16A	PR D93 092001	M.N. Achasov <i>et al.</i>	(SND Collab.)
ANASTASI	16B	PL B757 362	A. Anastasi <i>et al.</i>	(KLOE-2 Collab.)
KOZYREV	16	PL B760 314	E.A. Kozyrev <i>et al.</i>	(CMD-3 Collab.)
BABUSCI	15	PL B742 1	D. Babusci <i>et al.</i>	(KLOE-2 Collab.)
PDG	15	RPP 2015 at pdg.lbl.gov		(PDG Collab.)
LEES	14H	PR D89 092002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
PDG	14	CP C38 070001	K. Olive <i>et al.</i>	(PDG Collab.)
BABUSCI	13B	PL B720 111	D. Babusci <i>et al.</i>	(KLOE-2 Collab.)
BENAYOUN	13	EPJ C73 2453	M. Benayoun, P. David, L. DelBuono	(PARIN, BERLIN+)
LEES	13F	PR D87 052010	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13Q	PR D88 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ARCHILLI	12	PL B706 251	F. Archilli <i>et al.</i>	(KLOE-2 Collab.)
BENAYOUN	12	EPJ C72 1848	M. Benayoun <i>et al.</i>	
NIECKNIG	12	EPJ C72 2014	F. Niecknig, B. Kubis, S.P. Schneider	(BONN)
PDG	12	PR D86 010001	J. Beringer <i>et al.</i>	(PDG Collab.)
AKHMETSHIN	11	PL B695 412	R.R. Akhmetshin <i>et al.</i>	(CMD-2 Collab.)
ACHASOV	10A	PR D81 057102	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
BENAYOUN	10	EPJ C65 211	M. Benayoun <i>et al.</i>	
AMBROSINO	09C	PL B679 10	F. Ambrosino <i>et al.</i>	(KLOE Collab.)
AMBROSINO	09F	PL B681 5	F. Ambrosino <i>et al.</i>	(KLOE Collab.)
AKHMETSHIN	08	PL B669 217	R.R. Akhmetshin <i>et al.</i>	(CMD-2 Collab.)
AMBROSINO	08G	PL B669 223	F. Ambrosino <i>et al.</i>	(KLOE Collab.)
AULCHENKO	08	JETPL 88 85	V. Aulchenko <i>et al.</i>	(CMD-2 Collab.)

Translated from ZETFP 88 93.

FLOREZ-BAEZ	08	PR D78 077301	F.V. Florez-Baez, G. Lopez Castro
ACHASOV	07B	PR D76 077101	M.N. Achasov <i>et al.</i> (SND Collab.)
AMBROSINO	07	EPJ C49 473	F. Ambrosino <i>et al.</i> (KLOE Collab.)
AMBROSINO	07A	PL B648 267	F. Ambrosino <i>et al.</i> (KLOE Collab.)
DUBYNSKIY	07	PR D75 113001	S. Dubynskiy <i>et al.</i>
ACHASOV	06A	PR D74 014016	M.N. Achasov <i>et al.</i> (SND Collab.)
AKHMETSHIN	06	PL B642 203	R.R. Akhmetshin <i>et al.</i> (CMD-2 Collab.)
AKHMETSHIN	05	PL B605 26	R.R. Akhmetshin <i>et al.</i> (Novosibirsk CMD-2 Collab.)
AMBROSINO	05	PL B608 199	F. Ambrosino <i>et al.</i> (KLOE Collab.)
AUBERT,B	05J	PR D72 052008	B. Aubert <i>et al.</i> (BABAR Collab.)
AKHMETSHIN	04	PL B578 285	R.R. Akhmetshin <i>et al.</i> (Novosibirsk CMD-2 Collab.)
AUBERT,B	04N	PR D70 072004	B. Aubert <i>et al.</i> (BABAR Collab.)
ALOISIO	03	PL B561 55	A. Aloisio <i>et al.</i> (KLOE Collab.)
AULCHENKO	03B	JETP 97 24	V.M. Aulchenko <i>et al.</i> (Novosibirsk SND Collab.)
		Translated from ZETFP 124 28.	
ACHASOV	02	PR D65 032002	M.N. Achasov <i>et al.</i> (Novosibirsk SND Collab.)
ACHASOV	02D	JETPL 75 449	M.N. Achasov <i>et al.</i> (Novosibirsk SND Collab.)
		Translated from ZETFP 75 539.	
ALOISIO	02C	PL B536 209	A. Aloisio <i>et al.</i> (KLOE Collab.)
ALOISIO	02D	PL B537 21	A. Aloisio <i>et al.</i> (KLOE Collab.)
ALOISIO	02E	PL B541 45	A. Aloisio <i>et al.</i> (KLOE Collab.)
FISCHBACH	02	PL B526 355	E. Fischbach, A.W. Overhauser, B. Woodahl
GOKALP	02	JP G28 2783	A. Gokalp <i>et al.</i>
ACHASOV	01B	PL B504 275	M.N. Achasov <i>et al.</i> (Novosibirsk SND Collab.)
ACHASOV	01E	PR D63 072002	M.N. Achasov <i>et al.</i> (Novosibirsk SND Collab.)
ACHASOV	01F	PR D63 094007	N.N. Achasov, V.V. Gubin (Novosibirsk SND Collab.)
ACHASOV	01G	PRL 86 1698	M.N. Achasov <i>et al.</i> (Novosibirsk SND Collab.)
AITALA	01B	PRL 86 770	E.M. Aitala <i>et al.</i> (FNAL E791 Collab.)
AKHMETSHIN	01	PL B501 191	R.R. Akhmetshin <i>et al.</i> (Novosibirsk CMD-2 Collab.)
AKHMETSHIN	01B	PL B509 217	R.R. Akhmetshin <i>et al.</i> (Novosibirsk CMD-2 Collab.)
AKHMETSHIN	01C	PL B503 237	R.R. Akhmetshin <i>et al.</i> (Novosibirsk CMD-2 Collab.)
BENAYOUN	01	EPJ C22 503	M. Benayoun, H.B. O'Connell
ACHASOV	00	EPJ C12 25	M.N. Achasov <i>et al.</i> (Novosibirsk SND Collab.)
ACHASOV	00B	JETP 90 17	M.N. Achasov <i>et al.</i> (Novosibirsk SND Collab.)
		Translated from ZETFP 117 22.	
ACHASOV	00C	PL B474 188	M.N. Achasov <i>et al.</i> (Novosibirsk SND Collab.)
ACHASOV	00D	JETPL 72 282	M.N. Achasov <i>et al.</i> (Novosibirsk SND Collab.)
		Translated from ZETFP 72 411.	
ACHASOV	00E	NP B569 158	M.N. Achasov <i>et al.</i> (Novosibirsk SND Collab.)
ACHASOV	00F	PL B479 53	M.N. Achasov <i>et al.</i> (Novosibirsk SND Collab.)
ACHASOV	00H	PL B485 349	M.N. Achasov <i>et al.</i> (Novosibirsk SND Collab.)
AKHMETSHIN	00B	PL B473 337	R.R. Akhmetshin <i>et al.</i> (Novosibirsk CMD-2 Collab.)
AKHMETSHIN	00E	PL B491 81	R.R. Akhmetshin <i>et al.</i> (Novosibirsk CMD-2 Collab.)
AKHMETSHIN	00F	PL B494 26	R.R. Akhmetshin <i>et al.</i> (Novosibirsk CMD-2 Collab.)
AULCHENKO	00A	JETP 90 927	V.M. Aulchenko <i>et al.</i> (Novosibirsk SND Collab.)
		Translated from ZETFP 117 1067.	
BRAMON	00	PL B486 406	A. Bramon <i>et al.</i>
PDG	00	EPJ C15 1	D.E. Groom <i>et al.</i> (PDG Collab.)
ACHASOV	99	PL B449 122	M.N. Achasov <i>et al.</i>
ACHASOV	99C	PL B456 304	M.N. Achasov <i>et al.</i>
AKHMETSHIN	99B	PL B462 371	R.R. Akhmetshin <i>et al.</i> (Novosibirsk CMD-2 Collab.)
AKHMETSHIN	99C	PL B462 380	R.R. Akhmetshin <i>et al.</i> (Novosibirsk CMD-2 Collab.)
AKHMETSHIN	99D	PL B466 385	R.R. Akhmetshin <i>et al.</i> (Novosibirsk CMD-2 Collab.)
Also		PL B508 217 (errat.)	R.R. Akhmetshin <i>et al.</i> (Novosibirsk CMD-2 Collab.)
AKHMETSHIN	99F	PL B460 242	R.R. Akhmetshin <i>et al.</i> (Novosibirsk CMD-2 Collab.)
AULCHENKO	99	JETPL 69 97	V.M. Aulchenko <i>et al.</i>
		Translated from ZETFP 69 87.	
ACHASOV	98B	PL B438 441	M.N. Achasov <i>et al.</i> (Novosibirsk SND Collab.)
ACHASOV	98F	JETPL 68 573	M.N. Achasov <i>et al.</i> (Novosibirsk SND Collab.)
ACHASOV	98I	PL B440 442	M.N. Achasov <i>et al.</i>
AKHMETSHIN	98	PL B434 426	R.R. Akhmetshin <i>et al.</i> (CMD-2 Collab.)
AULCHENKO	98	PL B436 199	V.M. Aulchenko <i>et al.</i> (Novosibirsk SND Collab.)
BARBERIS	98	PL B432 436	D. Barberis <i>et al.</i> (Omega Expt.)
AKHMETSHIN	97B	PL B415 445	R.R. Akhmetshin <i>et al.</i> (NOVO, BOST, PITTP+)
AKHMETSHIN	97C	PL B415 452	R.R. Akhmetshin <i>et al.</i> (Novosibirsk CMD-2 Collab.)
BENAYOUN	96	ZPHY C72 221	M. Benayoun <i>et al.</i> (IPNP, NOVO)
AKHMETSHIN	95	PL B364 199	R.R. Akhmetshin <i>et al.</i> (Novosibirsk CMD-2 Collab.)
DOLINSKY	91	PRPL 202 99	S.I. Dolinsky <i>et al.</i> (NOVO)
KUHN	90	ZPHY C48 445	J.H. Kuhn <i>et al.</i> (MPIM)
ACHASOV	89	NP B315 465	N.N. Achasov, V.N. Ivanchenko
DOLINSKY	89	ZPHY C42 511	S.I. Dolinsky <i>et al.</i> (NOVO)
BARKOV	88	SJNP 47 248	L.M. Barkov <i>et al.</i> (NOVO)
		Translated from YAF 47 393.	

DOLINSKY	88	SJNP 48 277 Translated from YAF 48 442.	S.I. Dolinsky <i>et al.</i>	(NOVO)
DRUZHININ	87	ZPHY C37 1	V.P. Druzhinin <i>et al.</i>	(NOVO)
ARMSTRONG	86	PL 166B 245	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)
ATKINSON	86	ZPHY C30 521	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
BEBEK	86	PRL 56 1893	C. Bebek <i>et al.</i>	(CLEO Collab.)
DAVENPORT	86	PR D33 2519	T.F. Davenport	(TUFTS, ARIZ, FNAL, FSU, NDAM+)
DIJKSTRA	86	ZPHY C31 375	H. Dijkstra <i>et al.</i>	(ANIK, BRIS, CERN+)
FRAME	86	NP B276 667	D. Frame <i>et al.</i>	(GLAS)
GOLUBEV	86	SJNP 44 409 Translated from YAF 44 633.	V.B. Golubev <i>et al.</i>	(NOVO)
ALBRECHT	85D	PL 153B 343	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
GOLUBEV	85	SJNP 41 756	V.B. Golubev <i>et al.</i>	(NOVO)
DRUZHININ	84	PL 144B 136	V.P. Druzhinin <i>et al.</i>	(NOVO)
ARMSTRONG	83B	NP B224 193	T.A. Armstrong <i>et al.</i>	(BARI, BIRM, CERN+)
BARATE	83	PL 121B 449	R. Barate <i>et al.</i>	(SACL, LOIC, SHMP, IND)
KURDADZE	83C	JETPL 38 366	L.M. Kurdadze <i>et al.</i>	(NOVO)
ARENTON	82	Translated from ZETFP 38 306.		
PELLINEN	82	PR D25 2241	M.W. Arenton <i>et al.</i>	(ANL, ILL)
DAUM	81	PS 25 599	A. Pellinen, M. Roos	(HELS)
IVANOV	81	PL 100B 439	C. Daum <i>et al.</i>	(AMST, BRIS, CERN, CRAC+)
Also		PL 107B 297	P.M. Ivanov <i>et al.</i>	(NOVO)
VASSERMAN	81	Private Comm.	S.I. Eidelman	(NOVO)
Also		PL 99B 62	I.B. Vasserman <i>et al.</i>	(NOVO)
		SJNP 35 240	L.M. Kurdadze <i>et al.</i>	(NOVO)
CORDIER	80	Translated from YAF 35 352.		
CORDIER	79	NP B172 13	A. Cordier <i>et al.</i>	(LALO)
BUKIN	78B	PL 81B 389	A. Cordier <i>et al.</i>	(LALO)
BUKIN	78B	SJNP 27 521	A.D. Bokin <i>et al.</i>	(NOVO)
BUKIN	78C	Translated from YAF 27 985.		
		SJNP 27 516	A.D. Bokin <i>et al.</i>	(NOVO)
COOPER	78B	Translated from YAF 27 976.		
LOSTY	78	NP B146 1	A.M. Cooper <i>et al.</i>	(TATA, CERN, CDEF+)
AKERLOF	77	NP B133 38	M.J. Losty <i>et al.</i>	(CERN, AMST, NIJM+)
ANDREWS	77	PRL 39 861	C.W. Akerlof <i>et al.</i>	(FNAL, MICH, PURD)
BALDI	77	PRL 38 198	D.E. Andrews <i>et al.</i>	(ROCH)
CERRADA	77B	PL 68B 381	R. Baldi <i>et al.</i>	(GEVA)
COHEN	77	NP B126 241	M. Cerrada <i>et al.</i>	(AMST, CERN, NIJM+)
LAVEN	77	PRL 38 269	D. Cohen <i>et al.</i>	(ANL)
LYONS	77	NP B127 43	H. Laven <i>et al.</i>	(AACH3, BERL, CERN, LOIC+)
COSME	76	NP B125 207	L. Lyons, A.M. Cooper, A.G. Clark	(OXF)
KALBFLEISCH	76	PL 63B 352	G. Cosme <i>et al.</i>	(ORSAY)
KALBFLEISCH	76	PR D13 22	G.R. Kalbfleisch, R.C. Strand, J.W. Chapman	(BNL+)
PARROUR	76	PL 63B 357	G. Parrou <i>et al.</i>	(ORSAY)
PARROUR	76B	PL 63B 362	G. Parrou <i>et al.</i>	(ORSAY)
KALBFLEISCH	75	PR D11 987	G.R. Kalbfleisch, R.C. Strand, J.W. Chapman	(BNL+)
AYRES	74	PRL 32 1463	D.S. Ayres <i>et al.</i>	(ANL)
BESCH	74	NP B70 257	H.J. Besch <i>et al.</i>	(BONN)
COSME	74	PL 48B 155	G. Cosme <i>et al.</i>	(ORSAY)
COSME	74B	PL 48B 159	G. Cosme <i>et al.</i>	(ORSAY)
DEGROOT	74	NP B74 77	A.J. de Groot <i>et al.</i>	(AMST, NIJM)
AUGUSTIN	73	PRL 30 462	J.E. Augustin <i>et al.</i>	(ORSAY)
BALLAM	73	PR D7 3150	J. Ballam <i>et al.</i>	(SLAC, LBL)
BINNIE	73B	PR D8 2789	D.M. Binnie <i>et al.</i>	(LOIC, SHMP)
AGUILAR-...	72B	PR D6 29	M. Aguilar-Benitez <i>et al.</i>	(BNL)
ALVENSLEB...	72	PRL 28 66	H. Alvensleben <i>et al.</i>	(MIT, DESY)
BORENSTEIN	72	PR D5 1559	S.R. Borenstein <i>et al.</i>	(BNL, MICH)
COLLEY	72	NP B50 1	D.C. Colley <i>et al.</i>	(BIRM, GLAS)
BALAKIN	71	PL 34B 328	V.E. Balakin <i>et al.</i>	(NOVO)
CHATELUS	71	Thesis LAL 1247	Y. Chatelus	(STRB)
Also		PL 32B 416	J.C. Bizot <i>et al.</i>	(ORSAY)
HAYES	71	PR D4 899	S. Hayes <i>et al.</i>	(CORN)
STOTTLE...	71	Thesis ORO 2504 170	A.R. Stottlemeyer	(UMD)
BIZOT	70	PL 32B 416	J.C. Bizot <i>et al.</i>	(ORSAY)
Also		Liverpool Sym. 69	J.P. Perez-y-Jorba	
EARLES	70	PRL 25 1312	D.R. Earles <i>et al.</i>	(NEAS)
LINDSEY	66	PR 147 913	J.S. Lindsey, G. Smith	(LRL)
LONDON	66	PR 143 1034	G.W. London <i>et al.</i>	(BNL, SYRA) IGJPC
BADIER	65B	PL 17 337	J. Badier <i>et al.</i>	(EPOL, SACL, AMST)

LINDSEY 65 PRL 15 221 J.S. Lindsey, G.A. Smith (LRL)
LINDSEY 65 data included in LINDSEY 66.
SCHLEIN 63 PRL 10 368 P.E. Schlein *et al.* (UCLA) IGJP
