
 $I^G(J^{PC}) = 0^-(1^{--})$

See the Review on “Branching Ratios of $\psi(2S)$, $\chi_{c0,1,2}$ and $\eta_c(1S)$ ” before the $\chi_{c0}(1P)$ Listings.

$\psi(2S)$ MASS

OUR FIT includes measurements of $m_{\psi}(2S)$, $m_{\psi}(3770)$, and $m_{\psi}(3770) - m_{\psi}(2S)$.

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------------------------------|-----------|--|
| 3686.097±0.011 OUR FIT | | Error includes scale factor of 1.1. | | |
| 3686.097±0.010 OUR AVERAGE | | | | |
| 3686.099±0.004±0.009 | | ¹ ANASHIN | 15 KEDR | $e^+e^- \rightarrow$ hadrons |
| 3686.12 ± 0.06 ± 0.10 | 4k | AAIJ | 12H LHCb | $pp \rightarrow J/\psi\pi^+\pi^-X$ |
| 3685.95 ± 0.10 | 413 | ² ARTAMONOV 00 | OLYA | $e^+e^- \rightarrow$ hadrons |
| 3685.98 ± 0.09 ± 0.04 | | ³ ARMSTRONG 93B | E760 | $\bar{p}p \rightarrow e^+e^-$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 3686.08 ± 0.07 | 1301 | ⁴ AAIJ | 23AP LHCb | $B_s^0 \rightarrow J/\psi 2(\pi^+\pi^-)$ |
| 3686.114±0.007 ^{+0.011} _{-0.016} | | ⁵ ANASHIN | 12 KEDR | $e^+e^- \rightarrow$ hadrons |
| 3686.111±0.025±0.009 | | AULCHENKO 03 | KEDR | $e^+e^- \rightarrow$ hadrons |
| 3686.00 ± 0.10 | 413 | ⁶ ZHOLENTZ | 80 OLYA | e^+e^- |

¹ Supersedes AULCHENKO 03 and ANASHIN 12.

² Reanalysis of ZHOLENTZ 80 using new electron mass (COHEN 87) and radiative corrections (KURAEV 85).

³ Mass central value and systematic error recalculated by us according to Eq.(16) in ARMSTRONG 93B, using the value for the $J/\psi(1S)$ mass from AULCHENKO 03.

⁴ From a fit of a relativistic S -wave Breit-Wigner convolved with the detector resolution. The width of $\psi(2S)$ is constrained to the PDG 22 value. Systematic errors not evaluated.

⁵ From the scans in 2004 and 2006. ANASHIN 12 reports the value $3686.114 \pm 0.007 \pm 0.011^{+0.002}_{-0.012}$ MeV, where the third uncertainty is due to assumptions on the interference between the resonance and hadronic continuum. We combined the two systematic uncertainties.

⁶ Superseded by ARTAMONOV 00.

$m_{\psi(2S)} - m_{J/\psi(1S)}$

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|---|---------------------------|---------|--|
| 589.188±0.028 OUR AVERAGE | | | |
| 589.194±0.027±0.011 | ¹ AULCHENKO 03 | KEDR | $e^+e^- \rightarrow$ hadrons |
| 589.7 ± 1.2 | LEMOIGNE 82 | GOLI | $185\pi^-Be \rightarrow \gamma\mu^+\mu^-A$ |
| 589.07 ± 0.13 | ¹ ZHOLENTZ 80 | OLYA | e^+e^- |
| 588.7 ± 0.8 | LUTH 75 | MRK1 | |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 588 ± 1 | ² BAI | 98E BES | e^+e^- |

¹ Redundant with data in mass above.² Systematic errors not evaluated.

$\psi(2S)$ WIDTH

| <i>VALUE</i> (keV) | <i>EVTS</i> | <i>DOCUMENT ID</i> | <i>TECN</i> | <i>COMMENT</i> |
|---------------------------|-------------------------------------|--------------------|-------------|---|
| 293± 9 OUR FIT | Error includes scale factor of 1.2. | | | |
| 286±16 OUR AVERAGE | | | | |
| 358±88± 4 | | ABLIKIM 08B | BES2 | $e^+ e^- \rightarrow$ hadrons |
| 290±25± 4 | 2.7k | ANDREOTTI 07 | E835 | $p\bar{p} \rightarrow e^+ e^-$, $J/\psi X$ |
| 331±58± 2 | | ABLIKIM 06L | BES2 | $e^+ e^- \rightarrow$ hadrons |
| 264±27 | 1 BAI | | 02B | $e^+ e^-$ |
| 287±37±16 | | 2 ARMSTRONG 93B | E760 | $\bar{p}p \rightarrow e^+ e^-$ |

¹ From a simultaneous fit to the hadronic and $\mu^+ \mu^-$ cross section, assuming $\Gamma = \Gamma_h + \Gamma_e + \Gamma_\mu + \Gamma_\tau$ and lepton universality. Does not include vacuum polarization correction.

² The initial-state radiation correction reevaluated by ANDREOTTI 07 in its Ref. [4].

$\psi(2S)$ DECAY MODES

| Mode | Fraction (Γ_i/Γ) | Scale factor/ Confidence level |
|---|----------------------------------|-----------------------------------|
| Γ_1 hadrons | (97.85 ± 0.13) % | |
| Γ_2 virtual $\gamma \rightarrow$ hadrons | (1.79 ± 0.04) % | |
| Γ_3 ggg | (10.6 ± 1.6) % | |
| Γ_4 γgg | (1.03 ± 0.29) % | |
| Γ_5 light hadrons | (15.4 ± 1.5) % | |
| Γ_6 K_S^0 anything | (16.0 ± 1.1) % | |
| Γ_7 $e^+ e^-$ | (7.94 ± 0.22) $\times 10^{-3}$ | S=1.3 |
| Γ_8 $\mu^+ \mu^-$ | (8.0 ± 0.6) $\times 10^{-3}$ | |
| Γ_9 $\tau^+ \tau^-$ | (3.1 ± 0.4) $\times 10^{-3}$ | |

Decays into $J/\psi(1S)$ and anything

| | | | |
|---------------|------------------------|------------------------------------|-------|
| Γ_{10} | $J/\psi(1S)$ anything | (61.5 ± 0.7) % | S=1.3 |
| Γ_{11} | $J/\psi(1S)$ neutrals | (25.4 ± 0.5) % | S=1.6 |
| Γ_{12} | $J/\psi(1S)\pi^+\pi^-$ | (34.69 ± 0.34) % | S=1.1 |
| Γ_{13} | $J/\psi(1S)\pi^0\pi^0$ | (18.2 ± 0.5) % | S=1.6 |
| Γ_{14} | $J/\psi(1S)\eta$ | (3.37 ± 0.06) % | S=1.2 |
| Γ_{15} | $J/\psi(1S)\pi^0$ | (1.268 ± 0.032) $\times 10^{-3}$ | |

Hadronic decays

| | | | |
|---------------|---|----------------------------------|-------|
| Γ_{16} | $\pi^+ \pi^-$ | (7.8 ± 2.6) $\times 10^{-6}$ | |
| Γ_{17} | $\pi^+ \pi^- \pi^0$ | (2.01 ± 0.17) $\times 10^{-4}$ | S=1.7 |
| Γ_{18} | $\rho(770)\pi \rightarrow \pi^+ \pi^- \pi^0$ | (3.2 ± 1.2) $\times 10^{-5}$ | S=1.8 |
| Γ_{19} | $\rho(2150)\pi \rightarrow \pi^+ \pi^- \pi^0$ | (1.9 ± 0.4) $\times 10^{-4}$ | |
| Γ_{20} | $2(\pi^+ \pi^-)$ | (2.4 ± 0.6) $\times 10^{-4}$ | S=2.2 |
| Γ_{21} | $\rho^0 \pi^+ \pi^-$ | (2.2 ± 0.6) $\times 10^{-4}$ | S=1.4 |

| | | | |
|---------------|--|----------------------------------|--------|
| Γ_{22} | $2(\pi^+\pi^-)\pi^0$ | $(2.9 \pm 1.0) \times 10^{-3}$ | S=4.7 |
| Γ_{23} | $\rho\alpha_2(1320)$ | $(2.6 \pm 0.9) \times 10^{-4}$ | |
| Γ_{24} | $\pi^+\pi^-\pi^0\pi^0\pi^0$ | $(5.3 \pm 1.0) \times 10^{-3}$ | |
| Γ_{25} | $\rho^\pm\pi^\mp\pi^0\pi^0$ | $< 2.7 \times 10^{-3}$ | CL=90% |
| Γ_{26} | $\pi^+\pi^-4\pi^0$ | $(1.4 \pm 1.0) \times 10^{-3}$ | |
| Γ_{27} | $3(\pi^+\pi^-)$ | $(3.5 \pm 2.0) \times 10^{-4}$ | S=2.8 |
| Γ_{28} | $2(\pi^+\pi^-\pi^0)$ | $(4.8 \pm 1.5) \times 10^{-3}$ | |
| Γ_{29} | $3(\pi^+\pi^-)\pi^0$ | $(3.5 \pm 1.6) \times 10^{-3}$ | |
| Γ_{30} | $2(\pi^+\pi^-)3\pi^0$ | $(1.42 \pm 0.31) \%$ | |
| Γ_{31} | $\eta\pi^+\pi^-$ | $< 1.6 \times 10^{-4}$ | CL=90% |
| Γ_{32} | $\eta\pi^+\pi^-\pi^0$ | $(9.5 \pm 1.7) \times 10^{-4}$ | |
| Γ_{33} | $\eta 2(\pi^+\pi^-)$ | $(1.2 \pm 0.6) \times 10^{-3}$ | |
| Γ_{34} | $\eta\pi^+\pi^-\pi^0\pi^0$ | $< 4 \times 10^{-4}$ | CL=90% |
| Γ_{35} | $\eta\pi^+\pi^-3\pi^0$ | $< 2.1 \times 10^{-3}$ | CL=90% |
| Γ_{36} | $\eta 2(\pi^+\pi^-\pi^0)$ | $< 2.1 \times 10^{-3}$ | CL=90% |
| Γ_{37} | $\rho\eta$ | $(2.2 \pm 0.6) \times 10^{-5}$ | S=1.1 |
| Γ_{38} | $\eta'\pi^+\pi^-\pi^0$ | $(4.5 \pm 2.1) \times 10^{-4}$ | |
| Γ_{39} | $\eta'\rho$ | $(1.9 \pm 1.7) \times 10^{-5}$ | |
| Γ_{40} | $\omega\pi^0$ | $(2.1 \pm 0.6) \times 10^{-5}$ | |
| Γ_{41} | $\omega\pi^+\pi^-$ | $(7.3 \pm 1.2) \times 10^{-4}$ | S=2.1 |
| Γ_{42} | $\omega\pi^+\pi^-2\pi^0$ | $(8.7 \pm 2.4) \times 10^{-3}$ | |
| Γ_{43} | $b_1^\pm\pi^\mp$ | $(4.0 \pm 0.6) \times 10^{-4}$ | S=1.1 |
| Γ_{44} | $\omega f_2(1270)$ | $(2.2 \pm 0.4) \times 10^{-4}$ | |
| Γ_{45} | $\omega\pi^0\pi^0$ | $(1.11 \pm 0.35) \times 10^{-3}$ | |
| Γ_{46} | $\omega 3\pi^0$ | $< 8 \times 10^{-4}$ | CL=90% |
| Γ_{47} | $b_1^0\pi^0$ | $(2.4 \pm 0.6) \times 10^{-4}$ | |
| Γ_{48} | $\omega\eta$ | $< 1.1 \times 10^{-5}$ | CL=90% |
| Γ_{49} | $\omega\eta'$ | $(3.2 \pm 2.5) \times 10^{-5}$ | |
| Γ_{50} | $\phi\pi^0$ | $< 4 \times 10^{-7}$ | CL=90% |
| Γ_{51} | $\phi\pi^+\pi^-$ | $(1.18 \pm 0.26) \times 10^{-4}$ | S=1.5 |
| Γ_{52} | $\phi f_0(980) \rightarrow \pi^+\pi^-$ | $(7.5 \pm 3.3) \times 10^{-5}$ | S=1.6 |
| Γ_{53} | $\phi\eta$ | $(3.10 \pm 0.31) \times 10^{-5}$ | |
| Γ_{54} | $\eta\phi(2170), \phi(2170) \rightarrow \phi f_0(980), f_0 \rightarrow \pi^+\pi^-$ | $< 2.2 \times 10^{-6}$ | CL=90% |
| Γ_{55} | $\phi\eta'$ | $(1.54 \pm 0.20) \times 10^{-5}$ | |
| Γ_{56} | $\phi f_1(1285)$ | $(3.0 \pm 1.3) \times 10^{-5}$ | |
| Γ_{57} | $\phi\eta(1405) \rightarrow \phi\pi^+\pi^-\eta$ | $(8.5 \pm 1.7) \times 10^{-6}$ | |
| Γ_{58} | $\phi f'_2(1525)$ | $(4.4 \pm 1.6) \times 10^{-5}$ | |
| Γ_{59} | K^+K^- | $(7.5 \pm 0.5) \times 10^{-5}$ | |
| Γ_{60} | $K^+K^-\pi^+\pi^-$ | $(7.3 \pm 0.5) \times 10^{-4}$ | |
| Γ_{61} | $K^+K^-\pi^0$ | $(4.07 \pm 0.31) \times 10^{-5}$ | |
| Γ_{62} | $K_S^0 K_S^0$ | $< 4.6 \times 10^{-6}$ | |
| Γ_{63} | $K_S^0 K_L^0$ | $(5.34 \pm 0.33) \times 10^{-5}$ | |

| | | | |
|----------------|---|--|--------|
| Γ_{64} | $K_S^0 K_L^0 \pi^0$ | $< 3.0 \times 10^{-4}$ | CL=90% |
| Γ_{65} | $K^+ K^- \pi^0 \pi^0$ | $(2.6 \pm 1.3) \times 10^{-4}$ | |
| Γ_{66} | $K^+ K^- \pi^0 \pi^0 \pi^0$ | $(6.6 \pm 2.8) \times 10^{-4}$ | |
| Γ_{67} | $K_S^0 K^\pm \pi^\mp \pi^0 \pi^0$ | $(1.7 \pm 0.6) \times 10^{-3}$ | |
| Γ_{68} | $K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$ | $(2.2 \pm 0.4) \times 10^{-3}$ | |
| Γ_{69} | $K^+ K^- \pi^+ \pi^- \pi^0$ | $(1.26 \pm 0.09) \times 10^{-3}$ | |
| Γ_{70} | $\omega f_0(1710) \rightarrow \omega K^+ K^-$ | $(5.9 \pm 2.2) \times 10^{-5}$ | |
| Γ_{71} | $K^*(892)^0 K^- \pi^+ \pi^0 + \text{c.c.}$ | $(8.6 \pm 2.2) \times 10^{-4}$ | |
| Γ_{72} | $K^*(892)^+ K^- \pi^+ \pi^- + \text{c.c.}$ | $(9.6 \pm 2.8) \times 10^{-4}$ | |
| Γ_{73} | $K^*(892)^+ K^- \rho^0 + \text{c.c.}$ | $(7.3 \pm 2.6) \times 10^{-4}$ | |
| Γ_{74} | $K^*(892)^0 K^- \rho^+ + \text{c.c.}$ | $(6.1 \pm 1.8) \times 10^{-4}$ | |
| Γ_{75} | $K_S^0 K_S^0 \pi^+ \pi^-$ | $(2.2 \pm 0.4) \times 10^{-4}$ | |
| Γ_{76} | $K_S^0 K_L^0 \pi^0 \pi^0$ | $(1.3 \pm 0.6) \times 10^{-3}$ | |
| Γ_{77} | $K_S^0 K^*(892)^0 \pi^0 \pi^0$ | $(3.0 \pm 1.3) \times 10^{-4}$ | |
| Γ_{78} | $K_S^0 K^\pm \rho(770)^\mp \pi^0$ | $< 7 \times 10^{-4}$ | CL=90% |
| Γ_{79} | $K_S^0 K^\pm \pi^\mp \rho(770)^0$ | $< 7 \times 10^{-4}$ | CL=90% |
| Γ_{80} | $K^\mp K^*(892)^\pm \pi^0 \pi^0$ | $(7.0 \pm 2.9) \times 10^{-4}$ | |
| Γ_{81} | $K^*(892)^+ K^*(892)^- \pi^0$ | $(3.6 \pm 1.8) \times 10^{-3}$ | |
| Γ_{82} | $K_S^0 K_L^0 \eta$ | $(1.3 \pm 0.5) \times 10^{-3}$ | |
| Γ_{83} | $K^+ K^- \rho^0$ | $(2.2 \pm 0.4) \times 10^{-4}$ | |
| Γ_{84} | $K^*(892)^0 \bar{K}_2^*(1430)^0$ | $(1.9 \pm 0.5) \times 10^{-4}$ | |
| Γ_{85} | $K^+ K^- \pi^+ \pi^- \eta$ | $(1.3 \pm 0.7) \times 10^{-3}$ | |
| Γ_{86} | $K^+ K^- 2(\pi^+ \pi^-)$ | $(1.9 \pm 0.9) \times 10^{-3}$ | |
| Γ_{87} | $K^+ K^- 2(\pi^+ \pi^-) \pi^0$ | $(1.00 \pm 0.31) \times 10^{-3}$ | |
| Γ_{88} | $K^+ K^*(892)^- + \text{c.c.}$ | $(2.9 \pm 0.4) \times 10^{-5}$ | S=1.2 |
| Γ_{89} | $2(K^+ K^-)$ | $(6.3 \pm 1.3) \times 10^{-5}$ | |
| Γ_{90} | $2(K^+ K^-) \pi^0$ | $(1.10 \pm 0.28) \times 10^{-4}$ | |
| Γ_{91} | $K^+ K^- \phi$ | $(7.0 \pm 1.6) \times 10^{-5}$ | |
| Γ_{92} | $K_S^0 K_S^0 \phi$ | $(3.53 \pm 0.29) \times 10^{-5}$ | |
| Γ_{93} | $K_1(1270)^\pm K^\mp$ | $(1.00 \pm 0.28) \times 10^{-3}$ | |
| Γ_{94} | $K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.}$ | $(6.7 \pm 2.5) \times 10^{-4}$ | |
| Γ_{95} | $\eta K^+ K^-$, no $\eta \phi$ | $(3.49 \pm 0.17) \times 10^{-5}$ | |
| Γ_{96} | $\eta K^+ K^-$ | $< 2.6 \times 10^{-4}$ | CL=90% |
| Γ_{97} | $X(1750) \eta \rightarrow K^+ K^- \eta$ | $(4.8 \pm 2.8) \times 10^{-6}$ | |
| Γ_{98} | $K_1(1400)^\pm K^\mp$ | $< 3.1 \times 10^{-4}$ | CL=90% |
| Γ_{99} | $K_2^*(1430)^\pm K^\mp$ | $(7.1 \begin{array}{l} +1.3 \\ -0.9 \end{array}) \times 10^{-5}$ | |
| Γ_{100} | $K^*(892)^0 \bar{K}^0 + \text{c.c.}$ | $(1.09 \pm 0.20) \times 10^{-4}$ | |
| Γ_{101} | $\omega K^+ K^-$ | $(1.62 \pm 0.11) \times 10^{-4}$ | S=1.1 |
| Γ_{102} | $\omega K_S^0 K_S^0$ | $(7.0 \pm 0.5) \times 10^{-5}$ | |
| Γ_{103} | $\omega K^*(892)^+ K^- + \text{c.c.}$ | $(2.07 \pm 0.26) \times 10^{-4}$ | |
| Γ_{104} | $\omega K_2^*(1430)^+ K^- + \text{c.c.}$ | $(6.1 \pm 1.2) \times 10^{-5}$ | |
| Γ_{105} | $\omega \bar{K}^*(892)^0 K^0$ | $(1.68 \pm 0.30) \times 10^{-4}$ | |

| | | | |
|----------------|--|---|--------|
| Γ_{106} | $\omega \bar{K}_2^*(1430)^0 K^0$ | $(5.8 \pm 2.2) \times 10^{-5}$ | |
| Γ_{107} | $\omega X(1440) \rightarrow \omega K_S^0 K^- \pi^+ +$ c.c. | $(1.6 \pm 0.4) \times 10^{-5}$ | |
| Γ_{108} | $\omega X(1440) \rightarrow \omega K^+ K^- \pi^0$ | $(1.09 \pm 0.26) \times 10^{-5}$ | |
| Γ_{109} | $\omega f_1(1285) \rightarrow \omega K_S^0 K^- \pi^+ +$ c.c. | $(3.0 \pm 1.0) \times 10^{-6}$ | |
| Γ_{110} | $\omega f_1(1285) \rightarrow \omega K^+ K^- \pi^0$ | $(1.2 \pm 0.7) \times 10^{-6}$ | |
| Γ_{111} | $p \bar{p}$ | $(2.94 \pm 0.09) \times 10^{-4}$ | S=1.3 |
| Γ_{112} | $n \bar{n}$ | $(3.06 \pm 0.15) \times 10^{-4}$ | |
| Γ_{113} | $p \bar{p} \pi^0$ | $(1.53 \pm 0.07) \times 10^{-4}$ | |
| Γ_{114} | $N(940) \bar{p} + \text{c.c.} \rightarrow p \bar{p} \pi^0$ | $(6.4 \begin{array}{l} +1.8 \\ -1.3 \end{array}) \times 10^{-5}$ | |
| Γ_{115} | $N(1440) \bar{p} + \text{c.c.} \rightarrow p \bar{p} \pi^0$ | $(7.3 \begin{array}{l} +1.7 \\ -1.5 \end{array}) \times 10^{-5}$ | S=2.5 |
| Γ_{116} | $N(1520) \bar{p} + \text{c.c.} \rightarrow p \bar{p} \pi^0$ | $(6.4 \begin{array}{l} +2.3 \\ -1.8 \end{array}) \times 10^{-6}$ | |
| Γ_{117} | $N(1535) \bar{p} + \text{c.c.} \rightarrow p \bar{p} \pi^0$ | $(2.5 \pm 1.0) \times 10^{-5}$ | |
| Γ_{118} | $N(1650) \bar{p} + \text{c.c.} \rightarrow p \bar{p} \pi^0$ | $(3.8 \begin{array}{l} +1.4 \\ -1.7 \end{array}) \times 10^{-5}$ | |
| Γ_{119} | $N(1720) \bar{p} + \text{c.c.} \rightarrow p \bar{p} \pi^0$ | $(1.79 \begin{array}{l} +0.26 \\ -0.70 \end{array}) \times 10^{-5}$ | |
| Γ_{120} | $N(2300) \bar{p} + \text{c.c.} \rightarrow p \bar{p} \pi^0$ | $(2.6 \begin{array}{l} +1.2 \\ -0.7 \end{array}) \times 10^{-5}$ | |
| Γ_{121} | $N(2570) \bar{p} + \text{c.c.} \rightarrow p \bar{p} \pi^0$ | $(2.13 \begin{array}{l} +0.40 \\ -0.31 \end{array}) \times 10^{-5}$ | |
| Γ_{122} | $p \bar{p} \pi^+ \pi^-$ | $(6.0 \pm 0.4) \times 10^{-4}$ | |
| Γ_{123} | $p \bar{p} K^+ K^-$ | $(2.7 \pm 0.7) \times 10^{-5}$ | |
| Γ_{124} | $p \bar{p} \eta$ | $(6.0 \pm 0.4) \times 10^{-5}$ | |
| Γ_{125} | $N(1535) \bar{p} + \text{c.c.} \rightarrow p \bar{p} \eta$ | $(4.5 \begin{array}{l} +0.7 \\ -0.6 \end{array}) \times 10^{-5}$ | |
| Γ_{126} | $p \bar{p} \pi^+ \pi^- \pi^0$ | $(7.3 \pm 0.7) \times 10^{-4}$ | |
| Γ_{127} | $p \bar{p} \rho^0$ | $(5.0 \pm 2.2) \times 10^{-5}$ | |
| Γ_{128} | $p \bar{p} \omega$ | $(6.9 \pm 2.1) \times 10^{-5}$ | |
| Γ_{129} | $p \bar{p} \eta'$ | $(1.10 \pm 0.13) \times 10^{-5}$ | |
| Γ_{130} | $p \bar{p} \phi$ | $(6.1 \pm 0.6) \times 10^{-6}$ | |
| Γ_{131} | $\phi X(1835) \rightarrow p \bar{p} \phi$ | $< 1.82 \times 10^{-7}$ | CL=90% |
| Γ_{132} | $p \bar{n} \pi^- \text{ or c.c.}$ | $(2.48 \pm 0.17) \times 10^{-4}$ | |
| Γ_{133} | $p \bar{n} \pi^- \pi^0$ | $(3.2 \pm 0.7) \times 10^{-4}$ | |
| Γ_{134} | $\Lambda \bar{\Lambda}$ | $(3.81 \pm 0.13) \times 10^{-4}$ | S=1.4 |
| Γ_{135} | $\Lambda \bar{\Lambda} \pi^0$ | $(1.4 \pm 0.7) \times 10^{-6}$ | |
| Γ_{136} | $\Lambda \bar{\Lambda} \eta$ | $(2.43 \pm 0.32) \times 10^{-5}$ | |
| Γ_{137} | $\Lambda(1670) \bar{\Lambda} \rightarrow \Lambda \bar{\Lambda} \eta$ | $(1.3 \pm 0.7) \times 10^{-5}$ | |
| Γ_{138} | $\Lambda \bar{\Lambda} \eta'$ | $(7.3 \pm 1.0) \times 10^{-6}$ | |
| Γ_{139} | $\Lambda \bar{\Lambda} \omega(782)$ | $(3.3 \pm 0.4) \times 10^{-5}$ | |
| Γ_{140} | $\Lambda \bar{\Lambda} \pi^+ \pi^-$ | $(2.8 \pm 0.6) \times 10^{-4}$ | |
| Γ_{141} | $\Lambda \bar{p} K^+$ | $(1.00 \pm 0.14) \times 10^{-4}$ | |
| Γ_{142} | $\Lambda \bar{p} K^*(892)^+ + \text{c.c.}$ | $(6.3 \pm 0.7) \times 10^{-5}$ | |
| Γ_{143} | $\Lambda \bar{p} K^+ \pi^+ \pi^-$ | $(1.8 \pm 0.4) \times 10^{-4}$ | |

| | | | |
|----------------|---|------------------------------------|--------|
| Γ_{144} | $\bar{\Lambda} n K_S^0 + \text{c.c.}$ | $(8.1 \pm 1.8) \times 10^{-5}$ | |
| Γ_{145} | $\Delta^{++} \bar{\Delta}^{--}$ | $(1.28 \pm 0.35) \times 10^{-4}$ | |
| Γ_{146} | $\Lambda \bar{\Sigma}^+ \pi^- + \text{c.c.}$ | $(1.40 \pm 0.13) \times 10^{-4}$ | |
| Γ_{147} | $\Lambda \bar{\Sigma}^- \pi^+ + \text{c.c.}$ | $(1.54 \pm 0.14) \times 10^{-4}$ | |
| Γ_{148} | $\Lambda \bar{\Sigma}^0 + \text{c.c.}$ | $(1.6 \pm 0.7) \times 10^{-6}$ | |
| Γ_{149} | $\Lambda \bar{\Sigma}^0$ | | |
| Γ_{150} | $\Sigma^0 \bar{p} K^+ + \text{c.c.}$ | $(1.67 \pm 0.18) \times 10^{-5}$ | |
| Γ_{151} | $\Sigma^+ \bar{\Sigma}^-$ | $(2.43 \pm 0.10) \times 10^{-4}$ | S=1.4 |
| Γ_{152} | $\Sigma^0 \bar{\Sigma}^0$ | $(2.35 \pm 0.09) \times 10^{-4}$ | S=1.1 |
| Γ_{153} | $\Sigma^- \bar{\Sigma}^+$ | $(2.82 \pm 0.09) \times 10^{-4}$ | |
| Γ_{154} | $\Sigma^+ \bar{\Sigma}^- \eta$ | $(9.6 \pm 2.4) \times 10^{-6}$ | |
| Γ_{155} | $\Sigma^+ \bar{\Sigma}^- \omega$ | $(1.89 \pm 0.28) \times 10^{-5}$ | |
| Γ_{156} | $\Sigma^+ \bar{\Sigma}^- \phi$ | $(3.0 \pm 0.7) \times 10^{-6}$ | |
| Γ_{157} | $\Sigma(1385)^+ \bar{\Sigma}(1385)^-$ | $(8.5 \pm 0.7) \times 10^{-5}$ | |
| Γ_{158} | $\Sigma(1385)^- \bar{\Sigma}(1385)^+$ | $(8.5 \pm 0.8) \times 10^{-5}$ | |
| Γ_{159} | $\Sigma(1385)^0 \bar{\Sigma}(1385)^0$ | $(6.9 \pm 0.7) \times 10^{-5}$ | |
| Γ_{160} | $\Xi^- \bar{\Xi}^+$ | $(2.87 \pm 0.11) \times 10^{-4}$ | S=1.1 |
| Γ_{161} | $\Xi^0 \bar{\Xi}^0$ | $(2.3 \pm 0.4) \times 10^{-4}$ | S=4.2 |
| Γ_{162} | $\Xi(1530)^0 \bar{\Xi}(1530)^0$ | $(6.8 \pm 0.4) \times 10^{-5}$ | |
| Γ_{163} | $\Lambda \bar{\Xi}^+ K^- + \text{c.c.}$ | $(3.9 \pm 0.4) \times 10^{-5}$ | |
| Γ_{164} | $\Xi(1690)^- \bar{\Xi}^+ \rightarrow K^- \Lambda \bar{\Xi}^+ +$ | $(5.2 \pm 1.6) \times 10^{-6}$ | |
| Γ_{165} | $\Xi(1820)^- \bar{\Xi}^+ \rightarrow K^- \Lambda \bar{\Xi}^+ +$ c.c. | $(1.20 \pm 0.32) \times 10^{-5}$ | |
| Γ_{166} | $\Xi(1530)^- \bar{\Xi}(1530)^+$ | $(1.15 \pm 0.07) \times 10^{-4}$ | |
| Γ_{167} | $\Xi(1530)^- \bar{\Xi}^+$ | $(7.0 \pm 1.2) \times 10^{-6}$ | |
| Γ_{168} | $\Xi(1530)^0 \bar{\Xi}^0$ | $(5.3 \pm 0.5) \times 10^{-6}$ | |
| Γ_{169} | $\Sigma^0 \bar{\Xi}^+ K^- + \text{c.c.}$ | $(3.7 \pm 0.4) \times 10^{-5}$ | |
| Γ_{170} | $\Omega^- \bar{\Omega}^+$ | $(5.66 \pm 0.30) \times 10^{-5}$ | S=1.3 |
| Γ_{171} | $\eta_c \pi^+ \pi^- \pi^0$ | $< 1.0 \times 10^{-3}$ | CL=90% |
| Γ_{172} | $h_c(1P) \pi^0$ | $(7.4 \pm 0.5) \times 10^{-4}$ | |
| Γ_{173} | $\Lambda_c^+ \bar{p} e^+ e^- + \text{c.c.}$ | $< 1.7 \times 10^{-6}$ | CL=90% |
| Γ_{174} | $\Theta(1540) \bar{\Theta}(1540) \rightarrow K_S^0 p K^- \bar{n} + \text{c.c.}$ | [a] < 8.8 $\times 10^{-6}$ | CL=90% |
| Γ_{175} | $\Theta(1540) K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n}$ | [a] < 1.0 $\times 10^{-5}$ | CL=90% |
| Γ_{176} | $\Theta(1540) K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n$ | [a] < 7.0 $\times 10^{-6}$ | CL=90% |
| Γ_{177} | $\bar{\Theta}(1540) K^+ n \rightarrow K_S^0 \bar{p} K^+ n$ | [a] < 2.6 $\times 10^{-5}$ | CL=90% |
| Γ_{178} | $\bar{\Theta}(1540) K_S^0 p \rightarrow K_S^0 p K^- \bar{n}$ | [a] < 6.0 $\times 10^{-6}$ | CL=90% |

Radiative decays

| | | | |
|----------------|------------------------|----------------------------------|-------|
| Γ_{179} | $\gamma \chi_{c0}(1P)$ | $(9.77 \pm 0.23) \%$ | S=1.1 |
| Γ_{180} | $\gamma \chi_{c1}(1P)$ | $(9.75 \pm 0.27) \%$ | S=1.1 |
| Γ_{181} | $\gamma \chi_{c2}(1P)$ | $(9.36 \pm 0.23) \%$ | S=1.2 |
| Γ_{182} | $\gamma \eta_c(1S)$ | $(3.6 \pm 0.5) \times 10^{-3}$ | S=1.3 |

| | | | |
|----------------|---|----------------------------------|--------|
| Γ_{183} | $\gamma\eta_c(2S)$ | $(7 \pm 5) \times 10^{-4}$ | |
| Γ_{184} | $\gamma\pi^0$ | $(1.04 \pm 0.22) \times 10^{-6}$ | S=1.4 |
| Γ_{185} | $\gamma 2(\pi^+\pi^-)$ | $(4.0 \pm 0.6) \times 10^{-4}$ | |
| Γ_{186} | $\gamma 3(\pi^+\pi^-)$ | $< 1.7 \times 10^{-4}$ | CL=90% |
| Γ_{187} | $\gamma\eta'(958)$ | $(1.24 \pm 0.04) \times 10^{-4}$ | |
| Γ_{188} | $\gamma f_2(1270)$ | $(2.73 \pm 0.29) \times 10^{-4}$ | S=1.8 |
| Γ_{189} | $\gamma f_0(1370) \rightarrow \gamma K\bar{K}$ | $(3.1 \pm 1.7) \times 10^{-5}$ | |
| Γ_{190} | $\gamma f_0(1500)$ | $(9.3 \pm 1.9) \times 10^{-5}$ | |
| Γ_{191} | $\gamma f'_2(1525)$ | $(3.3 \pm 0.8) \times 10^{-5}$ | |
| Γ_{192} | $\gamma f_0(1710)$ | seen | |
| Γ_{193} | $\gamma f_0(1710) \rightarrow \gamma\pi\pi$ | $(3.5 \pm 0.6) \times 10^{-5}$ | |
| Γ_{194} | $\gamma f_0(1710) \rightarrow \gamma K\bar{K}$ | $(6.6 \pm 0.7) \times 10^{-5}$ | |
| Γ_{195} | $\gamma f_0(2100) \rightarrow \gamma\pi\pi$ | $(4.8 \pm 1.0) \times 10^{-6}$ | |
| Γ_{196} | $\gamma f_0(2200) \rightarrow \gamma K\bar{K}$ | $(3.2 \pm 1.0) \times 10^{-6}$ | |
| Γ_{197} | $\gamma f_J(2220) \rightarrow \gamma\pi\pi$ | $< 5.8 \times 10^{-6}$ | CL=90% |
| Γ_{198} | $\gamma f_J(2220) \rightarrow \gamma K\bar{K}$ | $< 9.5 \times 10^{-6}$ | CL=90% |
| Γ_{199} | $\gamma\eta$ | $(9.2 \pm 1.8) \times 10^{-7}$ | |
| Γ_{200} | $\gamma\eta\pi^+\pi^-$ | $(8.7 \pm 2.1) \times 10^{-4}$ | |
| Γ_{201} | $\gamma\eta(1405)$ | seen | |
| Γ_{202} | $\gamma\eta(1405) \rightarrow \gamma K\bar{K}\pi$ | $< 9 \times 10^{-5}$ | CL=90% |
| Γ_{203} | $\gamma\eta(1405) \rightarrow \gamma\eta\pi^+\pi^-$ | $(3.6 \pm 2.5) \times 10^{-5}$ | |
| Γ_{204} | $\gamma\eta(1405) \rightarrow \gamma f_0(980)\pi^0 \rightarrow \gamma\pi^+\pi^-\pi^0$ | $< 5.0 \times 10^{-7}$ | CL=90% |
| Γ_{205} | $\gamma\eta(1475)$ | seen | |
| Γ_{206} | $\gamma\eta(1475) \rightarrow \gamma K\bar{K}\pi$ | $< 1.4 \times 10^{-4}$ | CL=90% |
| Γ_{207} | $\gamma\eta(1475) \rightarrow \gamma\eta\pi^+\pi^-$ | $< 8.8 \times 10^{-5}$ | CL=90% |
| Γ_{208} | $\gamma K^{*0} K^+ \pi^- + \text{c.c.}$ | $(3.7 \pm 0.9) \times 10^{-4}$ | |
| Γ_{209} | $\gamma K^{*0} \bar{K}^{*0}$ | $(2.4 \pm 0.7) \times 10^{-4}$ | |
| Γ_{210} | $\gamma K_S^0 K^+ \pi^- + \text{c.c.}$ | $(2.6 \pm 0.5) \times 10^{-4}$ | |
| Γ_{211} | $\gamma K^+ K^- \pi^+ \pi^-$ | $(1.9 \pm 0.5) \times 10^{-4}$ | |
| Γ_{212} | $\gamma K^+ K^- 2(\pi^+ \pi^-)$ | $< 2.2 \times 10^{-4}$ | CL=90% |
| Γ_{213} | $\gamma 2(K^+ K^-)$ | $< 4 \times 10^{-5}$ | CL=90% |
| Γ_{214} | $\gamma p\bar{p}$ | $(3.9 \pm 0.5) \times 10^{-5}$ | S=2.0 |
| Γ_{215} | $\gamma f_2(1950) \rightarrow \gamma p\bar{p}$ | $(1.20 \pm 0.22) \times 10^{-5}$ | |
| Γ_{216} | $\gamma f_2(2150) \rightarrow \gamma p\bar{p}$ | $(7.2 \pm 1.8) \times 10^{-6}$ | |
| Γ_{217} | $\gamma X(1835) \rightarrow \gamma p\bar{p}$ | $(4.6 \pm 1.8) \times 10^{-6}$ | |
| Γ_{218} | $\gamma X \rightarrow \gamma p\bar{p}$ | [b] $< 2 \times 10^{-6}$ | CL=90% |
| Γ_{219} | $\gamma p\bar{p}\pi^+\pi^-$ | $(2.8 \pm 1.4) \times 10^{-5}$ | |
| Γ_{220} | $\gamma\gamma$ | $< 1.5 \times 10^{-4}$ | CL=90% |
| Γ_{221} | $\gamma\gamma J/\psi$ | $(3.1 \pm 1.0) \times 10^{-4}$ | |
| Γ_{222} | $e^+ e^- \eta'$ | $(1.90 \pm 0.26) \times 10^{-6}$ | |
| Γ_{223} | $e^+ e^- \eta_c(1S)$ | $(3.8 \pm 0.4) \times 10^{-5}$ | |

| | | |
|----------------|-------------------------|----------------------------------|
| Γ_{224} | $e^+ e^- \chi_{c0}(1P)$ | $(1.06 \pm 0.25) \times 10^{-3}$ |
| Γ_{225} | $e^+ e^- \chi_{c1}(1P)$ | $(8.5 \pm 0.7) \times 10^{-4}$ |
| Γ_{226} | $e^+ e^- \chi_{c2}(1P)$ | $(6.8 \pm 0.8) \times 10^{-4}$ |

Weak decays

| | | | |
|----------------|--|------------------------|--------|
| Γ_{227} | $D^0 e^+ e^- + \text{c.c.}$ | $< 1.4 \times 10^{-7}$ | CL=90% |
| Γ_{228} | $\Lambda_c^+ \bar{\Sigma}^- + \text{c.c.}$ | $< 1.4 \times 10^{-5}$ | CL=90% |

Other decays

| | | | | |
|----------------|-----------|---------|---|--------|
| Γ_{229} | invisible | < 1.6 | % | CL=90% |
|----------------|-----------|---------|---|--------|

[a] $\Theta(1540)$ is a hypothetical pentaquark state of $1.54 \text{ GeV}/c^2$ mass and a width of less than $25 \text{ MeV}/c^2$.

[b] For a narrow resonance in the range $2.2 < M(X) < 2.8 \text{ GeV}$.

CONSTRAINED FIT INFORMATION

A multiparticle fit to $\chi_{c1}(1P)$, $\chi_{c0}(1P)$, $\chi_{c2}(1P)$, and $\psi(2S)$ with 4 total widths, a partial width, 25 combinations of partial widths obtained from integrated cross section, and 86 branching ratios uses 253 measurements to determine 49 parameters. The overall fit has a $\chi^2 = 389.6$ for 204 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$.

| | | | | | | | | | | |
|-----------|-------|-------|-------|----------|----------|----------|-----------|-----------|-----------|-----------|
| x_8 | 3 | | | | | | | | | |
| x_9 | 0 | 0 | | | | | | | | |
| x_{12} | 21 | 13 | 2 | | | | | | | |
| x_{13} | 22 | 5 | 1 | 28 | | | | | | |
| x_{14} | 11 | 6 | 1 | 44 | 11 | | | | | |
| x_{111} | 0 | 0 | 0 | 3 | 2 | 1 | | | | |
| x_{179} | 0 | 0 | 0 | 2 | 1 | 1 | 0 | | | |
| x_{180} | 1 | 0 | 0 | 2 | 0 | 1 | 0 | 0 | | |
| x_{181} | 1 | 1 | 0 | 4 | 0 | 2 | 0 | 0 | 0 | |
| Γ | -85 | -4 | -1 | -29 | -31 | -15 | -4 | -1 | -1 | -2 |
| | x_7 | x_8 | x_9 | x_{12} | x_{13} | x_{14} | x_{111} | x_{179} | x_{180} | x_{181} |

FIT INFORMATION

A multiparticle fit to $\eta_c(1S)$, $J/\psi(1S)$, $\psi(2S)$, $h_c(1P)$, and B^\pm with the total width, 10 combinations of partial widths obtained from integrated cross section, and 38 branching ratios uses 113 measurements to determine 19 parameters. The overall fit has a $\chi^2 = 184.6$ for 94 degrees of freedom.

$\psi(2S)$ PARTIAL WIDTHS

$\Gamma(\text{hadrons})$

| VALUE (keV) | DOCUMENT ID | TECN | COMMENT | Γ_1 |
|--|-------------|------|----------------|------------|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 258 ± 26 | BAI | 02B | BES2 $e^+ e^-$ | |
| 224 ± 56 | LUTH | 75 | MRK1 $e^+ e^-$ | |

$\Gamma(e^+ e^-)$

| VALUE (keV) | DOCUMENT ID | TECN | COMMENT | Γ_7 |
|--|-------------------------------------|------|---|------------|
| 2.33 ± 0.04 OUR FIT | Error includes scale factor of 1.1. | | | |
| 2.29 ± 0.06 OUR AVERAGE | | | | |
| 2.23 ± 0.10 ± 0.02 | ¹ ABLIKIM | 15V | BES3 $e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$ | |
| 2.338 ± 0.037 ± 0.096 | ABLIKIM | 08B | BES2 $e^+ e^- \rightarrow \text{hadrons}$ | |
| 2.330 ± 0.036 ± 0.110 | ABLIKIM | 06L | BES2 $e^+ e^- \rightarrow \text{hadrons}$ | |
| 2.44 ± 0.21 | ² BAI | 02B | BES2 $e^+ e^-$ | |
| 2.14 ± 0.21 | ALEXANDER | 89 | RVUE See γ mini-review | |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 2.279 ± 0.015 ± 0.042 | ³ ANASHIN | 18 | KEDR $e^+ e^-$ | |
| 2.282 ± 0.015 ± 0.042 | ⁴ ANASHIN | 18 | KEDR $e^+ e^-$ | |
| 2.0 ± 0.3 | BRANDELIK | 79C | DASP $e^+ e^-$ | |
| 2.1 ± 0.3 | ⁵ LUTH | 75 | MRK1 $e^+ e^-$ | |

¹ ABLIKIM 15V reports $2.213 \pm 0.018 \pm 0.099$ keV from a measurement of $[\Gamma(\psi(2S) \rightarrow e^+ e^-)] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)]$ assuming $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.95 \pm 0.45) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.69 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² From a simultaneous fit to $e^+ e^-$, $\mu^+ \mu^-$, and hadronic channel, assuming $\Gamma_e = \Gamma_\mu = \Gamma_\tau / 0.38847$.

³ Combining $\Gamma_{e^+ e^-} \cdot B(\mu^+ \mu^-)$ from ANASHIN 18 with $\Gamma_{e^+ e^-} \cdot B(\text{hadrons})$ from ANASHIN 12 and assuming lepton universality.

⁴ From the sum of $\Gamma_{e^+ e^-} \cdot B(\text{hadrons})$ from ANASHIN 12, $\Gamma_{e^+ e^-} \cdot B(e^+ e^-)$ and $\Gamma_{e^+ e^-} \cdot B(\mu^+ \mu^-)$ from ANASHIN 18, and $\Gamma_{e^+ e^-} \cdot B(\tau^+ \tau^-)$ from ANASHIN 07.

⁵ From a simultaneous fit to $e^+ e^-$, $\mu^+ \mu^-$, and hadronic channels assuming $\Gamma(e^+ e^-) = \Gamma(\mu^+ \mu^-)$.

$\Gamma(\gamma\gamma)$

| VALUE (eV) | CL% | DOCUMENT ID | TECN | COMMENT | Γ_{220} |
|------------|-----|-------------|------|----------------|----------------|
| <43 | 90 | BRANDELIK | 79C | DASP $e^+ e^-$ | |

$\psi(2S) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

This combination of a partial width with the partial width into e^+e^- and with the total width is obtained from the integrated cross section into channel(*i*) in the e^+e^- annihilation. We list only data that have not been used to determine the partial width $\Gamma(i)$ or the branching ratio $\Gamma(i)/\text{total}$.

$\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_1\Gamma_7/\Gamma$

| VALUE (keV) | DOCUMENT ID | TECN | COMMENT |
|---|----------------------|------|--|
| 2.233±0.015±0.042 | ¹ ANASHIN | 12 | KEDR $e^+e^- \rightarrow \text{hadrons}$ |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | |
| 2.2 ± 0.4 | ABRAMS | 75 | MRK1 e^+e^- |

¹ ANASHIN 12 reports the value $2.233 \pm 0.015 \pm 0.037 \pm 0.020$ keV, where the third uncertainty is due to assumptions on the interference between the resonance and hadronic continuum. We combined the two systematic uncertainties.

$\Gamma(K_S^0 \text{anything}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_6\Gamma_7/\Gamma$

| VALUE (keV) | DOCUMENT ID | TECN | COMMENT |
|-----------------------------|-------------|------|---|
| 0.3738±0.0067±0.0200 | ABLIKIM | 21S | BES3 $e^+e^- \rightarrow K_S^0 \text{anything}$ |

$\Gamma(e^+e^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_7\Gamma_7/\Gamma$

| VALUE (eV) | DOCUMENT ID | TECN | COMMENT |
|---------------------|----------------------|------|---------------|
| 21.2±0.7±1.2 | ¹ ANASHIN | 18 | KEDR e^+e^- |

¹ From the average of nine scans of the $\psi(2S)$.

$\Gamma(\mu^+\mu^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_8\Gamma_7/\Gamma$

| VALUE (eV) | DOCUMENT ID | TECN | COMMENT |
|---------------------|----------------------|------|--|
| 19.3±0.3±0.5 | ¹ ANASHIN | 18 | KEDR $\psi(2S) \rightarrow \mu^+\mu^-$ |

¹ From the average of nine scans of the $\psi(2S)$.

$\Gamma(\tau^+\tau^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_9\Gamma_7/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|------|---------|
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | |

$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$

9.0 ± 2.6 79 ¹ ANASHIN 07 KEDR $e^+e^- \rightarrow \psi(2S) \rightarrow \tau^+\tau^-$

¹ Using $\psi(2S)$ total width of 337 ± 13 keV. Systematic errors not evaluated.

$\Gamma(J/\psi(1S)\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{12}\Gamma_7/\Gamma$

| VALUE (keV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------------|-------------------|-------------|------|---|
| 0.808±0.014 OUR FIT | | | | Error includes scale factor of 1.1. |
| 0.836±0.025 OUR AVERAGE | | | | Error includes scale factor of 1.3. See the ideogram below. |
| 0.78 $\pm 0.12 \pm 0.07$ | ¹ LEES | 23 | BABR | $e^+e^- \rightarrow \gamma_{ISR} \text{hadrons}$ |
| 0.837 $\pm 0.028 \pm 0.005$ | ² LEES | 12E | BABR | $10.6 e^+e^- \rightarrow 2\pi^+\pi^-\gamma$ |
| 0.852 $\pm 0.010 \pm 0.026$ | 19.5k ADAM | 06 | CLEO | $3.773 e^+e^- \rightarrow \gamma\psi(2S)$ |
| 0.68 ± 0.09 | ³ BAI | 98E | BES | e^+e^- |

$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$

0.93 $\pm 0.08 \pm 0.03$ 256 ⁴ AUBERT 07AU BABR $10.6 e^+e^- \rightarrow J/\psi\pi^+\pi^-\gamma$

0.755 $\pm 0.048 \pm 0.004$ 544 ⁵ AUBERT 05D BABR $10.6 e^+e^- \rightarrow \pi^+\pi^-\mu^+\mu^-\gamma$




¹ LEES 23 reports $[\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow K_S^0 K^\pm \pi^\mp)] = (4.14 \pm 0.55 \pm 0.29) \times 10^{-3}$ keV which we divide by our best value $B(J/\psi(1S) \rightarrow K_S^0 K^\pm \pi^\mp) = (5.3 \pm 0.5) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² LEES 12E reports $[\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \mu^+\mu^-)] = (49.9 \pm 1.3 \pm 1.0) \times 10^{-3}$ keV which we divide by our best value $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ The value of $\Gamma(e^+e^-)$ quoted in BAI 98E is derived using $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.4 \pm 2.6) \times 10^{-2}$ and $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1203 \pm 0.0038$. Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1181 \pm 0.0020$.

⁴ AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0)] = 0.0186 \pm 0.0012 \pm 0.0011$ keV which we divide by our best value $B(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0) = (2.00 \pm 0.07) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁵ AUBERT 05D reports $[\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \mu^+\mu^-)] = 0.0450 \pm 0.0018 \pm 0.0022$ keV which we divide by our best value $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Superseded by LEES 12E.



$\Gamma(J/\psi(1S)\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{13}\Gamma_7/\Gamma$

| VALUE (keV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|-------------------------------------|-------------|---|---------|
| 0.425±0.012 OUR FIT | Error includes scale factor of 1.4. | | | |
| 0.413±0.019 OUR AVERAGE | | | | |
| 0.45 ± 0.13 ± 0.02 | 1 LEES | 23 BABR | e ⁺ e ⁻ → γ _{ISR} hadrons | |
| 0.45 ± 0.12 ± 0.04 | 2 LEES | 23 BABR | e ⁺ e ⁻ → γ _{ISR} hadrons | |
| 0.411±0.008±0.018 | 3.6k ADAM | 06 CLEO | 3.773 e ⁺ e ⁻ → γψ(2S) | |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.51 ± 0.09 ± 0.02 | 142 3 LEES | 18E BABR | 10.6 e ⁺ e ⁻ → J/ψπ ⁰ π ⁰ γ | |
| 1 LEES 23 reports [Γ(ψ(2S) → J/ψ(1S)π ⁰ π ⁰) × Γ(ψ(2S) → e ⁺ e ⁻)/Γ _{total}] × [B(J/ψ(1S) → K ⁺ K ⁻ π ⁰) = (1.31 ± 0.35 ± 0.13) × 10 ⁻³ keV which we divide by our best value B(J/ψ(1S) → K ⁺ K ⁻ π ⁰) = (2.88 ± 0.12) × 10 ⁻³ . Our first error is their experiment's error and our second error is the systematic error from using our best value. | | | | |
| 2 LEES 23 reports [Γ(ψ(2S) → J/ψ(1S)π ⁰ π ⁰) × Γ(ψ(2S) → e ⁺ e ⁻)/Γ _{total}] × [B(J/ψ(1S) → K _S ⁰ K [±] π [∓]) = (2.36 ± 0.59 ± 0.24) × 10 ⁻³ keV which we divide by our best value B(J/ψ(1S) → K _S ⁰ K [±] π [∓]) = (5.3 ± 0.5) × 10 ⁻³ . Our first error is their experiment's error and our second error is the systematic error from using our best value. | | | | |
| 3 LEES 18E reports [Γ(ψ(2S) → J/ψ(1S)π ⁰ π ⁰) × Γ(ψ(2S) → e ⁺ e ⁻)/Γ _{total}] × [B(J/ψ(1S) → π ⁺ π ⁻ π ⁰) = 0.0101 ± 0.0015 ± 0.0011 keV which we divide by our best value B(J/ψ(1S) → π ⁺ π ⁻ π ⁰) = (2.00 ± 0.07) × 10 ⁻² . Our first error is their experiment's error and our second error is the systematic error from using our best value. | | | | |

 $\Gamma(J/\psi(1S)\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{14}\Gamma_7/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-------------------------------------|-------------|--|---------|
| 78.6±1.8 OUR FIT | Error includes scale factor of 1.1. | | | |
| 87 ± 9 OUR AVERAGE | | | | |
| 83 ± 25 ± 5 | 14 1 AUBERT | 07AU BABR | 10.6 e ⁺ e ⁻ → J/ψπ ⁺ π ⁻ π ⁰ γ | |
| 88 ± 6 ± 7 | 291 ± 24 ADAM | 06 CLEO | 3.773 e ⁺ e ⁻ → γψ(2S) | |
| 1 AUBERT 07AU quotes Γ _{ee} ^{ψ(2S)} · B(ψ(2S) → J/ψη) · B(J/ψ → μ ⁺ μ ⁻) · B(η → π ⁺ π ⁻ π ⁰) = 1.11 ± 0.33 ± 0.07 eV. | | | | |

 $\Gamma(J/\psi(1S)\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{15}\Gamma_7/\Gamma$

| VALUE (eV) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------|-----|------|-------------|---------|--|
| <8 | 90 | <37 | ADAM | 06 CLEO | 3.773 e ⁺ e ⁻ → γψ(2S) |

 $\Gamma(2(\pi^+\pi^-)\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{22}\Gamma_7/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------|------------|-------------|--|---------|
| 29.7±2.2±1.8 | 410 AUBERT | 07AU BABR | 10.6 e ⁺ e ⁻ → 2(π ⁺ π ⁻)π ⁰ γ | |

 $\Gamma(\pi^+\pi^-\pi^0\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{24}\Gamma_7/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------|----------|-------------|--|---------|
| 12.4±1.8±1.2 | 177 LEES | 18E BABR | 10.6 e ⁺ e ⁻ → π ⁺ π ⁻ 3π ⁰ γ | |

 $\Gamma(\pi^+\pi^-4\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{26}\Gamma_7/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------|---------|-------------|---|---------|
| 3.3±2.3±0.5 | 18 LEES | 21C BABR | e ⁺ e ⁻ → γ _{ISR} (π ⁺ π ⁻ 4π ⁰) | |

$$\Gamma(\rho^\pm \pi^\mp \pi^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{25} \Gamma_7/\Gamma$$

| VALUE (eV) | CL% | DOCUMENT ID | TECN | COMMENT |
|------------|-----|-------------|----------|--|
| <6.2 | 90 | LEES | 18E BABR | $10.6 e^+ e^- \rightarrow \pi^+ \pi^- 3\pi^0 \gamma$ |

$$\Gamma(2(\pi^+ \pi^- \pi^0)) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{28} \Gamma_7/\Gamma$$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------------------------|------|-------------|----------|--|
| 11.2 ± 3.3 ± 1.3 | 43 | AUBERT | 06D BABR | $10.6 e^+ e^- \rightarrow 2(\pi^+ \pi^- \pi^0) \gamma$ |

$$\Gamma(2(\pi^+ \pi^-)3\pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{30} \Gamma_7/\Gamma$$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------------------|------|-------------|---------|--|
| 33 ± 5 ± 5 | 14k | LEES | 21 BABR | $10.6 e^+ e^- \rightarrow 2(\pi^+ \pi^-)3\pi^0 \gamma$ |

$$\Gamma(\eta 2(\pi^+ \pi^-)) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{33} \Gamma_7/\Gamma$$

| VALUE (eV) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------------|-----|----------|-------------|---|---------|
| 2.87 ± 1.41 ± 0.01 | 16 | 1 AUBERT | 07AU BABR | $10.6 e^+ e^- \rightarrow 2(\pi^+ \pi^-) \eta \gamma$ | |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|----|----|-----|-------------------|---------|--|
| <7 | 90 | 14k | ² LEES | 21 BABR | $10.6 e^+ e^- \rightarrow 2(\pi^+ \pi^-)3\pi^0 \gamma$ |
|----|----|-----|-------------------|---------|--|

¹ AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow \eta 2(\pi^+ \pi^-)) \times \Gamma(\psi(2S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = 1.13 \pm 0.55 \pm 0.08$ eV which we divide by our best value $B(\eta \rightarrow 2\gamma) = (39.36 \pm 0.18) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² LEES 21 reports $[\Gamma(\psi(2S) \rightarrow \eta 2(\pi^+ \pi^-)) \times \Gamma(\psi(2S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 3\pi^0)] < 2.3$ eV which we divide by our best value $B(\eta \rightarrow 3\pi^0) = 32.57 \times 10^{-2}$.

$$\Gamma(\eta \pi^+ \pi^- \pi^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{34} \Gamma_7/\Gamma$$

| VALUE (eV) | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------|-----|-------------|----------|--|
| <0.85 | 90 | LEES | 18E BABR | $10.6 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \pi^0 \eta \gamma$ |

$$\Gamma(\eta \pi^+ \pi^- 3\pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{35} \Gamma_7/\Gamma$$

| VALUE (eV) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------|-----|-------------------|----------|---|
| <5 | 90 | ¹ LEES | 21C BABR | $e^+ e^- \rightarrow \gamma_{ISR} (\pi^+ \pi^- 3\pi^0 \gamma \gamma)$ |

¹ LEES 21C reports $[\Gamma(\psi(2S) \rightarrow \eta \pi^+ \pi^- 3\pi^0) \times \Gamma(\psi(2S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] < 1.9$ eV which we divide by our best value $B(\eta \rightarrow 2\gamma) = 39.36 \times 10^{-2}$.

$$\Gamma(\eta 2(\pi^+ \pi^- \pi^0)) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{36} \Gamma_7/\Gamma$$

| VALUE (eV) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------|-----|------|-------------------|---------|--|
| <5 | 90 | 14k | ¹ LEES | 21 BABR | $10.6 e^+ e^- \rightarrow 2(\pi^+ \pi^-)3\pi^0 \gamma$ |

¹ LEES 21 reports $[\Gamma(\psi(2S) \rightarrow \eta 2(\pi^+ \pi^- \pi^0)) \times \Gamma(\psi(2S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] < 1.9$ eV which we divide by our best value $B(\eta \rightarrow 2\gamma) = 39.36 \times 10^{-2}$.

$\Gamma(\omega\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{41}\Gamma_7/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|------|-------------|-----------|--|
| 3.01±0.84±0.02 | 37 | 1 AUBERT | 07AU BABR | $10.6 e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$ |

¹ AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow \omega\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = 2.69 \pm 0.73 \pm 0.16$ eV which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\omega\pi^+\pi^-2\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{42}\Gamma_7/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------|------|-------------|------|---|
| 20.2±5.6±0.1 | 14k | 1 LEES | 21 | $10.6 e^+e^- \rightarrow 2(\pi^+\pi^-)3\pi^0\gamma$ |

¹ LEES 21 reports $[\Gamma(\psi(2S) \rightarrow \omega\pi^+\pi^-2\pi^0) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = 18 \pm 4 \pm 3$ eV which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\omega\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{45}\Gamma_7/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|------|-------------|----------|--|
| 2.58±0.82±0.02 | 33 | 1 LEES | 18E BABR | $10.6 e^+e^- \rightarrow \pi^+\pi^-3\pi^0\gamma$ |

¹ LEES 18E reports $[\Gamma(\psi(2S) \rightarrow \omega\pi^0\pi^0) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = 2.3 \pm 0.7 \pm 0.2$ eV which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\omega 3\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{46}\Gamma_7/\Gamma$

| VALUE (eV) | CL% | DOCUMENT ID | TECN | COMMENT |
|----------------|-----|-------------|----------|---|
| <1.8 | 90 | 1 LEES | 21C BABR | $e^+e^- \rightarrow \gamma_{ISR}(\pi^+\pi^-4\pi^0)$ |

¹ LEES 21C reports $[\Gamma(\psi(2S) \rightarrow \omega 3\pi^0) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] < 1.6$ eV which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = 89.2 \times 10^{-2}$.

 $\Gamma(\phi\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{51}\Gamma_7/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|------|-------------|----------|--|
| 0.55±0.19±0.01 | 19 | 1 LEES | 12F BABR | $10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.57 \pm 0.23 \pm 0.01$ 10 ² AUBERT,BE 06D BABR $10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$

¹ LEES 12F reports $[\Gamma(\psi(2S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 0.27 \pm 0.09 \pm 0.02$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.1 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Superseded by LEES 12F. AUBERT,BE 06D reports $[\Gamma(\psi(2S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 0.28 \pm 0.11 \pm 0.02$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.1 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi f_0(980) \rightarrow \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{52} \Gamma_7/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------|------|-------------|----------|---|
| 0.346 ± 0.129 ± 0.004 | 12 | 1 LEES | 12F BABR | $10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-----------------------|-------|----------|-----------|---|
| 0.346 ± 0.168 ± 0.004 | 6 ± 3 | 2 AUBERT | 07AK BABR | $10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$ |
|-----------------------|-------|----------|-----------|---|

¹ LEES 12F reports $[\Gamma(\psi(2S) \rightarrow \phi f_0(980) \rightarrow \pi^+ \pi^-) \times \Gamma(\psi(2S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 0.17 \pm 0.06 \pm 0.02 \text{ eV}$ which we divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.1 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(\psi(2S) \rightarrow \phi f_0(980) \rightarrow \pi^+ \pi^-) \times \Gamma(\psi(2S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 0.17 \pm 0.08 \pm 0.02 \text{ eV}$ which we divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.1 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(K^+ K^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{59} \Gamma_7/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------|------|-------------|------|---------|
|------------|------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-----------------------|----|--------|----------|--------------------------------------|
| 0.147 ± 0.035 ± 0.005 | 66 | 1 LEES | 15J BABR | $e^+ e^- \rightarrow K^+ K^- \gamma$ |
| 0.197 ± 0.035 ± 0.005 | 66 | 2 LEES | 15J BABR | $e^+ e^- \rightarrow K^+ K^- \gamma$ |
| 0.35 ± 0.14 ± 0.03 | 11 | 3 LEES | 13Q BABR | $e^+ e^- \rightarrow K^+ K^- \gamma$ |

¹ $\sin\phi > 0$.

² $\sin\phi < 0$.

³ Interference with non-resonant $K^+ K^-$ production not taken into account.

 $\Gamma(K^+ K^- \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{60} \Gamma_7/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------------|------|-------------|----------|---|
| 1.92 ± 0.30 ± 0.06 | 133 | LEES | 12F BABR | $10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|--------------------|----|----------|-----------|---|
| 2.56 ± 0.42 ± 0.16 | 85 | 1 AUBERT | 07AK BABR | $10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$ |
|--------------------|----|----------|-----------|---|

¹ Superseded by LEES 12F.

 $\Gamma(K_S^0 K_L^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{64} \Gamma_7/\Gamma$

| VALUE (eV) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------|-----|------|-------------|----------|--|
| <0.7 | 90 | 8 | LEES | 17A BABR | $e^+ e^- \rightarrow K_S^0 K_L^0 \pi^0 \gamma$ |

 $\Gamma(K^+ K^- \pi^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{65} \Gamma_7/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------------|------|-------------|----------|---|
| 0.60 ± 0.31 ± 0.03 | 17 | LEES | 12F BABR | $10.6 e^+ e^- \rightarrow \pi^0 \pi^0 K^+ K^- \gamma$ |

 $\Gamma(K^+ K^- \pi^0 \pi^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{66} \Gamma_7/\Gamma$

| VALUE (eV) | DOCUMENT ID | TECN | COMMENT |
|---------------------------|-------------|---------|--|
| 1.54 ± 0.63 ± 0.15 | LEES | 23 BABR | $e^+ e^- \rightarrow \gamma_{ISR} \text{ hadrons}$ |

| $\Gamma(K_S^0 K^\pm \pi^\mp \pi^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ | | | | | $\Gamma_{67} \Gamma_7/\Gamma$ |
|--|------|---------------------|------|--|-------------------------------|
| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| 4.0±1.4±0.4 | | LEES | 23 | BABR $e^+ e^- \rightarrow \gamma_{ISR}$ hadrons | ■ |
| $\Gamma(K_S^0 K^\pm \pi^\mp \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ | | | | | $\Gamma_{68} \Gamma_7/\Gamma$ |
| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| 5.1±0.7±0.4 | | LEES | 23 | BABR $e^+ e^- \rightarrow \gamma_{ISR}$ hadrons | ■ |
| $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ | | | | | $\Gamma_{69} \Gamma_7/\Gamma$ |
| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| 4.4±1.3±0.3 | 32 | AUBERT | 07AU | BABR $10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \pi^0 \gamma$ | |
| $\Gamma(K_S^0 K_L^0 \pi^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ | | | | | $\Gamma_{76} \Gamma_7/\Gamma$ |
| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| 2.92±1.27±0.15 | 14 | LEES | 17A | BABR $e^+ e^- \rightarrow K_S^0 K_L^0 \pi^0 \pi^0 \gamma$ | |
| $\Gamma(K_S^0 K^*(892)^0 \pi^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ | | | | | $\Gamma_{77} \Gamma_7/\Gamma$ |
| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| 0.71±0.29±0.07 | | LEES | 23 | BABR $e^+ e^- \rightarrow \gamma_{ISR}$ hadrons | ■ |
| $\Gamma(K_S^0 K^\pm \rho(770)^\mp \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ | | | | | $\Gamma_{78} \Gamma_7/\Gamma$ |
| VALUE (eV) | CL% | DOCUMENT ID | TECN | COMMENT | |
| <1.6 | 90 | LEES | 23 | BABR $e^+ e^- \rightarrow \gamma_{ISR}$ hadrons | ■ |
| $\Gamma(K_S^0 K^\pm \rho(770)^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ | | | | | $\Gamma_{79} \Gamma_7/\Gamma$ |
| VALUE (eV) | CL% | DOCUMENT ID | TECN | COMMENT | |
| <1.6 | 90 | LEES | 23 | BABR $e^+ e^- \rightarrow \gamma_{ISR}$ hadrons | ■ |
| $\Gamma(K^*(892)^+ K^*(892)^- \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ | | | | | $\Gamma_{81} \Gamma_7/\Gamma$ |
| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| 8.46±4.05±0.90 | | LEES | 23 | BABR $e^+ e^- \rightarrow \gamma_{ISR}$ hadrons | ■ |
| $\Gamma(K^\mp K^*(892)^\pm \pi^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ | | | | | $\Gamma_{80} \Gamma_7/\Gamma$ |
| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| 1.62±0.66±0.15 | | LEES | 23 | BABR $e^+ e^- \rightarrow \gamma_{ISR}$ hadrons | ■ |
| $\Gamma(K_S^0 K_L^0 \eta) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ | | | | | $\Gamma_{82} \Gamma_7/\Gamma$ |
| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| 3.14±1.08±0.16 | 16 | LEES | 17A | BABR $e^+ e^- \rightarrow K_S^0 K_L^0 \eta \gamma$ | |
| $\Gamma(K^+ K^- \pi^+ \pi^- \eta) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ | | | | | $\Gamma_{85} \Gamma_7/\Gamma$ |
| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| 3.05±1.80±0.01 | 7 | ¹ AUBERT | 07AU | BABR $10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \eta \gamma$ | |

¹ AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \eta) \times \Gamma(\psi(2S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = 1.2 \pm 0.7 \pm 0.1$ eV which we divide by our best value $B(\eta \rightarrow 2\gamma) = (39.36 \pm 0.18) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

| $\Gamma(K^+ K^- 2(\pi^+ \pi^-)) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ | | | | $\Gamma_{86} \Gamma_7/\Gamma$ |
|---|------|-------------|------|---|
| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
| 4.4±2.1±0.3 | 26 | AUBERT | 06D | BABR $10.6 e^+ e^- \rightarrow K^+ K^- 2(\pi^+ \pi^-) \gamma$ |

| $\Gamma(2(K^+ K^-)) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ | | | | $\Gamma_{89} \Gamma_7/\Gamma$ |
|---|------|-------------|------|--|
| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
| 0.22±0.10±0.02 | 13 | LEES | 12F | BABR $10.6 e^+ e^- \rightarrow K^+ K^- K^+ K^- \gamma$ |

| $\Gamma(p\bar{p}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ | | | | $\Gamma_{111} \Gamma_7/\Gamma$ |
|---|------|-------------------------------------|------|--|
| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
| 0.686±0.024 OUR FIT | | Error includes scale factor of 1.2. | | |
| 0.63 ±0.05 OUR AVERAGE | | Error includes scale factor of 1.2. | | |
| 0.67 ±0.12 ±0.02 | 43 | ¹ LEES | 130 | BABR $e^+ e^- \rightarrow p\bar{p}\gamma$ |
| 0.74 ±0.07 ±0.04 | 142 | ² LEES | 13Y | BABR $e^+ e^- \rightarrow p\bar{p}\gamma$ |
| 0.579±0.038±0.036 | 2.7k | ANDREOTTI | 07 | E835 $p\bar{p} \rightarrow e^+ e^-$, $J/\psi X$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.70 ±0.17 ±0.03 | 22 | ³ AUBERT | 06B | BABR $e^+ e^- \rightarrow p\bar{p}\gamma$ |
| ¹ ISR photon reconstructed in the detector | | | | |
| ² ISR photon undetected | | | | |
| ³ Superseded by LEES 130 | | | | |

| $\Gamma(\Lambda\bar{\Lambda}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ | | | | $\Gamma_{134} \Gamma_7/\Gamma$ |
|---|------|-------------|------|--|
| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
| 1.5±0.4±0.1 | | AUBERT | 07BD | BABR $10.6 e^+ e^- \rightarrow \Lambda\bar{\Lambda}\gamma$ |

$\psi(2S)$ BRANCHING RATIOS

| $\Gamma(\text{hadrons})/\Gamma_{\text{total}}$ | | | | Γ_1/Γ |
|--|-------------------|------|----------------|-------------------|
| VALUE | DOCUMENT ID | TECN | COMMENT | |
| 0.9785±0.0013 OUR AVERAGE | | | | |
| 0.9779±0.0015 | ¹ BAI | 02B | BES2 $e^+ e^-$ | |
| 0.981 ±0.003 | ¹ LUTH | 75 | MRK1 $e^+ e^-$ | |

¹ Includes cascade decay into $J/\psi(1S)$.

| $\Gamma(\text{virtual } \gamma \rightarrow \text{hadrons})/\Gamma_{\text{total}}$ | | | | Γ_2/Γ |
|---|-------------------|------|----------------|-------------------|
| VALUE | DOCUMENT ID | TECN | COMMENT | |
| 0.0179±0.0004 | ¹ LIAO | 23 | RVUE $e^+ e^-$ | |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|---------------|---------------------|-----|----------------|
| 0.0166±0.0010 | ^{2,3} SETH | 04 | RVUE $e^+ e^-$ |
| 0.0199±0.0019 | ² BAI | 02B | BES2 $e^+ e^-$ |
| 0.029 ±0.004 | ² LUTH | 75 | MRK1 $e^+ e^-$ |

¹ Using $B(\psi(2S) \rightarrow \ell^+ \ell^-) = (0.794 \pm 0.017)\%$ and $R = 2.26 \pm 0.01$ determined by a fit to data from Mark-I, DM2, BESII, KEDR, and BESIII.

² Included in $\Gamma(\text{hadrons})/\Gamma_{\text{total}}$.

³ Using $B(\psi(2S) \rightarrow \ell^+ \ell^-) = (0.73 \pm 0.04)\%$ from RPP-2002 and $R = 2.28 \pm 0.04$ determined by a fit to data from BAI 00 and BAI 02C. Superseded by LIAO 23.

$\Gamma(ggg)/\Gamma_{\text{total}}$

| <u>VALUE (units 10^{-2})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|---------------------------------------|
| 10.58 ± 1.62 | 2.9 M | 1 LIBBY | 09 CLEO | $\psi(2S) \rightarrow \text{hadrons}$ |

¹ Calculated using $\Gamma(\gamma gg)/\Gamma(ggg) = 0.097 \pm 0.026 \pm 0.016$ from LIBBY 09, $B(\psi(2S) \rightarrow X J/\psi)$ relative and absolute branching fractions from MENDEZ 08, $B(\psi(2S) \rightarrow \gamma \eta_c)$ from MITCHELL 09, and $B(\psi(2S) \rightarrow \text{virtual } \gamma \rightarrow \text{hadrons})$, $B(\psi(2S) \rightarrow \gamma \chi_{cJ})$, and $B(\psi(2S) \rightarrow \ell^+ \ell^-)$ from PDG 08. The statistical error is negligible and the systematic error is largely uncorrelated with that of $\Gamma(\gamma gg)/\Gamma_{\text{total}}$ LIBBY 09 measurement.

 $\Gamma(\gamma gg)/\Gamma_{\text{total}}$

| <u>VALUE (units 10^{-2})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|--|
| 1.025 ± 0.288 | 200 k | 1 LIBBY | 09 CLEO | $\psi(2S) \rightarrow \gamma + \text{hadrons}$ |

¹ Calculated using $\Gamma(\gamma gg)/\Gamma(ggg) = 0.097 \pm 0.026 \pm 0.016$ from LIBBY 09. The statistical error is negligible and the systematic error is largely uncorrelated with that of $\Gamma(ggg)/\Gamma_{\text{total}}$ LIBBY 09 measurement.

 $\Gamma(\gamma gg)/\Gamma(ggg)$

| <u>VALUE (units 10^{-2})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|---|
| $9.7 \pm 2.6 \pm 1.6$ | 2.9 M | LIBBY | 09 CLEO | $\psi(2S) \rightarrow (\gamma +) \text{ hadrons}$ |

 $\Gamma(\text{light hadrons})/\Gamma_{\text{total}}$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-------------------------------------|--------------------|-------------|--------------------------------|
| 0.154 ± 0.015 | 1 MENDEZ | 08 CLEO | $e^+ e^- \rightarrow \psi(2S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|-------------------|--------|----------|--------------------------------|
| 0.169 ± 0.026 | 2 ADAM | 05A CLEO | $e^+ e^- \rightarrow \psi(2S)$ |
|-------------------|--------|----------|--------------------------------|

¹ Uses $B(\psi(2S) \rightarrow J/\psi X)$ from MENDEZ 08 and other branching fractions from PDG 07.

² Uses $B(J/\psi X)$ from ADAM 05A, $B(\chi_{cJ} \gamma)$, $B(\eta_c \gamma)$ from ATHAR 04 and $B(\ell^+ \ell^-)$ from PDG 04. Superseded by MENDEZ 08.

 $\Gamma(e^+ e^-)/\Gamma_{\text{total}}$

| <u>VALUE (units 10^{-4})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|-------------------------------------|-------------|----------------|
| $79.4 \pm 2.2 \text{ OUR FIT}$ | Error includes scale factor of 1.3. | | |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|---------|-----------|---------|-----------|
| 88 ± 13 | 1 FELDMAN | 77 RVUE | $e^+ e^-$ |
|---------|-----------|---------|-----------|

¹ From an overall fit assuming equal partial widths for $e^+ e^-$ and $\mu^+ \mu^-$. For a measurement of the ratio see the entry $\Gamma(\mu^+ \mu^-)/\Gamma(e^+ e^-)$ below. Includes LUTH 75, HILGER 75, BURMESTER 77.

 $\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$

| <u>VALUE (units 10^{-4})</u> | <u>DOCUMENT ID</u> |
|--|--------------------|
| $80 \pm 6 \text{ OUR FIT}$ | |

 $\Gamma(\mu^+ \mu^-)/\Gamma(e^+ e^-)$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|----------------|
| $1.00 \pm 0.08 \text{ OUR FIT}$ | | | |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|-------------|----------|----------|-----------|
| 0.89 ± 0.16 | BOYARSKI | 75C MRK1 | $e^+ e^-$ |
|-------------|----------|----------|-----------|

 Γ_3/Γ Γ_4/Γ Γ_5/Γ Γ_7/Γ Γ_8/Γ Γ_8/Γ_7

$\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}}$ Γ_9/Γ VALUE (units 10^{-4})DOCUMENT IDTECNCOMMENT**31 ±4 OUR FIT****30.8±2.1±3.8**1 ABLIKIM 06W BES $e^+e^- \rightarrow \psi(2S)$

¹ Computed using PDG 02 value of $B(\psi(2S) \rightarrow \text{hadrons}) = 0.9810 \pm 0.0030$ to estimate the total number of $\psi(2S)$ events.

— DECAYS INTO $J/\psi(1S)$ AND ANYTHING — $\Gamma(J/\psi(1S)\text{anything})/\Gamma_{\text{total}}$

$$\Gamma_{10}/\Gamma = (\Gamma_{12} + \Gamma_{13} + \Gamma_{14} + 0.343\Gamma_{180} + 0.195\Gamma_{181})/\Gamma$$

VALUE EVTS DOCUMENT ID TECN COMMENT**0.615 ±0.007 OUR FIT** Error includes scale factor of 1.3.**0.55 ±0.07 OUR AVERAGE**

| | |
|-----------------|---|
| 0.51 ± 0.12 | BRANDELIK 79C DASP $e^+e^- \rightarrow \mu^+\mu^-X$ |
| 0.57 ± 0.08 | ABRAMS 75B MRK1 $e^+e^- \rightarrow \mu^+\mu^-X$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | |
|--------------------------------|--|
| 0.644 $\pm 0.006 \pm 0.016$ | 1 ABLIKIM 21Z BES3 $e^+e^- \rightarrow \ell^+\ell^-X$ |
| 0.6254 $\pm 0.0016 \pm 0.0155$ | 1.1M 2 MENDEZ 08 CLEO $\psi(2S) \rightarrow \ell^+\ell^-X$ |
| 0.5950 $\pm 0.0015 \pm 0.0190$ | 151k ADAM 05A CLEO Repl. by MENDEZ 08 |


¹ From a fit to the $e^+e^- \rightarrow J/\psi X$ cross section between 3.645 and 3.891 GeV, with $\Gamma(ee)$ and Γ fixed to the PDG 20 values of the cross particle fit which are correlated to "OUR FIT" value for $B(\psi(2S) \rightarrow J/\psi X)$.

² Not independent from other measurements of MENDEZ 08.

 $\Gamma(e^+e^-)/\Gamma(J/\psi(1S)\text{anything})$ Γ_7/Γ_{10} VALUE (units 10^{-2}) EVTS DOCUMENT ID TECN COMMENT**1.291±0.035 OUR FIT** Error includes scale factor of 1.3.**1.28 ±0.04 OUR AVERAGE** Error includes scale factor of 1.6. See the ideogram below.

| | | |
|--------------------------|---------------|--|
| 1.22 $\pm 0.02 \pm 0.05$ | 5097 ± 73 | 1 ANDREOTTI 05 E835 $p\bar{p} \rightarrow \psi(2S) \rightarrow e^+e^-$ |
| 1.28 $\pm 0.03 \pm 0.02$ | | 1 AMBROGIANI 00A E835 $p\bar{p} \rightarrow \psi(2S)$ |
| 1.44 $\pm 0.08 \pm 0.02$ | | 1 ARMSTRONG 97 E760 $\bar{p}p \rightarrow \psi(2S)$ |

¹ Using $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0593 \pm 0.0010$.



$\Gamma(\mu^+ \mu^-)/\Gamma(J/\psi(1S)\text{anything})$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------------------|-------------|------|----------------|
| 0.0130±0.0010 OUR FIT | | | |
| 0.014 ± 0.003 | HILGER | 75 | SPEC $e^+ e^-$ |


$\Gamma(J/\psi(1S)\text{ neutrals})/\Gamma_{\text{total}}$

| VALUE | DOCUMENT ID |
|----------------------------|-------------------------------------|
| 0.254±0.005 OUR FIT | Error includes scale factor of 1.6. |

$\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------------|------|---|
| 0.3469±0.0034 OUR FIT | | | | Error includes scale factor of 1.1. |
| 0.348 ± 0.005 OUR AVERAGE | | | | Error includes scale factor of 1.3. See the ideogram below. |
| 0.3498±0.0002±0.0045 | 20M | ABLIKIM | 13R | BES3 $\psi(2S) \rightarrow J/\psi\pi^+\pi^-$ |
| 0.3504±0.0007±0.0077 | 565k | MENDEZ | 08 | CLEO $\psi(2S) \rightarrow \ell^+\ell^-\pi^+\pi^-$ |
| 0.323 ± 0.014 | | BAI | 02B | BES2 $e^+ e^-$ |
| 0.32 ± 0.04 | | ABRAMS | 75B | MRK1 $e^+ e^- \rightarrow J/\psi\pi^+\pi^-$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.3354±0.0014±0.0110 | 60k | ¹ ADAM | 05A | CLEO Repl. by MENDEZ 08 |

¹ Not independent from other values reported by ADAM 05A.



$\Gamma(e^+e^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$

Γ_7/Γ_{12}

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------------------|-------------------------------------|----------|----------|
| 0.0229±0.0006 OUR FIT | Error includes scale factor of 1.3. | | |
| 0.0252±0.0028±0.0011 | ¹ AUBERT | 02B BABR | e^+e^- |

¹ Using $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0593 \pm 0.0010$.

$\Gamma(\mu^+\mu^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$

Γ_8/Γ_{12}

| VALUE | DOCUMENT ID | TECN | COMMENT |
|----------------------------------|------------------------|----------|-------------------------------------|
| 0.0230±0.0017 OUR FIT | | | |
| 0.0228±0.0018 OUR AVERAGE | | | |
| 0.0230±0.0020±0.0012 | ¹ AAIJ | 16Y LHCb | $\Lambda_b^0 \rightarrow \psi(2S)X$ |
| 0.0216±0.0026±0.0014 | ² AUBERT | 02B BABR | e^+e^- |
| 0.0327±0.0077±0.0072 | ² GRIBUSHIN | 96 FMPS | 515 π^- Be $\rightarrow 2\mu X$ |

¹ Using $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033) \times 10^{-2}$.

² Using $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = (5.88 \pm 0.10) \times 10^{-2}$.

$\Gamma(\tau^+\tau^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$

Γ_9/Γ_{12}

| VALUE (units 10^{-3}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|--------|----------|
| 8.8 ±1.1 OUR FIT | | | |
| 8.73±1.39±1.57 | BAI | 02 BES | e^+e^- |


$\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma(J/\psi(1S)\text{anything})$

Γ_{12}/Γ_{10}

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-----------------|------|---|
| 0.564 ± 0.004 OUR FIT | | | | Error includes scale factor of 1.7. |
| 0.554 ± 0.008 OUR AVERAGE | | | | Error includes scale factor of 1.3. See the ideogram below. |
| 0.5604 ± 0.0009 ± 0.0062 | 565k | MENDEZ 08 | CLEO | $\psi(2S) \rightarrow \ell^+\ell^-\pi^+\pi^-$ |
| 0.525 ± 0.009 ± 0.022 | 4k | ANDREOTTI 05 | E835 | $\psi(2S) \rightarrow J/\psi X$ |
| 0.536 ± 0.007 ± 0.016 | 20k | 1,2 ABLIKIM 04B | BES | $\psi(2S) \rightarrow J/\psi X$ |
| 0.496 ± 0.037 | | ARMSTRONG 97 | E760 | $\bar{p}p \rightarrow \psi(2S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.5637 ± 0.0027 ± 0.0046 | 60k | ADAM 05A | CLEO | Repl. by MENDEZ 08 |

¹ From a fit to the J/ψ recoil mass spectra.

² ABLIKIM 04B quotes $B(\psi(2S) \rightarrow J/\psi X) / B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-)$.



$\Gamma(J/\psi(1S)\text{ neutrals})/\Gamma(J/\psi(1S)\pi^+\pi^-)$

$$\Gamma_{11}/\Gamma_{12} = (0.9761\Gamma_{13} + 0.719\Gamma_{14} + 0.343\Gamma_{180} + 0.195\Gamma_{181})/\Gamma_{12}$$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------------------|--------------|------|-------------------------------------|
| 0.732 ± 0.013 OUR FIT | | | Error includes scale factor of 1.7. |
| 0.73 ± 0.09 | TANENBAUM 76 | MRK1 | e^+e^- |

$\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma(J/\psi(1S)\text{anything})$ Γ_{13}/Γ_{10}

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-------------------------------------|-------------|----------------|--|
| 0.297 ± 0.005 OUR FIT | Error includes scale factor of 1.7. | | | |
| 0.320 ± 0.012 OUR AVERAGE | | | | |
| 0.300 ± 0.008 | ± 0.022 | 1655 ± 44 | ANDREOTTI 05 | $\psi(2S) \rightarrow J/\psi X$ |
| 0.328 ± 0.013 | ± 0.008 | | AMBROGIANI 00A | $p\bar{p} \rightarrow \psi(2S)$ |
| 0.323 ± 0.033 | | | ARMSTRONG 97 | $\bar{p}p \rightarrow \psi(2S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.2829 ± 0.0012 | ± 0.0056 | 61k | MENDEZ 08 | CLEO $\psi(2S) \rightarrow \ell^+\ell^-2\pi^0$ |
| 0.2776 ± 0.0025 | ± 0.0043 | 13.4k | ADAM 05A | CLEO Repl. by MENDEZ 08 |

$\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma(J/\psi(1S)\pi^+\pi^-)$ Γ_{13}/Γ_{12}

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-------------------------------------|-------------|---------------------------|--|
| 0.526 ± 0.013 OUR FIT | Error includes scale factor of 1.7. | | | |
| 0.513 ± 0.022 OUR AVERAGE | Error includes scale factor of 2.2. | | | |
| 0.5047 ± 0.0022 | ± 0.0102 | 61k | MENDEZ 08 | CLEO $\psi(2S) \rightarrow \ell^+\ell^-2\pi^0$ |
| 0.570 ± 0.009 | ± 0.026 | 14k | ¹ ABLIKIM 04B | BES $\psi(2S) \rightarrow J/\psi X$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.4924 ± 0.0047 | ± 0.0086 | 73k | ^{2,3} ADAM 05A | CLEO Repl. by MENDEZ 08 |
| 0.571 ± 0.018 | ± 0.044 | | ⁴ ANDREOTTI 05 | E835 $\psi(2S) \rightarrow J/\psi X$ |
| 0.53 ± 0.06 | | | TANENBAUM 76 | MRK1 e^+e^- |
| 0.64 ± 0.15 | | | ⁵ HILGER 75 | SPEC e^+e^- |

¹ From a fit to the J/ψ recoil mass spectra.

² Not independent from other values reported by ADAM 05A.

³ Using 13,217 $J/\psi\pi^0\pi^0$ and 60,010 $J/\psi\pi^+\pi^-$ events.

⁴ Not independent from other values reported by ANDREOTTI 05.

⁵ Ignoring the $J/\psi(1S)\eta$ and $J/\psi(1S)\gamma\gamma$ decays.

$\Gamma(J/\psi(1S)\eta)/\Gamma_{\text{total}}$ Γ_{14}/Γ


| VALUE (units 10^{-3}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|---|-------------|----------------------------|--|
| 33.7 ± 0.6 OUR FIT | Error includes scale factor of 1.2. | | | |
| 32.9 ± 1.7 OUR AVERAGE | Error includes scale factor of 2.1. See the ideogram below. | | | |
| 33.75 ± 0.17 | ± 0.86 | 68.2k | ABLIKIM 12M | $e^+e^- \rightarrow \ell^+\ell^-2\gamma$ |
| 29.8 ± 0.9 | ± 2.3 | 5.7k | BAI 04I | BES2 $\psi(2S) \rightarrow J/\psi\gamma\gamma$ |
| 25.5 ± 2.9 | | 386 | ¹ OREGLIA 80 | CBAL $e^+e^- \rightarrow J/\psi 2\gamma$ |
| 45 ± 12 | | 17 | ² BRANDELIK 79B | DASP $e^+e^- \rightarrow J/\psi 2\gamma$ |
| 42 ± 6 | | 164 | ² BARTEL 78B | CNTR e^+e^- |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 34.3 ± 0.4 | ± 0.9 | 18.4k | ³ MENDEZ 08 | CLEO $\psi(2S) \rightarrow \ell^+\ell^-\eta$ |
| 32.5 ± 0.6 | ± 1.1 | 2.8k | ⁴ ADAM 05A | CLEO Repl. by MENDEZ 08 |
| 43 ± 8 | | 44 | TANENBAUM 76 | MRK1 e^+e^- |

¹ Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1181 \pm 0.0020$.

² Recalculated by us using $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = 0.0588 \pm 0.0010$.

³ Not independent from other measurements of MENDEZ 08.

⁴ Not independent from other values reported by ADAM 05A.




$\Gamma(J/\psi(1S)\eta)/\Gamma(J/\psi(1S)\text{anything})$

Γ_{14}/Γ_{10}

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-------------|-----------------------------|---|---|
| 0.0549 ± 0.0009 OUR FIT | | | | Error includes scale factor of 1.2. |
| 0.058 ± 0.007 OUR AVERAGE | | | | Error includes scale factor of 1.4. See the ideogram below. |
| 0.050 ± 0.006 | ± 0.003 | 298 ± 20 | ANDREOTTI 05 E835 | $\psi(2S) \rightarrow J/\psi X$ |
| 0.072 ± 0.009 | | | AMBROGIANI 00A E835 | $p\bar{p} \rightarrow \psi(2S)$ |
| 0.061 ± 0.015 | | | ARMSTRONG 97 E760 | $\bar{p}p \rightarrow \psi(2S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| $0.0549 \pm 0.0006 \pm 0.0009$ | 18.4k | ¹ MENDEZ 08 CLEO | $\psi(2S) \rightarrow \ell^+ \ell^- \eta$ | |
| $0.0546 \pm 0.0010 \pm 0.0007$ | 2.8k | ADAM 05A CLEO | Repl. by MENDEZ 08 | |

¹ Not independent from other measurements of MENDEZ 08.



$\Gamma(J/\psi(1S)\eta)/\Gamma(J/\psi(1S)\pi^+\pi^-)$

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-------|-------------------------------------|------|---|
| 0.0972±0.0016 OUR FIT | | Error includes scale factor of 1.1. | | |
| 0.0979±0.0018 OUR AVERAGE | | | | |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.0979±0.0010±0.0015 | 18.4k | MENDEZ 08 | CLEO | $\psi(2S) \rightarrow \ell^+ \ell^- \eta$ |
| 0.098 ± 0.005 ± 0.010 | 2k | ¹ ABLIKIM 04B | BES | $\psi(2S) \rightarrow J/\psi X$ |
| 0.091 ± 0.021 | | ² HIMEL 80 | MRK2 | $e^+ e^- \rightarrow \psi(2S) X$ |
| 0.0968±0.0019±0.0013 | 2.8k | ³ ADAM 05A | CLEO | Repl. by MENDEZ 08 |
| 0.095 ± 0.007 ± 0.007 | | ⁴ ANDREOTTI 05 | E835 | $\psi(2S) \rightarrow J/\psi X$ |

¹ From a fit to the J/ψ recoil mass spectra.

² The value for $B(\psi(2S) \rightarrow J/\psi(1s)\eta)$ reported in HIMEL 80 is derived using $B(\psi(2S)) \rightarrow J/\psi(1S)\pi^+\pi^-) = (33 \pm 3)\%$ and $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.138 \pm 0.018$. Calculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = (0.1181 \pm 0.0020)$.

³ Not independent from other values reported by ADAM 05A.

⁴ Not independent from other values reported by ANDREOTTI 05.

$\Gamma(J/\psi(1S)\pi^0)/\Gamma_{\text{total}}$

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------------------|------|--|
| 12.68±0.32 OUR AVERAGE | | | | |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 12.6 ± 0.2 ± 0.3 | 4.1k | ABLIKIM 12M | BES3 | $e^+ e^- \rightarrow \ell^+ \ell^- 2\gamma$ |
| 13.3 ± 0.8 ± 0.3 | 530 | MENDEZ 08 | CLEO | $\psi(2S) \rightarrow \ell^+ \ell^- 2\gamma$ |
| 14.3 ± 1.4 ± 1.2 | 280 | BAI 04I | BES2 | $\psi(2S) \rightarrow J/\psi\gamma\gamma$ |
| 14 ± 6 | 7 | HIMEL 80 | MRK2 | $e^+ e^-$ |
| 9 ± 2 ± 1 | 23 | ¹ OREGLIA 80 | CBAL | $\psi(2S) \rightarrow J/\psi 2\gamma$ |
| 13 ± 1 ± 1 | 88 | ADAM 05A | CLEO | Repl. by MENDEZ 08 |

¹ Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$.

$$\Gamma(J/\psi(1S)\pi^0)/\Gamma(J/\psi(1S)\text{anything})$$

$$\Gamma_{15}/\Gamma_{10} = \Gamma_{15}/(\Gamma_{12} + \Gamma_{13} + \Gamma_{14} + 0.343\Gamma_{180} + 0.195\Gamma_{181})$$

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|------------------------|--|---------|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| $0.213 \pm 0.012 \pm 0.003$ | 527 | ¹ MENDEZ 08 | CLEO $e^+ e^- \rightarrow J/\psi \gamma \gamma$ | |
| $0.22 \pm 0.02 \pm 0.01$ | | ² ADAM 05A | CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow J/\psi \gamma \gamma$ | |

¹ Not independent from other values reported by MENDEZ 08. Supersedes ADAM 05A.

² Not independent from other values reported by ADAM 05A.

$$\Gamma(J/\psi(1S)\pi^0)/\Gamma(J/\psi(1S)\pi^+\pi^-)$$

$$\Gamma_{15}/\Gamma_{12}$$

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|------------------------|--|---------|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| $0.380 \pm 0.022 \pm 0.005$ | 527 | ¹ MENDEZ 08 | CLEO $e^+ e^- \rightarrow J/\psi \gamma \gamma$ | |
| $0.39 \pm 0.04 \pm 0.01$ | | ² ADAM 05A | CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow J/\psi \gamma \gamma$ | |

¹ Not independent from other values reported by MENDEZ 08. Supersedes ADAM 05A.

² Not independent from other values reported by ADAM 05A.

HADRONIC DECAYS

$$\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$$

$$\Gamma_{16}/\Gamma$$

| VALUE (units 10^{-5}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|-----|---------------------------|-------------|-------------------------------------|---------|
| 0.78 ± 0.26 OUR AVERAGE | | | | | |
| $0.76 \pm 0.25 \pm 0.06$ | 30 | ¹ METREVELI 12 | | $\psi(2S) \rightarrow \pi^+ \pi^-$ | |
| 8 ± 5 | | BRANDELIK 79c | DASP | $e^+ e^-$ | |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| <2.1 | 90 | DOBBS | 06A | CLEO $e^+ e^- \rightarrow \psi(2S)$ | |
| <5 | 90 | FELDMAN | 77 | MRK1 $e^+ e^-$ | |

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration. Using $\psi(3770) \rightarrow \pi^+ \pi^-$ for continuum subtraction.


$$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$$

$$\Gamma_{17}/\Gamma$$

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|--------------|--------------------------|------|--------------------------------------|
| 2.01 ± 0.17 OUR AVERAGE Error includes scale factor of 1.7. See the ideogram below. | | | | |
| $2.14 \pm 0.03^{+0.12}_{-0.11}$ | 7k | ¹ ABLIKIM 12H | BES3 | $e^+ e^- \rightarrow \psi(2S)$ |
| $1.81 \pm 0.18 \pm 0.19$ | 260 ± 19 | ² ABLIKIM 05J | BES2 | $e^+ e^- \rightarrow \psi(2S)$ |
| $1.88^{+0.16}_{-0.15} \pm 0.28$ | 194 | ADAM 05 | CLEO | $e^+ e^- \rightarrow \psi(2S)$ |
| 0.85 ± 0.46 | 4 | FRANKLIN 83 | MRK2 | $e^+ e^- \rightarrow \text{hadrons}$ |

¹ From $\psi(2S) \rightarrow \pi^+ \pi^- \pi^0$ events directly. The quoted systematic error includes a contribution of 4% (added in quadrature) from the uncertainty on the number of $\psi(2S)$ events.

² From a PW analysis of $\psi(2S) \rightarrow \pi^+ \pi^- \pi^0$.



$\Gamma(\rho(770)\pi \rightarrow \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$

Γ_{18}/Γ

| VALUE (units 10^{-4}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|-------------------------------------|------|----------------------|----------|---|
| 0.32±0.12 OUR AVERAGE | Error includes scale factor of 1.8. | | | | |
| 0.51±0.07±0.11 | | | ¹ ABLIKIM | 05J BES2 | $\psi(2S) \rightarrow \rho(770)\pi \rightarrow \pi^+ \pi^- \pi^0$ |
| 0.24 ^{+0.08} _{-0.07} ±0.02 | | 22 | ADAM | 05 CLEO | $e^+ e^- \rightarrow \psi(2S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | | |
|-------|----|---|---------------------|----|------|-----------|
| <0.83 | 90 | 1 | FRANKLIN | 83 | MRK2 | $e^+ e^-$ |
| <10 | 90 | | BARTEL | 76 | CNTR | $e^+ e^-$ |
| <10 | 90 | | ² ABRAMS | 75 | MRK1 | $e^+ e^-$ |

¹ From a PW analysis of $\psi(2S) \rightarrow \pi^+ \pi^- \pi^0$.

² Final state $\rho^0 \pi^0$.

$\Gamma(\rho(2150)\pi \rightarrow \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$

Γ_{19}/Γ

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|--|----------------------|----------|--|
| 1.94±0.25^{+1.15}_{-0.34} | ¹ ABLIKIM | 05J BES2 | $\psi(2S) \rightarrow \rho(2150)\pi \rightarrow \pi^+ \pi^- \pi^0$ |

¹ From a PW analysis of $\psi(2S) \rightarrow \pi^+ \pi^- \pi^0$.

$\Gamma(2(\pi^+\pi^-))/\Gamma_{\text{total}}$

Γ_{20}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------|-------------------------------------|-------------|------|---|
| 2.4±0.6 OUR AVERAGE | Error includes scale factor of 2.2. | | | |
| 2.2±0.2±0.2 | 308 | BRIERE | 05 | CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+\pi^-)$ |
| 4.5±1.0 | | TANENBAUM | 78 | MRK1 $e^+ e^-$ |

$\Gamma(\rho^0\pi^+\pi^-)/\Gamma_{\text{total}}$

Γ_{21}/Γ


| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------|-------------------------------------|-------------|------|---|
| 2.2±0.6 OUR AVERAGE | Error includes scale factor of 1.4. | | | |
| 2.0±0.2±0.4 | 285.5 | BRIERE | 05 | CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+\pi^-)$ |
| 4.2±1.5 | | TANENBAUM | 78 | MRK1 $e^+ e^-$ |

$\Gamma(2(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$

Γ_{22}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------------|---|---------------------|------|--|
| 29 ±10 OUR AVERAGE | Error includes scale factor of 4.7. See the ideogram below. | | | |
| 24.9± 0.7±3.6 | 2173 | ABLIKIM | 07D | BES2 $e^+ e^- \rightarrow \psi(2S)$ |
| 127 ±12 ±2 | 410 | ¹ AUBERT | 07AU | BABR $10.6 e^+ e^- \rightarrow 2(\pi^+\pi^-)\pi^0 \gamma$ |
| 26.1± 0.7±3.0 | 1703 | BRIERE | 05 | CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$ |
| 30 ± 8 | 42 | FRANKLIN | 83 | MRK2 $e^+ e^-$ |

¹AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+e^-)] = (297 \pm 22 \pm 18) \times 10^{-4}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+e^-) = 2.33 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.



$\Gamma(\rho a_2(1320))/\Gamma_{\text{total}}$

Γ_{23}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>CL%</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------|--------------------|-------------|---|
| 2.55 ± 0.73 ± 0.47 | | 112 ± 31 | BAI | 04C BES2 | $\psi(2S) \rightarrow 2(\pi^+ \pi^-) \pi^0$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| <2.3 | 90 | BAI | 98J BES | | $e^+ e^-$ |

$\Gamma(3(\pi^+ \pi^-))/\Gamma_{\text{total}}$

Γ_{27}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|-------------------------------------|-------------|---|
| 3.5 ± 2.0 OUR AVERAGE | | Error includes scale factor of 2.8. | | |
| 5.45 ± 0.42 ± 0.87 | 671 | ABLIKIM | 05H BES2 | $e^+ e^- \rightarrow \psi(2S) \rightarrow 3(\pi^+ \pi^-)$ |
| 1.5 ± 1.0 | | ¹ TANENBAUM 78 | MRK1 | $e^+ e^-$ |

¹ Assuming entirely strong decay.

$\Gamma(3(\pi^+ \pi^-)\pi^0)/\Gamma_{\text{total}}$

Γ_{29}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|------------------------------------|
| 35 ± 16 | 6 | FRANKLIN | 83 | MRK2 $e^+ e^- \rightarrow$ hadrons |

$\Gamma(\eta \pi^+ \pi^-)/\Gamma_{\text{total}}$

Γ_{31}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|--|
| <1.6 | 90 | BRIERE | 05 | CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+ \pi^-) \pi^0$ |

$\Gamma(\eta \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$

Γ_{32}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|---------------------|-------------|---|
| 9.5 ± 0.7 ± 1.5 | | ¹ BRIERE | 05 | CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow$ hadr |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 10.3 ± 0.8 ± 1.4 | 201.7 | ² BRIERE | 05 | CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow \eta 3\pi (\eta \rightarrow \gamma\gamma)$ |
| 8.1 ± 1.4 ± 1.6 | 50.0 | ² BRIERE | 05 | CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow \eta 3\pi (\eta \rightarrow 3\pi)$ |

¹ Average of $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow 3\pi$.

² Not independent from other values reported by BRIERE 05.

$\Gamma(\rho\eta)/\Gamma_{\text{total}}$

Γ_{37}/Γ

| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|-------------------------------------|-------------|-------------------------------------|
| 2.2 ± 0.6 OUR AVERAGE | | Error includes scale factor of 1.1. | | |
| 3.0 $^{+1.1}_{-0.9}$ ± 0.2 | 18 | ADAM | 05 | CLEO $e^+ e^- \rightarrow \psi(2S)$ |
| 1.78 $^{+0.67}_{-0.62}$ ± 0.17 | 13 | ABLIKIM | 04L BES | $e^+ e^- \rightarrow \psi(2S)$ |

$\Gamma(\eta' \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$

Γ_{38}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|--|
| 4.5 ± 1.6 ± 1.3 | 12.8 | BRIERE | 05 | CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow$ hadr |

$\Gamma(\eta' \rho)/\Gamma_{\text{total}}$ Γ_{39}/Γ

| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|--------------------------------|
| 1.87 $^{+1.64}_{-1.11}$ ± 0.33 | 2 | ABLIKIM | 04L BES | $e^+ e^- \rightarrow \psi(2S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-----------------------------|-----|----------------------|-----------|--------------------------------|
| 1.02 ± 0.11 ± 0.24 | 143 | ¹ ABLIKIM | 17AK BES3 | $e^+ e^- \rightarrow \psi(2S)$ |
| $0.569 \pm 0.128 \pm 0.236$ | 80 | ² ABLIKIM | 17AK BES3 | $e^+ e^- \rightarrow \psi(2S)$ |

¹ Destructive-interference solution of a partial wave analysis of the decay $\psi(2S) \rightarrow \pi^+ \pi^- \eta'$.

² Constructive-interference solution of a partial wave analysis of the decay $\psi(2S) \rightarrow \pi^+ \pi^- \eta'$.

 $\Gamma(\omega \pi^0)/\Gamma_{\text{total}}$ Γ_{40}/Γ


| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|--------------------------------|
| 2.1 ± 0.6 OUR AVERAGE | | | | |
| 2.5 $^{+1.2}_{-1.0}$ ± 0.2 | 14 | ADAM | 05 CLEO | $e^+ e^- \rightarrow \psi(2S)$ |
| $1.87^{+0.68}_{-0.62} \pm 0.28$ | 14 | ABLIKIM | 04L BES | $e^+ e^- \rightarrow \psi(2S)$ |

 $\Gamma(\omega \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{41}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------|---|-------------|---|
| 7.3 ± 1.2 OUR AVERAGE | | Error includes scale factor of 2.1. See the ideogram below. | | |
| 8.4 $\pm 0.5 \pm 1.2$ | 386 | ABLIKIM | 07D BES2 | $e^+ e^- \rightarrow \psi(2S)$ |
| 12.2 $\pm 2.2 \pm 0.7$ | 37 | ¹ AUBERT | 07AU BABR | $10.6 e^+ e^- \rightarrow \omega \pi^+ \pi^- \gamma$ |
| 8.2 $\pm 0.5 \pm 0.7$ | 391 | BRIERE | 05 CLEO | $e^+ e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+ \pi^-) \pi^0$ |
| 4.8 $\pm 0.6 \pm 0.7$ | 100 ± 22 | ² BAI | 03B BES | $\psi(2S) \rightarrow 2(\pi^+ \pi^-) \pi^0$ |

¹ AUBERT 07AU quotes $\Gamma_{ee}^{\psi(2S)} \cdot B(\psi(2S) \rightarrow \omega \pi^+ \pi^-) \cdot B(\omega \rightarrow 3\pi) = 2.69 \pm 0.73 \pm 0.16$ eV.

² Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.



$\Gamma(b_1^\pm\pi^\mp)/\Gamma_{\text{total}}$

Γ_{43}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-------------|--------------------|----------|---|
| 4.0 ± 0.6 OUR AVERAGE | | | | Error includes scale factor of 1.1. |
| 5.1 ± 0.6 ± 0.8 | 202 | ABLIKIM | 07D BES2 | $e^+e^- \rightarrow \psi(2S)$ |
| 4.18 ^{+0.43} _{-0.42} ± 0.92 | 170 | ADAM | 05 CLEO | $e^+e^- \rightarrow \psi(2S)$ |
| 3.2 ± 0.6 ± 0.5 | 61 ± 11 | ^{1,2} BAI | 03B BES | $\psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 5.2 ± 0.8 ± 1.0 | | ¹ BAI | 99C BES | Repl. by BAI 03B |

¹ Assuming $B(b_1 \rightarrow \omega\pi)=1$.

² Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.

$\Gamma(\omega f_2(1270))/\Gamma_{\text{total}}$

Γ_{44}/Γ

| VALUE (units 10^{-4}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------|------------------|----------|---|
| 2.2 ± 0.4 OUR AVERAGE | | | | | |
| 2.3 ± 0.5 ± 0.4 | | 57 | ABLIKIM | 07D BES2 | $e^+e^- \rightarrow \psi(2S)$ |
| 2.05 ^{+0.41} _{-0.38} ± 0.38 | | 62 ± 12 | BAI | 04C BES2 | $\psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| <1.5 | | 90 | ¹ BAI | 03B BES | $\psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$ |
| <1.7 | | 90 | BAI | 98J BES | Repl. by BAI 03B |

¹ Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.

$\Gamma(b_1^0\pi^0)/\Gamma_{\text{total}}$

Γ_{47}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|-------------|---------|-------------------------------|
| 2.35^{+0.47}_{-0.42} ± 0.40 | 45 | ADAM | 05 CLEO | $e^+e^- \rightarrow \psi(2S)$ |

| $\Gamma(\omega\eta)/\Gamma_{\text{total}}$ | | | | | Γ_{48}/Γ |
|---|------------|--------------------|-------------|-------------------------------------|----------------------|
| <u>VALUE (units 10^{-5})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
| <1.1 | 90 | ADAM | 05 | CLEO $e^+ e^- \rightarrow \psi(2S)$ | |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | | |
| <3.1 | 90 | ABLIKIM | 04K | BES $e^+ e^- \rightarrow \psi(2S)$ | |

| $\Gamma(\omega\eta')/\Gamma_{\text{total}}$ | | | | | Γ_{49}/Γ |
|---|-------------|----------------------|-------------|------------------------------------|----------------------|
| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
| $3.2^{+2.4}_{-2.0} \pm 0.7$ | 4 | ¹ ABLIKIM | 04K | BES $e^+ e^- \rightarrow \psi(2S)$ | |

¹ Calculated combining $\eta' \rightarrow \gamma\rho$ and $\eta\pi^+\pi^-$ channels.


| $\Gamma(\phi\pi^0)/\Gamma_{\text{total}}$ | | | | | Γ_{50}/Γ |
|---|------------|--------------------|-------------|-------------------------------------|----------------------|
| <u>VALUE (units 10^{-5})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
| <0.04 | 90 | ABLIKIM | 12L | BES3 $e^+ e^- \rightarrow \psi(2S)$ | |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | | |
| <0.7 | 90 | ADAM | 05 | CLEO $e^+ e^- \rightarrow \psi(2S)$ | |
| <0.4 | 90 | ABLIKIM | 04K | BES $e^+ e^- \rightarrow \psi(2S)$ | |

| $\Gamma(\phi\pi^+\pi^-)/\Gamma_{\text{total}}$ | | | | | Γ_{51}/Γ |
|---|----------------|---|-------------|---|----------------------|
| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
| 1.18 ± 0.26 OUR AVERAGE | | Error includes scale factor of 1.5. See the ideogram below. | | | |
| 2.3 $\pm 0.8 \pm 0.1$ | 19 ± 6 | LEES | 12F | BABR $10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$ | |
| 0.9 $\pm 0.2 \pm 0.1$ | 47.6 | BRIERE | 05 | CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$ | |
| 1.5 $\pm 0.2 \pm 0.2$ | 51.5 ± 8.3 | ¹ BAI | 03B | BES $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$ | |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | | |
| 2.45 $\pm 0.96 \pm 0.04$ | 10 ± 4 | ^{2,3} AUBERT | 07AK | BABR $10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$ | |

¹ Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.

² Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(\psi(2S) \rightarrow \phi\pi^+\pi^-)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (0.57 \pm 0.22 \pm 0.04) \times 10^{-3}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.33 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Using $B(\phi \rightarrow K^+ K^-) = (49.3 \pm 0.6)\%$.



$\Gamma(\phi f_0(980) \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}$

Γ_{52}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|----------------|-------------------------------------|-----------|---|
| 0.75±0.33 OUR AVERAGE | | Error includes scale factor of 1.6. | | |
| 1.5 ± 0.5 ± 0.1 | 12 ± 4 | LEES | 12F BABR | $10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$ |
| 0.6 ± 0.2 ± 0.1 | 18.4 ± 6.4 | ¹ BAI | 03B BES | $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 1.46±0.71±0.02 | 6 ± 3 | ^{2,3} AUBERT | 07AK BABR | $10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$ |

¹ Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.

² Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(\psi(2S) \rightarrow \phi f_0(980) \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+e^-)] = (0.34 \pm 0.16 \pm 0.04) \times 10^{-3}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+e^-) = 2.33 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Using $B(\phi \rightarrow K^+K^-) = (49.3 \pm 0.6)\%$.

$\Gamma(\phi\eta)/\Gamma_{\text{total}}$

Γ_{53}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------|------|-------------|----------|--------------------------------|
| 3.10±0.31 OUR AVERAGE | | | | |
| 3.14±0.23±0.23 | 0.2k | ABLIKIM | 12L BES3 | $e^+ e^- \rightarrow \psi(2S)$ |
| $2.0^{+1.5}_{-1.1} \pm 0.4$ | 6 | ADAM | 05 CLEO | $e^+ e^- \rightarrow \psi(2S)$ |
| 3.3 ± 1.1 ± 0.5 | 17 | ABLIKIM | 04K BES | $e^+ e^- \rightarrow \psi(2S)$ |

$\Gamma(\eta\phi(2170), \phi(2170) \rightarrow \phi f_0(980), f_0 \rightarrow \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{54}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-----|-------------|----------|---|
| $<2.2 \times 10^{-6}$ | 90 | ABLIKIM | 19I BES3 | $e^+ e^- \rightarrow \eta\phi f_0(980)$ |

$\Gamma(\phi\eta')/\Gamma_{\text{total}}$ Γ_{55}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|----------------------|-----------|--------------------------------|
| 1.54 ± 0.20 OUR AVERAGE | | | | |
| $1.51 \pm 0.16 \pm 0.12$ | 201 | ABLIKIM | 19BA BES3 | $e^+ e^- \rightarrow \psi(2S)$ |
| $3.1 \pm 1.4 \pm 0.7$ | 8 | ¹ ABLIKIM | 04K BES | $e^+ e^- \rightarrow \psi(2S)$ |

¹ Calculated combining $\eta' \rightarrow \gamma\rho$ and $\eta\pi^+\pi^-$ channels.

$\Gamma(\phi f_1(1285))/\Gamma_{\text{total}}$ Γ_{56}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|----------------------|-----------|--------------------------------|
| $3.0 \pm 0.4 \pm 1.3$ | 234 | ¹ ABLIKIM | 19BA BES3 | $e^+ e^- \rightarrow \psi(2S)$ |

¹ ABLIKIM 19BA reports $[\Gamma(\psi(2S) \rightarrow \phi f_1(1285))/\Gamma_{\text{total}}] \times [B(f_1(1285) \rightarrow \eta\pi^+\pi^-)] = (1.03 \pm 0.10 \pm 0.09) \times 10^{-5}$ which we divide by our best value $B(f_1(1285) \rightarrow \eta\pi^+\pi^-) = (35 \pm 15) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi\eta(1405) \rightarrow \phi\pi^+\pi^-\eta)/\Gamma_{\text{total}}$ Γ_{57}/Γ

| VALUE (units 10^{-6}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|-------------|-----------|--------------------------------|
| $8.46 \pm 1.37 \pm 0.92$ | 195 | ABLIKIM | 19BA BES3 | $e^+ e^- \rightarrow \psi(2S)$ |

$\Gamma(\phi f'_2(1525))/\Gamma_{\text{total}}$ Γ_{58}/Γ

| VALUE (units 10^{-4}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|-----|------------|-------------|------|-----------------------------------|
| $0.44 \pm 0.12 \pm 0.11$ | | 20 ± 6 | BAI | 04C | $\psi(2S) \rightarrow 2(K^+ K^-)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-------|----|-----|---------|----------------------------------|
| <0.45 | 90 | BAI | 98J BES | $e^+ e^- \rightarrow 2(K^+ K^-)$ |
|-------|----|-----|---------|----------------------------------|

$\Gamma(K^+ K^-)/\Gamma_{\text{total}}$ Γ_{59}/Γ

| VALUE (units 10^{-5}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|-----|------|------------------------|------|--------------------------------|
| $7.48 \pm 0.23 \pm 0.39$ | | 1.3k | ¹ METREVELI | 12 | $\psi(2S) \rightarrow K^+ K^-$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-----------------------|----|-------------|----------|--------------------------------------|
| 6.2 $\pm 1.5 \pm 0.2$ | 66 | 2,3 LEES | 15J BABR | $e^+ e^- \rightarrow K^+ K^- \gamma$ |
| 8.3 $\pm 1.5 \pm 0.2$ | 66 | 3,4 LEES | 15J BABR | $e^+ e^- \rightarrow K^+ K^- \gamma$ |
| 6.3 $\pm 0.6 \pm 0.3$ | | 5 DOBBS | 06A CLEO | $e^+ e^-$ |
| 10 ± 7 | | 5 BRANDELIK | 79C DASP | $e^+ e^-$ |
| < 5 | 90 | FELDMAN | 77 MRK1 | $e^+ e^-$ |

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

² $\sin\phi > 0$.

³ Using $\Gamma(\psi(2S) \rightarrow e^+ e^-) = (2.37 \pm 0.04)$ keV.

⁴ $\sin\phi < 0$.

⁵ Interference with non-resonant $K^+ K^-$ production not taken into account.

$\Gamma(K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{60}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-------|---------------------|-----------|--|
| 7.3 ± 0.5 OUR AVERAGE | | | | |
| 8.1 $\pm 1.3 \pm 0.3$ | 133 | LEES | 12F BABR | $10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$ |
| 7.1 $\pm 0.3 \pm 0.4$ | 817.2 | BRIERE | 05 CLEO | $e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$ |
| 16 ± 4 | | 1 TANENBAUM | 78 MRK1 | $e^+ e^-$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 11.0 $\pm 1.9 \pm 0.2$ | 85 | ² AUBERT | 07AK BABR | $10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$ |

¹ Assuming entirely strong decay.

² Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (2.56 \pm 0.42 \pm 0.16) \times 10^{-3}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.33 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{61}/Γ

| VALUE (units 10^{-5}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|------|-------------|----------|--------------------------------------|
| $4.07 \pm 0.16 \pm 0.26$ | 0.9k | | ABLIKIM | 12L BES3 | $e^+ e^- \rightarrow \psi(2S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| <8.9 | 90 | 1 | FRANKLIN | 83 MRK2 | $e^+ e^- \rightarrow \text{hadrons}$ |

$\Gamma(K_S^0 K_L^0)/\Gamma_{\text{total}}$ Γ_{62}/Γ

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|---------|-----------|
| <0.046 | 1 BAI | 04D BES | $e^+ e^-$ |

¹ Forbidden by CP.

$\Gamma(K_S^0 K_L^0)/\Gamma_{\text{total}}$ Γ_{63}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|--------------|------------------|----------|--|
| 5.34 ± 0.33 OUR AVERAGE | | | | |
| 5.28 $\pm 0.25 \pm 0.34$ | 478 ± 23 | 1 METREVELI | 12 | $\psi(2S) \rightarrow K_S^0 K_L^0$ |
| 5.8 $\pm 0.8 \pm 0.4$ | | DOBBS | 06A CLEO | $e^+ e^-$ |
| 5.24 $\pm 0.47 \pm 0.48$ | 156 ± 14 | ² BAI | 04B BES2 | $\psi(2S) \rightarrow K_S^0 K_L^0 \rightarrow \pi^+ \pi^- X$ |

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

² Using $B(K_S^0 \rightarrow \pi^+ \pi^-) = 0.6860 \pm 0.0027$.

$\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_{69}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|-------|---------------------|-----------|--|
| 12.6 ± 0.9 OUR AVERAGE | | | | |
| 18.9 $\pm 5.7 \pm 0.3$ | 32 | ¹ AUBERT | 07AU BABR | $10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \pi^0 \gamma$ |
| 11.7 $\pm 1.0 \pm 1.5$ | 597 | ABLIKIM | 06G BES2 | $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ |
| 12.7 $\pm 0.5 \pm 1.0$ | 711.6 | BRIERE | 05 CLEO | $e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ |

¹ AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (44 \pm 13 \pm 3) \times 10^{-4}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.33 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\omega f_0(1710) \rightarrow \omega K^+ K^-)/\Gamma_{\text{total}}$ Γ_{70}/Γ

| <u>VALUE</u> (units 10^{-5}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|-------------|--------------------|-------------|--|
| 5.9±2.0±0.9 | 19 | ABLIKIM | 06G BES2 | $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ |

 $\Gamma(K^*(892)^0 K^- \pi^+ \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{71}/Γ

| <u>VALUE</u> (units 10^{-4}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|-------------|--------------------|-------------|--|
| 8.6±1.3±1.8 | 238 | ABLIKIM | 06G BES2 | $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ |

 $\Gamma(K^*(892)^+ K^- \pi^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{72}/Γ

| <u>VALUE</u> (units 10^{-4}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|-------------|--------------------|-------------|--|
| 9.6±2.2±1.7 | 133 | ABLIKIM | 06G BES2 | $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ |

 $\Gamma(K^*(892)^+ K^- \rho^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{73}/Γ

| <u>VALUE</u> (units 10^{-4}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|-------------|--------------------|-------------|--|
| 7.3±2.2±1.4 | 78 | ABLIKIM | 06G BES2 | $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ |

 $\Gamma(K^*(892)^0 K^- \rho^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{74}/Γ

| <u>VALUE</u> (units 10^{-4}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|-------------|--------------------|-------------|--|
| 6.1±1.3±1.2 | 125 | ABLIKIM | 06G BES2 | $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ |

 $\Gamma(K_S^0 K_S^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{75}/Γ

| <u>VALUE</u> (units 10^{-4}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|-------------|--------------------|-------------|--------------------------------|
| 2.20±0.25±0.37 | 83 ± 9 | ABLIKIM | 050 BES2 | $e^+ e^- \rightarrow \psi(2S)$ |

 $\Gamma(K^+ K^- \rho^0)/\Gamma_{\text{total}}$ Γ_{83}/Γ

| <u>VALUE</u> (units 10^{-4}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|-------------|--------------------|-------------|--|
| 2.2±0.2±0.4 | 223.8 | BRIERE | 05 CLEO | $e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$ |

 $\Gamma(K^*(892)^0 \bar{K}_2^*(1430)^0)/\Gamma_{\text{total}}$ Γ_{84}/Γ

| <u>VALUE</u> (units 10^{-4}) | <u>CL%</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|------------|-------------|--------------------|-------------|--|
| 1.86±0.32±0.43 | | 93 ± 16 | BAI | 04C | $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.2 90 BAI 98J BES $e^+ e^-$

 $\Gamma(K^+ K^- \pi^+ \pi^- \eta)/\Gamma_{\text{total}}$ Γ_{85}/Γ

| <u>VALUE</u> (units 10^{-3}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|-------------|---------------------|-------------|--|
| 1.3±0.7±0.1 | 7 | ¹ AUBERT | 07AU BABR | $10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \eta \gamma$ |

¹AUBERT 07AU quotes $\Gamma_{ee}^{\psi(2S)} \cdot B(\psi(2S) \rightarrow 2(\pi^+ \pi^-) \eta) \cdot B(\eta \rightarrow \gamma \gamma) = 1.2 \pm 0.7 \pm 0.1 \text{ eV}$.

$\Gamma(K^+ K^- 2(\pi^+ \pi^-) \pi^0)/\Gamma_{\text{total}}$ Γ_{87}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|--------------------------------|
| $10.0 \pm 2.5 \pm 1.8$ | 65 | ABLIKIM | 07D BES2 | $e^+ e^- \rightarrow \psi(2S)$ |

 $\Gamma(K^+ K^*(892)^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{88}/Γ

| <u>VALUE (units 10^{-5})</u> | <u>CL %</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|-------------|-------------------------------------|-------------|----------------|
| 2.9 ± 0.4 OUR AVERAGE | | | Error includes scale factor of 1.2. | | |

3.18 $\pm 0.30^{+0.26}_{-0.31}$ 0.2k ABLIKIM 12L BES3 $e^+ e^- \rightarrow \psi(2S)$ 2.9 ± 0.4 9.6 ± 4.2 ABLIKIM 05I BES2 $e^+ e^- \rightarrow \psi(2S)$ 1.3 ± 0.3 7 ADAM 05 CLEO $e^+ e^- \rightarrow \psi(2S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<5.4 90 FRANKLIN 83 MRK2 $e^+ e^- \rightarrow \text{hadrons}$ $\Gamma(2(K^+ K^-))/\Gamma_{\text{total}}$ Γ_{89}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|---|
| 0.63 ± 0.13 OUR AVERAGE | | | | |
| 0.9 ± 0.4 ± 0.1 | 13 | LEES | 12F BABR | $10.6 e^+ e^- \rightarrow 2(K^+ K^-)\gamma$ |
| 0.6 ± 0.1 ± 0.1 | 59.2 | BRIERE | 05 CLEO | $e^+ e^- \rightarrow \psi(2S) \rightarrow 2(K^+ K^-)$ |

 $\Gamma(2(K^+ K^-)\pi^0)/\Gamma_{\text{total}}$ Γ_{90}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|--|
| $1.1 \pm 0.2 \pm 0.2$ | 44.7 | BRIERE | 05 CLEO | $e^+ e^- \rightarrow \psi(2S) \rightarrow 2(K^+ K^-)\pi^0$ |

 $\Gamma(K^+ K^- \phi)/\Gamma_{\text{total}}$ Γ_{91}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|----------------|--------------------|-------------|---|
| 0.70 ± 0.16 OUR AVERAGE | | | | |
| 0.8 ± 0.2 ± 0.1 | 36.8 | BRIERE | 05 CLEO | $e^+ e^- \rightarrow \psi(2S) \rightarrow 2(K^+ K^-)$ |
| 0.6 ± 0.2 ± 0.1 | 16.1 ± 5.0 | ¹ BAI | 03B BES | $\psi(2S) \rightarrow 2(K^+ K^-)$ |

¹ Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$. $\Gamma(K_S^0 K_S^0 \phi)/\Gamma_{\text{total}}$ Γ_{92}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|----------------------|-------------|--|
| $0.353 \pm 0.020 \pm 0.021$ | 687 | ¹ ABLIKIM | 23BA BES3 | $e^+ e^- \rightarrow \psi(2S) \rightarrow K_S^0 K_S^0 K^+ K^-$ |

¹ Solution with a constructive interference of the signal with the continuum background. $\Gamma(K_1(1270)^{\pm} K^{\mp})/\Gamma_{\text{total}}$ Γ_{93}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|----------------|
| $10.0 \pm 1.8 \pm 2.1$ | | ¹ BAI | 99C BES | $e^+ e^-$ |

¹ Assuming $B(K_1(1270) \rightarrow K\rho) = 0.42 \pm 0.06$

$\Gamma(K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{94}/Γ

| <u>VALUE</u> (units 10^{-4}) | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|--------------------|-------------|----------------|
| 6.7 ± 2.5 | TANENBAUM 78 | MRK1 | $e^+ e^-$ |

 $\Gamma(\eta K^+ K^-, \text{no } \eta\phi)/\Gamma_{\text{total}}$ Γ_{95}/Γ

| <u>VALUE</u> (units 10^{-5}) | <u>CL%</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|------------|-------------|--------------------|-------------|---|
| $3.49 \pm 0.09 \pm 0.15$ | | 1.8k | 1 ABLIKIM | 20F BES3 | $\psi(2S) \rightarrow K^+ K^- \gamma\gamma$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|--------------------------|------|-------------|----------|--|
| 3.08 $\pm 0.29 \pm 0.25$ | 0.3k | 1,2 ABLIKIM | 12L BES3 | $\psi(2S) \rightarrow K^+ K^- \gamma\gamma$ |
| <13 | 90 | BRIERE | 05 CLEO | $e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ |

¹ Excluding $\eta\phi$.

² Superseded by ABLIKIM 20F.

 $\Gamma(\eta K^+ K^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{96}\Gamma_7/\Gamma$

| <u>VALUE</u> (eV) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-------------------|------------|--------------------|-------------|--|
| <0.6 | 90 | 1 LEES | 23 BABR | $e^+ e^- \rightarrow \gamma_{ISR}$ hadrons |

¹ LEES 23 reports $[\Gamma(\psi(2S) \rightarrow \eta K^+ K^-) \times \Gamma(\psi(2S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 3\pi^0)] < 0.2$ eV which we divide by our best value $B(\eta \rightarrow 3\pi^0) = 32.57 \times 10^{-2}$.

 $\Gamma(X(1750)\eta \rightarrow K^+ K^- \eta)/\Gamma_{\text{total}}$ Γ_{97}/Γ

| <u>VALUE</u> (units 10^{-6}) | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|-------------------------------------|
| $4.8 \pm 1.0 \pm 2.6$ | ABLIKIM | 20F BES3 | $\psi(2S) \rightarrow K^+ K^- \eta$ |

 $\Gamma(K_1(1400)^{\pm} K^{\mp})/\Gamma_{\text{total}}$ Γ_{98}/Γ

| <u>VALUE</u> (units 10^{-4}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|------------|--------------------|-------------|----------------|
| <3.1 | 90 | 1 BAI | 99C BES | $e^+ e^-$ |

¹ Assuming $B(K_1(1400) \rightarrow K^* \pi) = 0.94 \pm 0.06$

 $\Gamma(K_2^*(1430)^{\pm} K^{\mp})/\Gamma_{\text{total}}$ Γ_{99}/Γ

| <u>VALUE</u> (units 10^{-5}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------|--------------------|-------------|--------------------------------|
| $7.12 \pm 0.62^{+1.13}_{-0.61}$ | 251 ± 22 | ABLIKIM | 12L BES3 | $e^+ e^- \rightarrow \psi(2S)$ |

 $\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{100}/Γ

| <u>VALUE</u> (units 10^{-5}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|----------------|--------------------|-------------|--------------------------------|
| 10.9 ± 2.0 OUR AVERAGE | | | | |
| $13.3^{+2.4}_{-2.8} \pm 1.7$ | 65.6 ± 9.0 | ABLIKIM | 05I BES2 | $e^+ e^- \rightarrow \psi(2S)$ |
| $9.2^{+2.7}_{-2.2} \pm 0.9$ | 25 | ADAM | 05 CLEO | $e^+ e^- \rightarrow \psi(2S)$ |

 $\Gamma(K^+ K^*(892)^- + \text{c.c.})/\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})$ Γ_{88}/Γ_{100}

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|--------------------------------|
| 0.16 ± 0.06 OUR AVERAGE | | | |
| $0.22^{+0.10}_{-0.14}$ | ABLIKIM | 05I BES2 | $e^+ e^- \rightarrow \psi(2S)$ |
| $0.14^{+0.08}_{-0.06}$ | ADAM | 05 CLEO | $e^+ e^- \rightarrow \psi(2S)$ |

$\Gamma(\omega K^+ K^-)/\Gamma_{\text{total}}$ Γ_{101}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|--|
| 1.62 ± 0.11 OUR AVERAGE | | | | Error includes scale factor of 1.1. |
| $1.56 \pm 0.04 \pm 0.11$ | 2.8k | ABLIKIM | 14G BES3 | $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ |
| $2.38 \pm 0.37 \pm 0.29$ | 78 | ABLIKIM | 06G BES2 | $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ |
| $1.9 \pm 0.3 \pm 0.3$ | 76.8 | BRIERE | 05 CLEO | $e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ |
| $1.5 \pm 0.3 \pm 0.2$ | 23 | ¹ BAI | 03B BES | $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ |

¹ Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

 $\Gamma(\omega K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{102}/Γ

| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|-------------|--------------------|-------------|--|
| $7.04 \pm 0.39 \pm 0.36$ | 1.5k | ABLIKIM | 21AL BES3 | $\psi(2S) \rightarrow \pi^+ \pi^- \pi^0 K_S^0 K_S^0$ |

 $\Gamma(\omega K^*(892)^+ K^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{103}/Γ

| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|-------------|--------------------|-------------|---|
| 20.7 ± 2.6 OUR AVERAGE | | | | |
| $18.9 \pm 2.9 \pm 2.2$ | 396 | ABLIKIM | 13M BES3 | $\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$ |
| $22.6 \pm 3.0 \pm 2.4$ | 535 | ABLIKIM | 13M BES3 | $\psi(2S) \rightarrow \omega K^+ K^- \pi^0$ |

 $\Gamma(\omega K_2^*(1430)^+ K^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{104}/Γ

| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|---|
| 6.1 ± 1.2 OUR AVERAGE | | | | |
| $6.39 \pm 1.50 \pm 0.78$ | 128 | ABLIKIM | 13M BES3 | $\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$ |
| $5.86 \pm 1.61 \pm 0.83$ | 143 | ABLIKIM | 13M BES3 | $\psi(2S) \rightarrow \omega K^+ K^- \pi^0$ |

 $\Gamma(\omega \bar{K}^*(892)^0 K^0)/\Gamma_{\text{total}}$ Γ_{105}/Γ

| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|---|
| $16.8 \pm 2.5 \pm 1.6$ | 356 | ABLIKIM | 13M BES3 | $\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$ |

 $\Gamma(\omega \bar{K}_2^*(1430)^0 K^0)/\Gamma_{\text{total}}$ Γ_{106}/Γ

| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|-------------|--------------------|-------------|---|
| $5.82 \pm 2.08 \pm 0.72$ | 116 | ABLIKIM | 13M BES3 | $\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$ |

 $\Gamma(\omega X(1440) \rightarrow \omega K_S^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{107}/Γ

| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|-------------|----------------------|-------------|---|
| $1.60 \pm 0.27 \pm 0.24$ | 109 | ¹ ABLIKIM | 13M BES3 | $\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$ |

¹ $X(1440)$ compatible with $\eta(1405)$ and $\eta(1475)$. A $f_1(1420)$ is also possible.

 $\Gamma(\omega X(1440) \rightarrow \omega K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{108}/Γ

| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|-------------|----------------------|-------------|---|
| $1.09 \pm 0.20 \pm 0.16$ | 82 | ¹ ABLIKIM | 13M BES3 | $\psi(2S) \rightarrow \omega K^+ K^- \pi^0$ |

¹ $X(1440)$ compatible with $\eta(1405)$ and $\eta(1475)$. A $f_1(1420)$ is also possible.

$\Gamma(\omega f_1(1285) \rightarrow \omega K_S^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{109}/Γ

| <u>VALUE</u> (units 10^{-5}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|----------------------|-------------|---|
| $0.302 \pm 0.098 \pm 0.027$ | 22 | ¹ ABLIKIM | 13M BES3 | $\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$ |

¹ Statistical significance 4.5σ . This measurement is equivalent to a limit of $< 0.478 \times 10^{-5}$ at 90% C.L.

 $\Gamma(\omega f_1(1285) \rightarrow \omega K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{110}/Γ

| <u>VALUE</u> (units 10^{-5}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|----------------------|-------------|---|
| $0.125 \pm 0.070 \pm 0.013$ | 10 | ¹ ABLIKIM | 13M BES3 | $\psi(2S) \rightarrow \omega K^+ K^- \pi^0$ |

¹ Statistical significance 3.2σ . This measurement is equivalent to a limit of $< 0.221 \times 10^{-5}$ at 90% C.L.

 $\Gamma(p\bar{p})/\Gamma_{\text{total}}$ Γ_{111}/Γ

| <u>VALUE</u> (units 10^{-4}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|-------------------------------------|
| 2.94 ± 0.09 OUR FIT | | | | Error includes scale factor of 1.3. |

 3.02 ± 0.08 OUR AVERAGE

| | | | | |
|--------------------------|------|--------------------|----------|---|
| 3.05 $\pm 0.02 \pm 0.12$ | 19k | ABLIKIM | 18T BES3 | $e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$ |
| 3.08 $\pm 0.05 \pm 0.18$ | 4.5k | ¹ DOBBS | 14 | $e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$ |
| 3.36 $\pm 0.09 \pm 0.25$ | 1.6k | ABLIKIM | 07C BES | $e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$ |
| 2.87 $\pm 0.12 \pm 0.15$ | 557 | PEDLAR | 05 CLEO | $e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$ |
| 1.4 ± 0.8 | 4 | BRANDELIK | 79C DASP | $e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$ |
| 2.3 ± 0.7 | | FELDMAN | 77 MRK1 | $e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$ |

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

 $\Gamma(p\bar{p})/\Gamma(J/\psi(1S)\pi^+\pi^-)$ Γ_{111}/Γ_{12}

| <u>VALUE</u> (units 10^{-4}) | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|--------------------|-------------|---|
| 8.49 ± 0.28 OUR FIT | | | Error includes scale factor of 1.3. |
| $6.98 \pm 0.49 \pm 0.97$ | BAI | 01 BES | $e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$ |

 $\Gamma(n\bar{n})/\Gamma_{\text{total}}$ Γ_{112}/Γ

| <u>VALUE</u> (units 10^{-4}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|-------------|--------------------|-------------|---|
| $3.06 \pm 0.06 \pm 0.14$ | 6k | ABLIKIM | 18T BES3 | $e^+ e^- \rightarrow \psi(2S) \rightarrow n\bar{n}$ |

 $\Gamma(p\bar{p}\pi^0)/\Gamma_{\text{total}}$ Γ_{113}/Γ

| <u>VALUE</u> (units 10^{-4}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|----------------------|-------------|---|
| 1.53 ± 0.07 OUR AVERAGE | | | | |
| 1.65 $\pm 0.03 \pm 0.15$ | 4.5k | ABLIKIM | 13A BES3 | $\psi(2S) \rightarrow p\bar{p}\pi^0$ |
| 1.54 $\pm 0.06 \pm 0.06$ | 948 | ALEXANDER | 10 CLEO | $\psi(2S) \rightarrow \pi^0 p\bar{p}$ |
| 1.32 $\pm 0.10 \pm 0.15$ | 256 | ¹ ABLIKIM | 05E BES2 | $e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\gamma\gamma$ |
| 1.4 ± 0.5 | 9 | FRANKLIN | 83 MRK2 | $e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\pi^0$ |

¹ Computed using $B(\pi^0 \rightarrow \gamma\gamma) = (98.80 \pm 0.03)\%$.

 $\Gamma(N(940)\bar{p} + \text{c.c.} \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}$ Γ_{114}/Γ

| <u>VALUE</u> (units 10^{-5}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|-------------|----------------------|-------------|--------------------------------------|
| $6.42 \pm 0.20 \pm 1.78$ | 1.9k | ¹ ABLIKIM | 13A BES3 | $\psi(2S) \rightarrow p\bar{p}\pi^0$ |

¹ From a fit of $\pi^0 p\bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.

$\Gamma(N(1440)\bar{p} + \text{c.c.} \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}$ Γ_{115}/Γ

| <u>VALUE</u> (units 10^{-5}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|-------------|--------------------|-------------|----------------|
|---------------------------------|-------------|--------------------|-------------|----------------|

7.3 ± 1.7 OUR AVERAGE Error includes scale factor of 2.5.

| | | | | |
|-------------------------------|------|------------------------|----------|---------------------------------------|
| 3.58 ± 0.25 ± 0.84 | 1.1k | ¹ ABLIKIM | 13A BES3 | $\psi(2S) \rightarrow p\bar{p}\pi^0$ |
| 8.1 ± 0.7 ± 0.3 | 474 | ² ALEXANDER | 10 CLEO | $\psi(2S) \rightarrow \pi^0 p\bar{p}$ |

¹ From a fit of $\pi^0 p\bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.

² From a fit of the $p\bar{p}$ and $p\pi^0$ mass distributions to a combination of $N(1440)\bar{p}$, a broad $p\bar{p}$ enhancement around 2100 MeV, and two other broad, unestablished resonances.

 $\Gamma(N(1520)\bar{p} + \text{c.c.} \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}$ Γ_{116}/Γ

| <u>VALUE</u> (units 10^{-5}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|-------------|--------------------|-------------|----------------|
|---------------------------------|-------------|--------------------|-------------|----------------|

| | | | | |
|-------------------------------|------|----------------------|----------|--------------------------------------|
| 0.64 ± 0.05 ± 0.17 | 0.2k | ¹ ABLIKIM | 13A BES3 | $\psi(2S) \rightarrow p\bar{p}\pi^0$ |
|-------------------------------|------|----------------------|----------|--------------------------------------|

¹ From a fit of $\pi^0 p\bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.

 $\Gamma(N(1535)\bar{p} + \text{c.c.} \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}$ Γ_{117}/Γ

| <u>VALUE</u> (units 10^{-5}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|-------------|--------------------|-------------|----------------|
|---------------------------------|-------------|--------------------|-------------|----------------|

| | | | | |
|-------------------------------|------|----------------------|----------|--------------------------------------|
| 2.47 ± 0.28 ± 0.97 | 0.7k | ¹ ABLIKIM | 13A BES3 | $\psi(2S) \rightarrow p\bar{p}\pi^0$ |
|-------------------------------|------|----------------------|----------|--------------------------------------|

¹ From a fit of $\pi^0 p\bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.

 $\Gamma(N(1650)\bar{p} + \text{c.c.} \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}$ Γ_{118}/Γ

| <u>VALUE</u> (units 10^{-5}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|-------------|--------------------|-------------|----------------|
|---------------------------------|-------------|--------------------|-------------|----------------|

| | | | | |
|-------------------------------|------|----------------------|----------|--------------------------------------|
| 3.76 ± 0.28 ± 1.66 | 1.1k | ¹ ABLIKIM | 13A BES3 | $\psi(2S) \rightarrow p\bar{p}\pi^0$ |
|-------------------------------|------|----------------------|----------|--------------------------------------|

¹ From a fit of $\pi^0 p\bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.

 $\Gamma(N(1720)\bar{p} + \text{c.c.} \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}$ Γ_{119}/Γ

| <u>VALUE</u> (units 10^{-5}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|-------------|--------------------|-------------|----------------|
|---------------------------------|-------------|--------------------|-------------|----------------|

| | | | | |
|-------------------------------|------|----------------------|----------|--------------------------------------|
| 1.79 ± 0.10 ± 0.71 | 0.5k | ¹ ABLIKIM | 13A BES3 | $\psi(2S) \rightarrow p\bar{p}\pi^0$ |
|-------------------------------|------|----------------------|----------|--------------------------------------|

¹ From a fit of $\pi^0 p\bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.

 $\Gamma(N(2300)\bar{p} + \text{c.c.} \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}$ Γ_{120}/Γ

| <u>VALUE</u> (units 10^{-5}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|-------------|--------------------|-------------|----------------|
|---------------------------------|-------------|--------------------|-------------|----------------|

| | | | | |
|-------------------------------|------|----------------------|----------|--------------------------------------|
| 2.62 ± 0.28 ± 0.64 | 0.9k | ¹ ABLIKIM | 13A BES3 | $\psi(2S) \rightarrow p\bar{p}\pi^0$ |
|-------------------------------|------|----------------------|----------|--------------------------------------|

¹ From a fit of $\pi^0 p\bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.

 $\Gamma(N(2570)\bar{p} + \text{c.c.} \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}$ Γ_{121}/Γ

| <u>VALUE</u> (units 10^{-5}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|-------------|--------------------|-------------|----------------|
|---------------------------------|-------------|--------------------|-------------|----------------|

| | | | | |
|-------------------------------|------|----------------------|----------|--------------------------------------|
| 2.13 ± 0.08 ± 0.30 | 0.8k | ¹ ABLIKIM | 13A BES3 | $\psi(2S) \rightarrow p\bar{p}\pi^0$ |
|-------------------------------|------|----------------------|----------|--------------------------------------|

¹ From a fit of $\pi^0 p\bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.

$\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{122}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|------------------------|-------------|--|
| 6.0 ± 0.4 OUR AVERAGE | | | | |
| $5.9 \pm 0.2 \pm 0.4$ | 904.5 | BRIERE | 05 | CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\pi^+\pi^-$ |
| 8 ± 2 | | ¹ TANENBAUM | 78 | MRK1 $e^+ e^-$ |

¹ Assuming entirely strong decay. $\Gamma(p\bar{p}K^+K^-)/\Gamma_{\text{total}}$ Γ_{123}/Γ

| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|--|
| $2.7 \pm 0.6 \pm 0.4$ | 30.1 | BRIERE | 05 | CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}K^+K^-$ |

 $\Gamma(p\bar{p}\eta)/\Gamma_{\text{total}}$ Γ_{124}/Γ

| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|----------------|------------------------|-------------|---|
| 6.0 ± 0.4 OUR AVERAGE | | | | |
| $6.4 \pm 0.2 \pm 0.6$ | 679 | ¹ ABLIKIM | 13S | BES3 $\psi(2S) \rightarrow \eta p\bar{p}$ |
| $5.6 \pm 0.6 \pm 0.3$ | 154 | ¹ ALEXANDER | 10 | CLEO $\psi(2S) \rightarrow \eta p\bar{p}$ |
| $5.8 \pm 1.1 \pm 0.7$ | 44.8 ± 8.5 | ² ABLIKIM | 05E | BES2 $e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\gamma\gamma$ |
| $8 \pm 3 \pm 3$ | 9.8 | BRIERE | 05 | CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\pi^0$ |

¹ With $N(1535)$ decaying to $p\eta$.² Computed using $B(\eta \rightarrow \gamma\gamma) = (39.43 \pm 0.26)\%$. $\Gamma(N(1535)\bar{p} + \text{c.c.} \rightarrow p\bar{p}\eta)/\Gamma_{\text{total}}$ Γ_{125}/Γ

| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|------------------------|-------------|---|
| 4.5 ± 0.7 OUR AVERAGE | | | | |
| $5.2 \pm 0.3 \pm 3.2$ | 527 | ¹ ABLIKIM | 13S | BES3 $\psi(2S) \rightarrow \eta p\bar{p}$ |
| $4.4 \pm 0.6 \pm 0.3$ | 123 | ² ALEXANDER | 10 | CLEO $\psi(2S) \rightarrow \eta p\bar{p}$ |

¹ With $N(1535)$ decaying to $p\eta$.² From a fit of the $p\bar{p}$ and $p\eta$ distributions to a combination of $N^*(1535)\bar{p}$ and a broad $p\bar{p}$ enhancement around 2100 MeV. $\Gamma(p\bar{p}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{126}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|---|
| $7.3 \pm 0.4 \pm 0.6$ | 434.9 | BRIERE | 05 | CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\pi^0$ |

 $\Gamma(p\bar{p}\rho^0)/\Gamma_{\text{total}}$ Γ_{127}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|--|
| $0.5 \pm 0.1 \pm 0.2$ | 61.1 | BRIERE | 05 | CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\pi^+\pi^-$ |

$\Gamma(p\bar{p}\omega)/\Gamma_{\text{total}}$ Γ_{128}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|------------------|--|
| 0.69±0.21 OUR AVERAGE | | | | |
| 0.6 ± 0.2 | ± 0.2 | 21.2 | BRIERE | 05 CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\pi^0$ |
| 0.8 ± 0.3 | ± 0.1 | 14.9 ± 0.1 | ¹ BAI | 03B BES $\psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\pi^0$ |

¹ Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.

 $\Gamma(p\bar{p}\eta')/\Gamma_{\text{total}}$ Γ_{129}/Γ

| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|----------------------|-------------|---------------------------------------|
| 1.10±0.10±0.08 | 491 | ¹ ABLIKIM | 19N BES3 | $\psi(2S) \rightarrow \eta' p\bar{p}$ |

¹ From the combination of $p\bar{p}\eta' \rightarrow p\bar{p}\pi^+\pi^-\eta$ and $p\bar{p}\eta' \rightarrow p\bar{p}\pi^+\pi^-\gamma$ channels.

 $\Gamma(p\bar{p}\phi)/\Gamma_{\text{total}}$ Γ_{130}/Γ

| <u>VALUE (units 10^{-6})</u> | <u>CL%</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|-------------|--------------------|-------------|---|
| 6.06±0.38±0.48 | | 753 | ABLIKIM | 19AO BES3 | $e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}K^+K^-$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-----|----|------------------|-----|------|---|
| <24 | 90 | BRIERE | 05 | CLEO | $e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}K^+K^-$ |
| <26 | 90 | ¹ BAI | 03B | BES | $\psi(2S) \rightarrow K^+K^-p\bar{p}$ |

¹ Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.

 $\Gamma(\phi X(1835) \rightarrow p\bar{p}\phi)/\Gamma_{\text{total}}$ Γ_{131}/Γ

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------------------|------------|--------------------|-------------|---|
| <1.82 × 10⁻⁷ | 90 | ABLIKIM | 19AO BES3 | $e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}K^+K^-$ |

 $\Gamma(p\bar{n}\pi^- \text{ or c.c.})/\Gamma_{\text{total}}$ Γ_{132}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|-------------------------------------|
| 2.48±0.17 OUR AVERAGE | | | | |
| 2.45 $\pm 0.11 \pm 0.21$ | 851 | ABLIKIM | 06I BES2 | $e^+ e^- \rightarrow p\pi^-X$ |
| 2.52 $\pm 0.12 \pm 0.22$ | 849 | ABLIKIM | 06I BES2 | $e^+ e^- \rightarrow \bar{p}\pi^+X$ |

 $\Gamma(p\bar{n}\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{133}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------|--------------------|-------------|------------------------------------|
| 3.18±0.50±0.50 | 135 ± 21 | ABLIKIM | 06I BES2 | $e^+ e^- \rightarrow p\pi^-\pi^0X$ |

 $\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$ Γ_{134}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>CL%</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|------------|---------------------|--------------------|---|----------------|
| 3.81±0.13 OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below. | | | | | |
| 3.97 $\pm 0.02 \pm 0.12$ | 31k | ABLIKIM | 17L BES3 | $e^+ e^- \rightarrow \Lambda\bar{\Lambda}$ | |
| 3.71 $\pm 0.05 \pm 0.15$ | 6.5k | ¹ DOBBS | 17 | $e^+ e^- \rightarrow \Lambda\bar{\Lambda}$ | |
| 3.39 $\pm 0.20 \pm 0.32$ | 337 | ABLIKIM | 07C BES | $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ | |
| 6.4 $\pm 1.8 \pm 0.1$ | | ² AUBERT | 07BD BABR | 10.6 $e^+ e^- \rightarrow \Lambda\bar{\Lambda}\gamma$ | |
| 3.28 $\pm 0.23 \pm 0.25$ | 208 | PEDLAR | 05 CLEO | $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ | |

• • • We do not use the following data for averages, fits, limits, etc. • • •


| | | | | |
|--------------------------|--------|----------------------|----|---|
| $3.75 \pm 0.09 \pm 0.23$ | $1.9k$ | ^{1,3} DOBBS | 14 | $e^+ e^- \rightarrow \Lambda \bar{\Lambda}$ |
| $1.81 \pm 0.20 \pm 0.27$ | 80 | ⁴ BAI | 01 | $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |
| < 4 | 90 | FELDMAN | 77 | $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² AUBERT 07BD reports $[\Gamma(\psi(2S) \rightarrow \Lambda \bar{\Lambda})/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (15 \pm 4 \pm 1) \times 10^{-4}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.33 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Superseded by DOBBS 17.

⁴ Estimated using $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$.



$\Gamma(\Lambda \bar{\Lambda} \pi^0)/\Gamma_{\text{total}}$

| VALUE (units 10^{-6}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT | |
|--|-----|------|----------------------|-----------|--|---|
| $1.42 \pm 0.39 \pm 0.59$ | | 23 | ¹ ABLIKIM | 22AP BES3 | $\psi(2S) \rightarrow p \bar{p} \pi^+ \pi^- \gamma \gamma$ | █ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-------|----|----------------------|----------|--|
| < 2.9 | 90 | ² ABLIKIM | 13F BES3 | $\psi(2S) \rightarrow p \bar{p} \pi^+ \pi^- \gamma \gamma$ |
| < 120 | 90 | ³ ABLIKIM | 07H BES2 | $e^+ e^- \rightarrow \psi(2S)$ |

¹ With a significance of 3.7σ . The corresponding 90% CL upper limit is 2.47×10^{-6} .

² Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\pi^0 \rightarrow \gamma \gamma) = 98.8\%$.

³ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\eta \rightarrow \gamma \gamma) = 39.4\%$.

$\Gamma(\Lambda\bar{\Lambda}\eta)/\Gamma_{\text{total}}$ Γ_{136}/Γ

| <u>VALUE</u> (units 10^{-5}) | <u>CL%</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
|---|------------|----------------------|--------------------|---|----------------|--|
| 2.43 ± 0.32 OUR AVERAGE | | | | | | |
| 2.34 $\pm 0.18 \pm 0.52$ | 218 | ABLIKIM | 22AP BES3 | $\psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$ | | |
| 2.48 $\pm 0.34 \pm 0.19$ | 60 | ¹ ABLIKIM | 13F BES3 | $\psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$ | | |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | | |
| <4.9 | 90 | ² ABLIKIM | 07H BES2 | $e^+e^- \rightarrow \psi(2S)$ | | |

¹ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\eta \rightarrow \gamma\gamma) = 39.31\%$.² Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$. $\Gamma(\Lambda\bar{\Lambda}\eta')/\Gamma_{\text{total}}$ Γ_{138}/Γ

| <u>VALUE</u> (units 10^{-6}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
|--|-------------|--------------------|-------------|--|--|
| $7.34 \pm 0.94 \pm 0.43$ | 218 | ABLIKIM | 23BV BES3 | $\psi(2S) \rightarrow p\bar{p}2(\pi^+\pi^-)\gamma(\gamma)$ | |

 $\Gamma(\Lambda\bar{\Lambda}\omega(782))/\Gamma_{\text{total}}$ Γ_{139}/Γ

| <u>VALUE</u> (units 10^{-5}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
|--|-------------|----------------------|-------------|-------------------------------|--|
| $3.30 \pm 0.34 \pm 0.29$ | 207 | ¹ ABLIKIM | 22AZ BES3 | $e^+e^- \rightarrow \psi(2S)$ | |
| 1 Using $B(\Lambda \rightarrow \pi^- p) = 0.639$ and $B(\omega \rightarrow \pi^+\pi^-\pi^0) = 0.893$. | | | | | |

 $\Gamma(\Lambda(1670)\bar{\Lambda} \rightarrow \Lambda\bar{\Lambda}\eta)/\Gamma_{\text{total}}$ Γ_{137}/Γ

| <u>VALUE</u> (units 10^{-5}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
|---|-------------|----------------------|-------------|---|--|
| $1.29 \pm 0.31 \pm 0.62$ | 116 | ¹ ABLIKIM | 22AP BES3 | $\psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$ | |
| 1 From a partial wave analysis of the $\Lambda\eta$ system. | | | | | |

 $\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{140}/Γ

| <u>VALUE</u> (units 10^{-4}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
|---|-------------|--------------------|-------------|--|--|
| $2.8 \pm 0.4 \pm 0.5$ | 73.4 | BRIERE | 05 | CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}2(\pi^+\pi^-)$ | |

 $\Gamma(\Lambda\bar{p}K^+)/\Gamma_{\text{total}}$ Γ_{141}/Γ

| <u>VALUE</u> (units 10^{-4}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
|---|-------------|--------------------|-------------|---|--|
| $1.0 \pm 0.1 \pm 0.1$ | 74.0 | BRIERE | 05 | CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}K^+\pi^-$ | |

 $\Gamma(\Lambda\bar{p}K^*(892)^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{142}/Γ

| <u>VALUE</u> (units 10^{-5}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
|---|-------------|--------------------|-------------|-------------------------------|--|
| $6.3 \pm 0.5 \pm 0.5$ | 1011 | ABLIKIM | 19AU BES3 | $e^+e^- \rightarrow \psi(2S)$ | |

 $\Gamma(\Lambda\bar{p}K^+\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{143}/Γ

| <u>VALUE</u> (units 10^{-4}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
|---|-------------|--------------------|-------------|---|--|
| $1.8 \pm 0.3 \pm 0.3$ | 45.8 | BRIERE | 05 | CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}K^+\pi^+\pi^-\pi^-$ | |

$\Gamma(\bar{\Lambda} n K_S^0 + \text{c.c.})/\Gamma_{\text{total}}$

Γ_{144}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|----------|------------------------------|
| 0.81±0.11±0.14 | 50 | 1 ABLIKIM | 08C BES2 | $e^+ e^- \rightarrow J/\psi$ |

¹ Using $B(\bar{\Lambda} \rightarrow \bar{p}\pi^+) = 63.9\%$ and $B(K_S^0 \rightarrow \pi^+\pi^-) = 69.2\%$.

$\Gamma(\Delta^{++} \bar{\Delta}^{--})/\Gamma_{\text{total}}$

Γ_{145}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|--------|---|
| 12.8±1.0±3.4 | 157 | 1 BAI | 01 BES | $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |

¹ Estimated using $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.310 \pm 0.028$.

$\Gamma(\Lambda \bar{\Sigma}^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$

Γ_{146}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|----------|---------------------------------------|
| 1.40±0.03±0.13 | 2.8k | ABLIKIM | 13W BES3 | $\psi(2S) \rightarrow \text{hadrons}$ |

$\Gamma(\Lambda \bar{\Sigma}^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$

Γ_{147}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|----------|---------------------------------------|
| 1.54±0.04±0.13 | 2.8k | ABLIKIM | 13W BES3 | $\psi(2S) \rightarrow \text{hadrons}$ |

$\Gamma(\Lambda \bar{\Sigma}^0 + \text{c.c.})/\Gamma_{\text{total}}$

Γ_{148}/Γ

| VALUE (units 10^{-6}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|----------|---|
| 1.60±0.31±0.59 | 60 | ABLIKIM | 21L BES3 | $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |

$\Gamma(\Lambda \bar{\Sigma}^0)/\Gamma_{\text{total}}$

Γ_{149}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | COMMENT |
|--|------|-------------|--|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 1.23±0.23±0.08 | 30 | 1 DOBBS | 17 $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\Sigma^0 \bar{p} K^+ + \text{c.c.})/\Gamma_{\text{total}}$

Γ_{150}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|----------|---|
| 1.67±0.13±0.12 | 276 | 1 ABLIKIM | 13D BES3 | $\psi(2S) \rightarrow \gamma \Lambda \bar{p} K^+$ |

¹ Using $B(\Lambda \rightarrow p\pi^-) = 63.9\%$, and $B(\Sigma^0 \rightarrow \Lambda\gamma) = 100\%$.

$\Gamma(\Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}$

Γ_{151}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------|------|-------------|------|-------------------------------------|
| 2.43±0.10 OUR AVERAGE | | | | Error includes scale factor of 1.4. |

| | | | | |
|----------------|------|--------------------|-----------|---|
| 2.52±0.04±0.09 | 5.4k | ABLIKIM | 21AT BES3 | $\psi(2S) \rightarrow p\pi^0 \bar{p}\pi^0$ |
| 2.31±0.06±0.10 | 1.9k | ¹ DOBBS | 17 | $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |
| 2.57±0.44±0.68 | 35 | PEDLAR | 05 CLEO | $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|----------------|-----|----------------------|----|---|
| 2.51±0.15±0.16 | 281 | ^{1,2} DOBBS | 14 | $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |
|----------------|-----|----------------------|----|---|

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² Superseded by DOBBS 17.

$\Gamma(\Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}$ Γ_{152}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
|--|-------------|-------------------------------------|-------------|---|--|
| 2.35 ± 0.09 OUR AVERAGE | | Error includes scale factor of 1.1. | | | |
| $2.44 \pm 0.03 \pm 0.11$ | 7k | ABLIKIM | 17L BES3 | $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ | |
| $2.22 \pm 0.05 \pm 0.11$ | 2.6k | ¹ DOBBS | 17 | $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ | |
| $2.35 \pm 0.36 \pm 0.32$ | 59 | ABLIKIM | 07C BES | $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ | |
| $2.63 \pm 0.35 \pm 0.21$ | 58 | PEDLAR | 05 CLEO | $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ | |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| $2.25 \pm 0.11 \pm 0.16$ | 439 | ^{1,2} DOBBS | 14 | $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ | |
| $1.2 \pm 0.4 \pm 0.4$ | 8 | ³ BAI | 01 BES | $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ | |

¹ Using CLEO-c data but not authored by the CLEO Collaboration.² Superseded by DOBBS 17.³ Estimated using $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$. $\Gamma(\Sigma^- \bar{\Sigma}^+)/\Gamma_{\text{total}}$ Γ_{153}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|-------------|--------------------|-------------|--|
| $2.82 \pm 0.04 \pm 0.08$ | 6.6k | ABLIKIM | 22AV BES3 | $\psi(2S) \rightarrow n\pi^- \bar{n}\pi^+$ |

 $\Gamma(\Sigma^+ \bar{\Sigma}^- \eta)/\Gamma_{\text{total}}$ Γ_{154}/Γ

| <u>VALUE (units 10^{-6})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|-------------|--------------------|-------------|---|
| $9.59 \pm 2.37 \pm 0.61$ | 21 | ABLIKIM | 22AY BES3 | $\psi(2S) \rightarrow \Sigma^+ \bar{\Sigma}^- \eta$ |

 $\Gamma(\Sigma^+ \bar{\Sigma}^- \omega)/\Gamma_{\text{total}}$ Γ_{155}/Γ

| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|-------------|--------------------|-------------|---|
| $1.89 \pm 0.18 \pm 0.21$ | 199 | ABLIKIM | 23BE BES3 | $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |

 $\Gamma(\Sigma^+ \bar{\Sigma}^- \phi)/\Gamma_{\text{total}}$ Γ_{156}/Γ

| <u>VALUE (units 10^{-6})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|-------------|--------------------|-------------|---|
| $2.96 \pm 0.54 \pm 0.41$ | 55 | ABLIKIM | 23BE BES3 | $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |

 $\Gamma(\Sigma(1385)^+ \bar{\Sigma}(1385)^-)/\Gamma_{\text{total}}$ Γ_{157}/Γ

| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|--|
| 8.5 ± 0.7 OUR AVERAGE | | | | |
| $8.4 \pm 0.5 \pm 0.5$ | 1.5k | ABLIKIM | 16L BES3 | $\psi(2S) \rightarrow \Sigma(1385)^+ \bar{\Sigma}(1385)^-$ |
| $11 \pm 3 \pm 3$ | 14 | ¹ BAI | 01 BES | $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |

¹ Estimated using $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$. $\Gamma(\Sigma(1385)^- \bar{\Sigma}(1385)^+)/\Gamma_{\text{total}}$ Γ_{158}/Γ

| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|--|
| $8.5 \pm 0.6 \pm 0.6$ | 1.4k | ABLIKIM | 16L BES3 | $\psi(2S) \rightarrow \Sigma(1385)^- \bar{\Sigma}(1385)^+$ |

 $\Gamma(\Sigma(1385)^0 \bar{\Sigma}(1385)^0)/\Gamma_{\text{total}}$ Γ_{159}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|-------------|--------------------|-------------|---|
| $0.69 \pm 0.05 \pm 0.05$ | 2.2k | ABLIKIM | 17E BES3 | $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |

$\Gamma(\Xi^-\Xi^+)/\Gamma_{\text{total}}$ Γ_{160}/Γ

| <i>VALUE</i> (units 10^{-4}) | <i>CL%</i> | <i>EVTS</i> | <i>DOCUMENT ID</i> | <i>TECN</i> | <i>COMMENT</i> |
|---|------------|-------------|--------------------|--|-------------------------------------|
| 2.87±0.11 OUR AVERAGE | | | | | |
| | | | | | Error includes scale factor of 1.1. |
| 3.03±0.05±0.14 | 3.6k | 1 DOBBS | 17 | $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ | |
| 2.78±0.05±0.14 | 5k | ABLIKIM | 16L BES3 | $\psi(2S) \rightarrow \Xi^-\Xi^+$ | |
| 3.03±0.40±0.32 | 67 | ABLIKIM | 07C BES | $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ | |
| 2.38±0.30±0.21 | 63 | PEDLAR | 05 CLEO | $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ | |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| 2.66±0.12±0.20 | 548 | 1,2 DOBBS | 14 | $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ | |
| 0.94±0.27±0.15 | 12 | 3 BAI | 01 BES | $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ | |
| <2 | 90 | FELDMAN | 77 MRK1 | $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ | |

¹ Using CLEO-c data but not authored by the CLEO Collaboration.² Superseded by DOBBS 17.³ Estimated using $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$. $\Gamma(\Xi^0\Xi^0)/\Gamma_{\text{total}}$ Γ_{161}/Γ

| <i>VALUE</i> (units 10^{-4}) | <i>EVTS</i> | <i>DOCUMENT ID</i> | <i>TECN</i> | <i>COMMENT</i> |
|---|-------------|--------------------|-------------|--|
| 2.3 ±0.4 OUR AVERAGE | | | | |
| | | | | Error includes scale factor of 4.2. |
| 2.73±0.03±0.13 | 11k | ABLIKIM | 17E BES3 | $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |
| 1.97±0.06±0.11 | 1.2k | 1 DOBBS | 17 | $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |
| 2.75±0.64±0.61 | 19 | PEDLAR | 05 CLEO | $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 2.02±0.19±0.15 | 112 | 1,2 DOBBS | 14 | $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |

¹ Using CLEO-c data but not authored by the CLEO Collaboration.² Superseded by DOBBS 17. $\Gamma(\Xi(1530)^0\Xi(1530)^0)/\Gamma_{\text{total}}$ Γ_{162}/Γ

| <i>VALUE</i> (units 10^{-5}) | <i>CL%</i> | <i>EVTS</i> | <i>DOCUMENT ID</i> | <i>TECN</i> | <i>COMMENT</i> |
|---|------------|-------------|--------------------|--|--|
| 6.77±0.14±0.39 | | | | | |
| | | 2951 | ABLIKIM | 21AO BES3 | $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| <32 | 90 | PEDLAR | 05 CLEO | $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ | |
| < 8.1 | 90 | 1 BAI | 01 BES | $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ | |

¹ Estimated using $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$. $\Gamma(\Lambda\Xi^+ K^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{163}/Γ

| <i>VALUE</i> (units 10^{-5}) | <i>EVTS</i> | <i>DOCUMENT ID</i> | <i>TECN</i> | <i>COMMENT</i> |
|---------------------------------|-------------|--------------------|-------------|---|
| 3.86±0.27±0.32 | | | | |
| | 236 | ABLIKIM | 15I BES3 | $e^+e^- \rightarrow \psi(2S) \rightarrow K^-\Lambda\Xi^+ + \text{c.c.}$ |

 $\Gamma(\Xi(1530)^-\Xi(1530)^+)/\Gamma_{\text{total}}$ Γ_{166}/Γ

| <i>VALUE</i> (units 10^{-5}) | <i>EVTS</i> | <i>DOCUMENT ID</i> | <i>TECN</i> | <i>COMMENT</i> |
|---------------------------------|-------------|--------------------|-------------|--|
| 11.45±0.40±0.59 | | | | |
| | 5k | ABLIKIM | 19AT BES3 | $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |

$\Gamma(\Xi(1530)^-\Xi^+)/\Gamma_{\text{total}}$ Γ_{167}/Γ

| <i>VALUE</i> (units 10^{-6}) | <i>EVTS</i> | <i>DOCUMENT ID</i> | <i>TECN</i> | <i>COMMENT</i> |
|---|-------------|--------------------|-------------|---|
| $7.0 \pm 1.1 \pm 0.4$ | 399 | ABLIKIM | 19AT BES3 | $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |

 $\Gamma(\Xi(1530)^0\Xi^0)/\Gamma_{\text{total}}$ Γ_{168}/Γ

| <i>VALUE</i> (units 10^{-5}) | <i>EVTS</i> | <i>DOCUMENT ID</i> | <i>TECN</i> | <i>COMMENT</i> |
|--|-------------|--------------------|-------------|---|
| $0.53 \pm 0.04 \pm 0.03$ | 278 | ABLIKIM | 21AO BES3 | $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |

 $\Gamma(\Xi(1690)^-\Xi^+ \rightarrow K^-\Lambda\Xi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{164}/Γ

| <i>VALUE</i> (units 10^{-6}) | <i>EVTS</i> | <i>DOCUMENT ID</i> | <i>TECN</i> | <i>COMMENT</i> |
|--|-------------|--------------------|-------------|--|
| $5.21 \pm 1.48 \pm 0.57$ | 74 | ABLIKIM | 15I | $e^+ e^- \rightarrow \psi(2S) \rightarrow K^-\Lambda\Xi^+ + \text{c.c.}$ |

 $\Gamma(\Xi(1820)^-\Xi^+ \rightarrow K^-\Lambda\Xi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{165}/Γ

| <i>VALUE</i> (units 10^{-6}) | <i>EVTS</i> | <i>DOCUMENT ID</i> | <i>TECN</i> | <i>COMMENT</i> |
|---|-------------|--------------------|-------------|--|
| $12.03 \pm 2.94 \pm 1.22$ | 136 | ABLIKIM | 15I | $e^+ e^- \rightarrow \psi(2S) \rightarrow K^-\Lambda\Xi^+ + \text{c.c.}$ |

 $\Gamma(\Sigma^0\Xi^+ K^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{169}/Γ

| <i>VALUE</i> (units 10^{-5}) | <i>EVTS</i> | <i>DOCUMENT ID</i> | <i>TECN</i> | <i>COMMENT</i> |
|--|-------------|--------------------|-------------|---|
| $3.67 \pm 0.33 \pm 0.28$ | 142 | ABLIKIM | 15I | $e^+ e^- \rightarrow \psi(2S) \rightarrow K^-\Sigma^0\Xi^+ + \text{c.c.}$ |

 $\Gamma(\Omega^-\bar{\Omega}^+)/\Gamma_{\text{total}}$ Γ_{170}/Γ

| <i>VALUE</i> (units 10^{-5}) | <i>CL%</i> | <i>EVTS</i> | <i>DOCUMENT ID</i> | <i>TECN</i> | <i>COMMENT</i> |
|---|------------|-------------|-------------------------------------|-------------|---|
| 5.66 ± 0.30 OUR AVERAGE | | | Error includes scale factor of 1.3. | | |
| 5.85 $\pm 0.12 \pm 0.25$ | | 4k | ¹ ABLIKIM | 21E BES3 | $\psi(2S) \rightarrow \Omega^-\bar{\Omega}^+ \rightarrow \Lambda K^-\bar{\Lambda}K^+$ |
| 5.2 $\pm 0.3 \pm 0.3$ | | 326 | ^{1,2} DOBBS | 17 | $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |

| | | | | | |
|-----------------------|--|----|------------------------|----------|---|
| 4.7 $\pm 0.9 \pm 0.5$ | | 27 | ^{1,2,3} DOBBS | 14 | $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |
| <15 | | 90 | ABLIKIM | 12Q BES2 | $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |
| <16 | | 90 | PEDLAR | 05 CLEO | $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |
| < 7.3 | | 90 | ⁴ BAI | 01 BES | $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ |

¹ Using $B(\Omega^- \rightarrow \Lambda K^-) = (67.8 \pm 0.7)\%$ and $B(\Lambda \rightarrow p\pi^-) = (63.9 \pm 0.5)\%$.² Using CLEO-c data but not authored by the CLEO Collaboration.³ Superseded by DOBBS 17.⁴ Estimated using $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.310 \pm 0.028$. $\Gamma(\eta_c\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{171}/Γ

| <i>VALUE</i> (units 10^{-3}) | <i>CL%</i> | <i>EVTS</i> | <i>DOCUMENT ID</i> | <i>TECN</i> | <i>COMMENT</i> |
|---------------------------------|------------|-------------|--------------------|-------------|--------------------------------|
| <1.0 | | 90 | PEDLAR | 07 CLEO | $e^+ e^- \rightarrow \psi(2S)$ |

$\Gamma(h_c(1P)\pi^0)/\Gamma_{\text{total}}$ Γ_{172}/Γ

| <u>VALUE</u> (units 10^{-4}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------------|----------------------|-------------|--|
| 7.4 ± 0.5 OUR AVERAGE | | | | |
| 7.32 ± 0.34 ± 0.41 | 46k | ABLIKIM | 22AQ | BES3 $\psi(2S) \rightarrow \pi^0$ hadrons |
| 9.0 ± 1.5 ± 1.3 | 3k | ¹ GE | 11 | CLEO $\psi(2S) \rightarrow \pi^0$ anything |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 8.4 ± 1.3 ± 1.0 | 11k | ² ABLIKIM | 10B | BES3 $\psi(2S) \rightarrow \pi^0 h_c$ |
| seen | 92^{+23}_{-22} | ADAMS | 09 | CLEO $\psi(2S) \rightarrow 2\pi^+ 2\pi^- 2\pi^0$ |
| seen | 1282 | DOBBS | 08A | CLEO $\psi(2S) \rightarrow \pi^0 \eta_c \gamma$ |
| seen | 168 ± 40 | ROSNER | 05 | CLEO $\psi(2S) \rightarrow \pi^0 \eta_c \gamma$ |

¹ Assuming a width $\Gamma(h_c(1P)) = 0.86$ MeV $\equiv \Gamma_0$, a measured dependence of the central value of $B = (7.6 + 1.4 \times \Gamma(h_c(1P)/\Gamma_0)) \times 10^{-4}$, and with a systematic error that accounts for the width variation range 0.43–1.29 MeV.

² Superseded by ABLIKIM 22AQ

 $\Gamma(\Lambda_c^+ \bar{p} e^+ e^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{173}/Γ

| <u>VALUE</u> | <u>CL%</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------|------------|-------------|--------------------|-------------|-------------------------------------|
| $<1.7 \times 10^{-6}$ | 90 | 450M | ABLIKIM | 18Q | BES3 $e^+ e^- \rightarrow \psi(2S)$ |

 $\Gamma(\Theta(1540) \bar{\Theta}(1540) \rightarrow K_S^0 p K^- \bar{n} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{174}/Γ

| <u>VALUE</u> (units 10^{-5}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|------------|--------------------|-------------|----------------|
| <0.88 | 90 | BAI | 04G | BES2 $e^+ e^-$ |

 $\Gamma(\Theta(1540) K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n})/\Gamma_{\text{total}}$ Γ_{175}/Γ

| <u>VALUE</u> (units 10^{-5}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|------------|--------------------|-------------|----------------|
| <1.0 | 90 | BAI | 04G | BES2 $e^+ e^-$ |

 $\Gamma(\Theta(1540) K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n)/\Gamma_{\text{total}}$ Γ_{176}/Γ

| <u>VALUE</u> (units 10^{-5}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|------------|--------------------|-------------|----------------|
| <0.70 | 90 | BAI | 04G | BES2 $e^+ e^-$ |

 $\Gamma(\bar{\Theta}(1540) K^+ n \rightarrow K_S^0 \bar{p} K^+ n)/\Gamma_{\text{total}}$ Γ_{177}/Γ

| <u>VALUE</u> (units 10^{-5}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|------------|--------------------|-------------|----------------|
| <2.6 | 90 | BAI | 04G | BES2 $e^+ e^-$ |

 $\Gamma(\bar{\Theta}(1540) K_S^0 p \rightarrow K_S^0 p K^- \bar{n})/\Gamma_{\text{total}}$ Γ_{178}/Γ

| <u>VALUE</u> (units 10^{-5}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|------------|--------------------|-------------|----------------|
| <0.60 | 90 | BAI | 04G | BES2 $e^+ e^-$ |

 RADIATIVE DECAYS

 $\Gamma(\gamma \chi_{c0}(1P))/\Gamma_{\text{total}}$ Γ_{179}/Γ

| <u>VALUE</u> (units 10^{-2}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|-------------|-------------------------------------|-------------|----------------|
| 9.77 ± 0.23 OUR FIT | | Error includes scale factor of 1.1. | | |

9.33 ± 0.26 OUR AVERAGE

| | | | | |
|-----------------------|------|---------|-----|-------------------------------------|
| 9.389 ± 0.014 ± 0.332 | 4.7M | ABLIKIM | 17U | BES3 $e^+ e^- \rightarrow \gamma X$ |
| 9.22 ± 0.11 ± 0.46 | 72k | ATHAR | 04 | CLEO $e^+ e^- \rightarrow \gamma X$ |

| | | | | | | |
|-----|-----------|-----------|-----------------------|----|------|--------------------------------|
| 9.9 | ± 0.5 | ± 0.8 | ¹ GAISER | 86 | CBAL | $e^+ e^- \rightarrow \gamma X$ |
| 7.2 | ± 2.3 | | ¹ BIDDICK | 77 | CNTR | $e^+ e^- \rightarrow \gamma X$ |
| 7.5 | ± 2.6 | | ¹ WHITAKER | 76 | MRK1 | $e^+ e^-$ |

¹ Angular distribution ($1+\cos^2\theta$) assumed.

$\Gamma(\gamma\chi_{c1}(1P))/\Gamma_{\text{total}}$

Γ_{180}/Γ

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------------------------------|------|-------------------------------------|
| 9.75 ± 0.27 OUR FIT | | Error includes scale factor of 1.1. | | |
| 9.54 ± 0.29 OUR AVERAGE | | | | |
| 9.905 $\pm 0.011 \pm 0.353$ | 5.0M | ABLIKIM | 17U | BES3 $e^+ e^- \rightarrow \gamma X$ |
| 9.07 $\pm 0.11 \pm 0.54$ | 76k | ATHAR | 04 | CLEO $e^+ e^- \rightarrow \gamma X$ |
| 9.0 $\pm 0.5 \pm 0.7$ | | ¹ GAISER | 86 | CBAL $e^+ e^- \rightarrow \gamma X$ |
| 7.1 ± 1.9 | | ² BIDDICK | 77 | CNTR $e^+ e^- \rightarrow \gamma X$ |

¹ Angular distribution ($1-0.189 \cos^2\theta$) assumed.

² Valid for isotropic distribution of the photon.

$\Gamma(\gamma\chi_{c0}(1P))/\Gamma(\gamma\chi_{c1}(1P))$

$\Gamma_{179}/\Gamma_{180}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|--------------------|------|-------------------------------------|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 1.02 $\pm 0.01 \pm 0.07$ | ¹ ATHAR | 04 | CLEO $e^+ e^- \rightarrow \gamma X$ |
| ¹ Not independent from ATHAR 04 measurements of $B(\gamma\chi_{cJ})$. | | | |


$\Gamma(\gamma\chi_{c2}(1P))/\Gamma_{\text{total}}$

Γ_{181}/Γ

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|---|------|-------------------------------------|
| 9.36 ± 0.23 OUR FIT | | Error includes scale factor of 1.2. | | |
| 9.42 ± 0.31 OUR AVERAGE | | Error includes scale factor of 1.3. See the ideogram below. | | |
| 9.621 $\pm 0.013 \pm 0.272$ | 4.2M | ABLIKIM | 17U | BES3 $e^+ e^- \rightarrow \gamma X$ |
| 9.33 $\pm 0.14 \pm 0.61$ | 79k | ATHAR | 04 | CLEO $e^+ e^- \rightarrow \gamma X$ |
| 8.0 $\pm 0.5 \pm 0.7$ | | ¹ GAISER | 86 | CBAL $e^+ e^- \rightarrow \gamma X$ |
| 7.0 ± 2.0 | | ² BIDDICK | 77 | CNTR $e^+ e^- \rightarrow \gamma X$ |

¹ Angular distribution ($1-0.052 \cos^2\theta$) assumed.

² Valid for isotropic distribution of the photon.



$$\frac{[\Gamma(\gamma\chi_{c0}(1P)) + \Gamma(\gamma\chi_{c1}(1P)) + \Gamma(\gamma\chi_{c2}(1P))]}{\Gamma_{\text{total}}} (\Gamma_{179} + \Gamma_{180} + \Gamma_{181})/\Gamma$$

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

$27.6 \pm 0.3 \pm 2.0$ ¹ ATHAR 04 CLEO $e^+ e^- \rightarrow \gamma X$

¹ Not independent from ATHAR 04 measurements of $B(\gamma\chi_{cJ})$.

$$\Gamma(\gamma\chi_{c0}(1P))/\Gamma(\gamma\chi_{c2}(1P))$$

$$\Gamma_{179}/\Gamma_{181}$$

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.99 \pm 0.02 \pm 0.08$ ¹ ATHAR 04 CLEO $e^+ e^- \rightarrow \gamma X$

¹ Not independent from ATHAR 04 measurements of $B(\gamma\chi_{cJ})$.

$$\Gamma(\gamma\chi_{c2}(1P))/\Gamma(\gamma\chi_{c1}(1P))$$

$$\Gamma_{181}/\Gamma_{180}$$

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.03 \pm 0.02 \pm 0.03$ ¹ ATHAR 04 CLEO $e^+ e^- \rightarrow \gamma X$

¹ Not independent from ATHAR 04 measurements of $B(\gamma\chi_{cJ})$.

$$\Gamma(\gamma\eta_c(1S))/\Gamma_{\text{total}}$$

$$\Gamma_{182}/\Gamma$$

VALUE (units 10^{-2}) EVTS DOCUMENT ID TECN COMMENT

0.36 ± 0.05 OUR FIT Error includes scale factor of 1.3.

0.34 ± 0.05 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.


$0.432 \pm 0.016 \pm 0.060$ MITCHELL 09 CLEO $e^+ e^- \rightarrow \gamma X$

$0.32 \pm 0.04 \pm 0.06$ ^{2.5k} ¹ ATHAR 04 CLEO $e^+ e^- \rightarrow \gamma X$

0.28 ± 0.06 ² GAISER 86 CBAL $e^+ e^- \rightarrow \gamma X$

¹ ATHAR 04 used $\Gamma_{\eta_c}(1S) = 24.8 \pm 4.9$ MeV to obtain this result.

² GAISER 86 used $\Gamma_{\eta_c}(1S) = 11.5 \pm 4.5$ MeV to obtain this result.



$\Gamma(\gamma\eta_c(2S))/\Gamma_{\text{total}}$

Γ_{183}/Γ

| VALUE (units 10^{-4}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-----------------------------|------|---|
| 7±2±4 | | ¹ ABLIKIM | 12G | $\psi(2S) \rightarrow \gamma K^0 K\pi, K\bar{K}\pi^0$ |
| < 8 | 90 | ² CRONIN-HEN..10 | CLEO | $\psi(2S) \rightarrow \gamma K\bar{K}\pi$ |
| <20 | 90 | ATHAR | 04 | $e^+ e^- \rightarrow \gamma X$ |
| 20–130 | 95 | EDWARDS | 82C | $e^+ e^- \rightarrow \gamma X$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 8 90 ² CRONIN-HENNESSY 10 CLEO $\psi(2S) \rightarrow \gamma K\bar{K}\pi$
<20 90 ATHAR 04 CLEO $e^+ e^- \rightarrow \gamma X$
20–130 95 EDWARDS 82C CBAL $e^+ e^- \rightarrow \gamma X$

¹ ABLIKIM 12G reports $[\Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}] \times [B(\eta_c(2S) \rightarrow K\bar{K}\pi)] = (1.30 \pm 0.20 \pm 0.30) \times 10^{-5}$ which we divide by our best value $B(\eta_c(2S) \rightarrow K\bar{K}\pi) = (1.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² CRONIN-HENNESSY 10 reports $[\Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}] \times [B(\eta_c(2S) \rightarrow K\bar{K}\pi)] < 14.5 \times 10^{-6}$ which we divide by our best value $B(\eta_c(2S) \rightarrow K\bar{K}\pi) = 1.9 \times 10^{-2}$. This measurement assumes $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$\Gamma(\gamma\pi^0)/\Gamma_{\text{total}}$

Γ_{184}/Γ

| VALUE (units 10^{-6}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-----|-----------------------|-------------------------------------|----------|------------------------------------|
| 1.04±0.22 OUR AVERAGE | | | Error includes scale factor of 1.4. | | |
| 0.95±0.16±0.05 | | 423 | ABLIKIM | 17X BES3 | $\psi(2S) \rightarrow \gamma\pi^0$ |
| 1.58±0.40±0.13 | | 37 | ABLIKIM | 10F BES3 | $\psi(2S) \rightarrow \gamma\pi^0$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| < 5 | 90 | PEDLAR | 09 | CLE3 | $\psi(2S) \rightarrow \gamma X$ |
| < 5400 | 95 | ¹ LIBERMAN | 75 | SPEC | e^+e^- |
| $< 1 \times 10^4$ | 90 | WIIK | 75 | DASP | e^+e^- |

¹ Restated by us using $B(\psi(2S) \rightarrow \mu^+\mu^-) = 0.0077$.

$\Gamma(\gamma 2(\pi^+\pi^-))/\Gamma_{\text{total}}$

Γ_{185}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|----------|-------------------------------|
| 39.6±2.8±5.0 | 583 | ABLIKIM | 07D BES2 | $e^+e^- \rightarrow \psi(2S)$ |

$\Gamma(\gamma 3(\pi^+\pi^-))/\Gamma_{\text{total}}$

Γ_{186}/Γ

| VALUE (units 10^{-5}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|----------|-------------------------------|
| <17 | 90 | ABLIKIM | 07D BES2 | $e^+e^- \rightarrow \psi(2S)$ |

$\Gamma(\gamma\eta'(958))/\Gamma_{\text{total}}$

Γ_{187}/Γ

| VALUE (units 10^{-4}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------------------------------|-----|------|----------------------|----------|--|
| 1.24 ±0.04 OUR AVERAGE | | | | | |
| 1.251±0.022±0.062 | | 56k | ABLIKIM | 17X BES3 | $\psi(2S) \rightarrow \gamma\pi^+\pi^-\eta,$ $\gamma\pi^0\pi^0\eta$ |
| 1.26 ±0.03 ±0.08 | | 2226 | ¹ ABLIKIM | 10F BES3 | $\psi(2S) \rightarrow 3\gamma\pi^+\pi^-,$ $2\gamma\pi^+\pi^-$ |
| 1.19 ±0.08 ±0.03 | | | PEDLAR | 09 CLE3 | $\psi(2S) \rightarrow \gamma X$ |
| 1.24 ±0.27 ±0.15 | | 23 | ABLIKIM | 06R BES2 | $e^+e^- \rightarrow \psi(2S)$ |
| 1.54 ±0.31 ±0.20 | | ~ 43 | BAI | 98F BES | $\psi(2S) \rightarrow \pi^+\pi^-2\gamma,$ $\pi^+\pi^-3\gamma$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 60 90 ² BRAUNSCH... 77 DASP e^+e^-

< 11 90 ³ BARTEL 76 CNTR e^+e^-

¹ Combining the results from $\eta' \rightarrow \pi^+\pi^-\eta$ and $\eta' \rightarrow \pi^+\pi^-\gamma$ decay modes.

² Restated by us using total decay width 228 keV.

³ The value is normalized to the branching ratio for $\Gamma(J/\psi(1S)\eta)/\Gamma_{\text{total}}$.

$\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$

Γ_{188}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|--------------|-------------------------------------|---------|---|
| 2.73±0.29 OUR AVERAGE | | Error includes scale factor of 1.8. | | |
| 2.84±0.15 ^{+0.03} _{-0.10} | 1.9k | 1,2 DOBBS | 15 | $\psi(2S) \rightarrow \gamma\pi\pi$ |
| 2.12±0.19±0.32 | | 3,4 BAI | 03C BES | $\psi(2S) \rightarrow \gamma\pi\pi$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 2.08±0.19±0.33 | 200.6 ± 18.8 | ³ BAI | 03C BES | $\psi(2S) \rightarrow \gamma\pi^+\pi^-$ |
| 2.90±1.08±1.07 | 29.9 ± 11.1 | ³ BAI | 03C BES | $\psi(2S) \rightarrow \gamma\pi^0\pi^0$ |

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² DOBBS 15 reports $[\Gamma(\psi(2S) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = (2.39 \pm 0.09 \pm 0.09) \times 10^{-4}$ which we divide by our best value $B(f_2(1270) \rightarrow \pi\pi) = (84.3^{+2.8}_{-1.0}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.

⁴ Combining the results from $\pi^+\pi^-$ and $\pi^0\pi^0$ decay modes.

$\Gamma(\gamma f_0(1370) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$

Γ_{189}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | COMMENT |
|---|------|-------------|--|
| $3.1 \pm 1.0 \pm 1.4$ | 175 | 1 DOBBS | $\psi(2S) \rightarrow \gamma K\bar{K}$ |

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\gamma f_0(1500))/\Gamma_{\text{total}}$

Γ_{190}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | COMMENT |
|---|------|-------------|-------------------------------------|
| $9.3 \pm 1.8 \pm 0.6$ | 274 | 1,2 DOBBS | $\psi(2S) \rightarrow \gamma\pi\pi$ |

¹ DOBBS 15 reports $[\Gamma(\psi(2S) \rightarrow \gamma f_0(1500))/\Gamma_{\text{total}}] \times [B(f_0(1500) \rightarrow \pi\pi)] = (3.2 \pm 0.6 \pm 0.2) \times 10^{-5}$ which we divide by our best value $B(f_0(1500) \rightarrow \pi\pi) = (34.5 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\gamma f'_2(1525))/\Gamma_{\text{total}}$

Γ_{191}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | COMMENT |
|---|------|-------------|--|
| $3.3 \pm 0.8 \pm 0.1$ | 136 | 1,2 DOBBS | $\psi(2S) \rightarrow \gamma K\bar{K}$ |

¹ DOBBS 15 reports $[\Gamma(\psi(2S) \rightarrow \gamma f'_2(1525))/\Gamma_{\text{total}}] \times [B(f'_2(1525) \rightarrow K\bar{K})] = (2.9 \pm 0.6 \pm 0.3) \times 10^{-5}$ which we divide by our best value $B(f'_2(1525) \rightarrow K\bar{K}) = (88.8 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\gamma f_0(1710) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$

Γ_{193}/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|----------------|-------------|---------|---|
| 3.5 ± 0.6 OUR AVERAGE | | | | |
| 3.6 $\pm 0.4 \pm 0.5$ | 290 | 1 DOBBS | 15 | $\psi(2S) \rightarrow \gamma\pi\pi$ |
| $3.01 \pm 0.41 \pm 1.24$ | 35.6 ± 4.8 | 2 BAI | 03C BES | $\psi(2S) \rightarrow \gamma\pi^+\pi^-$ |

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.

$\Gamma(\gamma f_0(1710) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$

Γ_{194}/Γ

| VALUE (units 10^{-5}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-----|----------------|---------------|---------|---|
| 6.6 ± 0.7 OUR AVERAGE | | | | | |
| 6.7 $\pm 0.6 \pm 0.6$ | | 375 | 1 DOBBS | 15 | $\psi(2S) \rightarrow \gamma K\bar{K}$ |
| $6.04 \pm 0.90 \pm 1.32$ | | 39.6 ± 5.9 | 2,3 BAI | 03C BES | $\psi(2S) \rightarrow \gamma K^+K^-$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| < 15.6 | | 90 | 6.8 ± 3.1 | 2,3 BAI | 03C BES $\psi(2S) \rightarrow \gamma K_S^0 K_S^0$ |

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² Includes unknown branching fractions to $K^+ K^-$ or $K_S^0 K_S^0$. We have multiplied the $K^+ K^-$ result by a factor of 2 and the $K_S^0 K_S^0$ result by a factor of 4 to obtain the $K \bar{K}$ result.

³ Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

$\Gamma(\gamma f_0(2100) \rightarrow \gamma \pi\pi)/\Gamma_{\text{total}}$

Γ_{195}/Γ

| VALUE (units 10^{-6}) | EVTS | DOCUMENT ID | COMMENT |
|---|------|--------------------|--------------------------------------|
| $4.8 \pm 0.5 \pm 0.9$ | 373 | ¹ DOBBS | $\psi(2S) \rightarrow \gamma \pi\pi$ |

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\gamma f_0(2200) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$

Γ_{196}/Γ

| VALUE (units 10^{-6}) | EVTS | DOCUMENT ID | COMMENT |
|---|------|--------------------|---|
| $3.2 \pm 0.6 \pm 0.8$ | 207 | ¹ DOBBS | $\psi(2S) \rightarrow \gamma K \bar{K}$ |

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\gamma f_J(2220) \rightarrow \gamma \pi\pi)/\Gamma_{\text{total}}$

Γ_{197}/Γ

| VALUE | CL% | DOCUMENT ID | COMMENT |
|--|-----|----------------------|--------------------------------------|
| $<5.8 \times 10^{-6}$ | 90 | ^{1,2} DOBBS | $\psi(2S) \rightarrow \gamma \pi\pi$ |

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² For $\Gamma = 20/50$ MeV, the 90% CL upper limits for $\pi^+ \pi^-$ and $\pi^0 \pi^0$ are $3.2/4.3 \times 10^{-6}$ and $2.6/4.0 \times 10^{-6}$, respectively.

$\Gamma(\gamma f_J(2220) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$

Γ_{198}/Γ

| VALUE | CL% | DOCUMENT ID | COMMENT |
|--|-----|----------------------|---|
| $<9.5 \times 10^{-6}$ | 90 | ^{1,2} DOBBS | $\psi(2S) \rightarrow \gamma K \bar{K}$ |

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² For $\Gamma = 20/50$ MeV, the 90% CL upper limits for $K^+ K^-$ and $K_S^0 K_S^0$ are $2.1/4.3 \times 10^{-6}$ and $3.7/5.5 \times 10^{-6}$, respectively.

$\Gamma(\gamma \eta)/\Gamma_{\text{total}}$

Γ_{199}/Γ

| VALUE (units 10^{-6}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-----|------|----------------------|----------|--|
| $0.92 \pm 0.18 \text{ OUR AVERAGE}$ | | | | | |
| $0.85 \pm 0.18 \pm 0.04$ | 382 | | ¹ ABLIKIM | 17X BES3 | $\psi(2S) \rightarrow \gamma \pi^+ \pi^- \pi^0$, $\gamma 3\pi^0$ |
| $1.38 \pm 0.48 \pm 0.09$ | 13 | | ¹ ABLIKIM | 10F BES3 | $\psi(2S) \rightarrow \gamma \pi^+ \pi^- \pi^0$, $\gamma 3\pi^0$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|---------|----|--------|-----|------|--|
| < 2 | 90 | PEDLAR | 09 | CLE3 | $\psi(2S) \rightarrow \gamma X$ |
| < 90 | 90 | BAI | 98F | BES | $\psi(2S) \rightarrow \pi^+ \pi^- 3\gamma$ |
| < 200 | 90 | YAMADA | 77 | DASP | $e^+ e^- \rightarrow 3\gamma$ |

¹ Combining the results from $\eta \rightarrow \pi^+ \pi^- \pi^0$ and $\eta \rightarrow 3\pi^0$ decay modes.

$\Gamma(\gamma \eta \pi^+ \pi^-)/\Gamma_{\text{total}}$

Γ_{200}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|-------------|----------|--|
| $8.71 \pm 1.25 \pm 1.64$ | 418 | ABLIKIM | 06R BES2 | $\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$ |

$\Gamma(\gamma\eta(1405) \rightarrow \gamma K\bar{K}\pi)/\Gamma_{\text{total}}$ Γ_{202}/Γ

| <u>VALUE</u> (units 10^{-4}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|----------------------|-------------|---|
| <0.9 | 90 | ABLIKIM | 06R | $\psi(2S) \rightarrow \gamma K_S^0 K^+ \pi^- + \text{c.c.}$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| <1.3 | 90 | ABLIKIM | 06R | $\psi(2S) \rightarrow \gamma K^+ K^- \pi^0$ |
| <1.2 | 90 | ¹ SCHARRE | 80 | $e^+ e^-$ |

¹ Includes unknown branching fraction $\eta(1405) \rightarrow K\bar{K}\pi$. $\Gamma(\gamma\eta(1405) \rightarrow \gamma\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{203}/Γ

| <u>VALUE</u> (units 10^{-4}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|-------------|--------------------|-------------|---|
| 0.36±0.25±0.05 | 10 | ABLIKIM | 06R | $\psi(2S) \rightarrow \gamma\eta\pi^+\pi^-$ |

 $\Gamma(\gamma\eta(1405) \rightarrow \gamma f_0(980)\pi^0 \rightarrow \gamma\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{204}/Γ

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------------------|------------|--------------------|-------------|--|
| <5.0 × 10⁻⁷ | 90 | ABLIKIM | 17AJ | $\psi(2S) \rightarrow \gamma\pi^+\pi^-\pi^0$ |

 $\Gamma(\gamma\eta(1475) \rightarrow \gamma K\bar{K}\pi)/\Gamma_{\text{total}}$ Γ_{206}/Γ

| <u>VALUE</u> (units 10^{-4}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|---|
| <1.4 | 90 | ABLIKIM | 06R | $\psi(2S) \rightarrow \gamma K^+ K^- \pi^0$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| <1.5 | 90 | ABLIKIM | 06R | $\psi(2S) \rightarrow \gamma K_S^0 K^+ \pi^- + \text{c.c.}$ |

 $\Gamma(\gamma\eta(1475) \rightarrow \gamma\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{207}/Γ

| <u>VALUE</u> (units 10^{-4}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|------------|--------------------|-------------|---|
| <0.88 | 90 | ABLIKIM | 06R | $\psi(2S) \rightarrow \gamma\eta\pi^+\pi^-$ |

 $\Gamma(\gamma K^{*0} K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{208}/Γ

| <u>VALUE</u> (units 10^{-5}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|-------------|--------------------|-------------|--------------------------------|
| 37.0±6.1±7.2 | 237 | ABLIKIM | 07D | $e^+ e^- \rightarrow \psi(2S)$ |

 $\Gamma(\gamma K^{*0} \bar{K}^{*0})/\Gamma_{\text{total}}$ Γ_{209}/Γ

| <u>VALUE</u> (units 10^{-5}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|-------------|--------------------|-------------|--------------------------------|
| 24.0±4.5±5.0 | 41 | ABLIKIM | 07D | $e^+ e^- \rightarrow \psi(2S)$ |

 $\Gamma(\gamma K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{210}/Γ

| <u>VALUE</u> (units 10^{-5}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|-------------|--------------------|-------------|--------------------------------|
| 25.6±3.6±3.6 | 115 | ABLIKIM | 07D | $e^+ e^- \rightarrow \psi(2S)$ |

 $\Gamma(\gamma K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{211}/Γ

| <u>VALUE</u> (units 10^{-5}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|-------------|--------------------|-------------|--------------------------------|
| 19.1±2.7±4.3 | 132 | ABLIKIM | 07D | $e^+ e^- \rightarrow \psi(2S)$ |

 $\Gamma(\gamma K^+ K^- 2(\pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{212}/Γ

| <u>VALUE</u> (units 10^{-5}) | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------------|------------|--------------------|-------------|--------------------------------|
| <22 | 90 | ABLIKIM | 07D | $e^+ e^- \rightarrow \psi(2S)$ |

| $\Gamma(\gamma 2(K^+ K^-))/\Gamma_{\text{total}}$ | | | | | Γ_{213}/Γ |
|---|------------|--------------------|-------------|----------------|--------------------------------|
| <u>VALUE (units 10^{-5})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
| <4 | 90 | ABLIKIM | 07D | BES2 | $e^+ e^- \rightarrow \psi(2S)$ |

| $\Gamma(\gamma p\bar{p})/\Gamma_{\text{total}}$ | | | | | Γ_{214}/Γ |
|---|-------------|-------------------------------------|-------------|----------------|--|
| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
| 3.9 ± 0.5 OUR AVERAGE | | Error includes scale factor of 2.0. | | | |
| 4.18 ± 0.26 ± 0.18 | 348 | ¹ ALEXANDER | 10 | CLEO | $\psi(2S) \rightarrow \gamma p\bar{p}$ |
| 2.9 ± 0.4 ± 0.4 | 142 | ABLIKIM | 07D | BES2 | $e^+ e^- \rightarrow \psi(2S)$ |

¹ From a fit of the $p\bar{p}$ mass distribution to a combination of $\gamma f_2(1950)$, $\gamma f_2(2150)$, and $\gamma p\bar{p}$ phase space, for $M(p\bar{p}) < 2.85$ GeV, and accounting for backgrounds from $\psi(2S) \rightarrow \pi^0 p\bar{p}$ and continuum.

| $\Gamma(\gamma f_2(1950) \rightarrow \gamma p\bar{p})/\Gamma_{\text{total}}$ | | | | | Γ_{215}/Γ |
|--|-------------|------------------------|-------------|----------------|--|
| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
| 1.2 ± 0.2 ± 0.1 | 111 | ¹ ALEXANDER | 10 | CLEO | $\psi(2S) \rightarrow \gamma p\bar{p}$ |

¹ From a fit of the $p\bar{p}$ mass distribution to a combination of $\gamma f_2(1950)$, $\gamma f_2(2150)$, and $\gamma p\bar{p}$ phase space, for $M(p\bar{p}) < 2.85$ GeV, and accounting for backgrounds from $\psi(2S) \rightarrow \pi^0 p\bar{p}$ and continuum.

| $\Gamma(\gamma f_2(2150) \rightarrow \gamma p\bar{p})/\Gamma_{\text{total}}$ | | | | | Γ_{216}/Γ |
|--|-------------|------------------------|-------------|----------------|--|
| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
| 0.72 ± 0.18 ± 0.03 | 73 | ¹ ALEXANDER | 10 | CLEO | $\psi(2S) \rightarrow \gamma p\bar{p}$ |

¹ From a fit of the $p\bar{p}$ mass distribution to a combination of $\gamma f_2(1950)$, $\gamma f_2(2150)$, and $\gamma p\bar{p}$ phase space, for $M(p\bar{p}) < 2.85$ GeV, and accounting for backgrounds from $\psi(2S) \rightarrow \pi^0 p\bar{p}$ and continuum.

| $\Gamma(\gamma X(1835) \rightarrow \gamma p\bar{p})/\Gamma_{\text{total}}$ | | | | | Γ_{217}/Γ |
|--|------------|--------------------|-------------|----------------|--|
| <u>VALUE (units 10^{-6})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
| 4.57 ± 0.36 ± 1.77 | | ABLIKIM | 12D | BES3 | $J/\psi \rightarrow \gamma p\bar{p}$ |
| <4.6 | 90 | ALEXANDER | 10 | CLEO | $\psi(2S) \rightarrow \gamma p\bar{p}$ |
| <5.4 | 90 | ABLIKIM | 07D | BES | $\psi(2S) \rightarrow \gamma p\bar{p}$ |

| $\Gamma(\gamma X \rightarrow \gamma p\bar{p})/\Gamma_{\text{total}}$ | | | | | Γ_{218}/Γ |
|--|------------|--------------------|-------------|----------------|--|
| For a narrow resonance in the range $2.2 < M(X) < 2.8$ GeV. | | | | | |
| <u>VALUE (units 10^{-6})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
| <2 | 90 | ALEXANDER | 10 | CLEO | $\psi(2S) \rightarrow \gamma p\bar{p}$ |

| $\Gamma(\gamma p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$ | | | | | Γ_{219}/Γ |
|---|-------------|--------------------|-------------|----------------|--------------------------------|
| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
| 2.8 ± 1.2 ± 0.7 | 17 | ABLIKIM | 07D | BES2 | $e^+ e^- \rightarrow \psi(2S)$ |

$\Gamma(\gamma\gamma J/\psi)/\Gamma_{\text{total}}$ Γ_{221}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|--------------------------------|
| $3.1 \pm 0.6 \pm 0.8$ | 1.1k | ABLIKIM | 120 BES3 | $e^+ e^- \rightarrow \psi(2S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|---------------|------|----------------------|----------|--|
| 3.2 \pm 0.6 | 1.1k | ¹ ABLIKIM | 17N BES3 | $\psi(2S) \rightarrow \gamma\gamma J/\psi$ |
|---------------|------|----------------------|----------|--|

¹ Uses $B(J/\psi \rightarrow e^+ e^-) = (5.971 \pm 0.032)\%$ and $B(J/\psi \rightarrow \mu^+ \mu^-) = (5.961 \pm 0.033)\%$. No systematic error estimation.

 $\Gamma(e^+ e^- \eta')/\Gamma_{\text{total}}$ Γ_{222}/Γ

| <u>VALUE (units 10^{-6})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|--|
| 1.90 ± 0.26 OUR AVERAGE | | | | |
| 1.99 \pm 0.33 \pm 0.12 | 57 | ABLIKIM | 18Z BES3 | $\psi(2S) \rightarrow \eta' e^+ e^-$, $\eta' \rightarrow \gamma\pi^+\pi^-$ |
| 1.79 \pm 0.38 \pm 0.11 | 20 | ABLIKIM | 18Z BES3 | $\psi(2S) \rightarrow \eta' e^+ e^-$, $\eta' \rightarrow \eta\pi^+\pi^-$ |

 $\Gamma(e^+ e^- \eta_c(1S))/\Gamma_{\text{total}}$ Γ_{223}/Γ

| <u>VALUE (units 10^{-5})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|-------------|----------------------|-------------|--------------------------------|
| $3.77 \pm 0.40 \pm 0.18$ | 3k | ¹ ABLIKIM | 22AX BES3 | $e^+ e^- \rightarrow \psi(2S)$ |

¹ From a fit to the recoil mass distribution of $e^+ e^-$ with inclusive $\eta_c(1S)$ decays.

 $\Gamma(e^+ e^- \chi_{c0}(1P))/\Gamma_{\text{total}}$ Γ_{224}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|----------------------|-------------|--|
| $10.6 \pm 2.4 \pm 0.7$ | 48 | ¹ ABLIKIM | 17I BES3 | $\psi(2S) \rightarrow e^+ e^- \gamma J/\psi$ |

¹ ABLIKIM 17I reports $(11.7 \pm 2.5 \pm 1.0) \times 10^{-4}$ from a measurement of $[\Gamma(\psi(2S) \rightarrow e^+ e^- \chi_{c0}(1P))/\Gamma_{\text{total}}] \times [B(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S))]$ assuming $B(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S)) = (1.27 \pm 0.06) \times 10^{-2}$, which we rescale to our best value $B(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S)) = (1.41 \pm 0.09) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(e^+ e^- \chi_{c0}(1P))/\Gamma(\gamma \chi_{c0}(1P))$ $\Gamma_{224}/\Gamma_{179}$

| <u>VALUE (units 10^{-3})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|----------------------|-------------|--|
| $9.4 \pm 1.9 \pm 0.6$ | 48 | ¹ ABLIKIM | 17I BES3 | $\psi(2S) \rightarrow e^+ e^- \gamma J/\psi$ |

¹ Uses $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) \times B(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S)) = (15.8 \pm 0.3 \pm 0.6) \times 10^{-4}$ from ABLIKIM 17N and accounts for common systematic errors.

 $\Gamma(e^+ e^- \chi_{c1}(1P))/\Gamma_{\text{total}}$ Γ_{225}/Γ

| <u>VALUE (units 10^{-4})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|----------------------|-------------|--|
| $8.5 \pm 0.6 \pm 0.3$ | 873 | ¹ ABLIKIM | 17I BES3 | $\psi(2S) \rightarrow e^+ e^- \gamma J/\psi$ |

¹ ABLIKIM 17I reports $(8.6 \pm 0.3 \pm 0.6) \times 10^{-4}$ from a measurement of $[\Gamma(\psi(2S) \rightarrow e^+ e^- \chi_{c1}(1P))/\Gamma_{\text{total}}] \times [B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S))]$ assuming $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (33.9 \pm 1.2) \times 10^{-2}$, which we rescale to our best value $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (34.3 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(e^+ e^- \chi_{c1}(1P))/\Gamma(\gamma \chi_{c1}(1P))$ $\Gamma_{225}/\Gamma_{180}$

| <u>VALUE</u> (units 10^{-3}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|----------------------|-------------|---|
| $8.3 \pm 0.3 \pm 0.4$ | 873 | ¹ ABLIKIM | 17I | BES3 $\psi(2S) \rightarrow e^+ e^- \gamma J/\psi$ |

¹ Uses $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) \times B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (351.8 \pm 1.0 \pm 12.0) \times 10^{-4}$ from ABLIKIM 17N and accounts for common systematic errors.

 $\Gamma(e^+ e^- \chi_{c2}(1P))/\Gamma_{\text{total}}$ Γ_{226}/Γ

| <u>VALUE</u> (units 10^{-4}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|----------------------|-------------|---|
| $6.8 \pm 0.7 \pm 0.3$ | 227 | ¹ ABLIKIM | 17I | BES3 $\psi(2S) \rightarrow e^+ e^- \gamma J/\psi$ |

¹ ABLIKIM 17I reports $(6.9 \pm 0.5 \pm 0.6) \times 10^{-4}$ from a measurement of $[\Gamma(\psi(2S) \rightarrow e^+ e^- \chi_{c2}(1P))/\Gamma_{\text{total}}] \times [B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S))]$ assuming $B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) = (19.2 \pm 0.7) \times 10^{-2}$, which we rescale to our best value $B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) = (19.5 \pm 0.8) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(e^+ e^- \chi_{c2}(1P))/\Gamma(\gamma \chi_{c2}(1P))$ $\Gamma_{226}/\Gamma_{181}$

| <u>VALUE</u> (units 10^{-3}) | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|----------------------|-------------|---|
| $6.6 \pm 0.5 \pm 0.4$ | 227 | ¹ ABLIKIM | 17I | BES3 $\psi(2S) \rightarrow e^+ e^- \gamma J/\psi$ |

¹ Uses $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) \times B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) = (199.6 \pm 0.8 \pm 7.0) \times 10^{-4}$ from ABLIKIM 17N and accounts for common systematic errors.

WEAK DECAYS $\Gamma(D^0 e^+ e^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{227}/Γ

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|------------|----------------------|-------------|-------------------------------------|
| $<1.4 \times 10^{-7}$ | 90 | ¹ ABLIKIM | 17AF | BES3 $e^+ e^- \rightarrow \psi(2S)$ |

¹ Using D^0 decays to $K^- \pi^+$, $K^- \pi^+ \pi^0$, and $K^- \pi^+ \pi^+ \pi^-$.

 $\Gamma(\Lambda_c^+ \bar{\Sigma}^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{228}/Γ

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|------------|----------------------|-------------|-------------------------------------|
| $<1.4 \times 10^{-5}$ | 90 | ¹ ABLIKIM | 23 | BES3 $e^+ e^- \rightarrow \psi(2S)$ |

¹ Using $\Lambda_c^+ \rightarrow p K^- \pi^+$ and $\bar{\Sigma}^- \rightarrow \bar{p} \pi^0$.

OTHER DECAYS $\Gamma(\text{invisible})/\Gamma(e^+ e^-)$ Γ_{229}/Γ_7

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------------|------------|--------------------|-------------|---------------------------------------|
| <2.0 | 90 | LEES | 13I | BABR $B \rightarrow K^{(*)} \psi(2S)$ |

 $\psi(2S)$ CROSS-PARTICLE BRANCHING RATIOS

For measurements involving $B(\psi(2S) \rightarrow \gamma \chi_{cJ}(1P)) \times B(\chi_{cJ}(1P) \rightarrow X)$ see the corresponding entries in the $\chi_{cJ}(1P)$ sections.

MULTIPOLE AMPLITUDE RATIOS IN RADIATIVE DECAYS

$\psi(2S) \rightarrow \gamma\chi_{cJ}(1P)$ and $\chi_{cJ} \rightarrow \gamma J/\psi(1S)$

$a_2(\chi_{c1})/a_2(\chi_{c2})$ Magnetic quadrupole transition amplitude ratio

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|----------------------|----------|---|
| 63 \pm 7 OUR AVERAGE | | | | |
| 61.7 \pm 8.3 | 253k | ¹ ABLIKIM | 17N BES3 | $\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$ |
| 67 \pm 19 -13 | 59k | ² ARTUSO | 09 CLEO | $\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$ |

¹ Statistical and systematic errors combined.

² Statistical and systematic errors combined. Using values from fits with floating $M2$ amplitudes $a_2(\chi_{c1})$, $a_2(\chi_{c2})$, $b_2(\chi_{c1})$, $b_2(\chi_{c2})$ and fixed $E3$ amplitudes of $a_3(\chi_{c2}) = b_3(\chi_{c2}) = 0$. Not independent of values for $a_2(\chi_{c1}(1P))$ and $a_2(\chi_{c2}(1P))$ from ARTUSO 09.

$b_2(\chi_{c2})/b_2(\chi_{c1})$ Magnetic quadrupole transition amplitude ratio

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|----------------------|----------|---|
| 60 \pm 31 OUR AVERAGE | | | | |
| 74 \pm 40 | 253k | ¹ ABLIKIM | 17N BES3 | $\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$ |
| 37 \pm 53 -47 | 59k | ² ARTUSO | 09 CLEO | $\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$ |

¹ Statistical and systematic errors combined. Derived from the reported measurement of $b_2(\chi_{c1})/b_2(\chi_{c2}) = 1.35 \pm 0.72$.

² Statistical and systematic errors combined. Using values from fits with floating $M2$ amplitudes $a_2(\chi_{c1})$, $a_2(\chi_{c2})$, $b_2(\chi_{c1})$, $b_2(\chi_{c2})$ and fixed $E3$ amplitudes of $a_3(\chi_{c2}) = b_3(\chi_{c2}) = 0$. Not independent of values for $b_2(\chi_{c1}(1P))$ and $b_2(\chi_{c2}(1P))$ from ARTUSO 09.

$\psi(2S)$ REFERENCES

| | | | |
|---------|---------------------|----------------------------|------------------|
| AAIJ | 23AP JHEP 2307 084 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| ABLIKIM | 23 CP C47 013002 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 23BA PR D108 052001 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 23BE PR D108 092011 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 23BV PR D108 112014 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| LEES | 23 PR D107 072001 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| LIAO | 23 PR D107 112007 | L. Liao <i>et al.</i> | |
| ABLIKIM | 22AP PR D106 072006 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 22AQ PR D106 072007 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 22AV JHEP 2212 016 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 22AX PR D106 112002 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 22AY PR D106 112007 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 22AZ PR D106 112011 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| PDG | 22 PTEP 2022 083C01 | R.L. Workman <i>et al.</i> | (PDG Collab.) |
| ABLIKIM | 21AL PR D104 092003 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 21AO PR D104 092012 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 21AT JHEP 2111 226 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 21E PRL 126 092002 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 21L PR D103 112004 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 21S PL B820 136576 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 21Z PRL 127 082002 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| LEES | 21 PR D103 092001 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| LEES | 21C PR D104 112004 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| ABLIKIM | 20F PR D101 032008 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| PDG | 20 PTEP 2020 083C01 | P.A. Zyla <i>et al.</i> | (PDG Collab.) |
| ABLIKIM | 19AO PR D99 112010 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 19AT PR D100 051101 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 19AU PR D100 052010 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 19BA PR D100 092003 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 19I PR D99 012014 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |

| | | | | |
|-----------------------------|---------------|-----------------------------------|------------------------------|---------------------|
| ABLIKIM | 19N | PR D99 032006 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 18Q | PR D97 091102 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 18T | PR D98 032006 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 18Z | PL B783 452 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ANASHIN | 18 | PL B781 174 | V.V. Anashin <i>et al.</i> | (KEDR Collab.) |
| LEES | 18E | PR D98 112015 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| ABLIKIM | 17AF | PR D96 111101 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 17AJ | PR D96 112008 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 17AK | PR D96 112012 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 17E | PL B770 217 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 17I | PRL 118 221802 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 17L | PR D95 052003 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 17N | PR D95 072004 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 17U | PR D96 032001 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 17X | PR D96 052003 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| DOBBS | 17 | PR D96 092004 | S. Dobbs <i>et al.</i> | (NWES, WAYN) |
| LEES | 17A | PR D95 052001 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| AAIJ | 16Y | JHEP 1605 132 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| ABLIKIM | 16L | PR D93 072003 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 15I | PR D91 092006 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 15V | PL B749 414 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ANASHIN | 15 | PL B749 50 | V.V. Anashin <i>et al.</i> | (KEDR Collab.) |
| DOBBS | 15 | PR D91 052006 | S. Dobbs <i>et al.</i> | (NWES) |
| LEES | 15J | PR D92 072008 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| ABLIKIM | 14G | PR D89 112006 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| DOBBS | 14 | PL B739 90 | S. Dobbs <i>et al.</i> | (NWES, WAYN) |
| ABLIKIM | 13A | PRL 110 022001 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 13D | PR D87 012007 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 13F | PR D87 052007 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 13M | PR D87 092006 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 13R | PR D88 032007 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 13S | PR D88 032010 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 13W | PR D88 112007 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| LEES | 13I | PR D87 112005 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| LEES | 13O | PR D87 092005 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| LEES | 13Q | PR D88 032013 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| LEES | 13Y | PR D88 072009 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| AAIJ | 12H | EPJ C72 1972 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| ABLIKIM | 12D | PRL 108 112003 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 12G | PRL 109 042003 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 12H | PL B710 594 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 12L | PR D86 072011 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 12M | PR D86 092008 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 12O | PRL 109 172002 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 12Q | CP C36 1040 | M. Ablikim <i>et al.</i> | (BES II Collab.) |
| ANASHIN | 12 | PL B711 280 | V.V. Anashin <i>et al.</i> | (KEDR Collab.) |
| LEES | 12E | PR D85 112009 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| LEES | 12F | PR D86 012008 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| METREVELI | 12 | PR D85 092007 | Z. Metreveli <i>et al.</i> | (NWES, FLOR, WAYN+) |
| GE | 11 | PR D84 032008 | J.Y. Ge <i>et al.</i> | (CLEO Collab.) |
| ABLIKIM | 10B | PRL 104 132002 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 10F | PRL 105 261801 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ALEXANDER | 10 | PR D82 092002 | J.P. Alexander <i>et al.</i> | (CLEO Collab.) |
| CRONIN-HEN... ¹⁰ | PR D81 052002 | D. Cronin-Hennessey <i>et al.</i> | (CLEO Collab.) | |
| ADAMS | 09 | PR D80 051106 | G.S. Adams <i>et al.</i> | (CLEO Collab.) |
| ARTUSO | 09 | PR D80 112003 | M. Artuso <i>et al.</i> | (CLEO Collab.) |
| LIBBY | 09 | PR D80 072002 | J. Libby <i>et al.</i> | (CLEO Collab.) |
| MITCHELL | 09 | PRL 102 011801 | R.E. Mitchell <i>et al.</i> | (CLEO Collab.) |
| PEDLAR | 09 | PR D79 111101 | T.K. Pedlar <i>et al.</i> | (CLEO Collab.) |
| ABLIKIM | 08B | PL B659 74 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 08C | PL B659 789 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| DOBBS | 08A | PRL 101 182003 | S. Dobbs <i>et al.</i> | (CLEO Collab.) |
| MENDEZ | 08 | PR D78 011102 | H. Mendez <i>et al.</i> | (CLEO Collab.) |
| PDG | 08 | PL B667 1 | C. Amsler <i>et al.</i> | (PDG Collab.) |
| ABLIKIM | 07C | PL B648 149 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 07D | PRL 99 011802 | M. Ablikim <i>et al.</i> | (BES II Collab.) |
| ABLIKIM | 07H | PR D76 092003 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ANASHIN | 07 | JETPL 85 347 | V.V. Anashin <i>et al.</i> | (KEDR Collab.) |

Translated from ZETFP 85 429.

| | | | | |
|------------|------|-----------------------------|-------------------------------|-------------------------|
| ANDREOTTI | 07 | PL B654 74 | M. Andreotti <i>et al.</i> | (Fermilab E835 Collab.) |
| AUBERT | 07AK | PR D76 012008 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 07AU | PR D76 092005 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| Also | | PR D77 119902E (errat.) | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 07BD | PR D76 092006 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| PDG | 07 | Unofficial 2007 WWW edition | | (PDG Collab.) |
| PEDLAR | 07 | PR D75 011102 | T.K. Pedlar <i>et al.</i> | (CLEO Collab.) |
| ABLIKIM | 06G | PR D73 052004 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 06I | PR D74 012004 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 06L | PRL 97 121801 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 06R | PR D74 072001 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 06W | PR D74 112003 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ADAM | 06 | PRL 96 082004 | N.E. Adam <i>et al.</i> | (CLEO Collab.) |
| AUBERT | 06B | PR D73 012005 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 06D | PR D73 052003 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,BE | 06D | PR D74 091103 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| DOBBS | 06A | PR D74 011105 | S. Dobbs <i>et al.</i> | (CLEO Collab.) |
| ABLIKIM | 05E | PR D71 072006 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 05H | PR D72 012002 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 05I | PL B614 37 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 05J | PL B619 247 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 05O | PL B630 21 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ADAM | 05 | PRL 94 012005 | N.E. Adam <i>et al.</i> | (CLEO Collab.) |
| ADAM | 05A | PRL 94 232002 | N.E. Adam <i>et al.</i> | (CLEO Collab.) |
| ANDREOTTI | 05 | PR D71 032006 | M. Andreotti <i>et al.</i> | (FNAL E835 Collab.) |
| AUBERT | 05D | PR D71 052001 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| BRIERE | 05 | PRL 95 062001 | R.A. Briere <i>et al.</i> | (CLEO Collab.) |
| PEDLAR | 05 | PR D72 051108 | T.K. Pedlar <i>et al.</i> | (CLEO Collab.) |
| ROSNER | 05 | PRL 95 102003 | J.L. Rosner <i>et al.</i> | (CLEO Collab.) |
| ABLIKIM | 04B | PR D70 012003 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 04K | PR D70 112003 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 04L | PR D70 112007 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ATHAR | 04 | PR D70 112002 | S.B. Athar <i>et al.</i> | (CLEO Collab.) |
| BAI | 04B | PRL 92 052001 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| BAI | 04C | PR D69 072001 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| BAI | 04D | PL B589 7 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| BAI | 04G | PR D70 012004 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| BAI | 04I | PR D70 012006 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| PDG | 04 | PL B592 1 | S. Eidelman <i>et al.</i> | (PDG Collab.) |
| SETH | 04 | PR D69 097503 | K.K. Seth | |
| AULCHENKO | 03 | PL B573 63 | V.M. Aulchenko <i>et al.</i> | (KEDR Collab.) |
| BAI | 03B | PR D67 052002 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| BAI | 03C | PR D67 032004 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| AUBERT | 02B | PR D65 031101 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| BAI | 02 | PR D65 052004 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| BAI | 02B | PL B550 24 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| BAI | 02C | PRL 88 101802 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| PDG | 02 | PR D66 010001 | K. Hagiwara <i>et al.</i> | (PDG Collab.) |
| BAI | 01 | PR D63 032002 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| AMBROGIANI | 00A | PR D62 032004 | M. Ambrogiani <i>et al.</i> | (FNAL E835 Collab.) |
| ARTAMONOV | 00 | PL B474 427 | A.S. Artamonov <i>et al.</i> | |
| BAI | 00 | PRL 84 594 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| BAI | 99C | PRL 83 1918 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| BAI | 98E | PR D57 3854 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| BAI | 98F | PR D58 097101 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| BAI | 98J | PRL 81 5080 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| ARMSTRONG | 97 | PR D55 1153 | T.A. Armstrong <i>et al.</i> | (E760 Collab.) |
| GРИBUSHIN | 96 | PR D53 4723 | A. Gribushin <i>et al.</i> | (E672 and E706 Collab.) |
| ARMSTRONG | 93B | PR D47 772 | T.A. Armstrong <i>et al.</i> | (FNAL E760 Collab.) |
| ALEXANDER | 89 | NP B320 45 | J.P. Alexander <i>et al.</i> | (LBL, MICH, SLAC) |
| COHEN | 87 | RMP 59 1121 | E.R. Cohen, B.N. Taylor | (RISC, NBS) |
| GAISER | 86 | PR D34 711 | J. Gaiser <i>et al.</i> | (Crystal Ball Collab.) |
| KURAEV | 85 | SJNP 41 466 | E.A. Kuraev, V.S. Fadin | (NOVO) |
| | | Translated from YAF 41 733. | | |
| FRANKLIN | 83 | PRL 51 963 | M.E.B. Franklin <i>et al.</i> | (LBL, SLAC) |
| EDWARDS | 82C | PRL 48 70 | C. Edwards <i>et al.</i> | (CIT, HARV, PRIN+) |
| LEMOIGNE | 82 | PL 113B 509 | Y. Lemoigne <i>et al.</i> | (SACL, LOIC, SHMP+) |
| HIMEL | 80 | PRL 44 920 | T. Himel <i>et al.</i> | (LBL, SLAC) |
| OREGLIA | 80 | PRL 45 959 | M.J. Oreglia <i>et al.</i> | (SLAC, CIT, HARV+) |
| SCHARRE | 80 | PL 97B 329 | D.L. Scharre <i>et al.</i> | (SLAC, LBL) |

| | | | | |
|-------------|-----|--------------------------------|--|---------------------|
| ZHOLENTZ | 80 | PL 96B 214 Also SJNP 34 814 | A.A. Zholents <i>et al.</i> A.A. Zholents <i>et al.</i> | (NOVO) (NOVO) |
| | | Translated from YAF 34 1471. | | |
| BRANDELIK | 79B | NP B160 426 | R. Brandelik <i>et al.</i> | (DASP Collab.) |
| BRANDELIK | 79C | ZPHY C1 233 | R. Brandelik <i>et al.</i> | (DASP Collab.) |
| BARTEL | 78B | PL 79B 492 | W. Bartel <i>et al.</i> | (DESY, HEIDP) |
| TANENBAUM | 78 | PR D17 1731 | W.M. Tanenbaum <i>et al.</i> | (SLAC, LBL) |
| BIDDICK | 77 | PRL 38 1324 | C.J. Biddick <i>et al.</i> | (UCSD, UMD, PAVI+) |
| BRAUNSCH... | 77 | PL 67B 249 | W. Braunschweig <i>et al.</i> | (DASP Collab.) |
| BURMESTER | 77 | PL 66B 395 | J. Burmester <i>et al.</i> | (DESY, HAMB, SIEG+) |
| FELDMAN | 77 | PRPL 33C 285 | G.J. Feldman, M.L. Perl | (LBL, SLAC) |
| YAMADA | 77 | Hamburg Conf. 69 | S. Yamada | (DASP Collab.) |
| BARTEL | 76 | PL 64B 483 | W. Bartel <i>et al.</i> | (DESY, HEIDP) |
| TANENBAUM | 76 | PRL 36 402 | W.M. Tanenbaum <i>et al.</i> | (SLAC, LBL) IG |
| WHITAKER | 76 | PRL 37 1596 | J.S. Whitaker <i>et al.</i> | (SLAC, LBL) |
| ABRAMS | 75 | Stanford Symp. 25 | G.S. Abrams | (LBL) |
| ABRAMS | 75B | PRL 34 1181 | G.S. Abrams <i>et al.</i> | (LBL, SLAC) |
| BOYARSKI | 75C | Palermo Conf. 54 | A.M. Boyarski <i>et al.</i> | (SLAC, LBL) |
| HILGER | 75 | PRL 35 625 | E. Hilger <i>et al.</i> | (STAN, PENN) |
| LIBERMAN | 75 | Stanford Symp. 55 | A.D. Liberman | (STAN) |
| LUTH | 75 | PRL 35 1124 | V. Luth <i>et al.</i> | (SLAC, LBL) JPC |
| WIJK | 75 | Stanford Symp. 69 | B.H. Wiik | (DESY) |