

$\psi(4040)$

$I^G(J^{PC}) = 0^-(1^{--})$

$\psi(4040)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
4039.6 ± 4.3	1 ABLIKIM	08D BES2	$e^+ e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
4034 ± 6	2 MO	10 RVUE	$e^+ e^- \rightarrow$ hadrons
4037 ± 2	3 SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
4040 ± 1	4 SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
4040 ± 10	BRANDELIK	78C DASP	$e^+ e^-$

¹ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (130 \pm 46)^\circ$.

² Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects.

³ From a fit to Crystal Ball (OSTERHELD 86) data.

⁴ From a fit to BES (BAI 02C) data.

$\psi(4040)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
84.5 ± 12.3	5 ABLIKIM	08D BES2	$e^+ e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
87 ± 11	6 MO	10 RVUE	$e^+ e^- \rightarrow$ hadrons
85 ± 10	7 SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
89 ± 6	8 SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
52 ± 10	BRANDELIK	78C DASP	$e^+ e^-$

⁵ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (130 \pm 46)^\circ$.

⁶ Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects.

⁷ From a fit to Crystal Ball (OSTERHELD 86) data.

⁸ From a fit to BES (BAI 02C) data.

$\psi(4040)$ DECAY MODES

Due to the complexity of the $c\bar{c}$ threshold region, in this listing, “seen” (“not seen”) means that a cross section for the mode in question has been measured at effective \sqrt{s} near this particle’s central mass value, more

(less) than 2σ above zero, without regard to any peaking behavior in \sqrt{s} or absence thereof. See mode listing(s) for details and references.

Mode	Fraction (Γ_i/Γ)	Confidence level
$\Gamma_1 e^+ e^-$	$(1.02 \pm 0.17) \times 10^{-5}$	
$\Gamma_2 D \bar{D}$	seen	
$\Gamma_3 D^0 \bar{D}^0$	seen	
$\Gamma_4 D^+ D^-$	seen	
$\Gamma_5 D^* \bar{D} + \text{c.c.}$	seen	
$\Gamma_6 D^*(2007)^0 \bar{D}^0 + \text{c.c.}$	seen	
$\Gamma_7 D^*(2010)^+ D^- + \text{c.c.}$	seen	
$\Gamma_8 D^* \bar{D}^*$	seen	
$\Gamma_9 D^*(2007)^0 \bar{D}^*(2007)^0$	seen	
$\Gamma_{10} D^*(2010)^+ D^*(2010)^-$	seen	
$\Gamma_{11} D \bar{D} \pi (\text{excl. } D^* \bar{D})$	not seen	
$\Gamma_{12} D^0 D^- \pi^+ + \text{c.c.} (\text{excl. } D^*(2010)^+ D^- + \text{c.c.})$	not seen	
$\Gamma_{13} D \bar{D}^* \pi (\text{excl. } D^* \bar{D}^*)$	not seen	
$\Gamma_{14} D^0 \bar{D}^{*-} \pi^+ + \text{c.c.} (\text{excl. } D^*(2010)^+ D^*(2010)^-)$	seen	
$\Gamma_{15} D_s^+ D_s^-$	seen	
$\Gamma_{16} \pi^+ \pi^+ \pi^- \pi^- \pi^0$	seen	
$\Gamma_{17} J/\psi(1S) \text{hadrons}$	seen	
$\Gamma_{18} J/\psi \pi^+ \pi^-$	$< 4 \times 10^{-3}$	90%
$\Gamma_{19} J/\psi \pi^0 \pi^0$	$< 2 \times 10^{-3}$	90%
$\Gamma_{20} J/\psi \eta$	$(5.2 \pm 0.7) \times 10^{-3}$	
$\Gamma_{21} J/\psi \pi^0$	$< 2.8 \times 10^{-4}$	90%
$\Gamma_{22} J/\psi \pi^+ \pi^- \pi^0$	$< 2 \times 10^{-3}$	90%
$\Gamma_{23} \chi_{c1} \gamma$	$< 3.4 \times 10^{-3}$	90%
$\Gamma_{24} \chi_{c2} \gamma$	$< 5 \times 10^{-3}$	90%
$\Gamma_{25} \chi_{c1} \pi^+ \pi^- \pi^0$	$< 1.1 \%$	90%
$\Gamma_{26} \chi_{c2} \pi^+ \pi^- \pi^0$	$< 3.2 \%$	90%
$\Gamma_{27} h_c(1P) \pi^+ \pi^-$	$< 3 \times 10^{-3}$	90%
$\Gamma_{28} \phi \pi^+ \pi^-$	$< 3 \times 10^{-3}$	90%
$\Gamma_{29} \Lambda \bar{\Lambda} \pi^+ \pi^-$	$< 2.9 \times 10^{-4}$	90%
$\Gamma_{30} \Lambda \bar{\Lambda} \pi^0$	$< 9 \times 10^{-5}$	90%
$\Gamma_{31} \Lambda \bar{\Lambda} \eta$	$< 3.0 \times 10^{-4}$	90%
$\Gamma_{32} \Lambda \bar{\Lambda}$	$< 6 \times 10^{-6}$	90%
$\Gamma_{33} \Sigma^+ \bar{\Sigma}^-$	$< 1.3 \times 10^{-4}$	90%
$\Gamma_{34} \Sigma^0 \bar{\Sigma}^0$	$< 7 \times 10^{-5}$	90%
$\Gamma_{35} \Xi^+ \bar{\Xi}^-$	$< 1.6 \times 10^{-4}$	90%

Γ_{36}	$\Xi^0 \Xi^0$	< 1.8	$\times 10^{-4}$	90%
Γ_{37}	$\Xi^- \Xi^+$	< 6	$\times 10^{-5}$	90%
Γ_{38}	$\mu^+ \mu^-$	(9 \pm 6)	$\times 10^{-6}$	

$\psi(4040)$ PARTIAL WIDTHS

$\Gamma(e^+ e^-)$

VALUE (keV)

0.86 \pm 0.07 OUR ESTIMATE

0.83 \pm 0.20

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.6 to 1.4

0.88 \pm 0.11

0.91 \pm 0.13

0.75 \pm 0.15

DOCUMENT ID

TECN

COMMENT

⁹ ABLIKIM 08D BES2 $e^+ e^- \rightarrow$ hadrons

¹⁰ MO 10 RVUE $e^+ e^- \rightarrow$ hadrons

¹¹ SETH 05A RVUE $e^+ e^- \rightarrow$ hadrons

¹² SETH 05A RVUE $e^+ e^- \rightarrow$ hadrons

BRANDELIK 78C DASP $e^+ e^-$

Γ_1

⁹ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (130 \pm 46)^\circ$.

¹⁰ Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects. Four sets of solutions are obtained with the same fit quality, mass and total width, but with different $e^+ e^-$ partial widths. We quote only the range of values.

¹¹ From a fit to Crystal Ball (OSTERHELD 86) data.

¹² From a fit to BES (BAI 02C) data.

$\Gamma(\mu^+ \mu^-)$

VALUE (keV)

0.73 \pm 0.48 \pm 0.12

DOCUMENT ID

TECN

COMMENT

^{13,14} ABLIKIM 20AG BES3 $e^+ e^- \rightarrow \mu^+ \mu^-$

¹³ From a fit to the $e^+ e^- \rightarrow \mu^+ \mu^-$ cross section between 3.8 and 4.6 GeV to the coherent sum of four resonant amplitudes assuming $\Gamma(\mu^+ \mu^-) = \Gamma(e^+ e^-)$.

¹⁴ From solution 1 of 8 with equal fit quality. Other solutions range from $0.58 \pm 0.52 \pm 0.10$ to $0.80 \pm 0.48 \pm 0.13$ keV.

Γ_{38}

$\psi(4040) \Gamma(i) \times \Gamma(e^+ e^-)/\Gamma(\text{total})$

$\Gamma(J/\psi \eta) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$

$\Gamma_{20} \Gamma_1 / \Gamma$

VALUE (eV)

DOCUMENT ID

TECN

COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.5 \pm 0.3

¹⁵ ABLIKIM 200 BES3 $e^+ e^- \rightarrow \eta J/\psi$

1.4 \pm 0.3

¹⁶ ABLIKIM 200 BES3 $e^+ e^- \rightarrow \eta J/\psi$

7.0 \pm 0.6

¹⁷ ABLIKIM 200 BES3 $e^+ e^- \rightarrow \eta J/\psi$

¹⁵ Solution 1 of three equivalent fit solutions using three resonant structures.

¹⁶ Solution 2 of three equivalent fit solutions using three resonant structures.

¹⁷ Solution 3 of three equivalent fit solutions using three resonant structures.

$\Gamma(\chi_{c1}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{23}\Gamma_1/\Gamma$			
<u>VALUE (eV)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2.9	90	18 HAN	15	BELL $10.58 e^+e^- \rightarrow \chi_{c1}\gamma$

18 Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.

$\Gamma(\chi_{c2}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{24}\Gamma_1/\Gamma$			
<u>VALUE (eV)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<4.6	90	19 HAN	15	BELL $10.58 e^+e^- \rightarrow \chi_{c2}\gamma$

19 Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.

$\Gamma(\Lambda\bar{\Lambda}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{32}\Gamma_1/\Gamma$			
<u>VALUE (eV)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<5.5 × 10 ⁻³	90	20 ABLIKIM	21AS BES3	$e^+e^- \rightarrow \psi(4040)$

20 From a measurement of the $e^+e^- \rightarrow \Lambda\bar{\Lambda}$ cross section between 3.5 and 4.6 GeV.

$\Gamma(\Xi^-\bar{\Xi}^+) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{37}\Gamma_1/\Gamma$			
<u>VALUE (eV)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.0519	90	21 ABLIKIM	23BK BES3	$e^+e^- \rightarrow \psi(4040)$

21 From a fit to $e^+e^- \rightarrow \Xi^-\bar{\Xi}^+$ cross sections.

$\psi(4040) \Gamma(i) \times \Gamma(e^+e^-)/\Gamma^2(\text{total})$

$\Gamma(J/\psi\eta)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{20}/\Gamma \times \Gamma_1/\Gamma$		
<u>VALUE (units 10⁻⁸)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
5.1 ± 1.4 ± 1.5	22 WANG	13B	BELL $e^+e^- \rightarrow J/\psi\eta\gamma$
12.8 ± 2.1 ± 1.9	23 WANG	13B	BELL $e^+e^- \rightarrow J/\psi\eta\gamma$

22 Solution I of two equivalent solutions in a fit using two interfering resonances. Mass and width fixed at 4039 MeV and 80 MeV, respectively.

23 Solution II of two equivalent solutions in a fit using two interfering resonances. Mass and width fixed at 4039 MeV and 80 MeV, respectively.

$\psi(4040)$ BRANCHING RATIOS

$\Gamma(e^+e^-)/\Gamma_{\text{total}}$	Γ_1/Γ		
<u>VALUE (units 10⁻⁵)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
~ 1.0	FELDMAN	77	MRK1 e^+e^-

$\Gamma(D^0\bar{D}^0)/\Gamma_{\text{total}}$	Γ_3/Γ		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
seen	AUBERT	09M	BABR $e^+e^- \rightarrow D^0\bar{D}^0\gamma$
seen	CRONIN-HEN..09	CLEO	$e^+e^- \rightarrow D^0\bar{D}^0$
seen	PAKHLOVA	08	BELL $e^+e^- \rightarrow D^0\bar{D}^0\gamma$

$\Gamma(D^+ D^-)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	AUBERT 09M	BABR	$e^+ e^- \rightarrow D^+ D^- \gamma$
seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^+ D^-$
seen	PAKHLOVA 08	BELL	$e^+ e^- \rightarrow D^+ D^- \gamma$

Γ_4/Γ

$\Gamma(D\bar{D})/\Gamma(D^*\bar{D} + \text{c.c.})$

VALUE	DOCUMENT ID	TECN	COMMENT
0.24±0.05±0.12	AUBERT 09M	BABR	$e^+ e^- \rightarrow \gamma D^{(*)}\bar{D}$

Γ_2/Γ_5

$\Gamma(D^0\bar{D}^0)/\Gamma(D^*(2007)^0\bar{D}^0 + \text{c.c.})$

VALUE	DOCUMENT ID	TECN	COMMENT
0.05±0.03	²⁴ GOLDHABER 77	MRK1	$e^+ e^-$

Γ_3/Γ_6

²⁴ Phase-space factor (p^3) explicitly removed.

$\Gamma(D^*(2007)^0\bar{D}^0 + \text{c.c.})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	AUBERT 09M	BABR	$e^+ e^- \rightarrow D^{*0}\bar{D}^0 \gamma$
seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^{*0}\bar{D}^0$

Γ_6/Γ

$\Gamma(D^*(2010)^+ D^- + \text{c.c.})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	²⁵ ZHUKOVA 18	BELL	$e^+ e^- \rightarrow D^{*+} D^- \gamma$
seen	AUBERT 09M	BABR	$e^+ e^- \rightarrow D^{*+} D^- \gamma$
seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^{*+} D^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
seen	PAKHLOVA 07	BELL	$e^+ e^- \rightarrow D^{*+} D^- \gamma$

Γ_7/Γ

²⁵ Supersedes PAKHLOVA 07.

$\Gamma(D^*(2010)^+ D^- + \text{c.c.})/\Gamma(D^*(2007)^0\bar{D}^0 + \text{c.c.})$

VALUE	DOCUMENT ID	TECN	COMMENT
0.95±0.09±0.10	AUBERT 09M	BABR	$e^+ e^- \rightarrow \gamma D^*\bar{D}$

Γ_7/Γ_6

$\Gamma(D^*\bar{D}^*)/\Gamma(D^*\bar{D} + \text{c.c.})$

VALUE	DOCUMENT ID	TECN	COMMENT
0.18±0.14±0.03	AUBERT 09M	BABR	$e^+ e^- \rightarrow \gamma D^{(*)}\bar{D}^{(*)}$

Γ_8/Γ_5

$\Gamma(D^*(2007)^0\bar{D}^*(2007)^0)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	AUBERT 09M	BABR	$e^+ e^- \rightarrow D^{*0}\bar{D}^{*0} \gamma$
seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^{*0}\bar{D}^{*0}$

Γ_9/Γ

$\Gamma(D^*(2007)^0\bar{D}^*(2007)^0)/\Gamma(D^*(2007)^0\bar{D}^0 + \text{c.c.})$

VALUE	DOCUMENT ID	TECN	COMMENT
32.0±12.0	²⁶ GOLDHABER 77	MRK1	$e^+ e^-$

Γ_9/Γ_6

²⁶ Phase-space factor (p^3) explicitly removed.

$\Gamma(D^*(2010)^+ D^*(2010)^-)/\Gamma_{\text{total}}$ Γ_{10}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
seen	27 ZHUKOVA 18	BELL	$e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$
seen	AUBERT 09M	BABR	$e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$
seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^{*+} D^{*-}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen PAKHLOVA 07 BELL $e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$

27 Supersedes PAKHLOVA 07.

 $\Gamma(D^0 D^- \pi^+ +\text{c.c. (excl. } D^*(2010)^+ D^- +\text{c.c.)})/\Gamma_{\text{total}}$ Γ_{12}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
not seen	PAKHLOVA 08A	BELL	$e^+ e^- \rightarrow D^0 D^- \pi^+ \gamma$

 $\Gamma(D \bar{D}^* \pi (\text{excl. } D^* \bar{D}^*))/\Gamma_{\text{total}}$ Γ_{13}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
not seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D \bar{D}^* \pi$

 $\Gamma(D^0 \bar{D}^{*-} \pi^+ +\text{c.c. (excl. } D^*(2010)^+ D^*(2010)^-))/\Gamma_{\text{total}}$ Γ_{14}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
seen	PAKHLOVA 09	BELL	$e^+ e^- \rightarrow D^0 \bar{D}^{*-} \pi^+ \gamma$

 $\Gamma(D_s^+ D_s^-)/\Gamma_{\text{total}}$ Γ_{15}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
seen	PAKHLOVA 11	BELL	$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$
seen	DEL-AMO-SA..10N	BABR	$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$
seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D_s^+ D_s^-$

 $\Gamma(\pi^+ \pi^+ \pi^- \pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_{16}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$(3.51 \pm 1.89 \pm 1.24) \times 10^{-5}$	28 ABLIKIM	21AW BES3	$e^+ e^- \rightarrow 2\pi^+ 2\pi^- \pi^0$

$(2.41 \pm 0.05 \pm 0.79) \times 10^{-2}$ 29 ABLIKIM 21AW BES3 $e^+ e^- \rightarrow 2\pi^+ 2\pi^- \pi^0$

28 Solution 1 of two solutions with equal fit quality. The significance of the $\psi(4040)$ signal is 3.6 σ .

29 Solution 2 of two solutions with equal fit quality. The significance of the $\psi(4040)$ signal is 3.6 σ .

 $\Gamma(J/\psi \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{18}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<4	90	COAN 06	CLEO	$3.97\text{--}4.06 e^+ e^- \rightarrow \text{hadrons}$

 $\Gamma(J/\psi \pi^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{19}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2	90	COAN 06	CLEO	$3.97\text{--}4.06 e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(J/\psi\eta)/\Gamma_{\text{total}}$ Γ_{20}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.2±0.5±0.5	30	ABLIKIM	12K BES3	$e^+e^- \rightarrow \ell^+\ell^-2\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<7 90 COAN 06 CLEO 3.97–4.06 $e^+e^- \rightarrow$ hadrons

30 ABLIKIM 12K measure $\sigma(e^+e^- \rightarrow J/\psi\eta) = 32.1 \pm 2.8 \pm 1.3$ pb. They assume the $\eta J/\psi$ fully originates from $\psi(4040)$ decays.

 $\Gamma(J/\psi\pi^0)/\Gamma_{\text{total}}$ Γ_{21}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.28	90	31 ABLIKIM	12K BES3	$e^+e^- \rightarrow \ell^+\ell^-2\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2 90 COAN 06 CLEO 3.97–4.06 $e^+e^- \rightarrow$ hadrons

31 ABLIKIM 12K measure $\sigma(e^+e^- \rightarrow J/\psi\pi^0) < 1.6$ pb. They assume the $\eta J/\psi$ fully originates from $\psi(4040)$ decays.

 $\Gamma(J/\psi\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{22}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2	90	COAN	06	CLEO 3.97–4.06 $e^+e^- \rightarrow$ hadrons

 $\Gamma(\chi_{c1}\gamma)/\Gamma_{\text{total}}$ Γ_{23}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<11	90	COAN	06	CLEO 3.97–4.06 $e^+e^- \rightarrow$ hadrons

 $\Gamma(\chi_{c2}\gamma)/\Gamma_{\text{total}}$ Γ_{24}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<17	90	COAN	06	CLEO 3.97–4.06 $e^+e^- \rightarrow$ hadrons

 $\Gamma(\chi_{c1}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{25}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<11	90	COAN	06	CLEO 3.97–4.06 $e^+e^- \rightarrow$ hadrons

 $\Gamma(\chi_{c2}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{26}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<32	90	COAN	06	CLEO 3.97–4.06 $e^+e^- \rightarrow$ hadrons

 $\Gamma(h_c(1P)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{27}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<3	90	32 PEDLAR	11 CLEO	$e^+e^- \rightarrow h_c(1P)\pi^+\pi^-$

32 From several values of \sqrt{s} near the peak of the $\psi(4040)$, PEDLAR 11 measures $\sigma(e^+e^- \rightarrow h_c(1P)\pi^+\pi^-) = 1.0 \pm 8.0 \pm 5.4 \pm 0.2$ pb, where the errors are statistical, systematic, and due to uncertainty in $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$, respectively.

$\Gamma(\phi\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{28}/Γ

<i>VALUE</i> (units 10^{-3})	<i>CL%</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
<3	90	COAN	06	CLEO 3.97–4.06 $e^+e^- \rightarrow$ hadrons

 $\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{29}/Γ

<i>VALUE</i> (units 10^{-4})	<i>CL%</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
<2.9	90	33 ABLIKIM	13Q	BES3 $e^+e^- \rightarrow \psi(4040)$

33 Assuming that interference effects between resonance and continuum can be neglected.

 $\Gamma(\Lambda\bar{\Lambda}\pi^0)/\Gamma_{\text{total}}$ Γ_{30}/Γ

<i>VALUE</i> (units 10^{-4})	<i>CL%</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
<0.9	90	34 ABLIKIM	13Q	BES3 $e^+e^- \rightarrow \psi(4040)$

34 Assuming that interference effects between resonance and continuum can be neglected.

 $\Gamma(\Lambda\bar{\Lambda}\eta)/\Gamma_{\text{total}}$ Γ_{31}/Γ

<i>VALUE</i> (units 10^{-4})	<i>CL%</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
<3.0	90	35 ABLIKIM	13Q	BES3 $e^+e^- \rightarrow \psi(4040)$

35 Assuming that interference effects between resonance and continuum can be neglected.

 $\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$ Γ_{33}/Γ

<i>VALUE</i> (units 10^{-4})	<i>CL%</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
<1.3	90	36 ABLIKIM	13Q	BES3 $e^+e^- \rightarrow \psi(4040)$

36 Assuming that interference effects between resonance and continuum can be neglected.

 $\Gamma(\Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}$ Γ_{34}/Γ

<i>VALUE</i> (units 10^{-4})	<i>CL%</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
<0.7	90	37 ABLIKIM	13Q	BES3 $e^+e^- \rightarrow \psi(4040)$

37 Assuming that interference effects between resonance and continuum can be neglected.

 $\Gamma(\Xi^+\bar{\Xi}^-)/\Gamma_{\text{total}}$ Γ_{35}/Γ

<i>VALUE</i> (units 10^{-4})	<i>CL%</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
<1.6	90	38 ABLIKIM	13Q	BES3 $e^+e^- \rightarrow \psi(4040)$

38 Assuming that interference effects between resonance and continuum can be neglected.

 $\Gamma(\Xi^0\bar{\Xi}^0)/\Gamma_{\text{total}}$ Γ_{36}/Γ

<i>VALUE</i> (units 10^{-4})	<i>CL%</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
<1.8	90	39 ABLIKIM	13Q	BES3 $e^+e^- \rightarrow \psi(4040)$

39 Assuming that interference effects between resonance and continuum can be neglected.

$\psi(4040)$ REFERENCES

ABLIKIM	23BK	JHEP 2311 228	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21AS	PR D104 L091104	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21AW	PR D104 112009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	20AG	PR D102 112009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	20O	PR D102 031101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ZHUKOVA	18	PR D97 012002	V. Zhukova <i>et al.</i>	(BELLE Collab.)
HAN	15	PR D92 012011	Y.L. Han <i>et al.</i>	(BELLE Collab.)
ABLIKIM	13Q	PR D87 112011	M. Ablikim <i>et al.</i>	(BESIII Collab.)
WANG	13B	PR D87 051101	X.L. Wang <i>et al.</i>	(BELLE Collab.)
ABLIKIM	12K	PR D86 071101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
PAKHLOVA	11	PR D83 011101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PEDLAR	11	PRL 107 041803	T. Pedlar <i>et al.</i>	(CLEO Collab.)
DEL-AMO-SA...	10N	PR D82 052004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
MO	10	PR D82 077501	X.H. Mo, C.Z. Yuan, P. Wang	(BHEP)
AUBERT	09M	PR D79 092001	B. Aubert <i>et al.</i>	(BABAR Collab.)
CRONIN-HEN...	09	PR D80 072001	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)
PAKHLOVA	09	PR D80 091101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
ABLIKIM	08D	PL B660 315	M. Ablikim <i>et al.</i>	(BES Collab.)
PAKHLOVA	08	PR D77 011103	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PAKHLOVA	08A	PRL 100 062001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PAKHLOVA	07	PRL 98 092001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
COAN	06	PRL 96 162003	T.E. Coan <i>et al.</i>	(CLEO Collab.)
SETH	05A	PR D72 017501	K.K. Seth	
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)
OSTERHELD	86	SLAC-PUB-4160	A. Osterheld <i>et al.</i>	(SLAC Crystal Ball Collab.)
BRANDELIK	78C	PL 76B 361	R. Brandelik <i>et al.</i>	(DASP Collab.)
Also		ZPHY C1 233	R. Brandelik <i>et al.</i>	(DASP Collab.)
FELDMAN	77	PRPL 33C 285	G.J. Feldman, M.L. Perl	(LBL, SLAC)
GOLDHABER	77	PL 69B 503	G. Goldhaber <i>et al.</i>	(Mark I Collab.)