

# $\Upsilon(2S)$

$$J^{PC} = 0^{-}(1^{-}-)$$

## $\Upsilon(2S)$ MASS

| <u>VALUE (MeV)</u>                                                            | <u>DOCUMENT ID</u>          | <u>TECN</u> | <u>COMMENT</u>               |
|-------------------------------------------------------------------------------|-----------------------------|-------------|------------------------------|
| <b>10023.4 ± 0.5</b>                                                          | <sup>1</sup> SHAMOV 23      | RVUE        | $e^+e^- \rightarrow$ hadrons |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● |                             |             |                              |
| 10022.7 ± 0.4                                                                 | <sup>2</sup> SHAMOV 23      | RVUE        | $e^+e^- \rightarrow$ hadrons |
| 10023.5 ± 0.5                                                                 | <sup>3,4</sup> ARTAMONOV 00 | MD1         | $e^+e^- \rightarrow$ hadrons |
| 10023.6 ± 0.5                                                                 | <sup>5,6</sup> BARU 86B     | MD1         | $e^+e^- \rightarrow$ hadrons |
| 10023.1 ± 0.4                                                                 | <sup>7</sup> BARBER 84      | ARG         | $e^+e^- \rightarrow$ hadrons |

<sup>1</sup> Reanalysis of MD1 data using the electron mass from COHEN 87, the radiative corrections from KURAEV 85 and interference effects.

<sup>2</sup> Obtained by reanalysing ARGUS and Crystal Ball data (BARBER 84), but not authored by the ARGUS and Crystal Ball collaboration.

<sup>3</sup> Reanalysis of BARU 86B using new electron mass (COHEN 87).

<sup>4</sup> Superseded by SHAMOV 23.

<sup>5</sup> Reanalysis of ARTAMONOV 84.

<sup>6</sup> Superseded by ARTAMONOV 00.

<sup>7</sup> Reanalysed by SHAMOV 23.

## $m\Upsilon(3S) - m\Upsilon(2S)$

| <u>VALUE (MeV)</u>          | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                   |
|-----------------------------|--------------------|-------------|----------------------------------|
| <b>331.50 ± 0.02 ± 0.13</b> | LEES 11C           | BABR        | $e^+e^- \rightarrow \pi^+\pi^-X$ |

## $\Upsilon(2S)$ WIDTH

| <u>VALUE (keV)</u>                 | <u>DOCUMENT ID</u>                                              |
|------------------------------------|-----------------------------------------------------------------|
| <b>31.98 ± 2.63 OUR EVALUATION</b> | See the Note on "Width Determinations of the $\Upsilon$ States" |

## $\Upsilon(2S)$ DECAY MODES

| Mode                                    | Fraction ( $\Gamma_i/\Gamma$ ) | Scale factor/<br>Confidence level |
|-----------------------------------------|--------------------------------|-----------------------------------|
| $\Gamma_1 \quad \Upsilon(1S)\pi^+\pi^-$ | (17.85 ± 0.26) %               |                                   |
| $\Gamma_2 \quad \Upsilon(1S)\pi^0\pi^0$ | ( 8.6 ± 0.4 ) %                |                                   |
| $\Gamma_3 \quad \tau^+\tau^-$           | ( 2.00 ± 0.21 ) %              |                                   |
| $\Gamma_4 \quad \mu^+\mu^-$             | ( 1.93 ± 0.17 ) %              | S=2.2                             |
| $\Gamma_5 \quad e^+e^-$                 | ( 1.91 ± 0.16 ) %              |                                   |
| $\Gamma_6 \quad \Upsilon(1S)\pi^0$      | < 4                            | $\times 10^{-5}$ CL=90%           |
| $\Gamma_7 \quad \Upsilon(1S)\eta$       | ( 2.9 ± 0.4 ) $\times 10^{-4}$ | S=2.0                             |
| $\Gamma_8 \quad J/\psi(1S)$ anything    | < 6                            | $\times 10^{-3}$ CL=90%           |
| $\Gamma_9 \quad J/\psi(1S)\eta_c$       | < 5.4                          | $\times 10^{-6}$ CL=90%           |
| $\Gamma_{10} \quad J/\psi(1S)\chi_{c0}$ | < 3.4                          | $\times 10^{-6}$ CL=90%           |

|               |                                                   |                            |                  |        |
|---------------|---------------------------------------------------|----------------------------|------------------|--------|
| $\Gamma_{11}$ | $J/\psi(1S)\chi_{c1}$                             | $< 1.2$                    | $\times 10^{-6}$ | CL=90% |
| $\Gamma_{12}$ | $J/\psi(1S)\chi_{c2}$                             | $< 2.0$                    | $\times 10^{-6}$ | CL=90% |
| $\Gamma_{13}$ | $J/\psi(1S)\eta_c(2S)$                            | $< 2.5$                    | $\times 10^{-6}$ | CL=90% |
| $\Gamma_{14}$ | $J/\psi(1S)X(3940)$                               | $< 2.0$                    | $\times 10^{-6}$ | CL=90% |
| $\Gamma_{15}$ | $J/\psi(1S)X(4160)$                               | $< 2.0$                    | $\times 10^{-6}$ | CL=90% |
| $\Gamma_{16}$ | $\chi_{c1}$ anything                              | $( 2.2 \pm 0.5 )$          | $\times 10^{-4}$ |        |
| $\Gamma_{17}$ | $\chi_{c1}(1P)^0 X_{tetra}$                       | $< 3.67$                   | $\times 10^{-5}$ | CL=90% |
| $\Gamma_{18}$ | $\chi_{c2}$ anything                              | $( 2.3 \pm 0.8 )$          | $\times 10^{-4}$ |        |
| $\Gamma_{19}$ | $\psi(2S)\eta_c$                                  | $< 5.1$                    | $\times 10^{-6}$ | CL=90% |
| $\Gamma_{20}$ | $\psi(2S)\chi_{c0}$                               | $< 4.7$                    | $\times 10^{-6}$ | CL=90% |
| $\Gamma_{21}$ | $\psi(2S)\chi_{c1}$                               | $< 2.5$                    | $\times 10^{-6}$ | CL=90% |
| $\Gamma_{22}$ | $\psi(2S)\chi_{c2}$                               | $< 1.9$                    | $\times 10^{-6}$ | CL=90% |
| $\Gamma_{23}$ | $\psi(2S)\eta_c(2S)$                              | $< 3.3$                    | $\times 10^{-6}$ | CL=90% |
| $\Gamma_{24}$ | $\psi(2S)X(3940)$                                 | $< 3.9$                    | $\times 10^{-6}$ | CL=90% |
| $\Gamma_{25}$ | $\psi(2S)X(4160)$                                 | $< 3.9$                    | $\times 10^{-6}$ | CL=90% |
| $\Gamma_{26}$ | $T_{c\bar{c}1}(3900)^+ T_{c\bar{c}1}(3900)^-$     | $< 1.0$                    | $\times 10^{-6}$ | CL=90% |
| $\Gamma_{27}$ | $T_{c\bar{c}1}(4200)^+ T_{c\bar{c}1}(4200)^-$     | $< 1.67$                   | $\times 10^{-5}$ | CL=90% |
| $\Gamma_{28}$ | $T_{c\bar{c}1}(3900)^\pm T_{c\bar{c}1}(4200)^\mp$ | $< 7.3$                    | $\times 10^{-6}$ | CL=90% |
| $\Gamma_{29}$ | $T_{c\bar{c}}(4050)^+ T_{c\bar{c}}(4050)^-$       | $< 1.35$                   | $\times 10^{-5}$ | CL=90% |
| $\Gamma_{30}$ | $T_{c\bar{c}}(4250)^+ T_{c\bar{c}}(4250)^-$       | $< 2.67$                   | $\times 10^{-5}$ | CL=90% |
| $\Gamma_{31}$ | $T_{c\bar{c}}(4050)^\pm T_{c\bar{c}}(4250)^\mp$   | $< 2.72$                   | $\times 10^{-5}$ | CL=90% |
| $\Gamma_{32}$ | $T_{c\bar{c}1}(4430)^+ T_{c\bar{c}1}(4430)^-$     | $< 2.03$                   | $\times 10^{-5}$ | CL=90% |
| $\Gamma_{33}$ | $T_{c\bar{c}}(4055)^\pm T_{c\bar{c}}(4055)^\mp$   | $< 1.11$                   | $\times 10^{-5}$ | CL=90% |
| $\Gamma_{34}$ | $T_{c\bar{c}}(4055)^\pm T_{c\bar{c}1}(4430)^\mp$  | $< 2.11$                   | $\times 10^{-5}$ | CL=90% |
| $\Gamma_{35}$ | $\bar{2}H$ anything                               | $( 2.78_{-0.26}^{+0.30} )$ | $\times 10^{-5}$ | S=1.2  |
| $\Gamma_{36}$ | hadrons                                           | $(94 \pm 11)$              | %                |        |
| $\Gamma_{37}$ | $ggg$                                             | $(58.8 \pm 1.2)$           | %                |        |
| $\Gamma_{38}$ | $\gamma gg$                                       | $( 1.87 \pm 0.28 )$        | %                |        |
| $\Gamma_{39}$ | $\phi K^+ K^-$                                    | $( 1.6 \pm 0.4 )$          | $\times 10^{-6}$ |        |
| $\Gamma_{40}$ | $\omega \pi^+ \pi^-$                              | $< 2.58$                   | $\times 10^{-6}$ | CL=90% |
| $\Gamma_{41}$ | $K^*(892)^0 K^- \pi^+ + \text{c.c.}$              | $( 2.3 \pm 0.7 )$          | $\times 10^{-6}$ |        |
| $\Gamma_{42}$ | $\phi f_2'(1525)$                                 | $< 1.33$                   | $\times 10^{-6}$ | CL=90% |
| $\Gamma_{43}$ | $\omega f_2(1270)$                                | $< 5.7$                    | $\times 10^{-7}$ | CL=90% |
| $\Gamma_{44}$ | $\rho(770) a_2(1320)$                             | $< 8.8$                    | $\times 10^{-7}$ | CL=90% |
| $\Gamma_{45}$ | $K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.}$    | $( 1.5 \pm 0.6 )$          | $\times 10^{-6}$ |        |
| $\Gamma_{46}$ | $K_1(1270)^\pm K^\mp$                             | $< 3.22$                   | $\times 10^{-6}$ | CL=90% |
| $\Gamma_{47}$ | $K_1(1400)^\pm K^\mp$                             | $< 8.3$                    | $\times 10^{-7}$ | CL=90% |
| $\Gamma_{48}$ | $b_1(1235)^\pm \pi^\mp$                           | $< 4.0$                    | $\times 10^{-7}$ | CL=90% |
| $\Gamma_{49}$ | $\rho \pi$                                        | $< 1.16$                   | $\times 10^{-6}$ | CL=90% |
| $\Gamma_{50}$ | $\pi^+ \pi^- \pi^0$                               | $< 8.0$                    | $\times 10^{-7}$ | CL=90% |
| $\Gamma_{51}$ | $\omega \pi^0$                                    | $< 1.63$                   | $\times 10^{-6}$ | CL=90% |
| $\Gamma_{52}$ | $\pi^+ \pi^- \pi^0 \pi^0$                         | $( 1.30 \pm 0.28 )$        | $\times 10^{-5}$ |        |
| $\Gamma_{53}$ | $K_S^0 K^+ \pi^- + \text{c.c.}$                   | $( 1.14 \pm 0.33 )$        | $\times 10^{-6}$ |        |

|               |                                                                        |                   |                  |        |
|---------------|------------------------------------------------------------------------|-------------------|------------------|--------|
| $\Gamma_{54}$ | $K^*(892)^0 \bar{K}^0 + \text{c.c.}$                                   | $< 4.22$          | $\times 10^{-6}$ | CL=90% |
| $\Gamma_{55}$ | $K^*(892)^- K^+ + \text{c.c.}$                                         | $< 1.45$          | $\times 10^{-6}$ | CL=90% |
| $\Gamma_{56}$ | $f_1(1285)$ anything                                                   | $(2.2 \pm 1.6)$   | $\times 10^{-3}$ |        |
| $\Gamma_{57}$ | $f_1(1285) X_{tetra}$                                                  | $< 6.47$          | $\times 10^{-5}$ | CL=90% |
| $\Gamma_{58}$ | $D_s^+ D_{s1}(2536)^-$                                                 |                   |                  |        |
| $\Gamma_{59}$ | $D_s^+ D_{s1}(2536)^-, D_{s1}^- \rightarrow$<br>$K^- D^*(2007)^0$      | $(1.6 \pm 0.4)$   | $\times 10^{-5}$ |        |
| $\Gamma_{60}$ | $D_s^+ D_{s1}(2536)^-, D_{s1}^- \rightarrow$<br>$K_S^0 D^*(2010)^-$    | $(8.4 \pm 2.3)$   | $\times 10^{-6}$ |        |
| $\Gamma_{61}$ | $D_s^{*+} D_{s1}(2536)^-$                                              |                   |                  |        |
| $\Gamma_{62}$ | $D_s^{*+} D_{s1}(2536)^-, D_{s1}^- \rightarrow$<br>$K^- D^*(2007)^0$   | $(1.4 \pm 0.4)$   | $\times 10^{-5}$ |        |
| $\Gamma_{63}$ | $D_s^{*+} D_{s1}(2536)^-, D_{s1}^- \rightarrow$<br>$K_S^0 D^*(2010)^-$ | $(8.2 \pm 3.1)$   | $\times 10^{-6}$ |        |
| $\Gamma_{64}$ | $D_s^+ D_{s2}^*(2573)^-$                                               |                   |                  |        |
| $\Gamma_{65}$ | $D_s^+ D_{s2}^*(2573)^-, D_{s2}^{*-} \rightarrow$<br>$K^- D^0$         | $(1.4 \pm 0.4)$   | $\times 10^{-5}$ |        |
| $\Gamma_{66}$ | $D_s^+ D_{s2}^*(2573)^-, D_{s2}^{*-} \rightarrow$<br>$K_S^0 D^-$       | $(6.9 \pm 3.0)$   | $\times 10^{-6}$ |        |
| $\Gamma_{67}$ | $D_s^{*+} D_{s2}^*(2573)^-$                                            |                   |                  |        |
| $\Gamma_{68}$ | $D_s^{*+} D_{s2}^*(2573)^-, D_{s2}^{*-} \rightarrow$<br>$K^- D^0$      | $(9 \pm 5)$       | $\times 10^{-6}$ |        |
| $\Gamma_{69}$ | $D_s^{*+} D_{s2}^*(2573)^-, D_{s2}^{*-} \rightarrow$<br>$K_S^0 D^-$    | $(5 \pm 6)$       | $\times 10^{-6}$ |        |
| $\Gamma_{70}$ | Sum of 100 exclusive modes                                             | $(2.90 \pm 0.30)$ | $\times 10^{-3}$ |        |

### Radiative decays

|               |                                                                               |                   |                  |        |
|---------------|-------------------------------------------------------------------------------|-------------------|------------------|--------|
| $\Gamma_{71}$ | $\gamma \chi_{b1}(1P)$                                                        | $(6.9 \pm 0.4)$   | %                |        |
| $\Gamma_{72}$ | $\gamma \chi_{b2}(1P)$                                                        | $(7.15 \pm 0.35)$ | %                |        |
| $\Gamma_{73}$ | $\gamma \chi_{b0}(1P)$                                                        | $(3.8 \pm 0.4)$   | %                |        |
| $\Gamma_{74}$ | $\gamma f_0(1710)$                                                            | $< 5.9$           | $\times 10^{-4}$ | CL=90% |
| $\Gamma_{75}$ | $\gamma f_2'(1525)$                                                           | $< 5.3$           | $\times 10^{-4}$ | CL=90% |
| $\Gamma_{76}$ | $\gamma f_2(1270)$                                                            | $< 2.41$          | $\times 10^{-4}$ | CL=90% |
| $\Gamma_{77}$ | $\gamma f_J(2220)$                                                            |                   |                  |        |
| $\Gamma_{78}$ | $\gamma \eta_c(1S)$                                                           | $< 2.7$           | $\times 10^{-5}$ | CL=90% |
| $\Gamma_{79}$ | $\gamma \chi_{c0}$                                                            | $< 1.0$           | $\times 10^{-4}$ | CL=90% |
| $\Gamma_{80}$ | $\gamma \chi_{c1}$                                                            | $< 3.6$           | $\times 10^{-6}$ | CL=90% |
| $\Gamma_{81}$ | $\gamma \chi_{c2}$                                                            | $< 1.5$           | $\times 10^{-5}$ | CL=90% |
| $\Gamma_{82}$ | $\gamma \chi_{c1}(3872)$                                                      | $< 2.3$           | $\times 10^{-5}$ | CL=90% |
| $\Gamma_{83}$ | $\gamma \chi_{c1}(3872), \chi_{c1} \rightarrow$<br>$\pi^+ \pi^- \pi^0 J/\psi$ | $< 2.4$           | $\times 10^{-6}$ | CL=90% |
| $\Gamma_{84}$ | $\gamma \chi_{c0}(3915) \rightarrow \omega J/\psi$                            | $< 2.8$           | $\times 10^{-6}$ | CL=90% |

|               |                                                                    |                 |                  |        |
|---------------|--------------------------------------------------------------------|-----------------|------------------|--------|
| $\Gamma_{85}$ | $\gamma\chi_{c1}(4140) \rightarrow \phi J/\psi$                    | $< 1.2$         | $\times 10^{-6}$ | CL=90% |
| $\Gamma_{86}$ | $\gamma X(4350) \rightarrow \phi J/\psi$                           | $< 1.3$         | $\times 10^{-6}$ | CL=90% |
| $\Gamma_{87}$ | $\gamma\eta_b(1S)$                                                 | $(5.5 \pm 1.1)$ | $\times 10^{-4}$ | S=1.2  |
| $\Gamma_{88}$ | $\gamma\eta_b(1S) \rightarrow \gamma$ Sum of 26 exclusive modes    | $< 3.7$         | $\times 10^{-6}$ | CL=90% |
| $\Gamma_{89}$ | $\gamma X_{b\bar{b}} \rightarrow \gamma$ Sum of 26 exclusive modes | $< 4.9$         | $\times 10^{-6}$ | CL=90% |
| $\Gamma_{90}$ | $\gamma X \rightarrow \gamma + \geq 4$ prongs                      | [a] $< 1.95$    | $\times 10^{-4}$ | CL=95% |
| $\Gamma_{91}$ | $\gamma A^0 \rightarrow \gamma$ hadrons                            | $< 8$           | $\times 10^{-5}$ | CL=90% |
| $\Gamma_{92}$ | $\gamma A^0 \rightarrow \gamma\mu^+\mu^-$                          | $< 8.3$         | $\times 10^{-6}$ | CL=90% |

**Lepton Family number (LF) violating modes**


|               |                    |    |         |                  |        |
|---------------|--------------------|----|---------|------------------|--------|
| $\Gamma_{93}$ | $e^\pm \tau^\mp$   | LF | $< 3.2$ | $\times 10^{-6}$ | CL=90% |
| $\Gamma_{94}$ | $\mu^\pm \tau^\mp$ | LF | $< 3.3$ | $\times 10^{-6}$ | CL=90% |

[a] 1.5 GeV <  $m_X$  < 5.0 GeV

**FIT INFORMATION**

An overall fit to 3 branching ratios uses 13 measurements to determine 2 parameters. The overall fit has a  $\chi^2 = 11.8$  for 11 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$ , in percent, from the fit to the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ .



**$\Upsilon(2S) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$**

|                                                                                    |                                                     |
|------------------------------------------------------------------------------------|-----------------------------------------------------|
| <b><math>\Gamma(\mu^+\mu^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}</math></b> | <b><math>\Gamma_4\Gamma_5/\Gamma</math></b>         |
| <u>VALUE (eV)</u>                                                                  | <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>       |
| <b>6.5±1.5±1.0</b>                                                                 | KOBEL    92    CBAL $e^+e^- \rightarrow \mu^+\mu^-$ |

|                                                                                                |                                                                                                   |
|------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------|
| <b><math>\Gamma(\Upsilon(1S)\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}</math></b> | <b><math>\Gamma_1\Gamma_5/\Gamma</math></b>                                                       |
| <u>VALUE (eV)</u>                                                                              | <u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>                                         |
| <b>105.4±1.0±4.2</b>                                                                           | 11.8k <sup>1</sup> AUBERT    08BP BABR    10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\ell^+\ell^-$ |

<sup>1</sup>Using  $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$  and  $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$ .

$\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{36}\Gamma_5/\Gamma$

| <u>VALUE (keV)</u>                           | <u>DOCUMENT ID</u>      | <u>TECN</u> | <u>COMMENT</u>                               |
|----------------------------------------------|-------------------------|-------------|----------------------------------------------|
| <b>0.577±0.009 OUR AVERAGE</b>               |                         |             |                                              |
| 0.581±0.004±0.009                            | <sup>1</sup> ROSNER     | 06 CLEO     | 10.0 e <sup>+</sup> e <sup>-</sup> → hadrons |
| 0.552±0.031±0.017                            | <sup>1</sup> BARU       | 96 MD1      | e <sup>+</sup> e <sup>-</sup> → hadrons      |
| 0.54 ±0.04 ±0.02                             | <sup>1</sup> JAKUBOWSKI | 88 CBAL     | e <sup>+</sup> e <sup>-</sup> → hadrons      |
| 0.58 ±0.03 ±0.04                             | <sup>2</sup> GILES      | 84B CLEO    | e <sup>+</sup> e <sup>-</sup> → hadrons      |
| 0.60 ±0.12 ±0.07                             | <sup>2</sup> ALBRECHT   | 82 DASP     | e <sup>+</sup> e <sup>-</sup> → hadrons      |
| 0.54 ±0.07 <sup>+0.09</sup> <sub>-0.05</sub> | <sup>2</sup> NICZYPORUK | 81C LENA    | e <sup>+</sup> e <sup>-</sup> → hadrons      |
| 0.41 ±0.18                                   | <sup>2</sup> BOCK       | 80 CNTR     | e <sup>+</sup> e <sup>-</sup> → hadrons      |

<sup>1</sup> Radiative corrections evaluated following KURAEV 85.  
<sup>2</sup> Radiative corrections reevaluated by BUCHMUELLER 88 following KURAEV 85.

**$\Upsilon(2S)$  PARTIAL WIDTHS**

$\Gamma(e^+e^-)$   $\Gamma_5$

| <u>VALUE (keV)</u>                | <u>DOCUMENT ID</u> |
|-----------------------------------|--------------------|
| <b>0.612±0.011 OUR EVALUATION</b> |                    |

**$\Upsilon(2S)$  BRANCHING RATIOS**

$\Gamma(\Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_1/\Gamma$

Abbreviation MM in the COMMENT field below stands for missing mass.

| <u>VALUE (units 10<sup>-2</sup>)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u>     | <u>TECN</u> | <u>COMMENT</u>                                                                                 |
|--------------------------------------|-------------|------------------------|-------------|------------------------------------------------------------------------------------------------|
| <b>17.85±0.26 OUR FIT</b>            |             |                        |             |                                                                                                |
| <b>17.92±0.26 OUR AVERAGE</b>        |             |                        |             |                                                                                                |
| 16.8 ±1.1 ±1.3                       | 906k        | <sup>1</sup> LEES      | 11C BABR    | e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> X                                |
| 17.80±0.05±0.37                      | 170k        | <sup>2</sup> LEES      | 11L BABR    | $\Upsilon(2S) \rightarrow \pi^+\pi^-\mu^+\mu^-$                                                |
| 18.02±0.02±0.61                      | 851k        | <sup>3</sup> BHARI     | 09 CLEO     | e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> MM                               |
| 17.22±0.17±0.75                      | 11.8k       | <sup>4</sup> AUBERT    | 08BP BABR   | e <sup>+</sup> e <sup>-</sup> → γπ <sup>+</sup> π <sup>-</sup> ℓ <sup>+</sup> ℓ <sup>-</sup>   |
| 19.2 ±0.2 ±1.0                       | 52.6k       | <sup>5</sup> ALEXANDER | 98 CLE2     | π <sup>+</sup> π <sup>-</sup> ℓ <sup>+</sup> ℓ <sup>-</sup> , π <sup>+</sup> π <sup>-</sup> MM |
| 18.1 ±0.5 ±1.0                       | 11.6k       | ALBRECHT               | 87 ARG      | e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> MM                               |
| 16.9 ±4.0                            |             | GELPHMAN               | 85 CBAL     | e <sup>+</sup> e <sup>-</sup> → e <sup>+</sup> e <sup>-</sup> π <sup>+</sup> π <sup>-</sup>    |
| 19.1 ±1.2 ±0.6                       |             | BESSION                | 84 CLEO     | π <sup>+</sup> π <sup>-</sup> MM                                                               |
| 18.9 ±2.6                            |             | FONSECA                | 84 CUSB     | e <sup>+</sup> e <sup>-</sup> → ℓ <sup>+</sup> ℓ <sup>-</sup> π <sup>+</sup> π <sup>-</sup>    |
| 21 ±7                                | 7           | NICZYPORUK             | 81B LENA    | e <sup>+</sup> e <sup>-</sup> → ℓ <sup>+</sup> ℓ <sup>-</sup> π <sup>+</sup> π <sup>-</sup>    |

<sup>1</sup> LEES 11C reports  $[\Gamma(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \Upsilon(2S)\text{anything})] = (1.78 \pm 0.02 \pm 0.11) \times 10^{-2}$  which we divide by our best value  $B(\Upsilon(3S) \rightarrow \Upsilon(2S)\text{anything}) = (10.6 \pm 0.8) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Using  $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$ .

<sup>3</sup> A weighted average of the inclusive and exclusive results.

<sup>4</sup> Using  $B(\Upsilon(2S) \rightarrow e^+e^-) = (1.91 \pm 0.16)\%$ ,  $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17)\%$  and,  $\Gamma_{ee}(\Upsilon(2S)) = 0.612 \pm 0.011$  keV.

<sup>5</sup> Using  $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.52 \pm 0.17)\%$  and  $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.07)\%$ .

$\Gamma(\Upsilon(1S)\pi^0\pi^0)/\Gamma_{\text{total}}$   $\Gamma_2/\Gamma$

| VALUE (units $10^{-2}$ )     | EVTS | DOCUMENT ID               | TECN | COMMENT                                     |
|------------------------------|------|---------------------------|------|---------------------------------------------|
| <b>8.6 ± 0.4 OUR AVERAGE</b> |      |                           |      |                                             |
| 8.43 ± 0.16 ± 0.42           | 38k  | <sup>1</sup> BHARI 09     | CLEO | $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$ |
| 9.2 ± 0.6 ± 0.8              | 275  | <sup>2</sup> ALEXANDER 98 | CLE2 | $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$ |
| 9.5 ± 1.9 ± 1.9              | 25   | ALBRECHT 87               | ARG  | $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$ |
| 8.0 ± 1.5                    |      | GELPHMAN 85               | CBAL | $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$ |
| 10.3 ± 2.3                   |      | FONSECA 84                | CUSB | $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$ |

<sup>1</sup> Authors assume  $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$ .

<sup>2</sup> Using  $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.52 \pm 0.17)\%$  and  $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.07)\%$ .

$\Gamma(\Upsilon(1S)\pi^0\pi^0)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$   $\Gamma_2/\Gamma_1$

| VALUE         | DOCUMENT ID           | TECN | COMMENT                           |
|---------------|-----------------------|------|-----------------------------------|
| 0.462 ± 0.037 | <sup>1</sup> BHARI 09 | CLEO | $e^+e^- \rightarrow \Upsilon(2S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup> Not independent of other values reported by BHARI 09.

$\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}}$   $\Gamma_3/\Gamma$

| VALUE (units $10^{-2}$ )       | EVTS | DOCUMENT ID            | TECN | COMMENT                                                    |
|--------------------------------|------|------------------------|------|------------------------------------------------------------|
| <b>2.00 ± 0.21 OUR AVERAGE</b> |      |                        |      |                                                            |
| 2.00 ± 0.12 ± 0.18             | 22k  | <sup>1</sup> BESSON 07 | CLEO | $e^+e^- \rightarrow \Upsilon(2S) \rightarrow \tau^+\tau^-$ |
| 1.7 ± 1.5 ± 0.6                |      | HAAS 84B               | CLEO | $e^+e^- \rightarrow \tau^+\tau^-$                          |

<sup>1</sup> BESSON 07 reports  $[\Gamma(\Upsilon(2S) \rightarrow \tau^+\tau^-)/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \mu^+\mu^-)] = 1.04 \pm 0.04 \pm 0.05$  which we multiply by our best value  $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$   $\Gamma_4/\Gamma$


| VALUE                              | CL% | EVTS | DOCUMENT ID              | TECN | COMMENT                                                     |
|------------------------------------|-----|------|--------------------------|------|-------------------------------------------------------------|
| <b>0.0193 ± 0.0017 OUR AVERAGE</b> |     |      |                          |      | Error includes scale factor of 2.2. See the ideogram below. |
| 0.0203 ± 0.0003 ± 0.0008           |     | 120k | ADAMS 05                 | CLEO | $e^+e^- \rightarrow \mu^+\mu^-$                             |
| 0.0122 ± 0.0028 ± 0.0019           |     |      | <sup>1</sup> KOBEL 92    | CBAL | $e^+e^- \rightarrow \mu^+\mu^-$                             |
| 0.0138 ± 0.0025 ± 0.0015           |     |      | KAARSBERG 89             | CSB2 | $e^+e^- \rightarrow \mu^+\mu^-$                             |
| 0.009 ± 0.006 ± 0.006              |     |      | <sup>2</sup> ALBRECHT 85 | ARG  | $e^+e^- \rightarrow \mu^+\mu^-$                             |
| 0.018 ± 0.008 ± 0.005              |     |      | HAAS 84B                 | CLEO | $e^+e^- \rightarrow \mu^+\mu^-$                             |

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.038      90      NICZYPORUK 81C LENA  $e^+e^- \rightarrow \mu^+\mu^-$

<sup>1</sup> Taking into account interference between the resonance and continuum.

<sup>2</sup> Re-evaluated using  $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 0.026$ .



**$\Gamma(\tau^+ \tau^-) / \Gamma(\mu^+ \mu^-)$   $\Gamma_3 / \Gamma_4$**

| VALUE                                      | EVTS | DOCUMENT ID | TECN | COMMENT                            |
|--------------------------------------------|------|-------------|------|------------------------------------|
| <b><math>1.04 \pm 0.04 \pm 0.05</math></b> | 22k  | BESSON 07   | CLEO | $e^+ e^- \rightarrow \Upsilon(2S)$ |

**$\Gamma(\Upsilon(1S)\pi^0) / \Gamma_{\text{total}}$   $\Gamma_6 / \Gamma$**

| VALUE (units $10^{-5}$ ) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|---------|
|--------------------------|-----|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

|       |    |                         |      |                                                   |
|-------|----|-------------------------|------|---------------------------------------------------|
| < 4   | 90 | <sup>1</sup> TAMPONI 13 | BELL | $e^+ e^- \rightarrow \Upsilon(1S)\pi^0$           |
| < 18  | 90 | <sup>2</sup> HE 08A     | CLEO | $e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$ |
| < 110 | 90 | ALEXANDER 98            | CLE2 | $e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$ |
| < 800 | 90 | LURZ 87                 | CBAL | $e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$ |

<sup>1</sup>TAMPONI 13 reports  $[\Gamma(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^0) / \Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-)] < 2.3 \times 10^{-4}$  which we multiply by our best value  $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-) = 17.85 \times 10^{-2}$ .

<sup>2</sup>Authors assume  $B(\Upsilon(1S) \rightarrow e^+ e^-) + B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = 4.96\%$ .

**$\Gamma(\Upsilon(1S)\pi^0) / \Gamma(\Upsilon(1S)\pi^+\pi^-)$   $\Gamma_6 / \Gamma_1$**

| VALUE (units $10^{-4}$ ) | CL% | DOCUMENT ID | TECN | COMMENT                                 |
|--------------------------|-----|-------------|------|-----------------------------------------|
| <b>&lt; 2.3</b>          | 90  | TAMPONI 13  | BELL | $e^+ e^- \rightarrow \Upsilon(1S)\pi^0$ |

$\Gamma(\Upsilon(1S)\eta)/\Gamma_{\text{total}}$

$\Gamma_7/\Gamma$

VALUE (units  $10^{-4}$ ) CL% EVTS DOCUMENT ID TECN COMMENT

**2.9 ± 0.4 OUR FIT** Error includes scale factor of 2.0.  
**2.9 ± 0.4 OUR AVERAGE** Error includes scale factor of 1.9. See the ideogram below.

|                                           |     |                   |     |      |                                               |
|-------------------------------------------|-----|-------------------|-----|------|-----------------------------------------------|
| 2.39 ± 0.31 ± 0.14                        | 112 | <sup>1</sup> LEES | 11L | BABR | $\Upsilon(2S) \rightarrow \ell^+ \ell^- \eta$ |
| 2.1 <sup>+0.7</sup> <sub>-0.6</sub> ± 0.3 | 14  | <sup>2</sup> HE   | 08A | CLEO | $e^+ e^- \rightarrow \ell^+ \ell^- \eta$      |

• • • We use the following data for averages but not for fits. • • •

|                    |     |                      |    |      |                                        |
|--------------------|-----|----------------------|----|------|----------------------------------------|
| 3.55 ± 0.32 ± 0.05 | 241 | <sup>3</sup> TAMPONI | 13 | BELL | $e^+ e^- \rightarrow \Upsilon(1S)\eta$ |
|--------------------|-----|----------------------|----|------|----------------------------------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

|       |    |                       |      |      |                                                                       |
|-------|----|-----------------------|------|------|-----------------------------------------------------------------------|
| < 9   | 90 | <sup>1,4</sup> AUBERT | 08BP | BABR | $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \pi^0 \ell^+ \ell^-$          |
| < 28  | 90 | ALEXANDER             | 98   | CLE2 | $e^+ e^- \rightarrow \ell^+ \ell^- \eta$                              |
| < 50  | 90 | ALBRECHT              | 87   | ARG  | $e^+ e^- \rightarrow \pi^+ \pi^- \ell^+ \ell^- \text{MM}$             |
| < 70  | 90 | LURZ                  | 87   | CBAL | $e^+ e^- \rightarrow \ell^+ \ell^- (\gamma\gamma, 3\pi^0)$            |
| < 100 | 90 | BESSON                | 84   | CLEO | $e^+ e^- \rightarrow \pi^+ \pi^- \ell^+ \ell^- \text{MM}$             |
| < 20  | 90 | FONSECA               | 84   | CUSB | $e^+ e^- \rightarrow \ell^+ \ell^- (\gamma\gamma, \pi^+ \pi^- \pi^0)$ |


<sup>1</sup> Using  $B(\Upsilon(1S) \rightarrow e^+ e^-) = (2.38 \pm 0.11)\%$  and  $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05)\%$ .

<sup>2</sup> Authors assume  $B(\Upsilon(1S) \rightarrow e^+ e^-) + B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = 4.96\%$ .

<sup>3</sup> TAMPONI 13 reports  $[\Gamma(\Upsilon(2S) \rightarrow \Upsilon(1S)\eta)/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+ \pi^-)] = (1.99 \pm 0.14 \pm 0.11) \times 10^{-3}$  which we multiply by our best value  $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+ \pi^-) = (17.85 \pm 0.26) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>4</sup> Using  $\Gamma_{ee}(\Upsilon(2S)) = 0.612 \pm 0.011$  keV.

WEIGHTED AVERAGE  
 2.9±0.4 (Error scaled by 1.9)





$\Gamma(\Upsilon(1S)\eta)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$   $\Gamma_7/\Gamma_1$

| VALUE (units $10^{-3}$ ) | CL% EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|----------|-------------|------|---------|
|--------------------------|----------|-------------|------|---------|

**1.64±0.25 OUR FIT** Error includes scale factor of 2.0.

**1.99±0.14±0.11** 241 TAMPONI 13 BELL  $e^+e^- \rightarrow \Upsilon(1S)\eta$

••• We do not use the following data for averages, fits, limits, etc. •••

1.35±0.17±0.08 <sup>1</sup> LEES 11L BABR  $\Upsilon(2S) \rightarrow (\pi^+\pi^-)(\gamma\gamma)\mu^+\mu^-$

< 5.2 90 <sup>2</sup> AUBERT 08BP BABR  $e^+e^- \rightarrow \gamma\pi^+\pi^-(\pi^0)\ell^+\ell^-$

<sup>1</sup> Not independent of other values reported by LEES 11L.

<sup>2</sup> Not independent of other values reported by AUBERT 08BP.

$\Gamma(\Upsilon(1S)\pi^0)/\Gamma(\Upsilon(1S)\eta)$   $\Gamma_6/\Gamma_7$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

••• We do not use the following data for averages, fits, limits, etc. •••

<0.13 90 TAMPONI 13 BELL  $e^+e^- \rightarrow \Upsilon(1S)\pi^0$

$\Gamma(J/\psi(1S) \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_8/\Gamma$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

<0.006 90 MASCHMANN 90 CBAL  $e^+e^- \rightarrow \text{hadrons}$

$\Gamma(J/\psi(1S)\eta_c)/\Gamma_{\text{total}}$   $\Gamma_9/\Gamma$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

<5.4 × 10<sup>-6</sup> 90 YANG 14 BELL  $e^+e^- \rightarrow J/\psi X$

$\Gamma(J/\psi(1S)\chi_{c0})/\Gamma_{\text{total}}$   $\Gamma_{10}/\Gamma$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

<3.4 × 10<sup>-6</sup> 90 YANG 14 BELL  $e^+e^- \rightarrow J/\psi X$

$\Gamma(J/\psi(1S)\chi_{c1})/\Gamma_{\text{total}}$   $\Gamma_{11}/\Gamma$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

<1.2 × 10<sup>-6</sup> 90 YANG 14 BELL  $e^+e^- \rightarrow J/\psi X$

$\Gamma(J/\psi(1S)\chi_{c2})/\Gamma_{\text{total}}$   $\Gamma_{12}/\Gamma$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

<2.0 × 10<sup>-6</sup> 90 YANG 14 BELL  $e^+e^- \rightarrow J/\psi X$

$\Gamma(J/\psi(1S)\eta_c(2S))/\Gamma_{\text{total}}$   $\Gamma_{13}/\Gamma$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

<2.5 × 10<sup>-6</sup> 90 YANG 14 BELL  $e^+e^- \rightarrow J/\psi X$

$\Gamma(J/\psi(1S)X(3940))/\Gamma_{\text{total}}$   $\Gamma_{14}/\Gamma$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

<2.0 × 10<sup>-6</sup> 90 YANG 14 BELL  $e^+e^- \rightarrow J/\psi X$

$\Gamma(J/\psi(1S)X(4160))/\Gamma_{\text{total}}$   $\Gamma_{15}/\Gamma$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

<2.0 × 10<sup>-6</sup> 90 YANG 14 BELL  $e^+e^- \rightarrow J/\psi X$

$\Gamma(\chi_{c1} \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_{16}/\Gamma$

| VALUE (units $10^{-4}$ )                   | EVTS | DOCUMENT ID | TECN | COMMENT                                           |
|--------------------------------------------|------|-------------|------|---------------------------------------------------|
| <b><math>2.24 \pm 0.44 \pm 0.20</math></b> | 376  | JIA         | 17   | BELL $\Upsilon(2S) \rightarrow \gamma J/\psi(1S)$ |

$\Gamma(\chi_{c1}(1P)^0 X_{tetra})/\Gamma_{\text{total}}$   $\Gamma_{17}/\Gamma$

| VALUE                                        | CL% | DOCUMENT ID      | TECN | COMMENT                                   |
|----------------------------------------------|-----|------------------|------|-------------------------------------------|
| <b><math>&lt; 36.7 \times 10^{-6}</math></b> | 90  | <sup>1</sup> JIA | 17A  | BELL $e^+ e^- \rightarrow \text{hadrons}$ |

<sup>1</sup> For a tetraquark state  $X_{tetra}$ , with mass in the range 1.16–2.46 GeV and width in the range 0–0.3 GeV. Measured 90% CL limits as a function of  $X_{tetra}$  mass and width range from  $4.4 \times 10^{-6}$  to  $36.7 \times 10^{-6}$ .

$\Gamma(\chi_{c2} \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_{18}/\Gamma$

| VALUE (units $10^{-4}$ )                   | DOCUMENT ID | TECN | COMMENT                                           |
|--------------------------------------------|-------------|------|---------------------------------------------------|
| <b><math>2.28 \pm 0.73 \pm 0.34</math></b> | JIA         | 17   | BELL $\Upsilon(2S) \rightarrow \gamma J/\psi(1S)$ |

$\Gamma(\psi(2S)\eta_c)/\Gamma_{\text{total}}$   $\Gamma_{19}/\Gamma$

| VALUE                                       | CL% | DOCUMENT ID | TECN | COMMENT                               |
|---------------------------------------------|-----|-------------|------|---------------------------------------|
| <b><math>&lt; 5.1 \times 10^{-6}</math></b> | 90  | YANG        | 14   | BELL $e^+ e^- \rightarrow \psi(2S) X$ |

$\Gamma(\psi(2S)\chi_{c0})/\Gamma_{\text{total}}$   $\Gamma_{20}/\Gamma$

| VALUE                                       | CL% | DOCUMENT ID | TECN | COMMENT                               |
|---------------------------------------------|-----|-------------|------|---------------------------------------|
| <b><math>&lt; 4.7 \times 10^{-6}</math></b> | 90  | YANG        | 14   | BELL $e^+ e^- \rightarrow \psi(2S) X$ |

$\Gamma(\psi(2S)\chi_{c1})/\Gamma_{\text{total}}$   $\Gamma_{21}/\Gamma$

| VALUE                                       | CL% | DOCUMENT ID | TECN | COMMENT                               |
|---------------------------------------------|-----|-------------|------|---------------------------------------|
| <b><math>&lt; 2.5 \times 10^{-6}</math></b> | 90  | YANG        | 14   | BELL $e^+ e^- \rightarrow \psi(2S) X$ |

$\Gamma(\psi(2S)\chi_{c2})/\Gamma_{\text{total}}$   $\Gamma_{22}/\Gamma$

| VALUE                                       | CL% | DOCUMENT ID | TECN | COMMENT                               |
|---------------------------------------------|-----|-------------|------|---------------------------------------|
| <b><math>&lt; 1.9 \times 10^{-6}</math></b> | 90  | YANG        | 14   | BELL $e^+ e^- \rightarrow \psi(2S) X$ |

$\Gamma(\psi(2S)\eta_c(2S))/\Gamma_{\text{total}}$   $\Gamma_{23}/\Gamma$

| VALUE                                       | CL% | DOCUMENT ID | TECN | COMMENT                               |
|---------------------------------------------|-----|-------------|------|---------------------------------------|
| <b><math>&lt; 3.3 \times 10^{-6}</math></b> | 90  | YANG        | 14   | BELL $e^+ e^- \rightarrow \psi(2S) X$ |

$\Gamma(\psi(2S)X(3940))/\Gamma_{\text{total}}$   $\Gamma_{24}/\Gamma$

| VALUE                                       | CL% | DOCUMENT ID | TECN | COMMENT                               |
|---------------------------------------------|-----|-------------|------|---------------------------------------|
| <b><math>&lt; 3.9 \times 10^{-6}</math></b> | 90  | YANG        | 14   | BELL $e^+ e^- \rightarrow \psi(2S) X$ |

$\Gamma(\psi(2S)X(4160))/\Gamma_{\text{total}}$   $\Gamma_{25}/\Gamma$

| VALUE                                       | CL% | DOCUMENT ID | TECN | COMMENT                               |
|---------------------------------------------|-----|-------------|------|---------------------------------------|
| <b><math>&lt; 3.9 \times 10^{-6}</math></b> | 90  | YANG        | 14   | BELL $e^+ e^- \rightarrow \psi(2S) X$ |

$\Gamma(T_{c\bar{c}1}(3900)^+ T_{c\bar{c}1}(3900)^-)/\Gamma_{\text{total}}$   $\Gamma_{26}/\Gamma$

| VALUE                                       | CL% | DOCUMENT ID      | TECN | COMMENT                                          |
|---------------------------------------------|-----|------------------|------|--------------------------------------------------|
| <b><math>&lt; 1.0 \times 10^{-6}</math></b> | 90  | <sup>1</sup> JIA | 18   | BELL $\Upsilon(2S) \rightarrow J/\psi \pi^\pm X$ |

<sup>1</sup> Assuming  $B(T_{c\bar{c}1}(3900)^\pm \rightarrow J/\psi \pi^\pm) = 1$ .

$\Gamma(T_{c\bar{c}1}(4200)^+ T_{c\bar{c}1}(4200)^-)/\Gamma_{\text{total}}$   $\Gamma_{27}/\Gamma$

| VALUE                  | CL% | DOCUMENT ID      | TECN | COMMENT                                          |
|------------------------|-----|------------------|------|--------------------------------------------------|
| $<16.7 \times 10^{-6}$ | 90  | <sup>1</sup> JIA | 18   | BELL $\Upsilon(1S) \rightarrow J/\psi \pi^\pm X$ |

<sup>1</sup> Assuming  $B(T_{c\bar{c}1}(4200)^\pm \rightarrow J/\psi \pi^\pm) = 1$

$\Gamma(T_{c\bar{c}1}(3900)^\pm T_{c\bar{c}1}(4200)^\mp)/\Gamma_{\text{total}}$   $\Gamma_{28}/\Gamma$

| VALUE                 | CL% | DOCUMENT ID      | TECN | COMMENT                                          |
|-----------------------|-----|------------------|------|--------------------------------------------------|
| $<7.3 \times 10^{-6}$ | 90  | <sup>1</sup> JIA | 18   | BELL $\Upsilon(2S) \rightarrow J/\psi \pi^\pm X$ |

<sup>1</sup> Assuming  $B(T_{c\bar{c}1}(4200)^\pm \rightarrow J/\psi \pi^\pm) = 1 = B(T_{c\bar{c}1}(3900)^\pm \rightarrow J/\psi \pi^\pm)$ .

$\Gamma(T_{c\bar{c}}(4050)^+ T_{c\bar{c}}(4050)^-)/\Gamma_{\text{total}}$   $\Gamma_{29}/\Gamma$

| VALUE                  | CL% | DOCUMENT ID      | TECN | COMMENT                                                 |
|------------------------|-----|------------------|------|---------------------------------------------------------|
| $<13.5 \times 10^{-6}$ | 90  | <sup>1</sup> JIA | 18   | BELL $\Upsilon(2S) \rightarrow \chi_{c1}(1P) \pi^\pm X$ |

<sup>1</sup> Assuming  $B(T_{c\bar{c}}(4050)^\pm \rightarrow \chi_{c1}(1P) \pi^\pm)$

$\Gamma(T_{c\bar{c}}(4250)^+ T_{c\bar{c}}(4250)^-)/\Gamma_{\text{total}}$   $\Gamma_{30}/\Gamma$

| VALUE                  | CL% | DOCUMENT ID      | TECN | COMMENT                                                 |
|------------------------|-----|------------------|------|---------------------------------------------------------|
| $<26.7 \times 10^{-6}$ | 90  | <sup>1</sup> JIA | 18   | BELL $\Upsilon(2S) \rightarrow \chi_{c1}(1P) \pi^\pm X$ |

<sup>1</sup> Assuming  $B(T_{c\bar{c}}(4250)^\pm \rightarrow \chi_{c1}(1P) \pi^\pm) = 1$

$\Gamma(T_{c\bar{c}}(4050)^\pm T_{c\bar{c}}(4250)^\mp)/\Gamma_{\text{total}}$   $\Gamma_{31}/\Gamma$

| VALUE                  | CL% | DOCUMENT ID      | TECN | COMMENT                                                 |
|------------------------|-----|------------------|------|---------------------------------------------------------|
| $<27.2 \times 10^{-6}$ | 90  | <sup>1</sup> JIA | 18   | BELL $\Upsilon(2S) \rightarrow \chi_{c1}(1P) \pi^\pm X$ |

<sup>1</sup> Assuming  $B(T_{c\bar{c}}(4050)^\pm \rightarrow \chi_{c1}(1P) \pi^\pm) = 1 = B(T_{c\bar{c}}(4250)^\pm \rightarrow \chi_{c1}(1P) \pi^\pm)$

$\Gamma(T_{c\bar{c}1}(4430)^+ T_{c\bar{c}1}(4430)^-)/\Gamma_{\text{total}}$   $\Gamma_{32}/\Gamma$

| VALUE                  | CL% | DOCUMENT ID      | TECN | COMMENT                                            |
|------------------------|-----|------------------|------|----------------------------------------------------|
| $<20.3 \times 10^{-6}$ | 90  | <sup>1</sup> JIA | 18   | BELL $\Upsilon(2S) \rightarrow \psi(2S) \pi^\pm X$ |

<sup>1</sup> Assuming  $B(T_{c\bar{c}1}(4430)^\pm \rightarrow \psi(2P) \pi^\pm) = 1$

$\Gamma(T_{c\bar{c}}(4055)^\pm T_{c\bar{c}}(4055)^\mp)/\Gamma_{\text{total}}$   $\Gamma_{33}/\Gamma$

| VALUE                  | CL% | DOCUMENT ID      | TECN | COMMENT                                            |
|------------------------|-----|------------------|------|----------------------------------------------------|
| $<11.1 \times 10^{-6}$ | 90  | <sup>1</sup> JIA | 18   | BELL $\Upsilon(2S) \rightarrow \psi(2S) \pi^\pm X$ |

<sup>1</sup> Assuming  $B(T_{c\bar{c}}(4055)^\pm \rightarrow \psi(2S) \pi^\pm) = 1$

$\Gamma(T_{c\bar{c}}(4055)^\pm T_{c\bar{c}1}(4430)^\mp)/\Gamma_{\text{total}}$   $\Gamma_{34}/\Gamma$

| VALUE                  | CL% | DOCUMENT ID      | TECN | COMMENT                                            |
|------------------------|-----|------------------|------|----------------------------------------------------|
| $<21.1 \times 10^{-6}$ | 90  | <sup>1</sup> JIA | 18   | BELL $\Upsilon(2S) \rightarrow \psi(2S) \pi^\pm X$ |

<sup>1</sup> Assuming  $B(T_{c\bar{c}}(4055)^\pm \rightarrow \psi(2S) \pi^\pm) = 1 = B(T_{c\bar{c}1}(4430)^\pm \rightarrow \psi(2S) \pi^\pm)$

$\Gamma(\overline{2H} \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_{35}/\Gamma$

| VALUE (units $10^{-5}$ ) | EVTS | DOCUMENT ID | TECN | COMMENT                                                |
|--------------------------|------|-------------|------|--------------------------------------------------------|
| $2.78^{+0.30}_{-0.26}$   |      |             |      | <b>OUR AVERAGE</b> Error includes scale factor of 1.2. |

2.64 ± 0.11 <sup>+0.26</sup>/<sub>-0.21</sub> LEES 14G BABR  $e^+e^- \rightarrow \overline{2H} X$

3.37 ± 0.50 ± 0.25 58 ASNER 07 CLEO  $e^+e^- \rightarrow \overline{2H} X$

**$\Gamma(g g g)/\Gamma_{\text{total}}$**   **$\Gamma_{37}/\Gamma$**

| <u>VALUE (units <math>10^{-2}</math>)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u>  | <u>TECN</u> | <u>COMMENT</u>                            |
|-------------------------------------------|-------------|---------------------|-------------|-------------------------------------------|
| <b>58.8±1.2</b>                           | 6M          | <sup>1</sup> BESSON | 06A CLEO    | $\Upsilon(2S) \rightarrow \text{hadrons}$ |

<sup>1</sup> Calculated using the value  $\Gamma(\gamma g g)/\Gamma(g g g) = (3.18 \pm 0.04 \pm 0.22 \pm 0.41)\%$  from BESSON 06A and PDG 08 values of  $B(\pi^+ \pi^- \Upsilon(1S)) = (18.1 \pm 0.4)\%$ ,  $B(\pi^0 \pi^0 \Upsilon(1S)) = (8.6 \pm 0.4)\%$ ,  $B(\mu^+ \mu^-) = (1.93 \pm 0.17)\%$ , and  $R_{\text{hadrons}} = 3.51$ . The statistical error is negligible and the systematic error is partially correlated with that of  $\Gamma(\gamma g g)/\Gamma_{\text{total}}$  measurement of BESSON 06A.

**$\Gamma(\gamma g g)/\Gamma(g g g)$**   **$\Gamma_{38}/\Gamma_{37}$**

| <u>VALUE (units <math>10^{-2}</math>)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                                       |
|-------------------------------------------|-------------|--------------------|-------------|------------------------------------------------------|
| <b>3.18±0.04±0.47</b>                     | 6M          | BESSON             | 06A CLEO    | $\Upsilon(2S) \rightarrow (\gamma +) \text{hadrons}$ |

**$\Gamma(\phi K^+ K^-)/\Gamma_{\text{total}}$**   **$\Gamma_{39}/\Gamma$**

| <u>VALUE (units <math>10^{-6}</math>)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                        |
|-------------------------------------------|-------------|--------------------|-------------|---------------------------------------|
| <b>1.58±0.33±0.18</b>                     | 58          | SHEN               | 12A BELL    | $\Upsilon(1S) \rightarrow 2(K^+ K^-)$ |

**$\Gamma(\omega \pi^+ \pi^-)/\Gamma_{\text{total}}$**   **$\Gamma_{40}/\Gamma$**

| <u>VALUE (units <math>10^{-6}</math>)</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                                  |
|-------------------------------------------|------------|--------------------|-------------|-------------------------------------------------|
| <b>&lt;2.58</b>                           | 90         | SHEN               | 12A BELL    | $\Upsilon(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$ |

**$\Gamma(K^*(892)^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$**   **$\Gamma_{41}/\Gamma$**

| <u>VALUE (units <math>10^{-6}</math>)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                                 |
|-------------------------------------------|-------------|--------------------|-------------|------------------------------------------------|
| <b>2.32±0.40±0.54</b>                     | 135         | SHEN               | 12A BELL    | $\Upsilon(1S) \rightarrow K^+ K^- \pi^+ \pi^-$ |

**$\Gamma(\phi f'_2(1525))/\Gamma_{\text{total}}$**   **$\Gamma_{42}/\Gamma$**

| <u>VALUE (units <math>10^{-6}</math>)</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                        |
|-------------------------------------------|------------|--------------------|-------------|---------------------------------------|
| <b>&lt;1.33</b>                           | 90         | SHEN               | 12A BELL    | $\Upsilon(1S) \rightarrow 2(K^+ K^-)$ |

**$\Gamma(\omega f_2(1270))/\Gamma_{\text{total}}$**   **$\Gamma_{43}/\Gamma$**

| <u>VALUE (units <math>10^{-6}</math>)</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                                  |
|-------------------------------------------|------------|--------------------|-------------|-------------------------------------------------|
| <b>&lt;0.57</b>                           | 90         | SHEN               | 12A BELL    | $\Upsilon(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$ |

**$\Gamma(\rho(770) a_2(1320))/\Gamma_{\text{total}}$**   **$\Gamma_{44}/\Gamma$**

| <u>VALUE (units <math>10^{-6}</math>)</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                                  |
|-------------------------------------------|------------|--------------------|-------------|-------------------------------------------------|
| <b>&lt;0.88</b>                           | 90         | SHEN               | 12A BELL    | $\Upsilon(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$ |

**$\Gamma(K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.})/\Gamma_{\text{total}}$**   **$\Gamma_{45}/\Gamma$**

| <u>VALUE (units <math>10^{-6}</math>)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                                 |
|-------------------------------------------|-------------|--------------------|-------------|------------------------------------------------|
| <b>1.53±0.52±0.19</b>                     | 32          | SHEN               | 12A BELL    | $\Upsilon(1S) \rightarrow K^+ K^- \pi^+ \pi^-$ |

**$\Gamma(K_1(1270)^\pm K^\mp)/\Gamma_{\text{total}}$**   **$\Gamma_{46}/\Gamma$**

| <u>VALUE (units <math>10^{-6}</math>)</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                                 |
|-------------------------------------------|------------|--------------------|-------------|------------------------------------------------|
| <b>&lt;3.22</b>                           | 90         | SHEN               | 12A BELL    | $\Upsilon(1S) \rightarrow K^+ K^- \pi^+ \pi^-$ |

| $\Gamma(K_1(1400)^\pm K^\mp)/\Gamma_{\text{total}}$                |              |                                                                                              |                                                                           |                |                                                    | $\Gamma_{47}/\Gamma$                       |
|--------------------------------------------------------------------|--------------|----------------------------------------------------------------------------------------------|---------------------------------------------------------------------------|----------------|----------------------------------------------------|--------------------------------------------|
| <u>VALUE (units <math>10^{-6}</math>)</u>                          | <u>CL%</u>   | <u>DOCUMENT ID</u>                                                                           | <u>TECN</u>                                                               | <u>COMMENT</u> |                                                    |                                            |
| <b>&lt;0.83</b>                                                    | 90           | SHEN                                                                                         | 12A                                                                       | BELL           | $\Upsilon(1S) \rightarrow K^+ K^- \pi^+ \pi^-$     |                                            |
| $\Gamma(b_1(1235)^\pm \pi^\mp)/\Gamma_{\text{total}}$              |              |                                                                                              |                                                                           |                |                                                    | $\Gamma_{48}/\Gamma$                       |
| <u>VALUE (units <math>10^{-6}</math>)</u>                          | <u>CL%</u>   | <u>DOCUMENT ID</u>                                                                           | <u>TECN</u>                                                               | <u>COMMENT</u> |                                                    |                                            |
| <b>&lt;0.40</b>                                                    | 90           | SHEN                                                                                         | 12A                                                                       | BELL           | $\Upsilon(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$    |                                            |
| $\Gamma(\rho\pi)/\Gamma_{\text{total}}$                            |              |                                                                                              |                                                                           |                |                                                    | $\Gamma_{49}/\Gamma$                       |
| <u>VALUE (units <math>10^{-6}</math>)</u>                          | <u>CL%</u>   | <u>DOCUMENT ID</u>                                                                           | <u>TECN</u>                                                               | <u>COMMENT</u> |                                                    |                                            |
| <b>&lt;1.16</b>                                                    | 90           | SHEN                                                                                         | 13                                                                        | BELL           | $\Upsilon(2S) \rightarrow \pi^+ \pi^- \pi^0$       |                                            |
| $\Gamma(\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$                  |              |                                                                                              |                                                                           |                |                                                    | $\Gamma_{50}/\Gamma$                       |
| <u>VALUE (units <math>10^{-6}</math>)</u>                          | <u>CL%</u>   | <u>DOCUMENT ID</u>                                                                           | <u>TECN</u>                                                               | <u>COMMENT</u> |                                                    |                                            |
| <b>&lt;0.80</b>                                                    | 90           | SHEN                                                                                         | 13                                                                        | BELL           | $\Upsilon(2S) \rightarrow \pi^+ \pi^- \pi^0$       |                                            |
| $\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$                        |              |                                                                                              |                                                                           |                |                                                    | $\Gamma_{51}/\Gamma$                       |
| <u>VALUE (units <math>10^{-6}</math>)</u>                          | <u>CL%</u>   | <u>DOCUMENT ID</u>                                                                           | <u>TECN</u>                                                               | <u>COMMENT</u> |                                                    |                                            |
| <b>&lt;1.63</b>                                                    | 90           | SHEN                                                                                         | 13                                                                        | BELL           | $\Upsilon(2S) \rightarrow \pi^+ \pi^- \pi^0 \pi^0$ |                                            |
| $\Gamma(\pi^+ \pi^- \pi^0 \pi^0)/\Gamma_{\text{total}}$            |              |                                                                                              |                                                                           |                |                                                    | $\Gamma_{52}/\Gamma$                       |
| <u>VALUE (units <math>10^{-6}</math>)</u>                          | <u>EVTS</u>  | <u>DOCUMENT ID</u>                                                                           | <u>TECN</u>                                                               | <u>COMMENT</u> |                                                    |                                            |
| <b><math>13.0 \pm 1.9 \pm 2.1</math></b>                           | $261 \pm 37$ | SHEN                                                                                         | 13                                                                        | BELL           | $\Upsilon(2S) \rightarrow \pi^+ \pi^- \pi^0 \pi^0$ |                                            |
| $\Gamma(K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$      |              |                                                                                              |                                                                           |                |                                                    | $\Gamma_{53}/\Gamma$                       |
| <u>VALUE (units <math>10^{-6}</math>)</u>                          | <u>CL%</u>   | <u>EVTS</u>                                                                                  | <u>DOCUMENT ID</u>                                                        | <u>TECN</u>    | <u>COMMENT</u>                                     |                                            |
| <b><math>1.14 \pm 0.30 \pm 0.13</math></b>                         | $40 \pm 10$  |                                                                                              | SHEN                                                                      | 13             | BELL                                               | $\Upsilon(2S) \rightarrow K_S^0 K^- \pi^+$ |
| <b>•••</b>                                                         |              |                                                                                              | ••• We do not use the following data for averages, fits, limits, etc. ••• |                |                                                    |                                            |
| <b>&lt;3.2</b>                                                     | 90           | <sup>1</sup> DOBBS                                                                           | 12A                                                                       |                |                                                    | $\Upsilon(2S) \rightarrow K_S^0 K^- \pi^+$ |
|                                                                    |              | <sup>1</sup> Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration. |                                                                           |                |                                                    |                                            |
| $\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ |              |                                                                                              |                                                                           |                |                                                    | $\Gamma_{54}/\Gamma$                       |
| <u>VALUE (units <math>10^{-6}</math>)</u>                          | <u>CL%</u>   | <u>DOCUMENT ID</u>                                                                           | <u>TECN</u>                                                               | <u>COMMENT</u> |                                                    |                                            |
| <b>&lt;4.22</b>                                                    | 90           | SHEN                                                                                         | 13                                                                        | BELL           | $\Upsilon(2S) \rightarrow K_S^0 K^- \pi^+$         |                                            |
| $\Gamma(K^*(892)^- K^+ + \text{c.c.})/\Gamma_{\text{total}}$       |              |                                                                                              |                                                                           |                |                                                    | $\Gamma_{55}/\Gamma$                       |
| <u>VALUE (units <math>10^{-6}</math>)</u>                          | <u>CL%</u>   | <u>DOCUMENT ID</u>                                                                           | <u>TECN</u>                                                               | <u>COMMENT</u> |                                                    |                                            |
| <b>&lt;1.45</b>                                                    | 90           | SHEN                                                                                         | 13                                                                        | BELL           | $\Upsilon(2S) \rightarrow K_S^0 K^- \pi^+$         |                                            |
| $\Gamma(f_1(1285) \text{ anything})/\Gamma_{\text{total}}$         |              |                                                                                              |                                                                           |                |                                                    | $\Gamma_{56}/\Gamma$                       |
| <u>VALUE (units <math>10^{-3}</math>)</u>                          | <u>EVTS</u>  | <u>DOCUMENT ID</u>                                                                           | <u>TECN</u>                                                               | <u>COMMENT</u> |                                                    |                                            |
| <b><math>2.20 \pm 1.50 \pm 0.63</math></b>                         | 2.9k         | JIA                                                                                          | 17A                                                                       | BELL           | $e^+ e^- \rightarrow \text{hadrons}$               |                                            |

$\Gamma(f_1(1285)X_{tetra})/\Gamma_{total}$   $\Gamma_{57}/\Gamma$

| VALUE                             | CL% | DOCUMENT ID      | TECN | COMMENT                                      |
|-----------------------------------|-----|------------------|------|----------------------------------------------|
| <b>&lt;64.7 × 10<sup>-6</sup></b> | 90  | <sup>1</sup> JIA | 17A  | BELL e <sup>+</sup> e <sup>-</sup> → hadrons |

<sup>1</sup> For a tetraquark state  $X_{tetra}$ , with mass in the range 1.16–2.46 GeV and width in the range 0–0.3 GeV. Measured 90% CL limits as a function of  $X_{tetra}$  mass and width range from  $7.8 \times 10^{-6}$  to  $64.7 \times 10^{-6}$ .

$\Gamma(D_s^+ D_{s1}(2536)^-, D_{s1}^- \rightarrow K^- D^*(2007)^0)/\Gamma_{total}$   $\Gamma_{59}/\Gamma$

| VALUE (units 10 <sup>-5</sup> ) | DOCUMENT ID | TECN | COMMENT                                         |
|---------------------------------|-------------|------|-------------------------------------------------|
| <b>1.6 ± 0.3 ± 0.2</b>          | GAO         | 23   | BELL e <sup>+</sup> e <sup>-</sup> at 10.52 GeV |

$\Gamma(D_s^+ D_{s1}(2536)^-, D_{s1}^- \rightarrow K_S^0 D^*(2010)^-)/\Gamma_{total}$   $\Gamma_{60}/\Gamma$

| VALUE (units 10 <sup>-5</sup> ) | DOCUMENT ID | TECN | COMMENT                                         |
|---------------------------------|-------------|------|-------------------------------------------------|
| <b>0.84 ± 0.18 ± 0.15</b>       | GAO         | 23   | BELL e <sup>+</sup> e <sup>-</sup> at 10.52 GeV |

$\Gamma(D_s^{*+} D_{s1}(2536)^-, D_{s1}^- \rightarrow K^- D^*(2007)^0)/\Gamma_{total}$   $\Gamma_{62}/\Gamma$

| VALUE (units 10 <sup>-5</sup> ) | DOCUMENT ID | TECN | COMMENT                                         |
|---------------------------------|-------------|------|-------------------------------------------------|
| <b>1.4 ± 0.4 ± 0.2</b>          | GAO         | 23   | BELL e <sup>+</sup> e <sup>-</sup> at 10.52 GeV |

$\Gamma(D_s^{*+} D_{s1}(2536)^-, D_{s1}^- \rightarrow K_S^0 D^*(2010)^-)/\Gamma_{total}$   $\Gamma_{63}/\Gamma$

| VALUE (units 10 <sup>-5</sup> ) | DOCUMENT ID | TECN | COMMENT                                         |
|---------------------------------|-------------|------|-------------------------------------------------|
| <b>0.82 ± 0.25 ± 0.19</b>       | GAO         | 23   | BELL e <sup>+</sup> e <sup>-</sup> at 10.52 GeV |

$\Gamma(D_s^+ D_{s2}^*(2573)^-, D_{s2}^{*-} \rightarrow K^- D^0)/\Gamma_{total}$   $\Gamma_{65}/\Gamma$

| VALUE (units 10 <sup>-5</sup> ) | DOCUMENT ID | TECN | COMMENT                                         |
|---------------------------------|-------------|------|-------------------------------------------------|
| <b>1.4 ± 0.4 ± 0.2</b>          | GAO         | 23   | BELL e <sup>+</sup> e <sup>-</sup> at 10.52 GeV |

$\Gamma(D_s^+ D_{s2}^*(2573)^-, D_{s2}^{*-} \rightarrow K_S^0 D^-)/\Gamma_{total}$   $\Gamma_{66}/\Gamma$

| VALUE (units 10 <sup>-5</sup> ) | DOCUMENT ID | TECN | COMMENT                                         |
|---------------------------------|-------------|------|-------------------------------------------------|
| <b>0.69 ± 0.20 ± 0.22</b>       | GAO         | 23   | BELL e <sup>+</sup> e <sup>-</sup> at 10.52 GeV |

$\Gamma(D_s^{*+} D_{s2}^*(2573)^-, D_{s2}^{*-} \rightarrow K^- D^0)/\Gamma_{total}$   $\Gamma_{68}/\Gamma$

| VALUE (units 10 <sup>-5</sup> ) | DOCUMENT ID | TECN | COMMENT                                         |
|---------------------------------|-------------|------|-------------------------------------------------|
| <b>0.9 ± 0.5 ± 0.2</b>          | GAO         | 23   | BELL e <sup>+</sup> e <sup>-</sup> at 10.52 GeV |

$\Gamma(D_s^{*+} D_{s2}^*(2573)^-, D_{s2}^{*-} \rightarrow K_S^0 D^-)/\Gamma_{total}$   $\Gamma_{69}/\Gamma$

| VALUE (units 10 <sup>-5</sup> ) | DOCUMENT ID | TECN | COMMENT                                         |
|---------------------------------|-------------|------|-------------------------------------------------|
| <b>0.54 ± 0.31 ± 0.47</b>       | GAO         | 23   | BELL e <sup>+</sup> e <sup>-</sup> at 10.52 GeV |

$\Gamma(\text{Sum of 100 exclusive modes})/\Gamma_{total}$   $\Gamma_{70}/\Gamma$

| VALUE (units 10 <sup>-2</sup> ) | DOCUMENT ID          | COMMENT                                |
|---------------------------------|----------------------|----------------------------------------|
| <b>0.29 ± 0.03</b>              | <sup>1,2</sup> DOBBS | 12A $\Upsilon(2S) \rightarrow$ hadrons |

<sup>1</sup> DOBBS 12A presents individual exclusive branching fractions or upper limits for 100 modes of four to ten pions, kaons, or protons.

<sup>2</sup> Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

| $\Gamma(\gamma\chi_{b1}(1P))/\Gamma_{\text{total}}$ |             |                    |             | $\Gamma_{71}/\Gamma$                       |
|-----------------------------------------------------|-------------|--------------------|-------------|--------------------------------------------|
| <u>VALUE</u>                                        | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                             |
| <b>0.069 ± 0.004 OUR AVERAGE</b>                    |             |                    |             |                                            |
| 0.0693 ± 0.0012 ± 0.0041                            | 407k        | ARTUSO             | 05 CLEO     | $e^+e^- \rightarrow \gamma X$              |
| 0.069 ± 0.005 ± 0.009                               |             | EDWARDS            | 99 CLE2     | $\Upsilon(2S) \rightarrow \gamma\chi(1P)$  |
| 0.091 ± 0.018 ± 0.022                               |             | ALBRECHT           | 85E ARG     | $e^+e^- \rightarrow \gamma\text{conv. } X$ |
| 0.065 ± 0.007 ± 0.012                               |             | NERNST             | 85 CBAL     | $e^+e^- \rightarrow \gamma X$              |
| 0.080 ± 0.017 ± 0.016                               |             | HAAS               | 84 CLEO     | $e^+e^- \rightarrow \gamma\text{conv. } X$ |
| 0.059 ± 0.014                                       |             | KLOPFEN...         | 83 CUSB     | $e^+e^- \rightarrow \gamma X$              |

| $\Gamma(\gamma\chi_{b2}(1P))/\Gamma_{\text{total}}$ |             |                    |             | $\Gamma_{72}/\Gamma$                       |
|-----------------------------------------------------|-------------|--------------------|-------------|--------------------------------------------|
| <u>VALUE</u>                                        | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                             |
| <b>0.0715 ± 0.0035 OUR AVERAGE</b>                  |             |                    |             |                                            |
| 0.0724 ± 0.0011 ± 0.0040                            | 410k        | ARTUSO             | 05 CLEO     | $e^+e^- \rightarrow \gamma X$              |
| 0.074 ± 0.005 ± 0.008                               |             | EDWARDS            | 99 CLE2     | $\Upsilon(2S) \rightarrow \gamma\chi(1P)$  |
| 0.098 ± 0.021 ± 0.024                               |             | ALBRECHT           | 85E ARG     | $e^+e^- \rightarrow \gamma\text{conv. } X$ |
| 0.058 ± 0.007 ± 0.010                               |             | NERNST             | 85 CBAL     | $e^+e^- \rightarrow \gamma X$              |
| 0.102 ± 0.018 ± 0.021                               |             | HAAS               | 84 CLEO     | $e^+e^- \rightarrow \gamma\text{conv. } X$ |
| 0.061 ± 0.014                                       |             | KLOPFEN...         | 83 CUSB     | $e^+e^- \rightarrow \gamma X$              |

| $\Gamma(\gamma\chi_{b0}(1P))/\Gamma_{\text{total}}$                           |             |                    |             | $\Gamma_{73}/\Gamma$                       |
|-------------------------------------------------------------------------------|-------------|--------------------|-------------|--------------------------------------------|
| <u>VALUE</u>                                                                  | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                             |
| <b>0.038 ± 0.004 OUR AVERAGE</b>                                              |             |                    |             |                                            |
| 0.0375 ± 0.0012 ± 0.0047                                                      | 198k        | ARTUSO             | 05 CLEO     | $e^+e^- \rightarrow \gamma X$              |
| 0.034 ± 0.005 ± 0.006                                                         |             | EDWARDS            | 99 CLE2     | $\Upsilon(2S) \rightarrow \gamma\chi(1P)$  |
| 0.064 ± 0.014 ± 0.016                                                         |             | ALBRECHT           | 85E ARG     | $e^+e^- \rightarrow \gamma\text{conv. } X$ |
| 0.036 ± 0.008 ± 0.009                                                         |             | NERNST             | 85 CBAL     | $e^+e^- \rightarrow \gamma X$              |
| 0.044 ± 0.023 ± 0.009                                                         |             | HAAS               | 84 CLEO     | $e^+e^- \rightarrow \gamma\text{conv. } X$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |             |                    |             |                                            |
| 0.035 ± 0.014                                                                 |             | KLOPFEN...         | 83 CUSB     | $e^+e^- \rightarrow \gamma X$              |

| $\Gamma(\gamma f_0(1710))/\Gamma_{\text{total}}$                                      |            |                       |             | $\Gamma_{74}/\Gamma$                        |
|---------------------------------------------------------------------------------------|------------|-----------------------|-------------|---------------------------------------------|
| <u>VALUE (units <math>10^{-5}</math>)</u>                                             | <u>CL%</u> | <u>DOCUMENT ID</u>    | <u>TECN</u> | <u>COMMENT</u>                              |
| <b>&lt;59</b>                                                                         | 90         | <sup>1</sup> ALBRECHT | 89 ARG      | $\Upsilon(2S) \rightarrow \gamma K^+ K^-$   |
| • • • We do not use the following data for averages, fits, limits, etc. • • •         |            |                       |             |                                             |
| < 5.9                                                                                 | 90         | <sup>2</sup> ALBRECHT | 89 ARG      | $\Upsilon(2S) \rightarrow \gamma\pi^+\pi^-$ |
| <sup>1</sup> Re-evaluated assuming $B(f_0(1710) \rightarrow K^+ K^-) = 0.19$ .        |            |                       |             |                                             |
| <sup>2</sup> Includes unknown branching ratio of $f_0(1710) \rightarrow \pi^+\pi^-$ . |            |                       |             |                                             |

| $\Gamma(\gamma f'_2(1525))/\Gamma_{\text{total}}$                                |            |                       |             | $\Gamma_{75}/\Gamma$                      |
|----------------------------------------------------------------------------------|------------|-----------------------|-------------|-------------------------------------------|
| <u>VALUE (units <math>10^{-5}</math>)</u>                                        | <u>CL%</u> | <u>DOCUMENT ID</u>    | <u>TECN</u> | <u>COMMENT</u>                            |
| <b>&lt;53</b>                                                                    | 90         | <sup>1</sup> ALBRECHT | 89 ARG      | $\Upsilon(2S) \rightarrow \gamma K^+ K^-$ |
| <sup>1</sup> Re-evaluated assuming $B(f'_2(1525) \rightarrow K\bar{K}) = 0.71$ . |            |                       |             |                                           |

| $\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$              |            |                       |             | $\Gamma_{76}/\Gamma$                        |
|---------------------------------------------------------------|------------|-----------------------|-------------|---------------------------------------------|
| <u>VALUE (units <math>10^{-5}</math>)</u>                     | <u>CL%</u> | <u>DOCUMENT ID</u>    | <u>TECN</u> | <u>COMMENT</u>                              |
| <b>&lt;24.1</b>                                               | 90         | <sup>1</sup> ALBRECHT | 89 ARG      | $\Upsilon(2S) \rightarrow \gamma\pi^+\pi^-$ |
| <sup>1</sup> Using $B(f_2(1270) \rightarrow \pi\pi) = 0.84$ . |            |                       |             |                                             |

**$\Gamma(\gamma f_J(2220))/\Gamma_{\text{total}}$**   **$\Gamma_{77}/\Gamma$**

| VALUE (units $10^{-5}$ ) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|---------|
|--------------------------|-----|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

|        |    |                       |    |                                               |
|--------|----|-----------------------|----|-----------------------------------------------|
| $<6.8$ | 90 | <sup>1</sup> ALBRECHT | 89 | ARG $\Upsilon(2S) \rightarrow \gamma K^+ K^-$ |
|--------|----|-----------------------|----|-----------------------------------------------|

<sup>1</sup> Includes unknown branching ratio of  $f_J(2220) \rightarrow K^+ K^-$ .

**$\Gamma(\gamma \eta_c(1S))/\Gamma_{\text{total}}$**   **$\Gamma_{78}/\Gamma$**

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

|                       |    |      |     |                                          |
|-----------------------|----|------|-----|------------------------------------------|
| $<2.7 \times 10^{-5}$ | 90 | WANG | 11B | BELL $\Upsilon(2S) \rightarrow \gamma X$ |
|-----------------------|----|------|-----|------------------------------------------|

**$\Gamma(\gamma \chi_{c0})/\Gamma_{\text{total}}$**   **$\Gamma_{79}/\Gamma$**

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

|                       |    |      |     |                                          |
|-----------------------|----|------|-----|------------------------------------------|
| $<1.0 \times 10^{-4}$ | 90 | WANG | 11B | BELL $\Upsilon(2S) \rightarrow \gamma X$ |
|-----------------------|----|------|-----|------------------------------------------|

**$\Gamma(\gamma \chi_{c1})/\Gamma_{\text{total}}$**   **$\Gamma_{80}/\Gamma$**

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

|                       |    |      |     |                                          |
|-----------------------|----|------|-----|------------------------------------------|
| $<3.6 \times 10^{-6}$ | 90 | WANG | 11B | BELL $\Upsilon(2S) \rightarrow \gamma X$ |
|-----------------------|----|------|-----|------------------------------------------|

**$\Gamma(\gamma \chi_{c2})/\Gamma_{\text{total}}$**   **$\Gamma_{81}/\Gamma$**

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

|                       |    |      |     |                                          |
|-----------------------|----|------|-----|------------------------------------------|
| $<1.5 \times 10^{-5}$ | 90 | WANG | 11B | BELL $\Upsilon(2S) \rightarrow \gamma X$ |
|-----------------------|----|------|-----|------------------------------------------|

**$\Gamma(\gamma \chi_{c1}(3872))/\Gamma_{\text{total}}$**   **$\Gamma_{82}/\Gamma$**

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

|                       |    |                   |     |                                          |
|-----------------------|----|-------------------|-----|------------------------------------------|
| $<2.3 \times 10^{-5}$ | 90 | <sup>1</sup> WANG | 11B | BELL $\Upsilon(2S) \rightarrow \gamma X$ |
|-----------------------|----|-------------------|-----|------------------------------------------|

<sup>1</sup> WANG 11B reports  $[\Gamma(\Upsilon(2S) \rightarrow \gamma \chi_{c1}(3872))/\Gamma_{\text{total}}] \times [B(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S))] < 0.8 \times 10^{-6}$  which we divide by our best value  $B(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S)) = 3.5 \times 10^{-2}$ .

**$\Gamma(\gamma \chi_{c1}(3872), \chi_{c1} \rightarrow \pi^+ \pi^- \pi^0 J/\psi)/\Gamma_{\text{total}}$**   **$\Gamma_{83}/\Gamma$**

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

|                       |    |      |     |                                          |
|-----------------------|----|------|-----|------------------------------------------|
| $<2.4 \times 10^{-6}$ | 90 | WANG | 11B | BELL $\Upsilon(2S) \rightarrow \gamma X$ |
|-----------------------|----|------|-----|------------------------------------------|

**$\Gamma(\gamma \chi_{c0}(3915) \rightarrow \omega J/\psi)/\Gamma_{\text{total}}$**   **$\Gamma_{84}/\Gamma$**

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

|                       |    |      |     |                                          |
|-----------------------|----|------|-----|------------------------------------------|
| $<2.8 \times 10^{-6}$ | 90 | WANG | 11B | BELL $\Upsilon(2S) \rightarrow \gamma X$ |
|-----------------------|----|------|-----|------------------------------------------|

**$\Gamma(\gamma \chi_{c1}(4140) \rightarrow \phi J/\psi)/\Gamma_{\text{total}}$**   **$\Gamma_{85}/\Gamma$**

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

|                       |    |      |     |                                          |
|-----------------------|----|------|-----|------------------------------------------|
| $<1.2 \times 10^{-6}$ | 90 | WANG | 11B | BELL $\Upsilon(2S) \rightarrow \gamma X$ |
|-----------------------|----|------|-----|------------------------------------------|

**$\Gamma(\gamma X(4350) \rightarrow \phi J/\psi)/\Gamma_{\text{total}}$**   **$\Gamma_{86}/\Gamma$**

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

|                       |    |      |     |                                          |
|-----------------------|----|------|-----|------------------------------------------|
| $<1.3 \times 10^{-6}$ | 90 | WANG | 11B | BELL $\Upsilon(2S) \rightarrow \gamma X$ |
|-----------------------|----|------|-----|------------------------------------------|



$\Gamma(\gamma\eta_b(1S))/\Gamma_{\text{total}}$   $\Gamma_{87}/\Gamma$

| VALUE (units $10^{-4}$ ) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|------|-------------|------|---------|
|--------------------------|-----|------|-------------|------|---------|

**5.5<sup>+1.1</sup><sub>-0.9</sub> OUR AVERAGE** Error includes scale factor of 1.2.

|                                             |  |     |        |    |                                          |
|---------------------------------------------|--|-----|--------|----|------------------------------------------|
| 6.1 <sup>+0.6+0.9</sup> <sub>-0.7-0.6</sub> |  | 29k | FULSOM | 18 | BELL $\Upsilon(2S) \rightarrow \gamma X$ |
|---------------------------------------------|--|-----|--------|----|------------------------------------------|

|                                               |  |             |                     |      |                                          |
|-----------------------------------------------|--|-------------|---------------------|------|------------------------------------------|
| 3.9 $\pm$ 1.1 <sup>+1.1</sup> <sub>-0.9</sub> |  | 13 $\pm$ 5k | <sup>1</sup> AUBERT | 09AQ | BABR $\Upsilon(2S) \rightarrow \gamma X$ |
|-----------------------------------------------|--|-------------|---------------------|------|------------------------------------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

|     |    |  |      |     |                                         |
|-----|----|--|------|-----|-----------------------------------------|
| <21 | 90 |  | LEES | 11J | BABR $\Upsilon(2S) \rightarrow X\gamma$ |
|-----|----|--|------|-----|-----------------------------------------|

|       |    |  |                        |    |                                          |
|-------|----|--|------------------------|----|------------------------------------------|
| < 8.4 | 90 |  | <sup>1</sup> BONVICINI | 10 | CLEO $\Upsilon(2S) \rightarrow \gamma X$ |
|-------|----|--|------------------------|----|------------------------------------------|

|       |    |  |                     |    |                                    |
|-------|----|--|---------------------|----|------------------------------------|
| < 5.1 | 90 |  | <sup>2</sup> ARTUSO | 05 | CLEO $e^+e^- \rightarrow \gamma X$ |
|-------|----|--|---------------------|----|------------------------------------|

<sup>1</sup> Assuming  $\Gamma_{\eta_b(1S)} = 10$  MeV.

<sup>2</sup> Superseded by BONVICINI 10.

$\Gamma(\gamma\eta_b(1S) \rightarrow \gamma \text{Sum of 26 exclusive modes})/\Gamma_{\text{total}}$   $\Gamma_{88}/\Gamma$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

|                       |    |          |    |                                                |
|-----------------------|----|----------|----|------------------------------------------------|
| <3.7 $\times 10^{-6}$ | 90 | SANDILYA | 13 | BELL $\Upsilon(2S) \rightarrow \gamma$ hadrons |
|-----------------------|----|----------|----|------------------------------------------------|

$\Gamma(\gamma X_{b\bar{b}} \rightarrow \gamma \text{Sum of 26 exclusive modes})/\Gamma_{\text{total}}$   $\Gamma_{89}/\Gamma$

| VALUE (units $10^{-6}$ ) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|------|-------------|------|---------|
|--------------------------|-----|------|-------------|------|---------|

|       |    |  |          |    |                                                |
|-------|----|--|----------|----|------------------------------------------------|
| < 4.9 | 90 |  | SANDILYA | 13 | BELL $\Upsilon(2S) \rightarrow \gamma$ hadrons |
|-------|----|--|----------|----|------------------------------------------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                                                   |  |    |                    |    |                                           |
|---------------------------------------------------|--|----|--------------------|----|-------------------------------------------|
| 46.2 <sup>+29.7</sup> <sub>-14.2</sub> $\pm$ 10.6 |  | 10 | <sup>1</sup> DOBBS | 12 | $\Upsilon(2S) \rightarrow \gamma$ hadrons |
|---------------------------------------------------|--|----|--------------------|----|-------------------------------------------|

<sup>1</sup> Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

$\Gamma(\gamma X \rightarrow \gamma + \geq 4 \text{ prongs})/\Gamma_{\text{total}}$   $\Gamma_{90}/\Gamma$   
 (1.5 GeV <  $m_X$  < 5.0 GeV)

| VALUE (units $10^{-4}$ ) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|---------|
|--------------------------|-----|-------------|------|---------|

|       |    |        |     |                                    |
|-------|----|--------|-----|------------------------------------|
| <1.95 | 95 | ROSNER | 07A | CLEO $e^+e^- \rightarrow \gamma X$ |
|-------|----|--------|-----|------------------------------------|

$\Gamma(\gamma A^0 \rightarrow \gamma \text{hadrons})/\Gamma_{\text{total}}$   $\Gamma_{91}/\Gamma$   
 (0.3 GeV <  $m_{A^0}$  < 7 GeV)

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

|                     |    |                   |     |                                                |
|---------------------|----|-------------------|-----|------------------------------------------------|
| <8 $\times 10^{-5}$ | 90 | <sup>1</sup> LEES | 11H | BABR $\Upsilon(2S) \rightarrow \gamma$ hadrons |
|---------------------|----|-------------------|-----|------------------------------------------------|

<sup>1</sup> For a narrow scalar or pseudoscalar,  $A^0$ , excluding known resonances, with mass in the range 0.3–7 GeV. Measured 90% CL limits as a function of  $m_{A^0}$  range from  $1 \times 10^{-6}$  to  $8 \times 10^{-5}$ .

$\Gamma(\gamma A^0 \rightarrow \gamma \mu^+ \mu^-)/\Gamma_{\text{total}}$   $\Gamma_{92}/\Gamma$

| VALUE (units $10^{-6}$ ) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|---------|
|--------------------------|-----|-------------|------|---------|

|      |    |                     |     |                                                              |
|------|----|---------------------|-----|--------------------------------------------------------------|
| <8.3 | 90 | <sup>1</sup> AUBERT | 09Z | BABR $e^+e^- \rightarrow A^0 \rightarrow \gamma \mu^+ \mu^-$ |
|------|----|---------------------|-----|--------------------------------------------------------------|

<sup>1</sup> For a narrow scalar or pseudoscalar,  $A^0$ , with mass in the range 212–9300 MeV, excluding  $J/\psi$  and  $\psi(2S)$ . Measured 90% CL limits as a function of  $m_{A^0}$  range from 0.26–8.3  $\times 10^{-6}$ .

LEPTON FAMILY NUMBER (*LF*) VIOLATING MODES

| $\Gamma(e^\pm \tau^\mp)/\Gamma_{\text{total}}$ |     |             |      |         | $\Gamma_{93}/\Gamma$                 |
|------------------------------------------------|-----|-------------|------|---------|--------------------------------------|
| VALUE (units $10^{-6}$ )                       | CL% | DOCUMENT ID | TECN | COMMENT |                                      |
| <b>&lt;3.2</b>                                 | 90  | LEES        | 10B  | BABR    | $e^+ e^- \rightarrow e^\pm \tau^\mp$ |

| $\Gamma(\mu^\pm \tau^\mp)/\Gamma_{\text{total}}$                              |     |             |      |         | $\Gamma_{94}/\Gamma$                   |
|-------------------------------------------------------------------------------|-----|-------------|------|---------|----------------------------------------|
| VALUE (units $10^{-6}$ )                                                      | CL% | DOCUMENT ID | TECN | COMMENT |                                        |
| <b>&lt; 3.3</b>                                                               | 90  | LEES        | 10B  | BABR    | $e^+ e^- \rightarrow \mu^\pm \tau^\mp$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |     |             |      |         |                                        |
| <14.4                                                                         | 95  | LOVE        | 08A  | CLEO    | $e^+ e^- \rightarrow \mu^\pm \tau^\mp$ |

$\Upsilon(2S)$  Cross-Particle Branching Ratios

$B(\Upsilon(2S) \rightarrow \pi^+ \pi^-) \times B(\Upsilon(3S) \rightarrow \Upsilon(2S) X)$

| VALUE (units $10^{-2}$ ) | EVTS | DOCUMENT ID | TECN | COMMENT                                  |
|--------------------------|------|-------------|------|------------------------------------------|
| <b>1.78±0.02±0.11</b>    | 906k | LEES        | 11C  | BABR $e^+ e^- \rightarrow \pi^+ \pi^- X$ |

$\Upsilon(2S)$  REFERENCES

|           |      |                |                              |                 |
|-----------|------|----------------|------------------------------|-----------------|
| GAO       | 23   | PR D108 112015 | B.S. Gao <i>et al.</i>       | (BELLE Collab.) |
| SHAMOV    | 23   | PL B839 137766 | A.G. Shamov, O.L. Rezanova   | (NOVO, NOVOU)   |
| FULSOM    | 18   | PRL 121 232001 | B.G. Fulsom <i>et al.</i>    | (BELLE Collab.) |
| JIA       | 18   | PR D97 112004  | S. Jia <i>et al.</i>         | (BELLE Collab.) |
| JIA       | 17   | PR D95 012001  | S. Jia <i>et al.</i>         | (BELLE Collab.) |
| JIA       | 17A  | PR D96 112002  | S. Jia <i>et al.</i>         | (BELLE Collab.) |
| LEES      | 14G  | PR D89 111102  | J.P. Lees <i>et al.</i>      | (BABAR Collab.) |
| YANG      | 14   | PR D90 112008  | S.D. Yang <i>et al.</i>      | (BELLE Collab.) |
| SANDILYA  | 13   | PRL 111 112001 | S. Sandilya <i>et al.</i>    | (BELLE Collab.) |
| SHEN      | 13   | PR D88 011102  | C.P. Shen <i>et al.</i>      | (BELLE Collab.) |
| TAMPONI   | 13   | PR D87 011104  | U. Tamponi <i>et al.</i>     | (BELLE Collab.) |
| DOBBS     | 12   | PRL 109 082001 | S. Dobbs <i>et al.</i>       |                 |
| DOBBS     | 12A  | PR D86 052003  | S. Dobbs <i>et al.</i>       |                 |
| SHEN      | 12A  | PR D86 031102  | S.P. Shen <i>et al.</i>      | (BELLE Collab.) |
| LEES      | 11C  | PR D84 011104  | J.P. Lees <i>et al.</i>      | (BABAR Collab.) |
| LEES      | 11H  | PRL 107 221803 | J.P. Lees <i>et al.</i>      | (BABAR Collab.) |
| LEES      | 11J  | PR D84 072002  | J.P. Lees <i>et al.</i>      | (BABAR Collab.) |
| LEES      | 11L  | PR D84 092003  | J.P. Lees <i>et al.</i>      | (BABAR Collab.) |
| WANG      | 11B  | PR D84 071107  | X.L. Wang <i>et al.</i>      | (BELLE Collab.) |
| BONVICINI | 10   | PR D81 031104  | G. Bonvicini <i>et al.</i>   | (CLEO Collab.)  |
| LEES      | 10B  | PRL 104 151802 | J.P. Lees <i>et al.</i>      | (BABAR Collab.) |
| AUBERT    | 09AQ | PRL 103 161801 | B. Aubert <i>et al.</i>      | (BABAR Collab.) |
| AUBERT    | 09Z  | PRL 103 081803 | B. Aubert <i>et al.</i>      | (BABAR Collab.) |
| BHARI     | 09   | PR D79 011103  | S.R. Bhari <i>et al.</i>     | (CLEO Collab.)  |
| AUBERT    | 08BP | PR D78 112002  | B. Aubert <i>et al.</i>      | (BABAR Collab.) |
| HE        | 08A  | PRL 101 192001 | Q. He <i>et al.</i>          | (CLEO Collab.)  |
| LOVE      | 08A  | PRL 101 201601 | W. Love <i>et al.</i>        | (CLEO Collab.)  |
| PDG       | 08   | PL B667 1      | C. Amsler <i>et al.</i>      | (PDG Collab.)   |
| ASNER     | 07   | PR D75 012009  | D.M. Asner <i>et al.</i>     | (CLEO Collab.)  |
| BESSON    | 07   | PRL 98 052002  | D. Besson <i>et al.</i>      | (CLEO Collab.)  |
| ROSNER    | 07A  | PR D76 117102  | J.L. Rosner <i>et al.</i>    | (CLEO Collab.)  |
| BESSON    | 06A  | PR D74 012003  | D. Besson <i>et al.</i>      | (CLEO Collab.)  |
| ROSNER    | 06   | PRL 96 092003  | J.L. Rosner <i>et al.</i>    | (CLEO Collab.)  |
| ADAMS     | 05   | PRL 94 012001  | G.S. Adams <i>et al.</i>     | (CLEO Collab.)  |
| ARTUSO    | 05   | PRL 94 032001  | M. Artuso <i>et al.</i>      | (CLEO Collab.)  |
| ARTAMONOV | 00   | PL B474 427    | A.S. Artamonov <i>et al.</i> |                 |
| EDWARDS   | 99   | PR D59 032003  | K.W. Edwards <i>et al.</i>   | (CLEO Collab.)  |
| ALEXANDER | 98   | PR D58 052004  | J.P. Alexander <i>et al.</i> | (CLEO Collab.)  |
| BARU      | 96   | PRPL 267 71    | S.E. Baru <i>et al.</i>      | (NOVO)          |

|                                                             |     |                         |                               |                              |
|-------------------------------------------------------------|-----|-------------------------|-------------------------------|------------------------------|
| KOBEL                                                       | 92  | ZPHY C53 193            | M. Kobel <i>et al.</i>        | (Crystal Ball Collab.)       |
| MASCHMANN                                                   | 90  | ZPHY C46 555            | W.S. Maschmann <i>et al.</i>  | (Crystal Ball Collab.)       |
| ALBRECHT                                                    | 89  | ZPHY C42 349            | H. Albrecht <i>et al.</i>     | (ARGUS Collab.)              |
| KAARSBERG                                                   | 89  | PRL 62 2077             | T.M. Kaarsberg <i>et al.</i>  | (CUSB Collab.)               |
| BUCHMUEL...                                                 | 88  | HE $e^+e^-$ Physics 412 | W. Buchmueller, S. Cooper     | (HANN, DESY, MIT)            |
| Editors: A. Ali and P. Soeding, World Scientific, Singapore |     |                         |                               |                              |
| JAKUBOWSKI                                                  | 88  | ZPHY C40 49             | Z. Jakubowski <i>et al.</i>   | (Crystal Ball Collab.) IGJPC |
| ALBRECHT                                                    | 87  | ZPHY C35 283            | H. Albrecht <i>et al.</i>     | (ARGUS Collab.)              |
| COHEN                                                       | 87  | RMP 59 1121             | E.R. Cohen, B.N. Taylor       | (RISC, NBS)                  |
| LURZ                                                        | 87  | ZPHY C36 383            | B. Lurz <i>et al.</i>         | (Crystal Ball Collab.)       |
| BARU                                                        | 86B | ZPHY C32 622 (errat.)   | S.E. Baru <i>et al.</i>       | (NOVO)                       |
| ALBRECHT                                                    | 85  | ZPHY C28 45             | H. Albrecht <i>et al.</i>     | (ARGUS Collab.)              |
| ALBRECHT                                                    | 85E | PL 160B 331             | H. Albrecht <i>et al.</i>     | (ARGUS Collab.)              |
| GELPHMAN                                                    | 85  | PR D32 2893             | D. Gelpman <i>et al.</i>      | (Crystal Ball Collab.)       |
| KURAEV                                                      | 85  | SJNP 41 466             | E.A. Kuraev, V.S. Fadin       | (NOVO)                       |
| Translated from YAF 41 733.                                 |     |                         |                               |                              |
| NERNST                                                      | 85  | PRL 54 2195             | R. Nernst <i>et al.</i>       | (Crystal Ball Collab.)       |
| ARTAMONOV                                                   | 84  | PL 137B 272             | A.S. Artamonov <i>et al.</i>  | (NOVO)                       |
| BARBER                                                      | 84  | PL 135B 498             | D.P. Barber <i>et al.</i>     |                              |
| BESSON                                                      | 84  | PR D30 1433             | D. Besson <i>et al.</i>       | (CLEO Collab.)               |
| FONSECA                                                     | 84  | NP B242 31              | V. Fonseca <i>et al.</i>      | (CUSB Collab.)               |
| GILES                                                       | 84B | PR D29 1285             | R. Giles <i>et al.</i>        | (CLEO Collab.)               |
| HAAS                                                        | 84  | PRL 52 799              | J. Haas <i>et al.</i>         | (CLEO Collab.)               |
| HAAS                                                        | 84B | PR D30 1996             | J. Haas <i>et al.</i>         | (CLEO Collab.)               |
| KLOPFEN...                                                  | 83  | PRL 51 160              | C. Klopfenstein <i>et al.</i> | (CUSB Collab.)               |
| ALBRECHT                                                    | 82  | PL 116B 383             | H. Albrecht <i>et al.</i>     | (DESY, DORT, HEIDH+)         |
| NICZYPORUK                                                  | 81B | PL 100B 95              | B. Niczyporuk <i>et al.</i>   | (LENA Collab.)               |
| NICZYPORUK                                                  | 81C | PL 99B 169              | B. Niczyporuk <i>et al.</i>   | (LENA Collab.)               |
| BOCK                                                        | 80  | ZPHY C6 125             | P. Bock <i>et al.</i>         | (HEIDP, MPIM, DESY, HAMB)    |