

$\Upsilon(3S)$	$I^G(J^{PC}) = 0^-(1^{--})$
----------------------------------	-----------------------------

 $\Upsilon(3S)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
10355.1±0.5	¹ SHAMOV 23	RVUE	$e^+ e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
10355.2±0.5	^{2,3} ARTAMONOV 00	MD1	$e^+ e^- \rightarrow$ hadrons
10355.3±0.5	^{4,5} BARU 86B	MD1	$e^+ e^- \rightarrow$ hadrons
¹ Reanalysis of MD1 data using the electron mass from COHEN 87, the radiative corrections from KURAEV 85 and interference effects.			
² Reanalysis of BARU 86B using new electron mass (COHEN 87).			
³ Superseded by SHAMOV 23.			
⁴ Reanalysis of ARTAMONOV 84.			
⁵ Superseded by ARTAMONOV 00.			

 $m_{\Upsilon(3S)} - m_{\Upsilon(2S)}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
331.50±0.02±0.13	LEES	11c BABR	$e^+ e^- \rightarrow \pi^+ \pi^- X$

 $\Upsilon(3S)$ WIDTH

VALUE (keV)	DOCUMENT ID
20.32±1.85 OUR EVALUATION	See the Note on "Width Determinations of the Υ States"

 $\Upsilon(3S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 $\Upsilon(2S)$ anything	(10.6 ± 0.8) %	
Γ_2 $\Upsilon(2S)\pi^+\pi^-$	(2.82± 0.18) %	S=1.6
Γ_3 $\Upsilon(2S)\pi^0\pi^0$	(1.85± 0.14) %	
Γ_4 $\Upsilon(2S)\gamma\gamma$	(5.0 ± 0.7) %	
Γ_5 $\Upsilon(2S)\pi^0$	< 5.1 × 10 ⁻⁴	CL=90%
Γ_6 $\Upsilon(1S)\pi^+\pi^-$	(4.37± 0.08) %	
Γ_7 $\Upsilon(1S)\pi^0\pi^0$	(2.20± 0.13) %	
Γ_8 $\Upsilon(1S)\eta$	< 1 × 10 ⁻⁴	CL=90%
Γ_9 $\Upsilon(1S)\pi^0$	< 7 × 10 ⁻⁵	CL=90%
Γ_{10} $h_b(1P)\pi^0$	< 1.2 × 10 ⁻³	CL=90%
Γ_{11} $h_b(1P)\pi^0 \rightarrow \gamma\eta_b(1S)\pi^0$	(4.3 ± 1.4) × 10 ⁻⁴	
Γ_{12} $h_b(1P)\pi^+\pi^-$	< 1.2 × 10 ⁻⁴	CL=90%

Γ_{13}	$\tau^+ \tau^-$	(2.29 ± 0.30) %	
Γ_{14}	$\mu^+ \mu^-$	(2.18 ± 0.21) %	S=2.1
Γ_{15}	$e^+ e^-$	(2.18 ± 0.20) %	
Γ_{16}	hadrons	(93 ± 12) %	
Γ_{17}	ggg	(35.7 ± 2.6) %	
Γ_{18}	γgg	(9.7 ± 1.8) $\times 10^{-3}$	
Γ_{19}	$\frac{1}{2}H$ anything	(2.33 ± 0.33) $\times 10^{-5}$	

Radiative decays

Γ_{20}	$\gamma \chi b_2(2P)$	(13.1 ± 1.6) %	S=3.4
Γ_{21}	$\gamma \chi b_1(2P)$	(12.6 ± 1.2) %	S=2.4
Γ_{22}	$\gamma \chi b_0(2P)$	(5.9 ± 0.6) %	S=1.4
Γ_{23}	$\gamma \chi b_2(1P)$	(10.0 ± 1.0) $\times 10^{-3}$	S=1.7
Γ_{24}	$\gamma \chi b_1(1P)$	(9 ± 5) $\times 10^{-4}$	S=1.8
Γ_{25}	$\gamma \chi b_0(1P)$	(2.7 ± 0.4) $\times 10^{-3}$	
Γ_{26}	$\gamma \eta_b(2S)$	< 6.2 $\times 10^{-4}$	CL=90%
Γ_{27}	$\gamma \eta_b(1S)$	(5.1 ± 0.7) $\times 10^{-4}$	
Γ_{28}	$\gamma A^0 \rightarrow \gamma$ hadrons	< 8 $\times 10^{-5}$	CL=90%
Γ_{29}	$\gamma X \rightarrow \gamma + \geq 4$ prongs	[a] < 2.2 $\times 10^{-4}$	CL=95%
Γ_{30}	$\gamma A^0 \rightarrow \gamma \mu^+ \mu^-$	< 5.5 $\times 10^{-6}$	CL=90%
Γ_{31}	$\gamma A^0 \rightarrow \gamma \tau^+ \tau^-$	[b] < 1.6 $\times 10^{-4}$	CL=90%

Lepton Family number (*LF*) violating modes

Γ_{32}	$e^\pm \tau^\mp$	<i>LF</i>	< 4.2 $\times 10^{-6}$	CL=90%
Γ_{33}	$e^\pm \mu^\mp$	<i>LF</i>	< 3.6 $\times 10^{-7}$	CL=90%
Γ_{34}	$\mu^\pm \tau^\mp$	<i>LF</i>	< 3.1 $\times 10^{-6}$	CL=90%

[a] $1.5 \text{ GeV} < m_X < 5.0 \text{ GeV}$

[b] For $m_{\tau^+ \tau^-}$ in the ranges 4.03–9.52 and 9.61–10.10 GeV.

$\Upsilon(3S) \Gamma(i) \Gamma(e^+ e^-)/\Gamma(\text{total})$

$\Gamma(\text{hadrons}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$		$\Gamma_{16}\Gamma_{15}/\Gamma$	
VALUE (keV)	DOCUMENT ID	TECN	COMMENT
0.414 ± 0.007 OUR AVERAGE			
0.413 $\pm 0.004 \pm 0.006$	ROSNER	06	CLEO $10.4 \text{ e}^+ \text{e}^- \rightarrow \text{hadrons}$
0.45 $\pm 0.03 \pm 0.03$	⁶ GILES	84B	CLEO $\text{e}^+ \text{e}^- \rightarrow \text{hadrons}$

⁶ Radiative corrections reevaluated by BUCHMUELLER 88 following KURAEV 85.

$\Gamma(\Upsilon(1S) \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$		$\Gamma_6 \Gamma_{15}/\Gamma$		
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
18.46 $\pm 0.27 \pm 0.77$	6.4k	⁷ AUBERT	08BP BABR	$e^+ e^- \rightarrow \gamma \pi^+ \pi^- \ell^+ \ell^-$
7 Using $B(\Upsilon(1S) \rightarrow e^+ e^-) = (2.38 \pm 0.11)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05)\%$.				

$\Upsilon(3S)$ PARTIAL WIDTHS $\Gamma(e^+ e^-)$ Γ_{15}

VALUE (keV)

DOCUMENT ID

0.443±0.008 OUR EVALUATION **$\Upsilon(3S)$ BRANCHING RATIOS** $\Gamma(\Upsilon(2S)\text{anything})/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE

EVTS

DOCUMENT ID

TECN

COMMENT

0.106 ±0.008 OUR AVERAGE

0.1023±0.0105	4625	8,9,10	BUTLER	94B	CLE2	$e^+ e^- \rightarrow \ell^+ \ell^- X$
0.111 ± 0.012	4891	9,10,11	BROCK	91	CLEO	$e^+ e^- \rightarrow \pi^+ \pi^- X,$ $\pi^+ \pi^- \ell^+ \ell^-$

⁸ Using $B(\Upsilon(2S) \rightarrow \Upsilon(1S) \gamma\gamma) = (0.038 \pm 0.007)\%$, and $B(\Upsilon(2S) \rightarrow \Upsilon(1S) \pi^0 \pi^0) = (1/2)B(\Upsilon(2S) \rightarrow \Upsilon(1S) \pi^+ \pi^-)$.

⁹ Using $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.06)\%$. With the assumption of $e\mu$ universality.

¹⁰ Using $B(\Upsilon(2S) \rightarrow \Upsilon(1S) \pi^+ \pi^-) = (18.5 \pm 0.8)\%$.

¹¹ Using $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.31 \pm 0.21)\%$, $B(\Upsilon(2S) \rightarrow \Upsilon(1S) \gamma\gamma) \times 2B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (0.188 \pm 0.035)\%$, and $B(\Upsilon(2S) \rightarrow \Upsilon(1S) \pi^0 \pi^0) \times 2B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (0.436 \pm 0.056)\%$. With the assumption of $e\mu$ universality.

 $\Gamma(\Upsilon(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_2/Γ VALUE (units 10^{-2})

EVTS

DOCUMENT ID

TECN

COMMENT

2.82±0.18 OUR AVERAGE Error includes scale factor of 1.6. See the ideogram below.

3.00±0.02±0.14	543k	LEES	11C	BABR	$e^+ e^- \rightarrow \pi^+ \pi^- X$
2.40±0.10±0.26	800	¹² AUBERT	08BP	BABR	$e^+ e^- \rightarrow \gamma \pi^+ \pi^- e^+ e^-$
3.12±0.49	980	^{13,14} BUTLER	94B	CLE2	$e^+ e^- \rightarrow \pi^+ \pi^- \ell^+ \ell^-$
2.13±0.38	974	¹⁵ BROCK	91	CLEO	$e^+ e^- \rightarrow \pi^+ \pi^- X,$ $\pi^+ \pi^- \ell^+ \ell^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.82±0.65±0.53 138 ¹⁵WU 93 CUSB $\Upsilon(3S) \rightarrow \pi^+ \pi^- \ell^+ \ell^-$


3.1 ±2.0 5 MAGERAS 82 CUSB $\Upsilon(3S) \rightarrow \pi^+ \pi^- \ell^+ \ell^-$

¹² Using $B(\Upsilon(1S) \rightarrow e^+ e^-) = (2.38 \pm 0.11)\%$, $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05)\%$, and $\Gamma_{ee}(\Upsilon(3S)) = 0.443 \pm 0.008$ keV.

¹³ From the exclusive mode.

¹⁴ Using $B(\Upsilon(2S) \rightarrow \Upsilon(1S) \gamma\gamma) = (0.038 \pm 0.007)\%$, and $B(\Upsilon(2S) \rightarrow \Upsilon(1S) \pi^0 \pi^0) = (1/2)B(\Upsilon(2S) \rightarrow \Upsilon(1S) \pi^+ \pi^-)$.

¹⁵ Using $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.31 \pm 0.21)\%$, $B(\Upsilon(2S) \rightarrow \Upsilon(1S) \gamma\gamma) \times 2B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (0.188 \pm 0.035)\%$, and $B(\Upsilon(2S) \rightarrow \Upsilon(1S) \pi^0 \pi^0) \times 2B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (0.436 \pm 0.056)\%$. With the assumption of $e\mu$ universality.



$\Gamma(\gamma(2S)\pi^0\pi^0)/\Gamma_{\text{total}}$

Γ_3/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.85±0.14 OUR AVERAGE				
1.82±0.09±0.12	4391	¹⁶ BHARI	09	CLEO $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
2.16±0.39		^{17,18} BUTLER	94B	CLE2 $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
1.7 ± 0.5 ± 0.2	10	¹⁹ HEINTZ	92	CSB2 $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$

¹⁶ Authors assume $B(\gamma(1S) \rightarrow e^+e^-) + B(\gamma(1S) \rightarrow \mu^+\mu^-) = 4.06\%$.

¹⁷ $B(\gamma(2S) \rightarrow \mu^+\mu^-) = (1.31 \pm 0.21)\%$ and assuming $e\mu$ universality.

¹⁸ From the exclusive mode.

¹⁹ $B(\gamma(2S) \rightarrow \mu^+\mu^-) = (1.44 \pm 0.10)\%$ and assuming $e\mu$ universality. Supersedes HEINTZ 91.

$\Gamma(\gamma(2S)\gamma\gamma)/\Gamma_{\text{total}}$

Γ_4/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0502±0.0069	20 BUTLER	94B	CLE2 $e^+e^- \rightarrow \ell^+\ell^- 2\gamma$

²⁰ From the exclusive mode.

$\Gamma(\gamma(2S)\pi^0)/\Gamma_{\text{total}}$

Γ_5/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.51	90	²¹ HE	08A	CLEO $e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$

²¹ Authors assume $B(\gamma(2S) \rightarrow e^+e^-) + B(\gamma(1S) \rightarrow \mu^+\mu^-) = 4.06\%$.

$\Gamma(\Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_6/Γ

Abbreviation MM in the *COMMENT* field below stands for missing mass.

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
4.37 ± 0.08 OUR AVERAGE				
4.32 ± 0.07 ± 0.13	90k	22 LEES	11L BABR	$\Upsilon(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$
4.46 ± 0.01 ± 0.13	190k	23 BHARI	09 CLEO	$e^+e^- \rightarrow \pi^+\pi^-$ MM
4.17 ± 0.06 ± 0.19	6.4k	24 AUBERT	08BP BABR	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\ell^+\ell^-$
4.52 ± 0.35	11830	25 BUTLER	94B CLE2	$e^+e^- \rightarrow \pi^+\pi^-X$, $\pi^+\pi^-\ell^+\ell^-$
4.46 ± 0.34 ± 0.50	451	25 WU	93 CUSB	$\Upsilon(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$
4.46 ± 0.30	11221	25 BROCK	91 CLEO	$e^+e^- \rightarrow \pi^+\pi^-X$, $\pi^+\pi^-\ell^+\ell^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
4.9 ± 1.0	22	GREEN	82 CLEO	$\Upsilon(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$
3.9 ± 1.3	26	MAGERAS	82 CUSB	$\Upsilon(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$

22 Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$.

23 A weighted average of the inclusive and exclusive results.

24 Using $B(\Upsilon(2S) \rightarrow e^+e^-) = (1.91 \pm 0.16)\%$, $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17)\%$, and $\Gamma_{ee}(\Upsilon(3S)) = 0.443 \pm 0.008$ keV.

25 Using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.06)\%$. With the assumption of $e\mu$ universality.

$\Gamma(\Upsilon(2S)\pi^+\pi^-)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$ Γ_2/Γ_6

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.577 ± 0.026 ± 0.060	800	26 AUBERT	08BP BABR	$e^+e^- \rightarrow \gamma\pi^+\pi^-\ell^+\ell^-$
26	Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$, $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$, $B(\Upsilon(2S) \rightarrow e^+e^-) = (1.91 \pm 0.16)\%$, and $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17)\%$. Not independent of other values reported by AUBERT 08BP.			

$\Gamma(\Upsilon(1S)\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.20 ± 0.13 OUR AVERAGE

2.24 ± 0.09 ± 0.11	6584	27 BHARI	09 CLEO	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
1.99 ± 0.34	56	28 BUTLER	94B CLE2	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
2.2 ± 0.4 ± 0.3	33	29 HEINTZ	92 CSB2	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$

27 Authors assume $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$.

28 Using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.06)\%$ and assuming $e\mu$ universality.

29 Using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.57 \pm 0.07)\%$ and assuming $e\mu$ universality. Supersedes HEINTZ 91.

$\Gamma(\Upsilon(1S)\pi^0\pi^0)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$ Γ_7/Γ_6

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.501 ± 0.043	30	BHARI	09 CLEO	$e^+e^- \rightarrow \Upsilon(3S)$
---------------	----	-------	---------	-----------------------------------

30 Not independent of other values reported by BHARI 09.

$\Gamma(\Upsilon(1S)\eta)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.1	90	31 LEES	11L BABR	$\Upsilon(3S) \rightarrow (\pi^+\pi^-)(\gamma\gamma)\ell^+\ell^-$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<0.8	90	31,32 AUBERT	08BP BABR	$e^+e^- \rightarrow \gamma\pi^+\pi^-\pi^0\ell^+\ell^-$
<0.18	90	33 HE	08A CLEO	$e^+e^- \rightarrow \ell^+\ell^-\eta$
<2.2	90	BROCK	91 CLEO	$e^+e^- \rightarrow \ell^+\ell^-\eta$

³¹ Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$, $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$.³² Using $\Gamma_{ee}(\Upsilon(3S)) = 0.443 \pm 0.008$ keV.³³ Authors assume $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$. $\Gamma(\Upsilon(1S)\eta)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$ Γ_8/Γ_6

VALUE (units 10^{-2})	CL%	DOCUMENT ID	TECN	COMMENT
<0.23	90	34 LEES	11L BABR	$\Upsilon(3S) \rightarrow (\pi^+\pi^-)(\gamma\gamma)\ell^+\ell^-$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<1.9	90	35 AUBERT	08BP BABR	$e^+e^- \rightarrow \gamma\pi^+\pi^-(\pi^0)\ell^+\ell^-$

³⁴ Not independent of other values reported by LEES 11L.³⁵ Not independent of other values reported by AUBERT 08BP. $\Gamma(\Upsilon(1S)\pi^0)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.07	90	36 HE	08A CLEO	$e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$
³⁶ Authors assume $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$.				

 $\Gamma(h_b(1P)\pi^0)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.2 \times 10^{-3}$	90	37 GE	11 CLEO	$\Upsilon(3S) \rightarrow \pi^0$ anything
³⁷ Assuming $M(h_b(1P)) = 9900$ MeV and $\Gamma(h_b(1P)) = 0$ MeV, and allowing $B(h_b(1P) \rightarrow \gamma\eta_b(1S))$ to vary from 0–100%.				

 $\Gamma(h_b(1P)\pi^0 \rightarrow \gamma\eta_b(1S)\pi^0)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$4.3 \pm 1.1 \pm 0.9$	LEES	11K BABR	$\Upsilon(3S) \rightarrow \eta_b\gamma\pi^0$

 $\Gamma(h_b(1P)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 1.2	90	38 LEES	11C BABR	$e^+e^- \rightarrow \pi^+\pi^-X$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<18		38 BUTLER	94B CLE2	$e^+e^- \rightarrow \pi^+\pi^-X$
<15		38 BROCK	91 CLEO	$e^+e^- \rightarrow \pi^+\pi^-X$

³⁸ For $M(h_b(1P)) = 9900$ MeV.

$\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.29±0.21±0.22	15k	39 BESSON	07	CLEO $e^+e^- \rightarrow \gamma(3S) \rightarrow \tau^+\tau^-$

³⁹ BESSON 07 reports $[\Gamma(\gamma(3S) \rightarrow \tau^+\tau^-)/\Gamma_{\text{total}}] / [B(\gamma(3S) \rightarrow \mu^+\mu^-)] = 1.05 \pm 0.08 \pm 0.05$ which we multiply by our best value $B(\gamma(3S) \rightarrow \mu^+\mu^-) = (2.18 \pm 0.21) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.


$\Gamma(\tau^+\tau^-)/\Gamma(\mu^+\mu^-)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.968±0.016 OUR AVERAGE				
0.966±0.008±0.014	2.2M	LEES	20E	BABR $e^+e^- \rightarrow \gamma(3S)$
1.05 ± 0.08 ± 0.05	15k	BESSON	07	CLEO $e^+e^- \rightarrow \gamma(3S)$

$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0218±0.0021 OUR AVERAGE		Error includes scale factor of 2.1. See the ideogram below.		
0.0239±0.0007±0.0010	81k	ADAMS	05	CLEO $e^+e^- \rightarrow \mu^+\mu^-$
0.0202±0.0019±0.0033		CHEN	89B	CLEO $e^+e^- \rightarrow \mu^+\mu^-$
0.0173±0.0015±0.0011		KAARSBERG	89	CSB2 $e^+e^- \rightarrow \mu^+\mu^-$
0.033 ± 0.013 ± 0.007	1096	ANDREWS	83	CLEO $e^+e^- \rightarrow \mu^+\mu^-$

WEIGHTED AVERAGE
0.0218±0.0021 (Error scaled by 2.1)



$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$

$\Gamma(ggg)/\Gamma_{\text{total}}$ Γ_{17}/Γ

<i>VALUE</i> (units 10^{-2})	<i>EVTS</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
35.7±2.6	3M	40 BESSON	06A CLEO	$\gamma(3S) \rightarrow \text{hadrons}$

⁴⁰ Calculated using BESSON 06A value of $\Gamma(\gamma gg)/\Gamma(ggg) = (2.72 \pm 0.06 \pm 0.32 \pm 0.37)\%$ and the PDG 08 values of $B(\gamma(2S) + \text{anything}) = (10.6 \pm 0.8)\%$, $B(\pi^+ \pi^- \gamma(1S)) = (4.40 \pm 0.10)\%$, $B(\pi^0 \pi^0 \gamma(1S)) = (2.20 \pm 0.13)\%$, $B(\gamma \chi b_2(2P)) = (13.1 \pm 1.6)\%$, $B(\gamma \chi b_1(2P)) = (12.6 \pm 1.2)\%$, $B(\gamma \chi b_0(2P)) = (5.9 \pm 0.6)\%$, $B(\gamma \chi b_0(1P)) = (0.30 \pm 0.11)\%$, $B(\mu^+ \mu^-) = (2.18 \pm 0.21)\%$, and $R_{\text{hadrons}} = 3.51$. The statistical error is negligible and the systematic error is partially correlated with $\Gamma(\gamma gg)/\Gamma_{\text{total}}$ BESSON 06A value.

 $\Gamma(\gamma gg)/\Gamma_{\text{total}}$ Γ_{18}/Γ

<i>VALUE</i> (units 10^{-2})	<i>EVTS</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
0.97±0.18	60k	41 BESSON	06A CLEO	$\gamma(3S) \rightarrow \gamma + \text{hadrons}$

⁴¹ Calculated using BESSON 06A values of $\Gamma(\gamma gg)/\Gamma(ggg) = (2.72 \pm 0.06 \pm 0.32 \pm 0.37)\%$ and $\Gamma(ggg)/\Gamma_{\text{total}}$. The statistical error is negligible and the systematic error is partially correlated with $\Gamma(ggg)/\Gamma_{\text{total}}$ BESSON 06A value.

 $\Gamma(\gamma gg)/\Gamma(ggg)$ Γ_{18}/Γ_{17}

<i>VALUE</i> (units 10^{-2})	<i>EVTS</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
2.72±0.06±0.49	3M	BESSON	06A CLEO	$\gamma(3S) \rightarrow (\gamma +) \text{hadrons}$


 $\Gamma(\overline{2}H \text{ anything})/\Gamma_{\text{total}}$ Γ_{19}/Γ

<i>VALUE</i> (units 10^{-5})	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
2.33±0.15^{+0.31}_{-0.28}	LEES	14G BABR	$e^+ e^- \rightarrow \overline{2}H X$

 $\Gamma(\gamma \chi b_2(2P))/\Gamma_{\text{total}}$ Γ_{20}/Γ

<i>VALUE</i>	<i>EVTS</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
0.131 ± 0.016 OUR AVERAGE		Error includes scale factor of 3.4. See the ideogram below.		
0.1579 ± 0.0017 ± 0.0073	568k	ARTUSO	05 CLEO	$e^+ e^- \rightarrow \gamma X$
0.111 ± 0.005 ± 0.004	10319	42 HEINTZ	92 CSB2	$e^+ e^- \rightarrow \gamma X$
0.135 ± 0.003 ± 0.017	30741	MORRISON	91 CLE2	$e^+ e^- \rightarrow \gamma X$


⁴² Supersedes NARAIN 91.



$\Gamma(\gamma\chi_{b1}(2P))/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{21}/Γ
0.126 ± 0.012 OUR AVERAGE	537k	ARTUSO	05	CLEO $e^+ e^- \rightarrow \gamma X$	
0.1454 ± 0.0018 ± 0.0073	11147	43 HEINTZ	92	CSB2 $e^+ e^- \rightarrow \gamma X$	
0.115 ± 0.005 ± 0.005	25759	MORRISON	91	CLE2 $e^+ e^- \rightarrow \gamma X$	
0.105 +0.003 -0.002 ± 0.013					


⁴³ Supersedes NARAIN 91.



$\Gamma(\gamma\chi_{b0}(2P))/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{22}/Γ
0.059 ±0.006 OUR AVERAGE		Error includes scale factor of 1.4. See the ideogram below.			
0.0677±0.0020±0.0065	225k	ARTUSO	05	CLEO $e^+ e^- \rightarrow \gamma X$	
0.060 ±0.004 ±0.006	4959	⁴⁴ HEINTZ	92	CSB2 $e^+ e^- \rightarrow \gamma X$	
0.049 ^{+0.003} _{-0.004} ±0.006	9903	MORRISON	91	CLE2 $e^+ e^- \rightarrow \gamma X$	

⁴⁴ Supersedes NARAIN 91.



$\Gamma(\gamma\chi_{b2}(1P))/\Gamma_{\text{total}}$

Γ_{23}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
10.0±1.0 OUR AVERAGE			Error includes scale factor of 1.7.		
$8.0 \pm 1.3 \pm 0.4$	126	45,46	KORNICER	11	CLEO $e^+ e^- \rightarrow \gamma\gamma\ell^+\ell^-$
$10.5 \pm 0.3 \pm 0.7$	9.7k	LEES		11J	BABR $\Upsilon(3S) \rightarrow X\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<19	90	47 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma + \text{hadrons}$
seen		48 HEINTZ	92 CSB2	$e^+ e^- \rightarrow \gamma\gamma\ell^+\ell^-$

⁴⁵ Assuming $B(\Upsilon(1S) \rightarrow \ell^+\ell^-) = (2.48 \pm 0.05)\%$.

⁴⁶ KORNICER 11 reports $[\Gamma(\Upsilon(3S) \rightarrow \gamma\chi_{b2}(1P))/\Gamma_{\text{total}}] \times [B(\chi_{b2}(1P) \rightarrow \gamma\Upsilon(1S))] = (1.435 \pm 0.162 \pm 0.169) \times 10^{-3}$ which we divide by our best value $B(\chi_{b2}(1P) \rightarrow \gamma\Upsilon(1S)) = (18.0 \pm 1.0) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴⁷ ASNER 08A reports $[\Gamma(\Upsilon(3S) \rightarrow \gamma\chi_{b2}(1P))/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P))] < 27.1 \times 10^{-2}$ which we multiply by our best value $B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P)) = 7.15 \times 10^{-2}$.

⁴⁸ HEINTZ 92, while unable to distinguish between different J states, measures $\sum_J B(\Upsilon(3S) \rightarrow \gamma\chi_{bJ}) \times B(\chi_{bJ} \rightarrow \gamma\Upsilon(1S)) = (1.7 \pm 0.4 \pm 0.6) \times 10^{-3}$ for $J = 0,1,2$ using inclusive $\Upsilon(1S)$ decays and $(1.2^{+0.4}_{-0.3} \pm 0.09) \times 10^{-3}$ for $J = 1,2$ using $\Upsilon(1S) \rightarrow \ell^+\ell^-$.

$\Gamma(\gamma\chi_{b1}(1P))/\Gamma_{\text{total}}$ Γ_{24}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT		
0.9±0.5 OUR AVERAGE			Error includes scale factor of 1.8.				
$1.5 \pm 0.4 \pm 0.1$	50	49,50	KORNICER	11	CLEO	$e^+ e^- \rightarrow \gamma\gamma\ell^+\ell^-$	
$0.5 \pm 0.3^{+0.2}_{-0.1}$			LEES	11J	BABR	$\gamma(3S) \rightarrow X\gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •							
<1.7	90	51	ASNER	08A	CLEO	$\gamma(3S) \rightarrow \gamma + \text{hadrons}$	
seen		52	HEINTZ	92	CSB2	$e^+ e^- \rightarrow \gamma\gamma\ell^+\ell^-$	

49 Assuming $B(\gamma(1S) \rightarrow \ell^+\ell^-) = (2.48 \pm 0.05)\%$.50 KORNICER 11 reports $[\Gamma(\gamma\chi_{b1}(1P))/\Gamma_{\text{total}}] \times [B(\chi_{b1}(1P) \rightarrow \gamma\gamma(1S))] = (5.38 \pm 1.20 \pm 0.95) \times 10^{-4}$ which we divide by our best value $B(\chi_{b1}(1P) \rightarrow \gamma\gamma(1S)) = (35.2 \pm 2.0) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.51 ASNER 08A reports $[\Gamma(\gamma\chi_{b1}(1P))/\Gamma_{\text{total}}] / [B(\gamma(2S) \rightarrow \gamma\chi_{b1}(1P))] < 2.5 \times 10^{-2}$ which we multiply by our best value $B(\gamma(2S) \rightarrow \gamma\chi_{b1}(1P)) = 6.9 \times 10^{-2}$.52 HEINTZ 92, while unable to distinguish between different J states, measures $\sum_J B(\gamma(3S) \rightarrow \gamma\chi_{bJ}) \times B(\chi_{bJ} \rightarrow \gamma\gamma(1S)) = (1.7 \pm 0.4 \pm 0.6) \times 10^{-3}$ for $J = 0,1,2$ using inclusive $\gamma(1S)$ decays and $(1.2^{+0.4}_{-0.3} \pm 0.09) \times 10^{-3}$ for $J = 1,2$ using $\gamma(1S) \rightarrow \ell^+\ell^-$. $\Gamma(\gamma\chi_{b0}(1P))/\Gamma_{\text{total}}$ Γ_{25}/Γ

VALUE (units 10^{-2})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT		
0.27±0.04 OUR AVERAGE							
$0.27 \pm 0.04 \pm 0.02$	2.3k	LEES	11J	BABR	$\gamma(3S) \rightarrow X\gamma$		
$0.30 \pm 0.04 \pm 0.10$	8.7k	ARTUSO	05	CLEO	$e^+ e^- \rightarrow \gamma X$		
• • • We do not use the following data for averages, fits, limits, etc. • • •							
<0.8	90	53	ASNER	08A	CLEO	$\gamma(3S) \rightarrow \gamma + \text{hadrons}$	

53 ASNER 08A reports $[\Gamma(\gamma\chi_{b0}(1P))/\Gamma_{\text{total}}] / [B(\gamma(2S) \rightarrow \gamma\chi_{b0}(1P))] < 21.9 \times 10^{-2}$ which we multiply by our best value $B(\gamma(2S) \rightarrow \gamma\chi_{b0}(1P)) = 3.8 \times 10^{-2}$.

 $\Gamma(\gamma\eta_b(2S))/\Gamma_{\text{total}}$ Γ_{26}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	
< 6.2	90		ARTUSO	05	CLEO	$e^+ e^- \rightarrow \gamma X$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<19	90		LEES	11J	BABR	$\gamma(3S) \rightarrow X\gamma$

 $\Gamma(\gamma\eta_b(1S))/\Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT		
5.1±0.7 OUR AVERAGE							
$7.1 \pm 1.8 \pm 1.3$	2.3 ± 0.5k	54	BONVICINI	10	CLEO	$\gamma(3S) \rightarrow \gamma X$	
$4.8 \pm 0.5 \pm 0.6$	19 ± 3k	54	AUBERT	09AQ	BABR	$\gamma(3S) \rightarrow \gamma X$	
• • • We do not use the following data for averages, fits, limits, etc. • • •							
<8.5	90		LEES	11J	BABR	$\gamma(3S) \rightarrow X\gamma$	
$4.8 \pm 0.5 \pm 1.2$	19 ± 3k	54,55	AUBERT	08V	BABR	$\gamma(3S) \rightarrow \gamma X$	
<4.3	90		56	ARTUSO	05	CLEO	
						$e^+ e^- \rightarrow \gamma X$	

⁵⁴ Assuming $\Gamma_{\eta_b(1S)} = 10$ MeV.

⁵⁵ Systematic error re-evaluated by AUBERT 09AQ.

⁵⁶ Superseded by BONVICINI 10.

$\Gamma(\gamma A^0 \rightarrow \gamma \text{hadrons})/\Gamma_{\text{total}}$
($0.3 \text{ GeV} < m_{A^0} < 7 \text{ GeV}$)

Γ_{28}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 8 \times 10^{-5}$	90	57 LEES	11H BABR	$\gamma(3S) \rightarrow \gamma \text{ hadrons}$

⁵⁷ For a narrow scalar or pseudoscalar, A^0 , excluding known resonances, with mass in the range $0.3\text{--}7$ GeV. Measured 90% CL limits as a function of m_{A^0} range from 1×10^{-6} to 8×10^{-5} .

$\Gamma(\gamma X \rightarrow \gamma + \geq 4 \text{ prongs})/\Gamma_{\text{total}}$
($1.5 \text{ GeV} < m_X < 5.0 \text{ GeV}$)

Γ_{29}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<2.2	95	ROSNER	07A CLEO	$e^+ e^- \rightarrow \gamma X$

$\Gamma(\gamma A^0 \rightarrow \gamma \mu^+ \mu^-)/\Gamma_{\text{total}}$

Γ_{30}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<5.5	90	58 AUBERT	09Z BABR	$e^+ e^- \rightarrow \gamma A^0 \rightarrow \gamma \mu^+ \mu^-$

⁵⁸ For a narrow scalar or pseudoscalar, A^0 , with mass in the range $212\text{--}9300$ MeV, excluding J/ψ and $\psi(2S)$. Measured 90% CL limits as a function of m_{A^0} range from $0.27\text{--}5.5 \times 10^{-6}$.

$\Gamma(\gamma A^0 \rightarrow \gamma \tau^+ \tau^-)/\Gamma_{\text{total}}$

Γ_{31}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.6 \times 10^{-4}$	90	59 AUBERT	09P BABR	$e^+ e^- \rightarrow \gamma A^0 \rightarrow \gamma \tau^+ \tau^-$

⁵⁹ For a narrow scalar or pseudoscalar, A^0 , with $M(\tau^+ \tau^-)$ in the ranges $4.03\text{--}9.52$ and $9.61\text{--}10.10$ GeV. Measured 90% CL limits as a function of $M(\tau^+ \tau^-)$ range from $1.5\text{--}16 \times 10^{-5}$.

— LEPTON FAMILY NUMBER (*LF*) VIOLATING MODES —

$\Gamma(e^\pm \tau^\mp)/\Gamma_{\text{total}}$

Γ_{32}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<4.2	90	LEES	10B BABR	$e^+ e^- \rightarrow e^\pm \tau^\mp$

$\Gamma(e^\pm \mu^\mp)/\Gamma_{\text{total}}$

Γ_{33}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 3.6 \times 10^{-7}$	90	LEES	22A BABR	$e^+ e^- \rightarrow e^\pm \mu^\mp$

$\Gamma(\mu^\pm \tau^\mp)/\Gamma_{\text{total}}$

Γ_{34}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 3.1	90	LEES	10B BABR	$e^+ e^- \rightarrow \mu^\pm \tau^\mp$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<20.3	95	LOVE	08A CLEO	$e^+ e^- \rightarrow \mu^\pm \tau^\mp$
-------	----	------	----------	--

$\tau(3S)$ REFERENCES

SHAMOV	23	PL B839 137766	A.G. Shamov, O.L. Rezanova	(NOVO, NOVOU)
LEES	22A	PRL 128 091804	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	20E	PRL 125 241801	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	14G	PR D89 111102	J.P. Lees <i>et al.</i>	(BABAR Collab.)
GE	11	PR D84 032008	J.Y. Ge <i>et al.</i>	(CLEO Collab.)
KORNICER	11	PR D83 054003	M. Kornicer <i>et al.</i>	(CLEO Collab.)
LEES	11C	PR D84 011104	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11H	PRL 107 221803	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11K	PR D84 091101	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11L	PR D84 092003	J.P. Lees <i>et al.</i>	(BABAR Collab.)
BONVICINI	10	PR D81 031104	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
LEES	10B	PRL 104 151802	J.P. Lees <i>et al.</i>	(BABAR Collab.)
AUBERT	09AQ	PRL 103 161801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09P	PRL 103 181801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09Z	PRL 103 081803	B. Aubert <i>et al.</i>	(BABAR Collab.)
BHARI	09	PR D79 011103	S.R. Bhari <i>et al.</i>	(CLEO Collab.)
ASNER	08A	PR D78 091103	D.M. Asner <i>et al.</i>	(CLEO Collab.)
AUBERT	08BP	PR D78 112002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08V	PRL 101 071801	B. Aubert <i>et al.</i>	(BABAR Collab.)
HE	08A	PRL 101 192001	Q. He <i>et al.</i>	(CLEO Collab.)
LOVE	08A	PRL 101 201601	W. Love <i>et al.</i>	(CLEO Collab.)
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
BESSON	07	PRL 98 052002	D. Besson <i>et al.</i>	(CLEO Collab.)
ROSNER	07A	PR D76 117102	J.L. Rosner <i>et al.</i>	(CLEO Collab.)
BESSON	06A	PR D74 012003	D. Besson <i>et al.</i>	(CLEO Collab.)
ROSNER	06	PRL 96 092003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)
ADAMS	05	PRL 94 012001	G.S. Adams <i>et al.</i>	(CLEO Collab.)
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>	
BUTLER	94B	PR D49 40	F. Butler <i>et al.</i>	(CLEO Collab.)
WU	93	PL B301 307	Q.W. Wu <i>et al.</i>	(CUSB Collab.)
HEINTZ	92	PR D46 1928	U. Heintz <i>et al.</i>	(CUSB II Collab.)
BROCK	91	PR D43 1448	I.C. Brock <i>et al.</i>	(CLEO Collab.)
HEINTZ	91	PRL 66 1563	U. Heintz <i>et al.</i>	(CUSB Collab.)
MORRISON	91	PRL 67 1696	R.J. Morrison <i>et al.</i>	(CLEO Collab.)
NARAIN	91	PRL 66 3113	M. Narain <i>et al.</i>	(CUSB Collab.)
CHEN	89B	PR D39 3528	W.Y. Chen <i>et al.</i>	(CLEO Collab.)
KAARSBERG	89	PRL 62 2077	T.M. Kaarsberg <i>et al.</i>	(CUSB Collab.)
BUCHMUEL...	88	HE e^+e^- Physics 412	W. Buchmueller, S. Cooper	(HANN, DESY, MIT)
Editors: A. Ali and P. Soeding, World Scientific, Singapore				
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)
BARU	86B	ZPHY C32 622 (errat.)	S.E. Baru <i>et al.</i>	(NOVO)
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)
		Translated from YAF 41 733.		
ARTAMONOV	84	PL 137B 272	A.S. Artamonov <i>et al.</i>	(NOVO)
GILES	84B	PR D29 1285	R. Giles <i>et al.</i>	(CLEO Collab.)
ANDREWS	83	PRL 50 807	D.E. Andrews <i>et al.</i>	(CLEO Collab.)
GREEN	82	PRL 49 617	J. Green <i>et al.</i>	(CLEO Collab.)
MAGERAS	82	PL 118B 453	G. Mageras <i>et al.</i>	(COLU, CORN, LSU+)