



$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+) \text{ Status: } ****$$

The parity has not actually been measured, but + is of course expected.

We have omitted some results that have been superseded by later experiments. See our earlier editions.

Ξ⁻ MASS

The fit uses the Ξ⁻, Ξ⁺, and Ξ⁰ masses and the Ξ⁻ - Ξ⁺ mass difference. It assumes that the Ξ⁻ and Ξ⁺ masses are the same.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1321.71 ± 0.07 OUR FIT				
1321.70 ± 0.08 ± 0.05	2478 ± 68	ABDALLAH	06E DLPH	from Z decays
• • •				We do not use the following data for averages, fits, limits, etc. • • •
1321.46 ± 0.34	632	DIBIANCA	75 DBC	4.9 GeV/c K⁻ d
1321.12 ± 0.41	268	WILQUET	72 HLBC	
1321.87 ± 0.51	195	¹ GOLDWASSER	70 HBC	5.5 GeV/c K⁻ p
1321.67 ± 0.52	6	CHIEN	66 HBC	6.9 GeV/c p̄p
1321.4 ± 1.1	299	LONDON	66 HBC	
1321.3 ± 0.4	149	PJERROU	65B HBC	
1321.1 ± 0.3	241	² BADIER	64 HBC	
1321.4 ± 0.4	517	² JAUNEAU	63D FBC	
1321.1 ± 0.65	62	² SCHNEIDER	63 HBC	

¹ GOLDWASSER 70 uses $m_\Lambda = 1115.58$ MeV.

² These masses have been increased 0.09 MeV because the Λ mass increased.

Ξ⁺ MASS

The fit uses the Ξ⁻, Ξ⁺, and Ξ⁰ masses and the Ξ⁻ - Ξ⁺ mass difference. It assumes that the Ξ⁻ and Ξ⁺ masses are the same.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1321.71 ± 0.07 OUR FIT				
1321.73 ± 0.08 ± 0.05	2256 ± 63	ABDALLAH	06E DLPH	from Z decays
• • •				We do not use the following data for averages, fits, limits, etc. • • •
1321.6 ± 0.8	35	VOTRUBA	72 HBC	10 GeV/c K⁺ p
1321.2 ± 0.4	34	STONE	70 HBC	
1320.69 ± 0.93	5	CHIEN	66 HBC	6.9 GeV/c p̄p

$$(m_{\Xi^-} - m_{\Xi^+}) / m_{\Xi^-}$$

A test of CPT invariance.

VALUE	DOCUMENT ID	TECN	COMMENT
(-2.5 ± 8.7) × 10⁻⁵	ABDALLAH	06E DLPH	from Z decays

Ξ^- MEAN LIFE

Measurements with an error $> 0.2 \times 10^{-10}$ s or with systematic errors not included have been omitted.

<u>VALUE (10^{-10} s)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.639±0.015 OUR AVERAGE				
1.65 ±0.07 ±0.12	2478 ± 68	ABDALLAH	06E DLPH	from Z decays
1.652±0.051	32k	BOURQUIN	84 SPEC	Hyperon beam
1.665±0.065	41k	BOURQUIN	79 SPEC	Hyperon beam
1.609±0.028	4286	HEMINGWAY	78 HBC	4.2 GeV/c $K^- p$
1.67 ±0.08		DIBIANCA	75 DBC	4.9 GeV/c $K^- d$
1.63 ±0.03	4303	BALTAY	74 HBC	1.75 GeV/c $K^- p$
1.73 ^{+0.08} _{-0.07}	680	MAYEUR	72 HLBC	2.1 GeV/c K^-
1.61 ±0.04	2610	DAUBER	69 HBC	
1.80 ±0.16	299	LONDON	66 HBC	
1.70 ±0.12	246	PJERROU	65B HBC	
1.69 ±0.07	794	HUBBARD	64 HBC	
1.86 ^{+0.15} _{-0.14}	517	JAUNEAU	63D FBC	

Ξ^+ MEAN LIFE

<u>VALUE (10^{-10} s)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.70±0.08±0.12				
	2256 ± 63	ABDALLAH	06E DLPH	from Z decays
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1.55 ^{+0.35} _{-0.20}	35	¹ VOTRUBA	72 HBC	10 GeV/c $K^+ p$
1.6 ±0.3	34	STONE	70 HBC	
1.9 ^{+0.7} _{-0.5}	12	¹ SHEN	67 HBC	
1.51±0.55	5	¹ CHIEN	66 HBC	6.9 GeV/c $\bar{p} p$

¹The error is statistical only.

$$(\tau_{\Xi^-} - \tau_{\Xi^+}) / \tau_{\Xi^-}$$

A test of CPT invariance.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.01±0.07	ABDALLAH	06E DLPH	from Z decays

Ξ^- MAGNETIC MOMENT

See the "Quark Model" review.

<u>VALUE (μ_N)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.6507±0.0025 OUR AVERAGE				
-0.6505±0.0025	4.36M	DURYEA	92 SPEC	800 GeV p Be
-0.661 ±0.036 ±0.036	44k	TROST	89 SPEC	$\Xi^- \sim 250$ GeV
-0.69 ±0.04	218k	RAMEIKA	84 SPEC	400 GeV p Be

• • • We do not use the following data for averages, fits, limits, etc. • • •

-0.674 ±0.021 ±0.020	122k	HO	90	SPEC	See
					DURYEY 92
-2.1 ±0.8	2436	COOL	74	OSPK	1.8 GeV/c K ⁻ p
-0.1 ±2.1	2724	BINGHAM	70B	OSPK	1.8 GeV/c K ⁻ p

Ξ^+ MAGNETIC MOMENT

See the "Quark Model" review.

VALUE (μ_N)	EVTS	DOCUMENT ID	TECN	COMMENT
+0.657±0.028±0.020	70k	HO	90	SPEC 800 GeV pBe

$$(\mu_{\Xi^-} + \mu_{\Xi^+}) / |\mu_{\Xi^-}|$$

A test of *CPT* invariance. We calculate this from the Ξ^- and Ξ^+ magnetic moments above.

VALUE	DOCUMENT ID
+0.01±0.05 OUR EVALUATION	

Ξ^- DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $\Lambda\pi^-$	(99.887±0.035) %	
Γ_2 $\Sigma^-\gamma$	(1.27 ±0.23) × 10 ⁻⁴	
Γ_3 $\Lambda e^- \bar{\nu}_e$	(5.63 ±0.31) × 10 ⁻⁴	
Γ_4 $\Lambda\mu^- \bar{\nu}_\mu$	(3.5 ^{+3.5} _{-2.2}) × 10 ⁻⁴	
Γ_5 $\Sigma^0 e^- \bar{\nu}_e$	(8.7 ±1.7) × 10 ⁻⁵	
Γ_6 $\Sigma^0 \mu^- \bar{\nu}_\mu$	< 8 × 10 ⁻⁴	90%
Γ_7 $\Xi^0 e^- \bar{\nu}_e$	< 2.59 × 10 ⁻⁴	90%

$\Delta S = 2$ forbidden (*S2*) modes

Γ_8 $n\pi^-$	<i>S2</i> < 1.9	× 10 ⁻⁵	90%
Γ_9 $ne^- \bar{\nu}_e$	<i>S2</i> < 3.2	× 10 ⁻³	90%
Γ_{10} $n\mu^- \bar{\nu}_\mu$	<i>S2</i> < 1.5	%	90%
Γ_{11} $p\pi^-\pi^-$	<i>S2</i> < 4	× 10 ⁻⁴	90%
Γ_{12} $p\pi^- e^- \bar{\nu}_e$	<i>S2</i> < 4	× 10 ⁻⁴	90%
Γ_{13} $p\pi^- \mu^- \bar{\nu}_\mu$	<i>S2</i> < 4	× 10 ⁻⁴	90%
Γ_{14} $p\mu^- \mu^-$	<i>L</i> < 4	× 10 ⁻⁸	90%

CONSTRAINED FIT INFORMATION

An overall fit to 4 branching ratios uses 5 measurements and one constraint to determine 5 parameters. The overall fit has a $\chi^2 = 1.0$ for 1 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-6			
x_3	-8	0		
x_4	-99	0	-1	
x_5	-5	0	0	0
	x_1	x_2	x_3	x_4

Ξ^- BRANCHING RATIOS

A number of early results have been omitted.

$\Gamma(\Sigma^- \gamma) / \Gamma(\Lambda \pi^-)$

Γ_2 / Γ_1

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.27 ± 0.24 OUR FIT				
1.27 ± 0.23 OUR AVERAGE				
1.22 ± 0.23 ± 0.06	211	¹ DUBBS	94 E761	Ξ^- 375 GeV
2.27 ± 1.02	9	BIAGI	87B SPEC	SPS hyperon beam

¹ DUBBS 94 also finds weak evidence that the asymmetry parameter α_γ is positive ($\alpha_\gamma = 1.0 \pm 1.3$).

$\Gamma(\Lambda e^- \bar{\nu}_e) / \Gamma(\Lambda \pi^-)$

Γ_3 / Γ_1

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.564 ± 0.031 OUR FIT				
0.564 ± 0.031	2857	BOURQUIN	83 SPEC	SPS hyperon beam
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.30 ± 0.13	11	THOMPSON	80 ASPK	Hyperon beam

$\Gamma(\Lambda \mu^- \bar{\nu}_\mu) / \Gamma(\Lambda \pi^-)$

Γ_4 / Γ_1

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.35^{+0.35}_{-0.22} OUR FIT					
0.35 ± 0.35		1	YEH	74 HBC	Effective denom.=2859
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 2.3	90	0	THOMPSON	80 ASPK	Effective denom.=1017
< 1.3			DAUBER	69 HBC	
< 12			BERGE	66 HBC	

$\Gamma(\Sigma^0 e^- \bar{\nu}_e)/\Gamma(\Lambda\pi^-)$ Γ_5/Γ_1

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.087±0.017 OUR FIT				
0.087±0.017	154	BOURQUIN 83	SPEC	SPS hyperon beam

$[\Gamma(\Lambda e^- \bar{\nu}_e) + \Gamma(\Sigma^0 e^- \bar{\nu}_e)]/\Gamma(\Lambda\pi^-)$ $(\Gamma_3+\Gamma_5)/\Gamma_1$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.651±0.031	3011	¹ BOURQUIN 83	SPEC	SPS hyperon beam
0.68 ±0.22	17	² DUCLOS 71	OSPK	

¹See the separate BOURQUIN 83 values for $\Gamma(\Lambda e^- \bar{\nu}_e)/\Gamma(\Lambda\pi^-)$ and $\Gamma(\Sigma^0 e^- \bar{\nu}_e)/\Gamma(\Lambda\pi^-)$ above.

²DUCLOS 71 cannot distinguish Σ^0 's from Λ 's. The Cabibbo theory predicts the Σ^0 rate is about a factor 6 smaller than the Λ rate.

$\Gamma(\Sigma^0 \mu^- \bar{\nu}_\mu)/\Gamma(\Lambda\pi^-)$ Γ_6/Γ_1

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<0.76	90	0	YEH 74	HBC	Effective denom.=3026

• • • We do not use the following data for averages, fits, limits, etc. • • •

<5		BERGE 66	HBC	
----	--	----------	-----	--

$\Gamma(\Xi^0 e^- \bar{\nu}_e)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.59 × 10⁻⁴	90	ABLIKIM 21AH	BES3	$J/\psi \rightarrow \Xi \bar{\Xi}$

$\Gamma(\Xi^0 e^- \bar{\nu}_e)/\Gamma(\Lambda\pi^-)$ Γ_7/Γ_1

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<2.3 × 10 ⁻³	90	YEH 74	HBC	Effective denom.=1000

$\Gamma(n\pi^-)/\Gamma(\Lambda\pi^-)$ Γ_8/Γ_1

$\Delta S=2$. Forbidden in first-order weak interaction.

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<0.019	90		BIAGI 82B	SPEC	SPS hyperon beam

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.0	90	0	YEH 74	HBC	Effective denom.=760
<1.1			DAUBER 69	HBC	
<5.0			FERRO-LUZZI 63	HBC	

$\Gamma(ne^- \bar{\nu}_e)/\Gamma(\Lambda\pi^-)$ Γ_9/Γ_1

$\Delta S=2$. Forbidden in first-order weak interaction.

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 3.2	90	0	YEH 74	HBC	Effective denom.=715

• • • We do not use the following data for averages, fits, limits, etc. • • •

<10	90		BINGHAM 65	RVUE	
-----	----	--	------------	------	--

$\Gamma(n\mu^-\bar{\nu}_\mu)/\Gamma(\Lambda\pi^-)$ Γ_{10}/Γ_1

$\Delta S=2$. Forbidden in first-order weak interaction.

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<15.3	90	0	YEH 74	HBC	Effective denom.=150

$\Gamma(p\pi^-\pi^-)/\Gamma(\Lambda\pi^-)$ Γ_{11}/Γ_1

$\Delta S=2$. Forbidden in first-order weak interaction.

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<3.7	90	0	YEH 74	HBC	Effective denom.=6200

$\Gamma(p\pi^-e^-\bar{\nu}_e)/\Gamma(\Lambda\pi^-)$ Γ_{12}/Γ_1

$\Delta S=2$. Forbidden in first-order weak interaction.

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<3.7	90	0	YEH 74	HBC	Effective denom.=6200

$\Gamma(p\pi^-\mu^-\bar{\nu}_\mu)/\Gamma(\Lambda\pi^-)$ Γ_{13}/Γ_1

$\Delta S=2$. Forbidden in first-order weak interaction.

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<3.7	90	0	YEH 74	HBC	Effective denom.=6200

$\Gamma(p\mu^-\mu^-)/\Gamma(\Lambda\pi^-)$ Γ_{14}/Γ_1

A $\Delta L=2$ decay, forbidden by total lepton number conservation.

VALUE (units 10^{-8})	CL%	DOCUMENT ID	TECN	COMMENT
<4.0	90	RAJARAM 05	HYCP	p Cu, 800 GeV

••• We do not use the following data for averages, fits, limits, etc. •••

<3.7 × 10⁴ 90 ¹LITTENBERG 92B HBC Uses YEH 74 data

¹This LITTENBERG 92B limit and the identical YEH 74 limits for the preceding three modes all result from nonobservance of any 3-prong decays of the Ξ^- . One could as well apply the limit to the *sum* of the four modes.

Ξ DECAY PARAMETERS

See the "Note on Baryon Decay Parameters" in the neutron Listings.

$\alpha(\Xi^-)\alpha_-(\Lambda)$

OUR FIT value is obtained from measurements of $\alpha(\Xi^-)$, $\alpha_-(\Lambda)$, and $\alpha(\Xi^-)\alpha_-(\Lambda)$.


VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

−0.291 ±0.004 **OUR FIT** Error includes scale factor of 1.8.

−0.294 ±0.005 **OUR AVERAGE** Error includes scale factor of 1.7. See the ideogram below.

−0.2963±0.0042	189k	LUK	00	E756	p Be, 800 GeV
−0.2894±0.0073	63k	¹ LUK	00	E756	p Be, 800 GeV
−0.303 ±0.004 ±0.004	192k	RAMEIKA	86	SPEC	400 GeV p Be
−0.257 ±0.020	11k	ASTON	85B	LASS	11 GeV/ c $K^- p$
−0.260 ±0.017	21k	BENSINGER	85	MPS	5 GeV/ c $K^- p$
−0.299 ±0.007	150k	BIAGI	82	SPEC	SPS hyperon beam
−0.315 ±0.026	9046	CLELAND	80C	ASPK	BNL hyperon beam
−0.239 ±0.021	6599	HEMINGWAY	78	HBC	4.2 GeV/ c $K^- p$
−0.243 ±0.025	4303	BALTAY	74	HBC	1.75 GeV/ c $K^- p$
−0.252 ±0.032	2436	COOL	74	OSPK	1.8 GeV/ c $K^- p$
−0.253 ±0.028	2781	DAUBER	69	HBC	

¹This LUK 00 value is for $\alpha(\Xi^+) \alpha_+(\bar{\Lambda})$. We assume CP conservation here by including it in the average for $\alpha(\Xi^-) \alpha_-(\Lambda)$. But see the second data block below for the CP test.



$\alpha(\Xi^-)$ for $\Xi^- \rightarrow \Lambda \pi^-$

OUR FIT value is obtained from measurements of $\alpha(\Xi^-)$, $\alpha_-(\Lambda)$, and $\alpha(\Xi^-) \alpha_-(\Lambda)$.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-0.390 ± 0.007 OUR FIT				Error includes scale factor of 2.0.
$-0.376 \pm 0.007 \pm 0.003$	73k	ABLIKIM	22AD BES3	$J/\psi \rightarrow \Xi \Xi \rightarrow \Lambda \bar{\Lambda} \pi \pi$
••• We do not use the following data for averages, fits, limits, etc. •••				
$-0.344 \pm 0.025 \pm 0.007$	5.4k	ABLIKIM	22BE BES3	$\psi(3686) \rightarrow \Xi \Xi \rightarrow \Lambda \bar{\Lambda} \pi \pi$

$\alpha(\Xi^+)$ for $\Xi^+ \rightarrow \bar{\Lambda} \pi^+$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$0.371 \pm 0.007 \pm 0.002$	73k	ABLIKIM	22AD BES3	$J/\psi \rightarrow \Xi \Xi \rightarrow \Lambda \bar{\Lambda} \pi \pi$
••• We do not use the following data for averages, fits, limits, etc. •••				
$0.355 \pm 0.025 \pm 0.002$	5.4k	ABLIKIM	22BE BES3	$\psi(3686) \rightarrow \Xi \Xi \rightarrow \Lambda \bar{\Lambda} \pi \pi$

$(\alpha + \bar{\alpha}) / (\alpha - \bar{\alpha})$ for $\Xi^- \rightarrow \Lambda \pi^-$, $\Xi^+ \rightarrow \bar{\Lambda} \pi^+$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$6 \pm 13 \pm 6$	73k	ABLIKIM	22AD BES3	$J/\psi \rightarrow \Xi \Xi \rightarrow \Lambda \bar{\Lambda} \pi \pi$
••• We do not use the following data for averages, fits, limits, etc. •••				
$-15 \pm 51 \pm 10$	5.4k	ABLIKIM	22BE BES3	$\psi(3686) \rightarrow \Xi \Xi \rightarrow \Lambda \bar{\Lambda} \pi \pi$

$$\frac{[\alpha(\Xi^-)\alpha_\Lambda - \alpha(\Xi^+)\alpha_\Lambda(\bar{\Lambda})]}{[\alpha(\Xi^-)\alpha_\Lambda + \alpha(\Xi^+)\alpha_\Lambda(\bar{\Lambda})]}$$

This is zero if CP is conserved. The α 's are the decay-asymmetry parameters for $\Xi^- \rightarrow \Lambda\pi^-$ and $\Lambda \rightarrow p\pi^-$ and for $\Xi^+ \rightarrow \bar{\Lambda}\pi^+$ and $\bar{\Lambda} \rightarrow \bar{p}\pi^+$.

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.0 ± 5.1 ± 4.4	158M	HOLMSTROM 04	HYCP	p Cu, 800 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
+120 ± 140	252k	LUK	00 E756	p Be, 800 GeV

ϕ ANGLE FOR $\Xi^- \rightarrow \Lambda\pi^-$

$$(\tan\phi = \beta/\gamma)$$

VALUE ($^\circ$)	EVTS	DOCUMENT ID	TECN	COMMENT
− 1.2 ± 1.0 OUR AVERAGE				Error includes scale factor of 1.4. See the ideogram below.
0.63 ± 1.09 ± 0.52	73k	¹ ABLIKIM 22AD	BES3	$J/\psi \rightarrow \Xi\Xi \rightarrow \Lambda\bar{\Lambda}\pi\pi$
− 2.39 ± 0.64 ± 0.64	144M	² HUANG 04	HYCP	p Cu, 800 GeV
− 1.61 ± 2.66 ± 0.37	1.35M	³ CHAKRAVO... 03	E756	p Be, 800 GeV
5 ± 10	11k	ASTON 85B	LASS	$K^- p$
14.7 ± 16.0	21k	⁴ BENSINGER 85	MPS	5 GeV/c $K^- p$
11 ± 9	4303	BALTAY 74	HBC	1.75 GeV/c $K^- p$
5 ± 16	2436	COOL 74	OSPK	1.8 GeV/c $K^- p$
− 14 ± 11	2781	DAUBER 69	HBC	Uses $\alpha_\Lambda = 0.647 \pm 0.020$
0 ± 12	1004	⁵ BERGE 66	HBC	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.32 ± 4.24 ± 0.17	5.4k	⁶ ABLIKIM 22BE	BES3	$\psi(3686) \rightarrow \Xi\Xi \rightarrow \Lambda\bar{\Lambda}\pi\pi$
− 26 ± 30	2724	BINGHAM 70B	OSPK	
0 ± 20.4	364	⁵ LONDON 66	HBC	Using $\alpha_\Lambda = 0.62$
54 ± 30	356	⁵ CARMONY 64B	HBC	

¹ Converted from radians to degrees. ABLIKIM 22AD reports a value of $0.011 \pm 0.019 \pm 0.009$ radians.


² From this result and α_Ξ , HUANG 04 gets $\beta_\Xi = -0.037 \pm 0.011 \pm 0.010$ and $\gamma_\Xi = 0.888 \pm 0.0004 \pm 0.006$. And the strong p-s phase difference for $\Lambda\pi^-$ scattering is $(4.6 \pm 1.4 \pm 1.2)^\circ$.

³ From this result and α_Ξ , CHAKRAVORTY 03 obtains $\beta_\Xi = -0.025 \pm 0.042 \pm 0.006$ and $\gamma_\Xi = 0.889 \pm 0.001 \pm 0.007$. And the strong p-s phase difference for $\Lambda\pi^-$ scattering is $(3.17 \pm 5.28 \pm 0.73)^\circ$.

⁴ BENSINGER 85 used $\alpha_\Lambda = 0.642 \pm 0.013$.

⁵ The errors have been multiplied by 1.2 due to approximations used for the Ξ polarization; see DAUBER 69 for a discussion.

⁶ Converted from radians to degrees. ABLIKIM 22BE reports a value of $0.023 \pm 0.074 \pm 0.003$ radians.



ϕ ANGLE FOR $\Xi^+ \rightarrow \bar{\Lambda} \pi^+$ ($\tan \phi = \beta/\gamma$)

VALUE ($^\circ$)	EVTS	DOCUMENT ID	TECN	COMMENT
$-1.20 \pm 1.09 \pm 0.40$	73k	¹ ABLIKIM 22AD	BES3	$J/\psi \rightarrow \Xi \bar{\Xi} \rightarrow \Lambda \bar{\Lambda} \pi \pi$
$-7.05 \pm 4.18 \pm 0.23$	5.4k	² ABLIKIM 22BE	BES3	$\psi(3686) \rightarrow \Xi \bar{\Xi} \rightarrow \Lambda \bar{\Lambda} \pi \pi$

••• We do not use the following data for averages, fits, limits, etc. •••

¹ Converted from radians to degrees. ABLIKIM 22AD reports a value of $-0.021 \pm 0.019 \pm 0.007$ radians.
² Converted from radians to degrees. ABLIKIM 22BE reports a value of $-0.123 \pm 0.073 \pm 0.004$ radians.

$\Delta\Phi_{CP} = (\Phi_- + \Phi_+)/2$

VALUE ($^\circ$)	EVTS	DOCUMENT ID	TECN	COMMENT
$-0.28 \pm 0.78 \pm 0.17$	73k	¹ ABLIKIM 22AD	BES3	$J/\psi \rightarrow \Xi \bar{\Xi} \rightarrow \Lambda \bar{\Lambda} \pi \pi$
$-2.86 \pm 2.98 \pm 0.17$	5.4k	² ABLIKIM 22BE	BES3	$\psi(3686) \rightarrow \Xi \bar{\Xi} \rightarrow \Lambda \bar{\Lambda} \pi \pi$

••• We do not use the following data for averages, fits, limits, etc. •••

¹ Converted from radians to degrees. ABLIKIM 22AD reports a value of $(-0.5 \pm 1.4 \pm 0.3) \times 10^{-2}$ radians.
² Converted from radians to degrees. ABLIKIM 22BE reports a value of $(-5.0 \pm 5.2 \pm 0.3) \times 10^{-2}$ radians.

g_A / g_V FOR $\Xi^- \rightarrow \Lambda e^- \bar{\nu}_e$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-0.25 ± 0.05	1992	¹ BOURQUIN 83	SPEC	SPS hyperon beam

¹ BOURQUIN 83 assumes that $g_2 = 0$. Also, the sign has been changed to agree with our conventions, given in the "Note on Baryon Decay Parameters" in the neutron Listings.

REFERENCES

We have omitted some papers that have been superseded by later experiments. See our earlier editions.

ABLIKIM	22AD	NAT 606 64	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22BE	PR D106 L091101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21AH	PR D104 072007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABDALLAH	06E	PL B639 179	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
RAJARAM	05	PRL 94 181801	D. Rajaram <i>et al.</i>	(FNAL HyperCP Collab.)
HOLMSTROM	04	PRL 93 262001	T. Holmstrom <i>et al.</i>	(FNAL HyperCP Collab.)
HUANG	04	PRL 93 011802	M. Huang <i>et al.</i>	(FNAL HyperCP Collab.)
CHAKRAVO...	03	PRL 91 031601	A. Chakravorty <i>et al.</i>	(FNAL E756 Collab.)
LUK	00	PRL 85 4860	K.B. Luk <i>et al.</i>	(FNAL E756 Collab.)
DUBBS	94	PRL 72 808	T. Dubbs <i>et al.</i>	(FNAL E761 Collab.)
DURYEA	92	PRL 68 768	J. Duryea <i>et al.</i>	(MINN, FNAL, MICH, RUTG)
LITTENBERG	92B	PR D46 892	L.S. Littenberg, R.E. Shrock	(BNL, STON)
HO	90	PRL 65 1713	P.M. Ho <i>et al.</i>	(MICH, FNAL, MINN, RUTG)
Also		PR D44 3402	P.M. Ho <i>et al.</i>	(MICH, FNAL, MINN, RUTG)
TROST	89	PR D40 1703	L.H. Trost <i>et al.</i>	(FNAL-715 Collab.)
BIAGI	87B	ZPHY C35 143	S.F. Biagi <i>et al.</i>	(BRIS, CERN, GEVA+)
RAMEIKA	86	PR D33 3172	R. Rameika <i>et al.</i>	(RUTG, MICH, WISC+)
ASTON	85B	PR D32 2270	D. Aston <i>et al.</i>	(SLAC, CARL, CNRC, CINC)
BENSINGER	85	NP B252 561	J.R. Bensinger <i>et al.</i>	(CHIC, ELMT, FNAL+)
BOURQUIN	84	NP B241 1	M.H. Bourquin <i>et al.</i>	(BRIS, GEVA, HEIDP+)
RAMEIKA	84	PRL 52 581	R. Rameika <i>et al.</i>	(RUTG, MICH, WISC+)
BOURQUIN	83	ZPHY C21 1	M.H. Bourquin <i>et al.</i>	(BRIS, GEVA, HEIDP+)
BIAGI	82	PL 112B 265	S.F. Biagi <i>et al.</i>	(BRIS, CAVE, GEVA+)
BIAGI	82B	PL 112B 277	S.F. Biagi <i>et al.</i>	(LOQM, GEVA, RL+)
CLELAND	80C	PR D21 12	W.E. Cleland <i>et al.</i>	(PITT, BNL)
THOMPSON	80	PR D21 25	J.A. Thompson <i>et al.</i>	(PITT, BNL)
BOURQUIN	79	PL 87B 297	M.H. Bourquin <i>et al.</i>	(BRIS, GEVA, HEIDP+)
HEMINGWAY	78	NP B142 205	R.J. Hemingway <i>et al.</i>	(CERN, ZEEM, NIJM+)
DIBIANCA	75	NP B98 137	F.A. Dibianca, R.J. Endorf	(CMU)
BALTAY	74	PR D9 49	C. Baltay <i>et al.</i>	(COLU, BING) J
COOL	74	PR D10 792	R.L. Cool <i>et al.</i>	(BNL)
Also		PRL 29 1630	R.L. Cool <i>et al.</i>	(BNL)
YEH	74	PR D10 3545	N. Yeh <i>et al.</i>	(BING, COLU)
MAYEUR	72	NP B47 333	C. Mayeur <i>et al.</i>	(BRUX, CERN, TUFTS, LOUC)
VOTRUBA	72	NP B45 77	M.F. Votruba, A. Safder, T.M. Ratcliffe	(BIRM+)
WILQUET	72	PL 42B 372	G. Wilquet <i>et al.</i>	(BRUX, CERN, TUFTS+)
DUCLOS	71	NP B32 493	J. Duclos <i>et al.</i>	(CERN)
BINGHAM	70B	PR D1 3010	G.M. Bingham <i>et al.</i>	(UCSD, WASH)
GOLDWASSER	70	PR D1 1960	E.L. Goldwasser, P.F. Schultz	(ILL)
STONE	70	PL 32B 515	S.L. Stone <i>et al.</i>	(ROCH)
DAUBER	69	PR 179 1262	P.M. Dauber <i>et al.</i>	(LRL) J
SHEN	67	PL 25B 443	B.C. Shen, A. Firestone, G. Goldhaber	(UCB+)
BERGE	66	PR 147 945	J.P. Berge <i>et al.</i>	(LRL)
CHIEN	66	PR 152 1171	C.Y. Chien <i>et al.</i>	(YALE, BNL)
LONDON	66	PR 143 1034	G.W. London <i>et al.</i>	(BNL, SYRA)
BINGHAM	65	PRSL 285 202	H.H. Bingham	(CERN)
PJERROU	65B	PRL 14 275	G.M. Pjerrou <i>et al.</i>	(UCLA)
Also		Thesis	G.M. Pjerrou	(UCLA)
BADIER	64	Dubna Conf. 1 593	J. Badier <i>et al.</i>	(EPOL, SACL, ZEEM)
CARMONY	64B	PRL 12 482	D.D. Carmony <i>et al.</i>	(UCLA) J
HUBBARD	64	PR 135 B183	J.R. Hubbard <i>et al.</i>	(LRL)
FERRO-LUZZI	63	PR 130 1568	M. Ferro-Luzzi <i>et al.</i>	(LRL)
JAUNEAU	63D	Siena Conf. 4	L. Jauneau <i>et al.</i>	(EPOL, CERN, LOUC+)
Also		PL 5 261	L. Jauneau <i>et al.</i>	(EPOL, CERN, LOUC+)
SCHNEIDER	63	PL 4 360	J. Schneider	(CERN)