



$$J = 1$$

See the related review(s):

[Z Boson](#)

### Z MASS

OUR AVERAGE is given by the weighted average of the combined CDF result and the combined LEP result, assuming no correlations between CDF and LEP. The combined LEP result,  $91.1876 \pm 0.0021$  GeV, is obtained using the fit procedure and correlations as determined by the LEP Electroweak Working Group (see the note “The Z boson” and ref. LEP-SLC 06). The LEP fit is performed using the Z mass and width, the Z hadronic pole cross section, the ratios of hadronic to leptonic partial widths, and the Z pole forward-backward lepton asymmetries. This set is believed to be most free of correlations.

The Z-boson mass listed here corresponds to the mass parameter in a Breit-Wigner distribution with mass dependent width. The value is 34 MeV greater than the real part of the position of the pole (in the energy-squared plane) in the Z-boson propagator. Also the LEP experiments have generally assumed a fixed value of the  $\gamma$ -Z interferences term based on the standard model. Keeping this term as free parameter leads to a somewhat larger error on the fitted Z mass. See ACCIARRI 00Q and ABBIENDI 04G for a detailed investigation of both these issues.

<u>VALUE (GeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>91.1880 ± 0.0020 OUR AVERAGE</b>				
91.1923 ± 0.0071		1 AALTONEN	22 CDF	$E_{cm}^{p\bar{p}} = 1.8$ TeV
91.1876 ± 0.0021		2 LEP-SLC	06 LEP	$E_{cm}^{ee} = 88-94$ GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
91.084 ± 0.107		3 ANDREEV	18A H1	$e^\pm p$
91.1872 ± 0.0033		4 ABBIENDI	04G OPAL	$E_{cm}^{ee} = \text{LEP1} +$ 130-209 GeV
91.272 ± 0.032 ± 0.033		5 ACHARD	04C L3	$E_{cm}^{ee} = 183-209$ GeV
91.1852 ± 0.0030	4.57M	6 ABBIENDI	01A OPAL	$E_{cm}^{ee} = 88-94$ GeV
91.1863 ± 0.0028	4.08M	7 ABREU	00F DLPH	$E_{cm}^{ee} = 88-94$ GeV
91.1898 ± 0.0031	3.96M	8 ACCIARRI	00C L3	$E_{cm}^{ee} = 88-94$ GeV
91.1875 ± 0.0039	3.97M	9 ACCIARRI	00Q L3	$E_{cm}^{ee} = \text{LEP1} +$ 130-189 GeV
91.1885 ± 0.0031	4.57M	10 BARATE	00C ALEP	$E_{cm}^{ee} = 88-94$ GeV
91.151 ± 0.008		11 MIYABAYASHI	95 TOPZ	$E_{cm}^{ee} = 57.8$ GeV
91.74 ± 0.28 ± 0.93	156	12 ALITTI	92B UA2	$E_{cm}^{p\bar{p}} = 630$ GeV
90.9 ± 0.3 ± 0.2	188	13 ABE	89C CDF	$E_{cm}^{p\bar{p}} = 1.8$ TeV
91.14 ± 0.12	480	14 ABRAMS	89B MRK2	$E_{cm}^{ee} = 89-93$ GeV
93.1 ± 1.0 ± 3.0	24	15 ALBAJAR	89 UA1	$E_{cm}^{p\bar{p}} = 546,630$ GeV

- <sup>1</sup> AALTONEN 22 analyse Z decays in the di-muon and di-electron channels using their full Run-II data set. They obtain Z mass values of  $91192.0 \pm 6.4(\text{stat.}) \pm 4.0(\text{syst.})$  MeV and  $91194.3 \pm 13.8(\text{stat.}) \pm 7.6(\text{syst.})$  MeV, respectively. Combining these results using the systematic uncertainty contributions and their correlations as given in AALTONEN 22, we obtain an average of  $91192.3 \pm 5.8(\text{stat.}) \pm 4.1(\text{syst.})$  MeV.
- <sup>2</sup> This result combines ABBIENDI 01A, ABREU 00F, ACCIARRI 00C, BARATE 00C, taking correlated uncertainties into account.
- <sup>3</sup> ANDREEV 18A obtain this result in a combined electroweak and QCD analysis using all deep-inelastic  $e^+p$  and  $e^-p$  neutral current and charged current scattering cross sections published by the H1 Collaboration, including data with longitudinally polarized lepton beams.
- <sup>4</sup> ABBIENDI 04G obtain this result using the S-matrix formalism for a combined fit to their cross section and asymmetry data at the Z peak and their data at 130–209 GeV. The authors have corrected the measurement for the 34 MeV shift with respect to the Breit-Wigner fits.
- <sup>5</sup> ACHARD 04C select  $e^+e^- \rightarrow Z\gamma$  events with hard initial-state radiation. Z decays to  $q\bar{q}$  and muon pairs are considered. The fit results obtained in the two samples are found consistent to each other and combined considering the uncertainty due to ISR modelling as fully correlated.
- <sup>6</sup> ABBIENDI 01A error includes approximately 2.3 MeV due to statistics and 1.8 MeV due to LEP energy uncertainty. This result is included in the LEP average LEP-SLC 06.
- <sup>7</sup> The error includes 1.6 MeV due to LEP energy uncertainty. This result is included in the LEP average LEP-SLC 06.
- <sup>8</sup> The error includes 1.8 MeV due to LEP energy uncertainty. This result is included in the LEP average LEP-SLC 06.
- <sup>9</sup> ACCIARRI 00Q interpret the s-dependence of the cross sections and lepton forward-backward asymmetries in the framework of the S-matrix formalism. They fit to their cross section and asymmetry data at high energies, using the results of S-matrix fits to Z-peak data (ACCIARRI 00C) as constraints. The 130–189 GeV data constrains the  $\gamma/Z$  interference term. The authors have corrected the measurement for the 34.1 MeV shift with respect to the Breit-Wigner fits. The error contains a contribution of  $\pm 2.3$  MeV due to the uncertainty on the  $\gamma Z$  interference.
- <sup>10</sup> BARATE 00C error includes approximately 2.4 MeV due to statistics, 0.2 MeV due to experimental systematics, and 1.7 MeV due to LEP energy uncertainty. This result is included in the LEP average LEP-SLC 06.
- <sup>11</sup> MIYABAYASHI 95 combine their low energy total hadronic cross-section measurement with the ACTON 93D data and perform a fit using an S-matrix formalism. As expected, this result is below the mass values obtained with the standard Breit-Wigner parametrization.
- <sup>12</sup> Enters fit through  $W/Z$  mass ratio given in the W Particle Listings. The ALITTI 92B systematic error ( $\pm 0.93$ ) has two contributions: one ( $\pm 0.92$ ) cancels in  $m_W/m_Z$  and one ( $\pm 0.12$ ) is noncancelling. These were added in quadrature.
- <sup>13</sup> First error of ABE 89 is combination of statistical and systematic contributions; second is mass scale uncertainty.
- <sup>14</sup> ABRAMS 89B uncertainty includes 35 MeV due to the absolute energy measurement.
- <sup>15</sup> ALBAJAR 89 result is from a total sample of 33  $Z \rightarrow e^+e^-$  events.

## Z WIDTH

OUR EVALUATION is obtained using the fit procedure and correlations as determined by the LEP Electroweak Working Group (see the note “The Z boson” and ref. LEP-SLC 06). Corrections as discussed in VOUTSINAS 20 and JANOT 20 are also included.

VALUE (GeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.4955 ± 0.0023 OUR EVALUATION</b>				
2.4955 ± 0.0023		<sup>1</sup> JANOT	20	

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.4955 ± 0.0023		<sup>2</sup> VOUTSINAS	20					
2.4952 ± 0.0023		LEP-SLC	06			$E_{cm}^{ee} = 88-94$ GeV		
2.4943 ± 0.0041		<sup>3</sup> ABBIENDI	04G	OPAL		$E_{cm}^{ee} = \text{LEP1} +$ 130–209 GeV		
2.4948 ± 0.0041	4.57M	<sup>4</sup> ABBIENDI	01A	OPAL		$E_{cm}^{ee} = 88-94$ GeV		
2.4876 ± 0.0041	4.08M	<sup>5</sup> ABREU	00F	DLPH		$E_{cm}^{ee} = 88-94$ GeV		
2.5024 ± 0.0042	3.96M	<sup>6</sup> ACCIARRI	00C	L3		$E_{cm}^{ee} = 88-94$ GeV		
2.5025 ± 0.0041	3.97M	<sup>7</sup> ACCIARRI	00Q	L3		$E_{cm}^{ee} = \text{LEP1} +$ 130–189 GeV		
2.4951 ± 0.0043	4.57M	<sup>8</sup> BARATE	00C	ALEP		$E_{cm}^{ee} = 88-94$ GeV		
2.50 ± 0.21 ± 0.06		<sup>9</sup> ABREU	96R	DLPH		$E_{cm}^{ee} = 91.2$ GeV		
3.8 ± 0.8 ± 1.0	188	ABE	89C	CDF		$E_{cm}^{pp} = 1.8$ TeV		
2.42 $\begin{smallmatrix} +0.45 \\ -0.35 \end{smallmatrix}$	480	<sup>10</sup> ABRAMS	89B	MRK2		$E_{cm}^{ee} = 89-93$ GeV		
2.7 $\begin{smallmatrix} +1.2 \\ -1.0 \end{smallmatrix}$ ± 1.3	24	<sup>11</sup> ALBAJAR	89	UA1		$E_{cm}^{pp} = 546,630$ GeV		
2.7 ± 2.0 ± 1.0	25	<sup>12</sup> ANSARI	87	UA2		$E_{cm}^{pp} = 546,630$ GeV		

<sup>1</sup> JANOT 20 applies a correction to LEP-SLC 06 using an updated Bhabha cross section calculation. This result also includes a correction to account for correlated luminosity bias as presented in VOUTSINAS 20.

<sup>2</sup> VOUTSINAS 20 applies a correction to LEP-SLC 06 to account for correlated luminosity bias.

<sup>3</sup> ABBIENDI 04G obtain this result using the S-matrix formalism for a combined fit to their cross section and asymmetry data at the Z peak and their data at 130–209 GeV. The authors have corrected the measurement for the 1 MeV shift with respect to the Breit-Wigner fits.

<sup>4</sup> ABBIENDI 01A error includes approximately 3.6 MeV due to statistics, 1 MeV due to event selection systematics, and 1.3 MeV due to LEP energy uncertainty.

<sup>5</sup> The error includes 1.2 MeV due to LEP energy uncertainty.

<sup>6</sup> The error includes 1.3 MeV due to LEP energy uncertainty.

<sup>7</sup> ACCIARRI 00Q interpret the s-dependence of the cross sections and lepton forward-backward asymmetries in the framework of the S-matrix formalism. They fit to their cross section and asymmetry data at high energies, using the results of S-matrix fits to Z-peak data (ACCIARRI 00C) as constraints. The 130–189 GeV data constrains the  $\gamma/Z$  interference term. The authors have corrected the measurement for the 0.9 MeV shift with respect to the Breit-Wigner fits.

<sup>8</sup> BARATE 00C error includes approximately 3.8 MeV due to statistics, 0.9 MeV due to experimental systematics, and 1.3 MeV due to LEP energy uncertainty.

<sup>9</sup> ABREU 96R obtain this value from a study of the interference between initial and final state radiation in the process  $e^+ e^- \rightarrow Z \rightarrow \mu^+ \mu^-$ .

<sup>10</sup> ABRAMS 89B uncertainty includes 50 MeV due to the miniSAM background subtraction error.

<sup>11</sup> ALBAJAR 89 result is from a total sample of 33  $Z \rightarrow e^+ e^-$  events.

<sup>12</sup> Quoted values of ANSARI 87 are from direct fit. Ratio of Z and W production gives either  $\Gamma(Z) < (1.09 \pm 0.07) \times \Gamma(W)$ , CL = 90% or  $\Gamma(Z) = (0.82^{+0.19}_{-0.14} \pm 0.06) \times \Gamma(W)$ . Assuming Standard-Model value  $\Gamma(W) = 2.65$  GeV then gives  $\Gamma(Z) < 2.89 \pm 0.19$  or  $= 2.17^{+0.50}_{-0.37} \pm 0.16$ .

**Z DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level
$\Gamma_1$ $e^+ e^-$	( 3.3632 $\pm$ 0.0042 ) %	
$\Gamma_2$ $\mu^+ \mu^-$	( 3.3662 $\pm$ 0.0066 ) %	
$\Gamma_3$ $\tau^+ \tau^-$	( 3.3696 $\pm$ 0.0083 ) %	
$\Gamma_4$ $\ell^+ \ell^-$	[a] ( 3.3658 $\pm$ 0.0023 ) %	
$\Gamma_5$ $\mu^+ \mu^- \mu^+ \mu^-$		
$\Gamma_6$ $\ell^+ \ell^- \ell^+ \ell^-$	[b] ( 4.55 $\pm$ 0.17 ) $\times 10^{-6}$	
$\Gamma_7$ invisible	(20.000 $\pm$ 0.055 ) %	
$\Gamma_8$ hadrons	(69.911 $\pm$ 0.056 ) %	
$\Gamma_9$ $(u\bar{u} + c\bar{c})/2$	(11.6 $\pm$ 0.6 ) %	
$\Gamma_{10}$ $(d\bar{d} + s\bar{s} + b\bar{b})/3$	(15.6 $\pm$ 0.4 ) %	
$\Gamma_{11}$ $c\bar{c}$	(12.03 $\pm$ 0.21 ) %	
$\Gamma_{12}$ $b\bar{b}$	(15.12 $\pm$ 0.05 ) %	
$\Gamma_{13}$ $b\bar{b}b\bar{b}$	( 3.6 $\pm$ 1.3 ) $\times 10^{-4}$	
$\Gamma_{14}$ $g g g$	< 1.1	% CL=95%
$\Gamma_{15}$ $\pi^0 \gamma$	< 2.01	$\times 10^{-5}$ CL=95%
$\Gamma_{16}$ $\eta \gamma$	< 5.1	$\times 10^{-5}$ CL=95%
$\Gamma_{17}$ $\rho^0 \gamma$	< 4.0	$\times 10^{-6}$ CL=95%
$\Gamma_{18}$ $\omega \gamma$	< 3.9	$\times 10^{-6}$ CL=95%
$\Gamma_{19}$ $\eta'(958) \gamma$	< 4.2	$\times 10^{-5}$ CL=95%
$\Gamma_{20}$ $\phi \gamma$	< 7	$\times 10^{-7}$ CL=95%
$\Gamma_{21}$ $\gamma \gamma$	< 1.46	$\times 10^{-5}$ CL=95%
$\Gamma_{22}$ $\pi^0 \pi^0$	< 1.52	$\times 10^{-5}$ CL=95%
$\Gamma_{23}$ $\gamma \gamma \gamma$	< 2.2	$\times 10^{-6}$ CL=95%
$\Gamma_{24}$ $\pi^\pm W^\mp$	[c] < 7	$\times 10^{-5}$ CL=95%
$\Gamma_{25}$ $\rho^\pm W^\mp$	[c] < 8.3	$\times 10^{-5}$ CL=95%
$\Gamma_{26}$ $J/\psi(1S) X$	( 3.51 $^{+0.23}_{-0.25}$ ) $\times 10^{-3}$	S=1.1
$\Gamma_{27}$ $J/\psi(1S) \gamma$	< 1.2	$\times 10^{-6}$ CL=95%
$\Gamma_{28}$ $\psi(2S) X$	( 1.60 $\pm$ 0.29 ) $\times 10^{-3}$	
$\Gamma_{29}$ $\psi(2S) \gamma$	< 2.4	$\times 10^{-6}$ CL=95%
$\Gamma_{30}$ $J/\psi(1S) \ell^+ \ell^-$		
$\Gamma_{31}$ $J/\psi(1S) J/\psi(1S)$	< 2.2	$\times 10^{-6}$ CL=95%
$\Gamma_{32}$ $\chi_{c1}(1P) X$	( 2.9 $\pm$ 0.7 ) $\times 10^{-3}$	
$\Gamma_{33}$ $\chi_{c2}(1P) X$	< 3.2	$\times 10^{-3}$ CL=90%
$\Gamma_{34}$ $\Upsilon(1S) X + \Upsilon(2S) X$ $+ \Upsilon(3S) X$	( 1.0 $\pm$ 0.5 ) $\times 10^{-4}$	
$\Gamma_{35}$ $\Upsilon(1S) X$	< 4.4	$\times 10^{-5}$ CL=95%
$\Gamma_{36}$ $\Upsilon(1S) \gamma$	< 1.1	$\times 10^{-6}$ CL=95%
$\Gamma_{37}$ $\Upsilon(2S) X$	< 1.39	$\times 10^{-4}$ CL=95%
$\Gamma_{38}$ $\Upsilon(2S) \gamma$	< 1.3	$\times 10^{-6}$ CL=95%
$\Gamma_{39}$ $\Upsilon(3S) X$	< 9.4	$\times 10^{-5}$ CL=95%

$\Gamma_{40}$	$\Upsilon(3S)\gamma$		$< 2.4$	$\times 10^{-6}$	CL=95%
$\Gamma_{41}$	$\Upsilon(1, 2, 3S)\Upsilon(1, 2, 3S)$		$< 1.5$	$\times 10^{-6}$	CL=95%
$\Gamma_{42}$	$D^0\gamma$		$< 2.2$	$\times 10^{-3}$	CL=95%
$\Gamma_{43}$	$(D^0/\bar{D}^0)X$		(20.7 $\pm$ 2.0 ) %		
$\Gamma_{44}$	$D^\pm X$		(12.2 $\pm$ 1.7 ) %		
$\Gamma_{45}$	$D^*(2010)^\pm X$	[c]	(11.4 $\pm$ 1.3 ) %		
$\Gamma_{46}$	$D_{s1}(2536)^\pm X$		( 3.6 $\pm$ 0.8 ) $\times 10^{-3}$		
$\Gamma_{47}$	$D_{sJ}(2573)^\pm X$		( 5.8 $\pm$ 2.2 ) $\times 10^{-3}$		
$\Gamma_{48}$	$D^{*l}(2629)^\pm X$		searched for		
$\Gamma_{49}$	$BX$				
$\Gamma_{50}$	$B^*X$				
$\Gamma_{51}$	$B^+X$	[d]	( 6.08 $\pm$ 0.13 ) %		
$\Gamma_{52}$	$B_s^0X$	[d]	( 1.59 $\pm$ 0.13 ) %		
$\Gamma_{53}$	$B_c^+X$		searched for		
$\Gamma_{54}$	$\Lambda_c^+X$		( 1.54 $\pm$ 0.33 ) %		
$\Gamma_{55}$	$\Xi_c^0X$		seen		
$\Gamma_{56}$	$\Xi_b X$		seen		
$\Gamma_{57}$	$b$ -baryon $X$	[d]	( 1.38 $\pm$ 0.22 ) %		
$\Gamma_{58}$	anomalous $\gamma$ + hadrons	[e]	$< 3.2$	$\times 10^{-3}$	CL=95%
$\Gamma_{59}$	$e^+e^-\gamma$	[e]	$< 5.2$	$\times 10^{-4}$	CL=95%
$\Gamma_{60}$	$\mu^+\mu^-\gamma$	[e]	$< 5.6$	$\times 10^{-4}$	CL=95%
$\Gamma_{61}$	$\tau^+\tau^-\gamma$	[e]	$< 7.3$	$\times 10^{-4}$	CL=95%
$\Gamma_{62}$	$\ell^+\ell^-\gamma\gamma$	[f]	$< 6.8$	$\times 10^{-6}$	CL=95%
$\Gamma_{63}$	$q\bar{q}\gamma\gamma$	[f]	$< 5.5$	$\times 10^{-6}$	CL=95%
$\Gamma_{64}$	$\nu\bar{\nu}\gamma\gamma$	[f]	$< 3.1$	$\times 10^{-6}$	CL=95%
$\Gamma_{65}$	$e^\pm\mu^\mp$	LF	[c] $< 2.62$	$\times 10^{-7}$	CL=95%
$\Gamma_{66}$	$e^\pm\tau^\mp$	LF	[c] $< 5.0$	$\times 10^{-6}$	CL=95%
$\Gamma_{67}$	$\mu^\pm\tau^\mp$	LF	[c] $< 6.5$	$\times 10^{-6}$	CL=95%
$\Gamma_{68}$	$p e$	L,B	$< 1.8$	$\times 10^{-6}$	CL=95%
$\Gamma_{69}$	$p\mu$	L,B	$< 1.8$	$\times 10^{-6}$	CL=95%

[a]  $\ell$  indicates each type of lepton ( $e$ ,  $\mu$ , and  $\tau$ ), not sum over them.

[b] Here  $\ell$  indicates  $e$  or  $\mu$ .

[c] The value is for the sum of the charge states or particle/antiparticle states indicated.

[d] This value is updated using the product of (i) the  $Z \rightarrow b\bar{b}$  fraction from this listing and (ii) the  $b$ -hadron fraction in an unbiased sample of weakly decaying  $b$ -hadrons produced in  $Z$ -decays provided by the Heavy Flavor Averaging Group (HFLAV, [http://www.slac.stanford.edu/xorg/hflav/osc/PDG\\_2009/#FRACZ](http://www.slac.stanford.edu/xorg/hflav/osc/PDG_2009/#FRACZ)).

[e] See the Particle Listings below for the  $\gamma$  energy range used in this measurement.

[f] For  $m_{\gamma\gamma} = (60 \pm 5)$  GeV.

## Z PARTIAL WIDTHS

### $\Gamma(e^+e^-)$

$\Gamma_1$

For the LEP experiments, this parameter is not directly used in the overall fit but is derived using the fit results; see the note “The Z boson” and ref. LEP-SLC 06.

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>83.91±0.12 OUR FIT</b>				
83.66±0.20	137.0k	ABBIENDI	01A OPAL	$E_{cm}^{ee} = 88-94$ GeV
83.54±0.27	117.8k	ABREU	00F DLPH	$E_{cm}^{ee} = 88-94$ GeV
84.16±0.22	124.4k	ACCIARRI	00C L3	$E_{cm}^{ee} = 88-94$ GeV
83.88±0.19		BARATE	00C ALEP	$E_{cm}^{ee} = 88-94$ GeV
82.89±1.20±0.89		<sup>1</sup> ABE	95J SLD	$E_{cm}^{ee} = 91.31$ GeV

<sup>1</sup> ABE 95J obtain this measurement from Bhabha events in a restricted fiducial region to improve systematics. They use the values 91.187 and 2.489 GeV for the Z mass and total decay width to extract this partial width.

### $\Gamma(\mu^+\mu^-)$

$\Gamma_2$

This parameter is not directly used in the overall fit but is derived using the fit results; see the note “The Z boson” and ref. LEP-SLC 06.

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>83.99±0.18 OUR FIT</b>				
84.03±0.30	182.8k	ABBIENDI	01A OPAL	$E_{cm}^{ee} = 88-94$ GeV
84.48±0.40	157.6k	ABREU	00F DLPH	$E_{cm}^{ee} = 88-94$ GeV
83.95±0.44	113.4k	ACCIARRI	00C L3	$E_{cm}^{ee} = 88-94$ GeV
84.02±0.28		BARATE	00C ALEP	$E_{cm}^{ee} = 88-94$ GeV

### $\Gamma(\tau^+\tau^-)$

$\Gamma_3$

This parameter is not directly used in the overall fit but is derived using the fit results; see the note “The Z boson” and ref. LEP-SLC 06.

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>84.08±0.22 OUR FIT</b>				
83.94±0.41	151.5k	ABBIENDI	01A OPAL	$E_{cm}^{ee} = 88-94$ GeV
83.71±0.58	104.0k	ABREU	00F DLPH	$E_{cm}^{ee} = 88-94$ GeV
84.23±0.58	103.0k	ACCIARRI	00C L3	$E_{cm}^{ee} = 88-94$ GeV
84.38±0.31		BARATE	00C ALEP	$E_{cm}^{ee} = 88-94$ GeV

### $\Gamma(\ell^+\ell^-)$

$\Gamma_4$

$\ell$  indicates each type of lepton ( $e$ ,  $\mu$ , and  $\tau$ ), not sum over them.

In our fit  $\Gamma(\ell^+\ell^-)$  is defined as the partial Z width for the decay into a pair of massless charged leptons. This parameter is not directly used in the 5-parameter fit assuming lepton universality but is derived using the fit results. See the note “The Z boson” and ref. LEP-SLC 06.

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>83.984 ± 0.086 OUR FIT</b>				
83.82 ± 0.15	471.3k	ABBIENDI	01A OPAL	$E_{cm}^{ee} = 88-94$ GeV
83.85 ± 0.17	379.4k	ABREU	00F DLPH	$E_{cm}^{ee} = 88-94$ GeV
84.14 ± 0.17	340.8k	ACCIARRI	00C L3	$E_{cm}^{ee} = 88-94$ GeV
84.02 ± 0.15	500k	BARATE	00C ALEP	$E_{cm}^{ee} = 88-94$ GeV

### $\Gamma(\text{invisible})$

**$\Gamma_7$**

The  $Z$  boson also decays to final states invisible in any detector, for example, the decay to a neutrino pair as predicted in the Standard Model. Measurements of  $\Gamma(\text{invisible})$  fall into two categories: direct or indirect. Direct measurements look for final states with missing energy, missing momentum, or missing mass, corresponding to the invisible decay of a produced  $Z$  boson, including single-photon final states which arise from initial-state radiation. The indirect determination is based on  $Z$  lineshape analyses performed at the LEP collider, where the invisible decay width is calculated by subtracting all visible partial decay widths from the total decay width of the  $Z$  boson. Within the framework of the Standard Model these two determinations should be identical, but not in non-SM scenarios.

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>499.2 ± 1.5 OUR AVERAGE</b>				
523 ± 3 ± 16		<sup>1</sup> TUMASYAN	23E CMS	$E_{cm}^{pp} = 13$ TeV
499.0 ± 1.5		<sup>2</sup> LEP-SLC	06 LEP	$E_{cm}^{ee} = 88-94$ GeV
498 ± 12 ± 12	1791	<sup>3</sup> ACCIARRI	98G L3	$E_{cm}^{ee} = 88-94$ GeV
539 ± 26 ± 17	410	<sup>3</sup> AKERS	95C OPAL	$E_{cm}^{ee} = 88-94$ GeV
450 ± 34 ± 34	258	<sup>3</sup> BUSKULIC	93L ALEP	$E_{cm}^{ee} = 88-94$ GeV
540 ± 80 ± 40	52	<sup>3</sup> ADEVA	92 L3	$E_{cm}^{ee} = 88-94$ GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
498.1 ± 2.6		<sup>4</sup> ABBIENDI	01A OPAL	$E_{cm}^{ee} = 88-94$ GeV
498.1 ± 3.2		<sup>4</sup> ABREU	00F DLPH	$E_{cm}^{ee} = 88-94$ GeV
499.1 ± 2.9		<sup>4</sup> ACCIARRI	00C L3	$E_{cm}^{ee} = 88-94$ GeV
499.1 ± 2.5		<sup>4</sup> BARATE	00C ALEP	$E_{cm}^{ee} = 88-94$ GeV

<sup>1</sup> TUMASYAN 23E analyses leptonic  $Z$  decay modes, with the invisible  $Z$  decay identified by missing momentum.

<sup>2</sup> The LEP Collaborations perform a combined fit to their line-shape results and determine this quantity as a difference between the total width and the sum of all the visible widths, assuming lepton universality. This result combines ABBIENDI 01A, ABREU 00F, ACCIARRI 00C, BARATE 00C, taking correlated uncertainties into account.

<sup>3</sup> This analysis selects single-photon events arising from initial state radiation.

<sup>4</sup> This is an indirect determination of  $\Gamma(\text{invisible})$  from a fit to the visible  $Z$  decay modes. It is included in the determination of the LEP average LEP-SLC 06 reported above.

### $\Gamma(\text{hadrons})$

**$\Gamma_8$**

This parameter is not directly used in the 5-parameter fit assuming lepton universality, but is derived using the fit results. See the note “The  $Z$  boson” and ref. LEP-SLC 06.

<u>VALUE (MeV)</u>	<u>EVTs</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1744.4 ± 2.0 OUR FIT</b>				
1745.4 ± 3.5	4.10M	ABBIENDI	01A OPAL	$E_{cm}^{ee} = 88-94$ GeV
1738.1 ± 4.0	3.70M	ABREU	00F DLPH	$E_{cm}^{ee} = 88-94$ GeV
1751.1 ± 3.8	3.54M	ACCIARRI	00C L3	$E_{cm}^{ee} = 88-94$ GeV
1744.0 ± 3.4	4.07M	BARATE	00C ALEP	$E_{cm}^{ee} = 88-94$ GeV

## Z BRANCHING RATIOS

OUR FIT is obtained using the fit procedure and correlations as determined by the LEP Electroweak Working Group (see the note “The Z boson” and ref. LEP-SLC 06).

$\Gamma(\mu^+ \mu^-)/\Gamma(e^+ e^-)$				$\Gamma_2/\Gamma_1$
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>1.0001 ± 0.0024 OUR AVERAGE</b>				
0.9974 ± 0.0050	<sup>1</sup> AABOUD	17Q ATLS	$E_{cm}^{pp} = 7$ TeV	
1.0009 ± 0.0028	<sup>2</sup> LEP-SLC	06	$E_{cm}^{ee} = 88-94$ GeV	

<sup>1</sup> AABOUD 17Q make a precise determination of  $Z \rightarrow ee$  and  $Z \rightarrow \mu\mu$  production in the lepton pseudo-rapidity range  $|\eta| < 2.5$  and determine the ratio of the Z branching fractions  $B(Z \rightarrow ee)/B(Z \rightarrow \mu\mu) = 1.0026 \pm 0.0013 \pm 0.0048 = 1.0026 \pm 0.0050$ .

<sup>2</sup> This parameter is not directly used in the overall fit but is derived using the fit results; see the note “The Z boson” and ref. LEP-SLC 06.

$\Gamma(\tau^+ \tau^-)/\Gamma(e^+ e^-)$				$\Gamma_3/\Gamma_1$
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>1.0020 ± 0.0032 OUR AVERAGE</b>				
1.02 ± 0.06	<sup>1</sup> AAIJ	18AR LHCB	$E_{cm}^{pp} = 8$ TeV	
1.0019 ± 0.0032	<sup>2</sup> LEP-SLC	06	$E_{cm}^{ee} = 88-94$ GeV	

<sup>1</sup> AAIJ 18AR obtain the result from the ratio of the measured  $pp \rightarrow Z + X$  cross sections in the corresponding Z decay channels.

<sup>2</sup> This parameter is not directly used in the overall fit but is derived using the fit results; see the note “The Z boson” and ref. LEP-SLC 06.

$\Gamma(\tau^+ \tau^-)/\Gamma(\mu^+ \mu^-)$				$\Gamma_3/\Gamma_2$
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>1.0010 ± 0.0026 OUR AVERAGE</b>				
1.01 ± 0.05	<sup>1</sup> AAIJ	18AR LHCB	$E_{cm}^{pp} = 8$ TeV	
1.0010 ± 0.0026	<sup>2</sup> LEP-SLC	06	$E_{cm}^{ee} = 88-94$ GeV	

<sup>1</sup> AAIJ 18AR obtain the result from the ratio of the measured  $pp \rightarrow Z + X$  cross sections in the corresponding Z decay channels.

<sup>2</sup> This parameter is not directly used in the overall fit but is derived using the fit results; see the note “The Z boson” and ref. LEP-SLC 06.



$\Gamma(\ell^+ \ell^- \ell^+ \ell^-) / \Gamma_{\text{total}}$   $\Gamma_6 / \Gamma$

Here  $\ell$  indicates either  $e$  or  $\mu$ . The branching fractions in this node are given within the phase-space defined by the requirements that (i) the 4-lepton invariant mass is between 80 GeV and 100 GeV, and (ii) any opposite-sign same-flavor lepton pair has a di-lepton invariant mass larger than 4 GeV.

<u>VALUE (units <math>10^{-6}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>4.55 ± 0.17 OUR AVERAGE</b>				
4.41 ± 0.13 ± 0.27		<sup>1</sup> AAD	21AQ ATLS	$E_{\text{cm}}^{pp} = 13$ TeV
4.70 ± 0.32 ± 0.25		<sup>2</sup> AABOUD	19N ATLS	$E_{\text{cm}}^{pp} = 13$ TeV
4.83 $^{+0.23+0.35}_{-0.22-0.32}$	509	<sup>3</sup> SIRUNYAN	18BT CMS	$E_{\text{cm}}^{pp} = 13$ TeV
4.9 $^{+0.8+0.4}_{-0.7-0.2}$	39	<sup>4</sup> KHACHATRYAN	16CC CMS	$E_{\text{cm}}^{pp} = 13$ TeV
4.31 ± 0.34 ± 0.17	172	AAD	14N ATLS	$E_{\text{cm}}^{pp} = 7, 8$ TeV
4.6 $^{+1.0}_{-0.9} \pm 0.2$	28	<sup>5</sup> CHATRCHYAN	12BN CMS	$E_{\text{cm}}^{pp} = 7$ TeV

<sup>1</sup> AAD 21AQ analyze differential cross-sections in four-lepton events. Based on the measured cross section in the  $Z \rightarrow 4\ell$  channel, a branching fraction of  $B(Z \rightarrow 4\ell) = (4.41 \pm 0.13 \pm 0.23 \pm 0.09 \pm 0.12) \times 10^{-6}$  is obtained, where the uncertainties are statistical, systematic, theory and luminosity, respectively.

<sup>2</sup> AABOUD 19N reports  $(4.70 \pm 0.32 \pm 0.21 \pm 0.14) \times 10^{-6}$ , where the uncertainties are statistical, systematic, and luminosity. We have combined the latter two in quadrature.

<sup>3</sup> SIRUNYAN 18BT report the  $Z \rightarrow 4\ell$  branching fraction =  $(4.83^{+0.23+0.32}_{-0.22-0.29} \pm 0.08 \pm 0.12) \times 10^{-6}$ , where the uncertainties are statistical, systematic, due to theory, and luminosity. The last three have been added in quadrature to obtain the total systematic error.

<sup>4</sup> KHACHATRYAN 16CC reports  $(4.9^{+0.8+0.3+0.2+0.1}_{-0.7-0.2-0.1-0.1}) \times 10^{-6}$  value, where the uncertainties are statistical, systematic, theory, and due to luminosity. We have combined uncertainties in quadrature.

<sup>5</sup> CHATRCHYAN 12BN reports  $(4.2^{+0.9}_{-0.8} \pm 0.2) \times 10^{-6}$  value. Their result (both central value and uncertainties) is scaled up by 10% to account for the different phase-space definition used here (see RAINBOLT 19).

$\Gamma(\text{hadrons}) / \Gamma(e^+ e^-)$   $\Gamma_8 / \Gamma_1$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>20.804 ± 0.050 OUR FIT</b>				
20.902 ± 0.084	137.0k	<sup>1</sup> ABBIENDI	01A OPAL	$E_{\text{cm}}^{ee} = 88-94$ GeV
20.88 ± 0.12	117.8k	ABREU	00F DLPH	$E_{\text{cm}}^{ee} = 88-94$ GeV
20.816 ± 0.089	124.4k	ACCIARRI	00C L3	$E_{\text{cm}}^{ee} = 88-94$ GeV
20.677 ± 0.075		<sup>2</sup> BARATE	00C ALEP	$E_{\text{cm}}^{ee} = 88-94$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

27.0 $^{+11.7}_{-8.8}$	12	<sup>3</sup> ABRAMS	89D MRK2	$E_{\text{cm}}^{ee} = 89-93$ GeV
------------------------	----	---------------------	----------	----------------------------------

<sup>1</sup> ABBIENDI 01A error includes approximately 0.067 due to statistics, 0.040 due to event selection systematics, 0.027 due to the theoretical uncertainty in  $t$ -channel prediction, and 0.014 due to LEP energy uncertainty.

<sup>2</sup> BARATE 00C error includes approximately 0.062 due to statistics, 0.033 due to experimental systematics, and 0.026 due to the theoretical uncertainty in  $t$ -channel prediction.

<sup>3</sup> ABRAMS 89D have included both statistical and systematic uncertainties in their quoted errors.

## $\Gamma(\text{hadrons})/\Gamma(\mu^+\mu^-)$ $\Gamma_8/\Gamma_2$

OUR FIT is obtained using the fit procedure and correlations as determined by the LEP Electroweak Working Group (see the note “The Z boson” and ref. LEP-SLC 06).

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>20.785 ± 0.033 OUR FIT</b>				
20.811 ± 0.058	182.8k	<sup>1</sup> ABBIENDI	01A OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
20.65 ± 0.08	157.6k	ABREU	00F DLPH	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
20.861 ± 0.097	113.4k	ACCIARRI	00C L3	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
20.799 ± 0.056		<sup>2</sup> BARATE	00C ALEP	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
18.9 $\begin{smallmatrix} +7.1 \\ -5.3 \end{smallmatrix}$	13	<sup>3</sup> ABRAMS	89D MRK2	$E_{\text{cm}}^{ee} = 89\text{--}93$ GeV

<sup>1</sup> ABBIENDI 01A error includes approximately 0.050 due to statistics and 0.027 due to event selection systematics.

<sup>2</sup> BARATE 00C error includes approximately 0.053 due to statistics and 0.021 due to experimental systematics.

<sup>3</sup> ABRAMS 89D have included both statistical and systematic uncertainties in their quoted errors.

## $\Gamma(\text{hadrons})/\Gamma(\tau^+\tau^-)$ $\Gamma_8/\Gamma_3$

OUR FIT is obtained using the fit procedure and correlations as determined by the LEP Electroweak Working Group (see the note “The Z boson” and ref. LEP-SLC 06).

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>20.764 ± 0.045 OUR FIT</b>				
20.832 ± 0.091	151.5k	<sup>1</sup> ABBIENDI	01A OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
20.84 ± 0.13	104.0k	ABREU	00F DLPH	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
20.792 ± 0.133	103.0k	ACCIARRI	00C L3	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
20.707 ± 0.062		<sup>2</sup> BARATE	00C ALEP	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
15.2 $\begin{smallmatrix} +4.8 \\ -3.9 \end{smallmatrix}$	21	<sup>3</sup> ABRAMS	89D MRK2	$E_{\text{cm}}^{ee} = 89\text{--}93$ GeV

<sup>1</sup> ABBIENDI 01A error includes approximately 0.055 due to statistics and 0.071 due to event selection systematics.

<sup>2</sup> BARATE 00C error includes approximately 0.054 due to statistics and 0.033 due to experimental systematics.

<sup>3</sup> ABRAMS 89D have included both statistical and systematic uncertainties in their quoted errors.

## $\Gamma(\text{hadrons})/\Gamma(\ell^+\ell^-)$ $\Gamma_8/\Gamma_4$

$\ell$  indicates each type of lepton ( $e$ ,  $\mu$ , and  $\tau$ ), not sum over them.

Our fit result is obtained requiring lepton universality.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>20.767 ± 0.025 OUR FIT</b>				
20.823 ± 0.044	471.3k	<sup>1</sup> ABBIENDI	01A OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
20.730 ± 0.060	379.4k	ABREU	00F DLPH	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
20.810 ± 0.060	340.8k	ACCIARRI	00C L3	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
20.725 ± 0.039	500k	<sup>2</sup> BARATE	00C ALEP	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
18.9 $\begin{smallmatrix} +3.6 \\ -3.2 \end{smallmatrix}$	46	ABRAMS	89B MRK2	$E_{\text{cm}}^{ee} = 89\text{--}93$ GeV

- <sup>1</sup> ABBIENDI 01A error includes approximately 0.034 due to statistics and 0.027 due to event selection systematics.  
<sup>2</sup> BARATE 00C error includes approximately 0.033 due to statistics, 0.020 due to experimental systematics, and 0.005 due to the theoretical uncertainty in  $t$ -channel prediction.

### $\Gamma((u\bar{u}+c\bar{c})/2)/\Gamma(\text{hadrons})$

$\Gamma_9/\Gamma_8$

This quantity is the branching ratio of  $Z \rightarrow$  “up-type” quarks to  $Z \rightarrow$  hadrons. Except ACKERSTAFF 97T the values of  $Z \rightarrow$  “up-type” and  $Z \rightarrow$  “down-type” branchings are extracted from measurements of  $\Gamma(\text{hadrons})$ , and  $\Gamma(Z \rightarrow \gamma + \text{jets})$  where  $\gamma$  is a high-energy ( $>5$  or  $7$  GeV) isolated photon. As the experiments use different procedures and slightly different values of  $M_Z$ ,  $\Gamma(\text{hadrons})$  and  $\alpha_s$  in their extraction procedures, our average has to be taken with caution.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.166±0.009 OUR AVERAGE</b>			
0.172 <sup>+0.011</sup> <sub>-0.010</sub>	<sup>1</sup> ABBIENDI	04E OPAL	$E_{\text{cm}}^{ee} = 91.2$ GeV
0.160±0.019±0.019	<sup>2</sup> ACKERSTAFF	97T OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
0.137 <sup>+0.038</sup> <sub>-0.054</sub>	<sup>3</sup> ABREU	95X DLPH	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
0.137±0.033	<sup>4</sup> ADRIANI	93 L3	$E_{\text{cm}}^{ee} = 91.2$ GeV

- <sup>1</sup> ABBIENDI 04E select photons with energy  $> 7$  GeV and use  $\Gamma(\text{hadrons}) = 1744.4 \pm 2.0$  MeV and  $\alpha_s = 0.1172 \pm 0.002$  to obtain  $\Gamma_u = 300^{+19}_{-18}$  MeV.  
<sup>2</sup> ACKERSTAFF 97T measure  $\Gamma_{u\bar{u}}/(\Gamma_{d\bar{d}}+\Gamma_{u\bar{u}}+\Gamma_{s\bar{s}}) = 0.258 \pm 0.031 \pm 0.032$ . To obtain this branching ratio authors use  $R_c+R_b = 0.380 \pm 0.010$ . This measurement is fully negatively correlated with the measurement of  $\Gamma_{d\bar{d},s\bar{s}}/(\Gamma_{d\bar{d}}+\Gamma_{u\bar{u}}+\Gamma_{s\bar{s}})$  given in the next data block.  
<sup>3</sup> ABREU 95X use  $M_Z = 91.187 \pm 0.009$  GeV,  $\Gamma(\text{hadrons}) = 1725 \pm 12$  MeV and  $\alpha_s = 0.123 \pm 0.005$ . To obtain this branching ratio we divide their value of  $C_{2/3} = 0.91^{+0.25}_{-0.36}$  by their value of  $(3C_{1/3} + 2C_{2/3}) = 6.66 \pm 0.05$ .  
<sup>4</sup> ADRIANI 93 use  $M_Z = 91.181 \pm 0.022$  GeV,  $\Gamma(\text{hadrons}) = 1742 \pm 19$  MeV and  $\alpha_s = 0.125 \pm 0.009$ . To obtain this branching ratio we divide their value of  $C_{2/3} = 0.92 \pm 0.22$  by their value of  $(3C_{1/3} + 2C_{2/3}) = 6.720 \pm 0.076$ .

### $\Gamma((d\bar{d}+s\bar{s}+b\bar{b})/3)/\Gamma(\text{hadrons})$

$\Gamma_{10}/\Gamma_8$

This quantity is the branching ratio of  $Z \rightarrow$  “down-type” quarks to  $Z \rightarrow$  hadrons. Except ACKERSTAFF 97T the values of  $Z \rightarrow$  “up-type” and  $Z \rightarrow$  “down-type” branchings are extracted from measurements of  $\Gamma(\text{hadrons})$ , and  $\Gamma(Z \rightarrow \gamma + \text{jets})$  where  $\gamma$  is a high-energy ( $>5$  or  $7$  GeV) isolated photon. As the experiments use different procedures and slightly different values of  $M_Z$ ,  $\Gamma(\text{hadrons})$  and  $\alpha_s$  in their extraction procedures, our average has to be taken with caution.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.223±0.006 OUR AVERAGE</b>			
0.218±0.007	<sup>1</sup> ABBIENDI	04E OPAL	$E_{\text{cm}}^{ee} = 91.2$ GeV
0.230±0.010±0.010	<sup>2</sup> ACKERSTAFF	97T OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
0.243 <sup>+0.036</sup> <sub>-0.026</sub>	<sup>3</sup> ABREU	95X DLPH	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
0.243±0.022	<sup>4</sup> ADRIANI	93 L3	$E_{\text{cm}}^{ee} = 91.2$ GeV

- <sup>1</sup> ABBIENDI 04E select photons with energy  $> 7$  GeV and use  $\Gamma(\text{hadrons}) = 1744.4 \pm 2.0$  MeV and  $\alpha_s = 0.1172 \pm 0.002$  to obtain  $\Gamma_d = 381 \pm 12$  MeV.
- <sup>2</sup> ACKERSTAFF 97T measure  $\Gamma_{d\bar{d},s\bar{s}}/(\Gamma_{d\bar{d}}+\Gamma_{u\bar{u}}+\Gamma_{s\bar{s}}) = 0.371 \pm 0.016 \pm 0.016$ . To obtain this branching ratio authors use  $R_c+R_b = 0.380 \pm 0.010$ . This measurement is fully negatively correlated with the measurement of  $\Gamma_{u\bar{u}}/(\Gamma_{d\bar{d}} + \Gamma_{u\bar{u}} + \Gamma_{s\bar{s}})$  presented in the previous data block.
- <sup>3</sup> ABREU 95X use  $M_Z = 91.187 \pm 0.009$  GeV,  $\Gamma(\text{hadrons}) = 1725 \pm 12$  MeV and  $\alpha_s = 0.123 \pm 0.005$ . To obtain this branching ratio we divide their value of  $C_{1/3} = 1.62^{+0.24}_{-0.17}$  by their value of  $(3C_{1/3} + 2C_{2/3}) = 6.66 \pm 0.05$ .
- <sup>4</sup> ADRIANI 93 use  $M_Z = 91.181 \pm 0.022$  GeV,  $\Gamma(\text{hadrons}) = 1742 \pm 19$  MeV and  $\alpha_s = 0.125 \pm 0.009$ . To obtain this branching ratio we divide their value of  $C_{1/3} = 1.63 \pm 0.15$  by their value of  $(3C_{1/3} + 2C_{2/3}) = 6.720 \pm 0.076$ .

### $R_c = \Gamma(c\bar{c})/\Gamma(\text{hadrons})$

$\Gamma_{11}/\Gamma_8$

OUR FIT is obtained by a simultaneous fit to several  $c$ - and  $b$ -quark measurements as explained in the note "The Z boson" and ref. LEP-SLC 06.

The Standard Model predicts  $R_c = 0.1723$  for  $m_t = 174.3$  GeV and  $M_H = 150$  GeV.

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.1721±0.0030 OUR FIT</b>			
0.1744±0.0031±0.0021	<sup>1</sup> ABE	05F SLD	$E_{cm}^{ee}=91.28$ GeV
0.1665±0.0051±0.0081	<sup>2</sup> ABREU	00 DLPH	$E_{cm}^{ee}=88-94$ GeV
0.1698±0.0069	<sup>3</sup> BARATE	00B ALEP	$E_{cm}^{ee}=88-94$ GeV
0.180 ±0.011 ±0.013	<sup>4</sup> ACKERSTAFF	98E OPAL	$E_{cm}^{ee}=88-94$ GeV
0.167 ±0.011 ±0.012	<sup>5</sup> ALEXANDER	96R OPAL	$E_{cm}^{ee}=88-94$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.1623±0.0085±0.0209	<sup>6</sup> ABREU	95D DLPH	$E_{cm}^{ee}=88-94$ GeV

- <sup>1</sup> ABE 05F use hadronic Z decays collected during 1996–98 to obtain an enriched sample of  $c\bar{c}$  events using a double tag method. The single  $c$ -tag is obtained with a neural network trained to perform flavor discrimination using as input several signatures (corrected secondary vertex mass, vertex decay length, multiplicity and total momentum of the hemisphere). A multitag approach is used, defining 4 regions of the output value of the neural network and  $R_c$  is extracted from a simultaneous fit to the count rates of the 4 different tags. The quoted systematic error includes an uncertainty of  $\pm 0.0006$  due to the uncertainty on  $R_b$ .
- <sup>2</sup> ABREU 00 obtain this result properly combining the measurement from the  $D^{*+}$  production rate ( $R_c = 0.1610 \pm 0.0104 \pm 0.0077 \pm 0.0043$  (BR)) with that from the overall charm counting ( $R_c = 0.1692 \pm 0.0047 \pm 0.0063 \pm 0.0074$  (BR)) in  $c\bar{c}$  events. The systematic error includes an uncertainty of  $\pm 0.0054$  due to the uncertainty on the charmed hadron branching fractions.
- <sup>3</sup> BARATE 00B use exclusive decay modes to independently determine the quantities  $R_c \times f(c \rightarrow X)$ ,  $X=D^0, D^+, D_s^+$ , and  $\Lambda_c$ . Estimating  $R_c \times f(c \rightarrow \Xi_c/\Omega_c) = 0.0034$ , they simply sum over all the charm decays to obtain  $R_c = 0.1738 \pm 0.0047 \pm 0.0088 \pm 0.0075$ (BR). This is combined with all previous ALEPH measurements (BARATE 98T and BUSKULIC 94G,  $R_c = 0.1681 \pm 0.0054 \pm 0.0062$ ) to obtain the quoted value.
- <sup>4</sup> ACKERSTAFF 98E use an inclusive/exclusive double tag. In one jet  $D^{*\pm}$  mesons are exclusively reconstructed in several decay channels and in the opposite jet a slow pion (opposite charge inclusive  $D^{*\pm}$ ) tag is used. The  $b$  content of this sample is measured by the simultaneous detection of a lepton in one jet and an inclusively reconstructed  $D^{*\pm}$  meson in the opposite jet. The systematic error includes an uncertainty of  $\pm 0.006$  due to the external branching ratios.

- <sup>5</sup> ALEXANDER 96R obtain this value via direct charm counting, summing the partial contributions from  $D^0$ ,  $D^+$ ,  $D_s^+$ , and  $\Lambda_c^+$ , and assuming that strange-charmed baryons account for the 15% of the  $\Lambda_c^+$  production. An uncertainty of  $\pm 0.005$  due to the uncertainties in the charm hadron branching ratios is included in the overall systematics.
- <sup>6</sup> ABREU 95D perform a maximum likelihood fit to the combined  $p$  and  $p_T$  distributions of single and dilepton samples. The second error includes an uncertainty of  $\pm 0.0124$  due to models and branching ratios.

$$R_b = \Gamma(b\bar{b})/\Gamma(\text{hadrons}) \qquad \Gamma_{12}/\Gamma_8$$

OUR FIT is obtained by a simultaneous fit to several  $c$ - and  $b$ -quark measurements as explained in the note "The Z boson" and ref. LEP-SLC 06.

The Standard Model predicts  $R_b=0.21581$  for  $m_t=174.3$  GeV and  $M_H=150$  GeV.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.21629 ± 0.00066 OUR FIT</b>			
0.21594 ± 0.00094 ± 0.00075	<sup>1</sup> ABE	05F SLD	$E_{cm}^{ee}=91.28$ GeV
0.2174 ± 0.0015 ± 0.0028	<sup>2</sup> ACCIARRI	00 L3	$E_{cm}^{ee}=89-93$ GeV
0.2178 ± 0.0011 ± 0.0013	<sup>3</sup> ABBIENDI	99B OPAL	$E_{cm}^{ee}=88-94$ GeV
0.21634 ± 0.00067 ± 0.00060	<sup>4</sup> ABREU	99B DLPH	$E_{cm}^{ee}=88-94$ GeV
0.2159 ± 0.0009 ± 0.0011	<sup>5</sup> BARATE	97F ALEP	$E_{cm}^{ee}=88-94$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.2145 ± 0.0089 ± 0.0067	<sup>6</sup> ABREU	95D DLPH	$E_{cm}^{ee}=88-94$ GeV
0.219 ± 0.006 ± 0.005	<sup>7</sup> BUSKULIC	94G ALEP	$E_{cm}^{ee}=88-94$ GeV
0.251 ± 0.049 ± 0.030	<sup>8</sup> JACOBSEN	91 MRK2	$E_{cm}^{ee}=91$ GeV

- <sup>1</sup> ABE 05F use hadronic Z decays collected during 1996–98 to obtain an enriched sample of  $b\bar{b}$  events using a double tag method. The single  $b$ -tag is obtained with a neural network trained to perform flavor discrimination using as input several signatures (corrected secondary vertex mass, vertex decay length, multiplicity and total momentum of the hemisphere; the key tag is obtained requiring the secondary vertex corrected mass to be above the  $D$ -meson mass). ABE 05F obtain  $R_b = 0.21604 \pm 0.00098 \pm 0.00074$  where the systematic error includes an uncertainty of  $\pm 0.00012$  due to the uncertainty on  $R_c$ . The value reported here is obtained properly combining with ABE 98D. The quoted systematic error includes an uncertainty of  $\pm 0.00012$  due to the uncertainty on  $R_c$ .
- <sup>2</sup> ACCIARRI 00 obtain this result using a double-tagging technique, with a high  $p_T$  lepton tag and an impact parameter tag in opposite hemispheres.
- <sup>3</sup> ABBIENDI 99B tag  $Z \rightarrow b\bar{b}$  decays using leptons and/or separated decay vertices. The  $b$ -tagging efficiency is measured directly from the data using a double-tagging technique.
- <sup>4</sup> ABREU 99B obtain this result combining in a multivariate analysis several tagging methods (impact parameter and secondary vertex reconstruction, complemented by event shape variables). For  $R_c$  different from its Standard Model value of 0.172,  $R_b$  varies as  $-0.024 \times (R_c - 0.172)$ .
- <sup>5</sup> BARATE 97F combine the lifetime-mass hemisphere tag (BARATE 97E) with event shape information and lepton tag to identify  $Z \rightarrow b\bar{b}$  candidates. They further use  $c$ - and  $uds$ -selection tags to identify the background. For  $R_c$  different from its Standard Model value of 0.172,  $R_b$  varies as  $-0.019 \times (R_c - 0.172)$ .
- <sup>6</sup> ABREU 95D perform a maximum likelihood fit to the combined  $p$  and  $p_T$  distributions of single and dilepton samples. The second error includes an uncertainty of  $\pm 0.0023$  due to models and branching ratios.
- <sup>7</sup> BUSKULIC 94G perform a simultaneous fit to the  $p$  and  $p_T$  spectra of both single and dilepton events.
- <sup>8</sup> JACOBSEN 91 tagged  $b\bar{b}$  events by requiring coincidence of  $\geq 3$  tracks with significant impact parameters using vertex detector. Systematic error includes lifetime and decay uncertainties ( $\pm 0.014$ ).

### $\Gamma(b\bar{b}b\bar{b})/\Gamma(\text{hadrons})$

$\Gamma_{13}/\Gamma_8$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>5.2 \pm 1.9</math> OUR AVERAGE</b>			
$3.6 \pm 1.7 \pm 2.7$	<sup>1</sup> ABBIENDI	01G OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
$6.0 \pm 1.9 \pm 1.4$	<sup>2</sup> ABREU	99U DLPH	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV

<sup>1</sup> ABBIENDI 01G use a sample of four-jet events from hadronic  $Z$  decays. To enhance the  $b\bar{b}b\bar{b}$  signal, at least three of the four jets are required to have a significantly detached secondary vertex.

<sup>2</sup> ABREU 99U force hadronic  $Z$  decays into 3 jets to use all the available phase space and require a  $b$  tag for every jet. This decay mode includes primary and secondary  $4b$  production, e.g, from gluon splitting to  $b\bar{b}$ .

### $\Gamma(gg)/\Gamma(\text{hadrons})$

$\Gamma_{14}/\Gamma_8$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>&lt; 1.6 \times 10^{-2}</math></b>	95	<sup>1</sup> ABREU	96S DLPH	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV

<sup>1</sup> This branching ratio is slightly dependent on the jet-finder algorithm. The value we quote is obtained using the JADE algorithm, while using the DURHAM algorithm ABREU 96S obtain an upper limit of  $1.5 \times 10^{-2}$ .

### $\Gamma(\pi^0\gamma)/\Gamma_{\text{total}}$

$\Gamma_{15}/\Gamma$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>&lt; 2.01 \times 10^{-5}</math></b>	95	AALTONEN	14E CDF	$E_{\text{cm}}^{p\bar{p}} = 1.96$ TeV
$< 5.2 \times 10^{-5}$	95	<sup>1</sup> ACCIARRI	95G L3	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
$< 5.5 \times 10^{-5}$	95	ABREU	94B DLPH	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
$< 2.1 \times 10^{-4}$	95	DECAMP	92 ALEP	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
$< 1.4 \times 10^{-4}$	95	AKRAWY	91F OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV

<sup>1</sup> This limit is for both decay modes  $Z \rightarrow \pi^0\gamma/\gamma\gamma$  which are indistinguishable in ACCIARRI 95G.

### $\Gamma(\eta\gamma)/\Gamma_{\text{total}}$

$\Gamma_{16}/\Gamma$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 7.6 \times 10^{-5}$	95	ACCIARRI	95G L3	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
$< 8.0 \times 10^{-5}$	95	ABREU	94B DLPH	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
<b><math>&lt; 5.1 \times 10^{-5}</math></b>	95	DECAMP	92 ALEP	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
$< 2.0 \times 10^{-4}$	95	AKRAWY	91F OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV

### $\Gamma(\rho^0\gamma)/\Gamma_{\text{total}}$

$\Gamma_{17}/\Gamma$

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>&lt; 4.0 \times 10^{-6}</math></b>	95	12.5k	<sup>1</sup> AABOUD	18AU ATLS	$E_{\text{cm}}^{p\bar{p}} = 13$ TeV

<sup>1</sup> AABOUD 18AU search for the  $Z \rightarrow \rho\gamma$  decay mode where the  $\rho$  is identified through its decay  $\rho \rightarrow \pi^+\pi^-$ . In the data corresponding to  $32.3 \text{ fb}^{-1}$ , 12,583 events are selected for  $635 < m(\pi^+\pi^-) < 915$  MeV. See erratum AABOUD 23A.

### $\Gamma(\omega\gamma)/\Gamma_{\text{total}}$

$\Gamma_{18}/\Gamma$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>&lt; 3.9 \times 10^{-6}</math></b>	95	AAD	23BS ATLS	$E_{\text{cm}}^{p\bar{p}} = 13$ TeV
$< 6.5 \times 10^{-4}$	95	ABREU	94B DLPH	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\Gamma(\eta'(958)\gamma)/\Gamma_{\text{total}}$   $\Gamma_{19}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.2 \times 10^{-5}$	95	DECAMP 92	ALEP	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV

$\Gamma(\phi\gamma)/\Gamma_{\text{total}}$   $\Gamma_{20}/\Gamma$

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$<7 \times 10^{-7}$	95	3.3k	<sup>1</sup> AABOUD 18AU	ATLS	$E_{\text{cm}}^{pp} = 13$ TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<8.3 \times 10^{-6}$	95	1.0k	<sup>2</sup> AABOUD 16K	ATLS	$E_{\text{cm}}^{pp} = 13$ TeV
-----------------------	----	------	-------------------------	------	-------------------------------

<sup>1</sup> AABOUD 18AU search for the  $Z \rightarrow \phi\gamma$  decay mode where the  $\phi$  is identified through its decay  $\phi \rightarrow K^+K^-$ . In the data corresponding to  $32.3 \text{ fb}^{-1}$ , 3,364 events are selected for  $1012 < m(K^+K^-) < 1028$  MeV. See erratum AABOUD 23A.

<sup>2</sup> AABOUD 16K search for the  $Z \rightarrow \phi\gamma$  decay mode where the  $\phi$  is identified through its decay into  $K^+K^-$ . In the data corresponding to a total luminosity of  $2.7 \text{ fb}^{-1}$ , 1065 events are selected and their  $K^+K^-\gamma$  invariant mass spectrum is analyzed.

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$   $\Gamma_{21}/\Gamma$

This decay would violate the Landau-Yang theorem.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.46 \times 10^{-5}$	95	AALTONEN 14E	CDF	$E_{\text{cm}}^{pp} = 1.96$ TeV
$<5.2 \times 10^{-5}$	95	<sup>1</sup> ACCIARRI 95G	L3	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
$<5.5 \times 10^{-5}$	95	ABREU 94B	DLPH	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
$<1.4 \times 10^{-4}$	95	AKRAWY 91F	OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV

<sup>1</sup> This limit is for both decay modes  $Z \rightarrow \pi^0\gamma/\gamma\gamma$  which are indistinguishable in ACCIARRI 95G.

$\Gamma(\pi^0\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{22}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.52 \times 10^{-5}$	95	AALTONEN 14E	CDF	$E_{\text{cm}}^{pp} = 1.96$ TeV

$\Gamma(\gamma\gamma\gamma)/\Gamma_{\text{total}}$   $\Gamma_{23}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.2 \times 10^{-6}$	95	AAD 16L	ATLS	$E_{\text{cm}}^{pp} = 8$ TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.0 \times 10^{-5}$	95	<sup>1</sup> ACCIARRI 95C	L3	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
$<1.7 \times 10^{-5}$	95	<sup>1</sup> ABREU 94B	DLPH	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
$<6.6 \times 10^{-5}$	95	AKRAWY 91F	OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV

<sup>1</sup> Limit derived in the context of composite  $Z$  model.

$\Gamma(\pi^\pm W^\mp)/\Gamma_{\text{total}}$   $\Gamma_{24}/\Gamma$

The value is for the sum of the charge states indicated.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7 \times 10^{-5}$	95	DECAMP 92	ALEP	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV

$\Gamma(\rho^\pm W^\mp)/\Gamma_{\text{total}}$   $\Gamma_{25}/\Gamma$

The value is for the sum of the charge states indicated.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8.3 \times 10^{-5}$	95	DECAMP 92	ALEP	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV

$\Gamma(J/\psi(1S)X)/\Gamma_{\text{total}}$   $\Gamma_{26}/\Gamma$

VALUE (units  $10^{-3}$ )    EVTS    DOCUMENT ID    TECN    COMMENT

**3.51<sup>+0.23</sup><sub>-0.25</sub> OUR AVERAGE** Error includes scale factor of 1.1.

3.21 ± 0.21 <sup>+0.19</sup> <sub>-0.28</sub>	553	<sup>1</sup> ACCIARRI	99F L3	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
3.9 ± 0.2 ± 0.3	511	<sup>2</sup> ALEXANDER	96B OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
3.73 ± 0.39 ± 0.36	153	<sup>3</sup> ABREU	94P DLPH	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV

<sup>1</sup> ACCIARRI 99F combine  $\mu^+\mu^-$  and  $e^+e^- J/\psi(1S)$  decay channels. The branching ratio for prompt  $J/\psi(1S)$  production is measured to be  $(2.1 \pm 0.6 \pm 0.4_{-0.2}^{+0.4}(\text{theor.})) \times 10^{-4}$ .

<sup>2</sup> ALEXANDER 96B identify  $J/\psi(1S)$  from the decays into lepton pairs.  $(4.8 \pm 2.4)\%$  of this branching ratio is due to prompt  $J/\psi(1S)$  production (ALEXANDER 96N).

<sup>3</sup> Combining  $\mu^+\mu^-$  and  $e^+e^-$  channels and taking into account the common systematic errors.  $(7.7_{-5.4}^{+6.3})\%$  of this branching ratio is due to prompt  $J/\psi(1S)$  production.

$\Gamma(J/\psi(1S)\gamma)/\Gamma_{\text{total}}$   $\Gamma_{27}/\Gamma$

VALUE    CL%    DOCUMENT ID    TECN    COMMENT

**<1.2 × 10<sup>-6</sup>**    95    AAD    23CD ATLS     $E_{\text{cm}}^{pp} = 13$  TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.4 × 10 <sup>-6</sup>	95	<sup>1</sup> SIRUNYAN	19AJ CMS	$E_{\text{cm}}^{pp} = 13$ TeV
<2.3 × 10 <sup>-6</sup>	95	<sup>2</sup> AABOUD	18BL ATLS	$E_{\text{cm}}^{pp} = 13$ TeV
<2.6 × 10 <sup>-6</sup>	95	<sup>3</sup> AAD	15I ATLS	$E_{\text{cm}}^{pp} = 8$ TeV

<sup>1</sup> SIRUNYAN 19AJ study  $Z \rightarrow J/\psi\gamma$  with  $J/\psi \rightarrow \mu^+\mu^-$ . Candidate events are selected by requiring a pair of oppositely charged muons and a well isolated photon. The leading (subleading) muon is required to have a transverse momentum larger than 20 GeV (4 GeV), while the photon must have a transverse energy larger than 33 GeV. Requiring the invariant mass of the  $\mu\mu$  ( $\mu\mu\gamma$ ) system in the range 3.0 to 3.2 (81 to 101) GeV, selects 183 data events which is consistent with the expected background. The 95% C.L. limit on the  $Z$  branching fraction is obtained assuming the  $J/\psi$  to be unpolarized.

<sup>2</sup> AABOUD 18BL study  $Z \rightarrow J/\psi\gamma$  in 13 TeV  $pp$  interactions. Two triggers were used: isolated photon of  $p_T > 35(25)$  GeV and a muon with  $p_T > 18(24)$  GeV. The  $J/\psi$  is detected via its dimuon decay and it is required that the azimuthal angle between the photon and the  $J/\psi$  in the plane transverse to the beam direction is  $> \pi/2$ . The number of observed/expected background events is 92/89 ± 6 in the dimuon mass range 2.9–3.3 GeV leading to the quoted 95% C.L. limit.

<sup>3</sup> AAD 15I use events with the highest  $p_T$  muon in the pair required to have  $p_T > 20$  GeV, the dimuon mass required to be within 0.2 GeV of the  $J/\psi(1S)$  mass and its transverse momentum required to be  $> 36$  GeV. The photon is also required to have its  $p_T > 36$  GeV.

$\Gamma(\psi(2S)X)/\Gamma_{\text{total}}$   $\Gamma_{28}/\Gamma$

VALUE (units  $10^{-3}$ )    EVTS    DOCUMENT ID    TECN    COMMENT

**1.60 ± 0.29 OUR AVERAGE**

1.6 ± 0.5 ± 0.3	39	<sup>1</sup> ACCIARRI	97J L3	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
1.6 ± 0.3 ± 0.2	46.9	<sup>2</sup> ALEXANDER	96B OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
1.60 ± 0.73 ± 0.33	5.4	<sup>3</sup> ABREU	94P DLPH	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV



- <sup>1</sup> ACCIARRI 97J measure this branching ratio via the decay channel  $\psi(2S) \rightarrow \ell^+ \ell^-$  ( $\ell = \mu, e$ ).
- <sup>2</sup> ALEXANDER 96B measure this branching ratio via the decay channel  $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$ , with  $J/\psi \rightarrow \ell^+ \ell^-$ .
- <sup>3</sup> ABREU 94P measure this branching ratio via decay channel  $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$ , with  $J/\psi \rightarrow \mu^+ \mu^-$ .

$\Gamma(\psi(2S)\gamma)/\Gamma_{\text{total}}$					$\Gamma_{29}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<2.4 \times 10^{-6}$	95	AAD	23CD ATLS	$E_{\text{cm}}^{pp} = 13 \text{ TeV}$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<4.5 \times 10^{-6}$	95	<sup>1</sup> AABOUD	18BL ATLS	$E_{\text{cm}}^{pp} = 13 \text{ TeV}$
-----------------------	----	---------------------	-----------	---------------------------------------

<sup>1</sup> AABOUD 18BL study  $Z \rightarrow \psi(2S)\gamma$  in 13 TeV  $pp$  interactions. Two triggers were used: isolated photon of  $p_T > 35(25) \text{ GeV}$  and a muon with  $p_T > 18(24) \text{ GeV}$ . The  $\psi(2S)$  is detected via its dimuon decay and it is required that the azimuthal angle between the photon and the  $\psi(2S)$  in the plane transverse to the beam direction is  $> \pi/2$ . The number of observed/expected background events is  $43/42 \pm 5$  in the dimuon mass range 3.5–3.9 GeV leading to the quoted 95% C.L. limit.

$\Gamma(J/\psi(1S)\ell^+\ell^-)/\Gamma(\mu^+\mu^-\mu^+\mu^-)$				$\Gamma_{30}/\Gamma_5$
VALUE	DOCUMENT ID	TECN	COMMENT	
$0.67 \pm 0.18 \pm 0.05$	<sup>1</sup> SIRUNYAN	18DZ CMS	$pp$ at 13 TeV	

<sup>1</sup> SIRUNYAN 18DZ observe the decay  $Z \rightarrow \Psi \ell^+ \ell^-$  in  $pp$  collisions at  $\sqrt{s} = 13 \text{ TeV}$ , where  $\Psi$  includes  $J/\psi$  as well as  $\psi(2S) \rightarrow J/\psi X$ , and  $\ell^+ \ell^-$  represents an electron or muon pair while the  $J/\psi$  is detected via its  $\mu^+ \mu^-$  decay channel. To reduce systematic errors they determine the ratio of the branching fraction of this decay to that of  $Z \rightarrow \mu^+ \mu^- \mu^+ \mu^-$  within phase-space cuts imposed on lepton transverse momentum and pseudo rapidity, dilepton invariant mass, and  $J/\psi$  transverse momentum. The number of selected  $\Psi \mu^+ \mu^-$  ( $\Psi e^+ e^-$ ) candidate events is 29 (18). Analyzing the  $\mu^+ \mu^-$  and  $\mu^+ \mu^- \ell^+ \ell^-$  invariant mass distributions, a yield of  $13.0 \pm 3.9$  ( $11.2 \pm 3.4$ ) events for the  $\Psi \mu^+ \mu^-$  ( $\Psi e^+ e^-$ ) mode is obtained. The ratio of the branching fractions is determined as  $0.67 \pm 0.18 \pm 0.05$  within the selected phase-space cuts. Assuming extrapolation to full phase space cancels in the ratio, and using their measured value of  $B(Z \rightarrow \mu^+ \mu^- \mu^+ \mu^-) = (1.20 \pm 0.08) \times 10^{-6}$ , they estimate  $B(Z \rightarrow J/\psi \ell^+ \ell^-) = 8 \times 10^{-7}$ .

$\Gamma(J/\psi(1S)J/\psi(1S))/\Gamma_{\text{total}}$					$\Gamma_{31}/\Gamma$
VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$<2.2 \times 10^{-6}$	95	189	<sup>1</sup> SIRUNYAN	19BR CMS	$E_{\text{cm}}^{pp} = 13 \text{ TeV}$

<sup>1</sup> SIRUNYAN 19BR search for  $Z$  decays to a pair of  $J/\psi$  mesons in the channel  $J/\psi \rightarrow \mu^+ \mu^-$ . The invariant masses of the higher/lower- $p_T$   $J/\psi$  candidates have to be within 0.1/0.15 GeV of the nominal  $J/\psi$  mass. A total of 189 events are selected in the 40–140 GeV 4-muon invariant mass range. An un-binned extended maximum likelihood fit leads to the 95% C.L. upper limit, obtained assuming the  $J/\psi$  mesons to be unpolarised.

$\Gamma(\chi_{c1}(1P)X)/\Gamma_{\text{total}}$					$\Gamma_{32}/\Gamma$
VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>2.9±0.7 OUR AVERAGE</b>					
$2.7 \pm 0.6 \pm 0.5$	33	<sup>1</sup> ACCIARRI 97J	L3	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV	
$5.0 \pm 2.1^{+1.5}_{-0.9}$	6.4	<sup>2</sup> ABREU 94P	DLPH	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV	

<sup>1</sup> ACCIARRI 97J measure this branching ratio via the decay channel  $\chi_{c1} \rightarrow J/\psi + \gamma$ , with  $J/\psi \rightarrow \ell^+ \ell^-$  ( $\ell = \mu, e$ ). The  $M(\ell^+ \ell^- \gamma) - M(\ell^+ \ell^-)$  mass difference spectrum is fitted with two gaussian shapes for  $\chi_{c1}$  and  $\chi_{c2}$ .

<sup>2</sup> This branching ratio is measured via the decay channel  $\chi_{c1} \rightarrow J/\psi + \gamma$ , with  $J/\psi \rightarrow \mu^+ \mu^-$ .

$\Gamma(\chi_{c2}(1P)X)/\Gamma_{\text{total}}$					$\Gamma_{33}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<b>&lt;3.2 × 10<sup>-3</sup></b>					
	90	<sup>1</sup> ACCIARRI 97J	L3	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV	

<sup>1</sup> ACCIARRI 97J derive this limit via the decay channel  $\chi_{c2} \rightarrow J/\psi + \gamma$ , with  $J/\psi \rightarrow \ell^+ \ell^-$  ( $\ell = \mu, e$ ). The  $M(\ell^+ \ell^- \gamma) - M(\ell^+ \ell^-)$  mass difference spectrum is fitted with two gaussian shapes for  $\chi_{c1}$  and  $\chi_{c2}$ .

$\Gamma(\Upsilon(1S)X + \Upsilon(2S)X + \Upsilon(3S)X)/\Gamma_{\text{total}}$					$\Gamma_{34}/\Gamma = (\Gamma_{35} + \Gamma_{37} + \Gamma_{39})/\Gamma$
VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>1.0±0.4±0.22</b>					
	6.4	<sup>1</sup> ALEXANDER 96F	OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV	

<sup>1</sup> ALEXANDER 96F identify the  $\Upsilon$  (which refers to any of the three lowest bound states) through its decay into  $e^+ e^-$  and  $\mu^+ \mu^-$ . The systematic error includes an uncertainty of  $\pm 0.2$  due to the production mechanism.

$\Gamma(\Upsilon(1S)X)/\Gamma_{\text{total}}$					$\Gamma_{35}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<b>&lt;4.4 × 10<sup>-5</sup></b>					
	95	<sup>1</sup> ACCIARRI 99F	L3	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV	

<sup>1</sup> ACCIARRI 99F search for  $\Upsilon(1S)$  through its decay into  $\ell^+ \ell^-$  ( $\ell = e$  or  $\mu$ ).

$\Gamma(\Upsilon(1S)\gamma)/\Gamma_{\text{total}}$					$\Gamma_{36}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<b>&lt;1.1 × 10<sup>-6</sup></b>					
	95	AAD	23CD ATLS	$E_{\text{cm}}^{pp} = 13$ TeV	

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<2.8 \times 10^{-6}$	95	<sup>1</sup> AABOUD 18BL	ATLS	$E_{\text{cm}}^{pp} = 13$ TeV
$<3.4 \times 10^{-6}$	95	<sup>2</sup> AAD 15I	ATLS	$E_{\text{cm}}^{pp} = 8$ TeV

<sup>1</sup> AABOUD 18BL study  $Z \rightarrow \Upsilon(1S)\gamma$  in 13 TeV  $pp$  interactions. Two triggers were used: isolated photon of  $p_T > 35(25)$  GeV and a muon with  $p_T > 18(24)$  GeV. The  $\Upsilon(1S)$  is detected via its dimuon decay and it is required that the azimuthal angle between the photon and the  $\Upsilon(1S)$  in the plane transverse to the beam direction is  $> \pi/2$ . The number of observed/expected background events is  $115/126 \pm 8$  in the dimuon mass range 9.0–10.0 GeV leading to the quoted 95% C.L. limit.

<sup>2</sup> AAD 15I use events with the highest  $p_T$  muon in the pair required to have  $p_T > 20$  GeV, the dimuon mass required to be in the range 8–12 GeV and it's transverse momentum required to be  $> 36$  GeV. The photon is also required to have it's  $p_T > 36$  GeV.

$\Gamma(\Upsilon(2S)X)/\Gamma_{\text{total}}$   $\Gamma_{37}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<13.9 \times 10^{-5}$	95	<sup>1</sup> ACCIARRI 97R	L3	$E_{\text{cm}}^{\text{ee}} = 88\text{--}94$ GeV

<sup>1</sup> ACCIARRI 97R search for  $\Upsilon(2S)$  through its decay into  $\ell^+\ell^-$  ( $\ell = e$  or  $\mu$ ).

$\Gamma(\Upsilon(2S)\gamma)/\Gamma_{\text{total}}$   $\Gamma_{38}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.3 \times 10^{-6}$	95	AAD	23CD ATLS	$E_{\text{cm}}^{\text{pp}} = 13$ TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<1.7 \times 10^{-6}$	95	<sup>1</sup> AABOUD	18BL ATLS	$E_{\text{cm}}^{\text{pp}} = 13$ TeV
$<6.5 \times 10^{-6}$	95	<sup>2</sup> AAD	15I ATLS	$E_{\text{cm}}^{\text{pp}} = 8$ TeV

<sup>1</sup> AABOUD 18BL study  $Z \rightarrow \Upsilon(2S)\gamma$  in 13 TeV  $pp$  interactions. Two triggers were used: isolated photon of  $p_{\mathcal{T}} > 35(25)$  GeV and a muon with  $p_{\mathcal{T}} > 18(24)$  GeV. The  $\Upsilon(2S)$  is detected via its dimuon decay and it is required that the azimuthal angle between the photon and the  $\Upsilon(2S)$  in the plane transverse to the beam direction is  $> \pi/2$ . The number of observed/expected background events is  $106/121 \pm 8$  in the dimuon mass range 9.5–10.5 GeV leading to the quoted 95% C.L. limit.

<sup>2</sup> AAD 15I use events with the highest  $p_{\mathcal{T}}$  muon in the pair required to have  $p_{\mathcal{T}} > 20$  GeV, the dimuon mass required to be in the range 8–12 GeV and it's transverse momentum required to be  $> 36$  GeV. The photon is also required to have it's  $p_{\mathcal{T}} > 36$  GeV.

$\Gamma(\Upsilon(3S)X)/\Gamma_{\text{total}}$   $\Gamma_{39}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<9.4 \times 10^{-5}$	95	<sup>1</sup> ACCIARRI 97R	L3	$E_{\text{cm}}^{\text{ee}} = 88\text{--}94$ GeV

<sup>1</sup> ACCIARRI 97R search for  $\Upsilon(3S)$  through its decay into  $\ell^+\ell^-$  ( $\ell = e$  or  $\mu$ ).

$\Gamma(\Upsilon(3S)\gamma)/\Gamma_{\text{total}}$   $\Gamma_{40}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.4 \times 10^{-6}$	95	AAD	23CD ATLS	$E_{\text{cm}}^{\text{pp}} = 13$ TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<4.8 \times 10^{-6}$	95	<sup>1</sup> AABOUD	18BL ATLS	$E_{\text{cm}}^{\text{pp}} = 13$ TeV
$<5.4 \times 10^{-6}$	95	<sup>2</sup> AAD	15I ATLS	$E_{\text{cm}}^{\text{pp}} = 8$ TeV

<sup>1</sup> AABOUD 18BL study  $Z \rightarrow \Upsilon(3S)\gamma$  in 13 TeV  $pp$  interactions. Two triggers were used: isolated photon of  $p_{\mathcal{T}} > 35(25)$  GeV and a muon with  $p_{\mathcal{T}} > 18(24)$  GeV. The  $\Upsilon(3S)$  is detected via its dimuon decay and it is required that the azimuthal angle between the photon and the  $\Upsilon(3S)$  in the plane transverse to the beam direction is  $> \pi/2$ . The number of observed/expected background events is  $112/113 \pm 8$  in the dimuon mass range 10.0–11.0 GeV leading to the quoted 95% C.L. limit.

<sup>2</sup> AAD 15I use events with the highest  $p_{\mathcal{T}}$  muon in the pair required to have  $p_{\mathcal{T}} > 20$  GeV, the dimuon mass required to be in the range 8–12 GeV and it's transverse momentum required to be  $> 36$  GeV. The photon is also required to have it's  $p_{\mathcal{T}} > 36$  GeV.

$\Gamma(\Upsilon(1, 2, 3S)\Upsilon(1, 2, 3S))/\Gamma_{\text{total}}$   $\Gamma_{41}/\Gamma$

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$<1.5 \times 10^{-6}$	95	106	<sup>1</sup> SIRUNYAN 19BR	CMS	$E_{\text{cm}}^{\text{pp}} = 13$ TeV

<sup>1</sup> SIRUNYAN 19BR search for  $Z$  decays to a pair of  $\Upsilon$  mesons in the channel  $\Upsilon \rightarrow \mu^+\mu^-$ . The invariant mass of the  $\Upsilon$  candidates has to be in the range of 8.5 to 11 GeV. A total of 106 events are selected in the 20–140 GeV 4-muon invariant mass range. An un-binned extended maximum likelihood fit leads to the 95% C.L. upper limit, obtained assuming the  $\Upsilon$  mesons to be unpolarised.

$\Gamma(D^0\gamma)/\Gamma(\mu^+\mu^-)$   $\Gamma_{42}/\Gamma_2$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.4 \times 10^{-2}$	95	<sup>1</sup> AAIJ	23AMLHCB	$E_{cm}^{pp} = 13$ TeV

<sup>1</sup> AAIJ 23AM also quotes the branching fraction limit  $B(Z \rightarrow D^0\gamma) < 2.1 \times 10^{-3}$ , using the known  $Z \rightarrow \mu\mu$  branching fraction.

$\Gamma((D^0/\bar{D}^0)X)/\Gamma(\text{hadrons})$   $\Gamma_{43}/\Gamma_8$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$0.296 \pm 0.019 \pm 0.021$	369	<sup>1</sup> ABREU	93i	DLPH $E_{cm}^{ee} = 88-94$ GeV

<sup>1</sup> The  $(D^0/\bar{D}^0)$  states in ABREU 93i are detected by the  $K\pi$  decay mode. This is a corrected result (see the erratum of ABREU 93i).

$\Gamma(D^\pm X)/\Gamma(\text{hadrons})$   $\Gamma_{44}/\Gamma_8$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$0.174 \pm 0.016 \pm 0.018$	539	<sup>1</sup> ABREU	93i	DLPH $E_{cm}^{ee} = 88-94$ GeV

<sup>1</sup> The  $D^\pm$  states in ABREU 93i are detected by the  $K\pi\pi$  decay mode. This is a corrected result (see the erratum of ABREU 93i).

$\Gamma(D^*(2010)^\pm X)/\Gamma(\text{hadrons})$   $\Gamma_{45}/\Gamma_8$

The value is for the sum of the charge states indicated.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.163 \pm 0.019</math> OUR AVERAGE</b>		Error includes scale factor of 1.3.		
$0.155 \pm 0.010 \pm 0.013$	358	<sup>1</sup> ABREU	93i	DLPH $E_{cm}^{ee} = 88-94$ GeV
$0.21 \pm 0.04$	362	<sup>2</sup> DECAMP	91J	ALEP $E_{cm}^{ee} = 88-94$ GeV

<sup>1</sup>  $D^*(2010)^\pm$  in ABREU 93i are reconstructed from  $D^0\pi^\pm$ , with  $D^0 \rightarrow K^-\pi^+$ . The new CLEO II measurement of  $B(D^{*\pm} \rightarrow D^0\pi^\pm) = (68.1 \pm 1.6)\%$  is used. This is a corrected result (see the erratum of ABREU 93i).

<sup>2</sup> DECAMP 91J report  $B(D^*(2010)^+ \rightarrow D^0\pi^+) B(D^0 \rightarrow K^-\pi^+) \Gamma(D^*(2010)^\pm X) / \Gamma(\text{hadrons}) = (5.11 \pm 0.34) \times 10^{-3}$ . They obtained the above number assuming  $B(D^0 \rightarrow K^-\pi^+) = (3.62 \pm 0.34 \pm 0.44)\%$  and  $B(D^*(2010)^+ \rightarrow D^0\pi^+) = (55 \pm 4)\%$ . We have rescaled their original result of  $0.26 \pm 0.05$  taking into account the new CLEO II branching ratio  $B(D^*(2010)^+ \rightarrow D^0\pi^+) = (68.1 \pm 1.6)\%$ .

$\Gamma(D_{s1}(2536)^\pm X)/\Gamma(\text{hadrons})$   $\Gamma_{46}/\Gamma_8$

$D_{s1}(2536)^\pm$  is an expected orbitally-excited state of the  $D_s$  meson.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$0.52 \pm 0.09 \pm 0.06$	92	<sup>1</sup> HEISTER	02B	ALEP $E_{cm}^{ee} = 88-94$ GeV

<sup>1</sup> HEISTER 02B reconstruct this meson in the decay modes  $D_{s1}(2536)^\pm \rightarrow D^{*\pm}K^0$  and  $D_{s1}(2536)^\pm \rightarrow D^{*0}K^\pm$ . The quoted branching ratio assumes that the decay width of the  $D_{s1}(2536)$  is saturated by the two measured decay modes.

$\Gamma(D_{sJ}(2573)^\pm X)/\Gamma(\text{hadrons})$   $\Gamma_{47}/\Gamma_8$

$D_{sJ}(2573)^\pm$  is an expected orbitally-excited state of the  $D_s$  meson.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$0.83 \pm 0.29^{+0.07}_{-0.13}$	64	<sup>1</sup> HEISTER	02B	ALEP $E_{cm}^{ee} = 88-94$ GeV

<sup>1</sup> HEISTER 02B reconstruct this meson in the decay mode  $D_{s2}^*(2573)^\pm \rightarrow D^0K^\pm$ . The quoted branching ratio assumes that the detected decay mode represents 45% of the full decay width.

### $\Gamma(D^{*'}(2629)^{\pm}X)/\Gamma(\text{hadrons})$

$\Gamma_{48}/\Gamma_8$

$D^{*'}(2629)^{\pm}$  is a predicted radial excitation of the  $D^*(2010)^{\pm}$  meson.

VALUE	DOCUMENT ID	TECN	COMMENT
<b>searched for</b>	<sup>1</sup> ABBIENDI	01N OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV

<sup>1</sup> ABBIENDI 01N searched for the decay mode  $D^{*'}(2629)^{\pm} \rightarrow D^{*\pm}\pi^+\pi^-$  with  $D^{*+} \rightarrow D^0\pi^+$ , and  $D^0 \rightarrow K^-\pi^+$ . They quote a 95% CL limit for  $Z \rightarrow D^{*'}(2629)^{\pm} \times B(D^{*'}(2629)^{\pm} \rightarrow D^{*+}\pi^+\pi^-) < 3.1 \times 10^{-3}$ .

### $\Gamma(B^*X)/[\Gamma(BX) + \Gamma(B^*X)]$

$\Gamma_{50}/(\Gamma_{49} + \Gamma_{50})$

As the experiments assume different values of the  $b$ -baryon contribution, our average should be taken with caution.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.75 ± 0.04 OUR AVERAGE</b>				
0.760 ± 0.036 ± 0.083		<sup>1</sup> ACKERSTAFF	97M OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
0.771 ± 0.026 ± 0.070		<sup>2</sup> BUSKULIC	96D ALEP	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
0.72 ± 0.03 ± 0.06		<sup>3</sup> ABREU	95R DLPH	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
0.76 ± 0.08 ± 0.06	1378	<sup>4</sup> ACCIARRI	95B L3	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV

<sup>1</sup> ACKERSTAFF 97M use an inclusive  $B$  reconstruction method and assume a  $(13.2 \pm 4.1)\%$   $b$ -baryon contribution. The value refers to a  $b$ -flavored meson mixture of  $B_u$ ,  $B_d$ , and  $B_s$ .

<sup>2</sup> BUSKULIC 96D use an inclusive reconstruction of  $B$  hadrons and assume a  $(12.2 \pm 4.3)\%$   $b$ -baryon contribution. The value refers to a  $b$ -flavored mixture of  $B_u$ ,  $B_d$ , and  $B_s$ .

<sup>3</sup> ABREU 95R use an inclusive  $B$ -reconstruction method and assume a  $(10 \pm 4)\%$   $b$ -baryon contribution. The value refers to a  $b$ -flavored meson mixture of  $B_u$ ,  $B_d$ , and  $B_s$ .

<sup>4</sup> ACCIARRI 95B assume a 9.4%  $b$ -baryon contribution. The value refers to a  $b$ -flavored mixture of  $B_u$ ,  $B_d$ , and  $B_s$ .

### $\Gamma(B^+X)/\Gamma(\text{hadrons})$

$\Gamma_{51}/\Gamma_8$

"OUR EVALUATION" is obtained using our current values for  $f(\bar{b} \rightarrow B^+)$  and  $R_b = \Gamma(b\bar{b})/\Gamma(\text{hadrons})$ . We calculate  $\Gamma(B^+X)/\Gamma(\text{hadrons}) = R_b \times f(\bar{b} \rightarrow B^+)$ .

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.0869 ± 0.0019 OUR EVALUATION</b>	(Produced by HFLAV)		
<b>0.0887 ± 0.0030</b>	<sup>1</sup> ABDALLAH	03K DLPH	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV

<sup>1</sup> ABDALLAH 03K measure the production fraction of  $B^+$  mesons in hadronic  $Z$  decays  $f(B^+) = (40.99 \pm 0.82 \pm 1.11)\%$ . The value quoted here is obtained multiplying this production fraction by our value of  $R_b = \Gamma(\bar{b}b)/\Gamma(\text{hadrons})$ .

### $\Gamma(B_s^0X)/\Gamma(\text{hadrons})$

$\Gamma_{52}/\Gamma_8$

"OUR EVALUATION" is obtained using our current values for  $f(\bar{b} \rightarrow B_s^0)$  and  $R_b = \Gamma(b\bar{b})/\Gamma(\text{hadrons})$ . We calculate  $\Gamma(B_s^0X)/\Gamma(\text{hadrons}) = R_b \times f(\bar{b} \rightarrow B_s^0)$ .

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.0227 ± 0.0019 OUR EVALUATION</b>	(Produced by HFLAV)		
seen	<sup>1</sup> ABREU	92M DLPH	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
seen	<sup>2</sup> ACTON	92N OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
seen	<sup>3</sup> BUSKULIC	92E ALEP	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV

- <sup>1</sup> ABREU 92M reported value is  $\Gamma(B_S^0 X) * B(B_S^0 \rightarrow D_S \mu \nu_\mu X) * B(D_S \rightarrow \phi \pi) / \Gamma(\text{hadrons}) = (18 \pm 8) \times 10^{-5}$ .
- <sup>2</sup> ACTON 92N find evidence for  $B_S^0$  production using  $D_S$ - $\ell$  correlations, with  $D_S^+ \rightarrow \phi \pi^+$  and  $K^*(892) K^+$ . Assuming  $R_b$  from the Standard Model and averaging over the  $e$  and  $\mu$  channels, authors measure the product branching fraction to be  $f(\bar{b} \rightarrow B_S^0) \times B(B_S^0 \rightarrow D_S^- \ell^+ \nu_\ell X) \times B(D_S^- \rightarrow \phi \pi^-) = (3.9 \pm 1.1 \pm 0.8) \times 10^{-4}$ .
- <sup>3</sup> BUSKULIC 92E find evidence for  $B_S^0$  production using  $D_S$ - $\ell$  correlations, with  $D_S^+ \rightarrow \phi \pi^+$  and  $K^*(892) K^+$ . Using  $B(D_S^+ \rightarrow \phi \pi^+) = (2.7 \pm 0.7)\%$  and summing up the  $e$  and  $\mu$  channels, the weighted average product branching fraction is measured to be  $B(\bar{b} \rightarrow B_S^0) \times B(B_S^0 \rightarrow D_S^- \ell^+ \nu_\ell X) = 0.040 \pm 0.011^{+0.010}_{-0.012}$ .

### $\Gamma(B_c^+ X) / \Gamma(\text{hadrons})$

$\Gamma_{53} / \Gamma_8$

VALUE	DOCUMENT ID	TECN	COMMENT
searched for	<sup>1</sup> ACKERSTAFF 98O	OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
searched for	<sup>2</sup> ABREU 97E	DLPH	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
searched for	<sup>3</sup> BARATE 97H	ALEP	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV

- <sup>1</sup> ACKERSTAFF 98O searched for the decay modes  $B_c \rightarrow J/\psi \pi^+$ ,  $J/\psi a_1^+$ , and  $J/\psi \ell^+ \nu_\ell$ , with  $J/\psi \rightarrow \ell^+ \ell^-$ ,  $\ell = e, \mu$ . The number of candidates (background) for the three decay modes is 2 ( $0.63 \pm 0.2$ ), 0 ( $1.10 \pm 0.22$ ), and 1 ( $0.82 \pm 0.19$ ) respectively. Interpreting the  $2 B_c \rightarrow J/\psi \pi^+$  candidates as signal, they report  $\Gamma(B_c^+ X) \times B(B_c \rightarrow J/\psi \pi^+) / \Gamma(\text{hadrons}) = (3.8^{+5.0}_{-2.4} \pm 0.5) \times 10^{-5}$ . Interpreted as background, the 90% CL bounds are  $\Gamma(B_c^+ X) * B(B_c \rightarrow J/\psi \pi^+) / \Gamma(\text{hadrons}) < 1.06 \times 10^{-4}$ ,  $\Gamma(B_c^+ X) * B(B_c \rightarrow J/\psi a_1^+) / \Gamma(\text{hadrons}) < 5.29 \times 10^{-4}$ ,  $\Gamma(B_c^+ X) * B(B_c \rightarrow J/\psi \ell^+ \nu_\ell) / \Gamma(\text{hadrons}) < 6.96 \times 10^{-5}$ .
- <sup>2</sup> ABREU 97E searched for the decay modes  $B_c \rightarrow J/\psi \pi^+$ ,  $J/\psi \ell^+ \nu_\ell$ , and  $J/\psi (3\pi)^+$ , with  $J/\psi \rightarrow \ell^+ \ell^-$ ,  $\ell = e, \mu$ . The number of candidates (background) for the three decay modes is 1 (1.7), 0 (0.3), and 1 (2.3) respectively. They report the following 90% CL limits:  $\Gamma(B_c^+ X) * B(B_c \rightarrow J/\psi \pi^+) / \Gamma(\text{hadrons}) < (1.05\text{--}0.84) \times 10^{-4}$ ,  $\Gamma(B_c^+ X) * B(B_c \rightarrow J/\psi \ell \nu_\ell) / \Gamma(\text{hadrons}) < (5.8\text{--}5.0) \times 10^{-5}$ ,  $\Gamma(B_c^+ X) * B(B_c \rightarrow J/\psi (3\pi)^+) / \Gamma(\text{hadrons}) < 1.75 \times 10^{-4}$ , where the ranges are due to the predicted  $B_c$  lifetime (0.4–1.4) ps.
- <sup>3</sup> BARATE 97H searched for the decay modes  $B_c \rightarrow J/\psi \pi^+$  and  $J/\psi \ell^+ \nu_\ell$  with  $J/\psi \rightarrow \ell^+ \ell^-$ ,  $\ell = e, \mu$ . The number of candidates (background) for the two decay modes is 0 (0.44) and 2 (0.81) respectively. They report the following 90% CL limits:  $\Gamma(B_c^+ X) * B(B_c \rightarrow J/\psi \pi^+) / \Gamma(\text{hadrons}) < 3.6 \times 10^{-5}$  and  $\Gamma(B_c^+ X) * B(B_c \rightarrow J/\psi \ell^+ \nu_\ell) / \Gamma(\text{hadrons}) < 5.2 \times 10^{-5}$ .

### $\Gamma(\Lambda_c^+ X) / \Gamma(\text{hadrons})$

$\Gamma_{54} / \Gamma_8$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.022 ± 0.005 OUR AVERAGE</b>			
0.024 ± 0.005 ± 0.006	<sup>1</sup> ALEXANDER 96R	OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
0.021 ± 0.003 ± 0.005	<sup>2</sup> BUSKULIC 96Y	ALEP	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV

<sup>1</sup> ALEXANDER 96R measure  $R_b \times f(b \rightarrow \Lambda_c^+ X) \times B(\Lambda_c^+ \rightarrow pK^- \pi^+) = (0.122 \pm 0.023 \pm 0.010)\%$  in hadronic  $Z$  decays; the value quoted here is obtained using our best value  $B(\Lambda_c^+ \rightarrow pK^- \pi^+) = (5.0 \pm 1.3)\%$ . The first error is the total experiment's error and the second error is the systematic error due to the branching fraction uncertainty.

<sup>2</sup> BUSKULIC 96Y obtain the production fraction of  $\Lambda_c^+$  baryons in hadronic  $Z$  decays  $f(b \rightarrow \Lambda_c^+ X) = 0.110 \pm 0.014 \pm 0.006$  using  $B(\Lambda_c^+ \rightarrow pK^- \pi^+) = (4.4 \pm 0.6)\%$ ; we have rescaled using our best value  $B(\Lambda_c^+ \rightarrow pK^- \pi^+) = (5.0 \pm 1.3)\%$  obtaining  $f(b \rightarrow \Lambda_c^+ X) = 0.097 \pm 0.013 \pm 0.025$  where the first error is their total experiment's error and the second error is the systematic error due to the branching fraction uncertainty. The value quoted here is obtained multiplying this production fraction by our value of  $R_b = \Gamma(b\bar{b})/\Gamma(\text{hadrons})$ .

### $\Gamma(\Xi_c^0 X)/\Gamma(\text{hadrons})$

$\Gamma_{55}/\Gamma_8$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen <sup>1</sup> ABDALLAH 05C DLPH  $E_{\text{cm}}^{\text{ee}} = 88\text{--}94$  GeV

<sup>1</sup> ABDALLAH 05C searched for the charmed strange baryon  $\Xi_c^0$  in the decay channel  $\Xi_c^0 \rightarrow \Xi^- \pi^+$  ( $\Xi^- \rightarrow \Lambda \pi^-$ ). The production rate is measured to be  $f_{\Xi_c^0} \times B(\Xi_c^0 \rightarrow \Xi^- \pi^+) = (4.7 \pm 1.4 \pm 1.1) \times 10^{-4}$  per hadronic  $Z$  decay.

### $\Gamma(\Xi_b X)/\Gamma(\text{hadrons})$

$\Gamma_{56}/\Gamma_8$

Here  $\Xi_b$  is used as a notation for the strange  $b$ -baryon states  $\Xi_b^-$  and  $\Xi_b^0$ .

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen <sup>1</sup> ABDALLAH 05C DLPH  $E_{\text{cm}}^{\text{ee}} = 88\text{--}94$  GeV

seen <sup>2</sup> BUSKULIC 96T ALEP  $E_{\text{cm}}^{\text{ee}} = 88\text{--}94$  GeV

seen <sup>3</sup> ABREU 95V DLPH  $E_{\text{cm}}^{\text{ee}} = 88\text{--}94$  GeV

<sup>1</sup> ABDALLAH 05C searched for the beauty strange baryon  $\Xi_b$  in the inclusive semileptonic decay channel  $\Xi_b \rightarrow \Xi^- \ell^- \bar{\nu}_\ell X$ . Evidence for the  $\Xi_b$  production is seen from the observation of  $\Xi^\mp$  production accompanied by a lepton of the same sign. From the excess of "right-sign" pairs  $\Xi^\mp \ell^\mp$  compared to "wrong-sign" pairs  $\Xi^\mp \ell^\pm$  the production rate is measured to be  $B(b \rightarrow \Xi_b) \times B(\Xi_b \rightarrow \Xi^- \ell^- X) = (3.0 \pm 1.0 \pm 0.3) \times 10^{-4}$  per lepton species, averaged over electrons and muons.

<sup>2</sup> BUSKULIC 96T investigate  $\Xi$ -lepton correlations and find a significant excess of "right-sign" pairs  $\Xi^\mp \ell^\mp$  compared to "wrong-sign" pairs  $\Xi^\mp \ell^\pm$ . This excess is interpreted as evidence for  $\Xi_b$  semileptonic decay. The measured product branching ratio is  $B(b \rightarrow \Xi_b) \times B(\Xi_b \rightarrow X_c X \ell^- \bar{\nu}_\ell) \times B(X_c \rightarrow \Xi^- X') = (5.4 \pm 1.1 \pm 0.8) \times 10^{-4}$  per lepton species, averaged over electrons and muons, with  $X_c$  a charmed baryon.

<sup>3</sup> ABREU 95V observe an excess of "right-sign" pairs  $\Xi^\mp \ell^\mp$  compared to "wrong-sign" pairs  $\Xi^\mp \ell^\pm$  in jets: this excess is interpreted as evidence for the beauty strange baryon  $\Xi_b$  production, with  $\Xi_b \rightarrow \Xi^- \ell^- \bar{\nu}_\ell X$ . They find that the probability for this signal to come from non  $b$ -baryon decays is less than  $5 \times 10^{-4}$  and that  $\Lambda_b$  decays can account for less than 10% of these events. The  $\Xi_b$  production rate is then measured to be  $B(b \rightarrow \Xi_b) \times B(\Xi_b \rightarrow \Xi^- \ell^- X) = (5.9 \pm 2.1 \pm 1.0) \times 10^{-4}$  per lepton species, averaged over electrons and muons.

### $\Gamma(b\text{-baryon } X)/\Gamma(\text{hadrons})$

$\Gamma_{57}/\Gamma_8$

“OUR EVALUATION” is obtained using our current values for  $f(b \rightarrow b\text{-baryon})$  and  $R_b = \Gamma(b\bar{b})/\Gamma(\text{hadrons})$ . We calculate  $\Gamma(b\text{-baryon } X)/\Gamma(\text{hadrons}) = R_b \times f(b \rightarrow b\text{-baryon})$ .

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>0.0197 ± 0.0032 OUR EVALUATION</b>		(Produced by HFLAV)		
<b>0.0221 ± 0.0015 ± 0.0058</b>		<sup>1</sup> BARATE	98V ALEP	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV

<sup>1</sup> BARATE 98V use the overall number of identified protons in  $b$ -hadron decays to measure  $f(b \rightarrow b\text{-baryon}) = 0.102 \pm 0.007 \pm 0.027$ . They assume  $\text{BR}(b\text{-baryon} \rightarrow pX) = (58 \pm 6)\%$  and  $\text{BR}(B_s^0 \rightarrow pX) = (8.0 \pm 4.0)\%$ . The value quoted here is obtained multiplying this production fraction by our value of  $R_b = \Gamma(b\bar{b})/\Gamma(\text{hadrons})$ .

### $\Gamma(\text{anomalous } \gamma + \text{hadrons})/\Gamma_{\text{total}}$

$\Gamma_{58}/\Gamma$

Limits on additional sources of prompt photons beyond expectations for final-state bremsstrahlung.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 3.2 × 10<sup>-3</sup></b>	95	<sup>1</sup> AKRAWY	90J OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV

<sup>1</sup> AKRAWY 90J report  $\Gamma(\gamma X) < 8.2$  MeV at 95%CL. They assume a three-body  $\gamma q\bar{q}$  distribution and use  $E(\gamma) > 10$  GeV.

### $\Gamma(e^+ e^- \gamma)/\Gamma_{\text{total}}$

$\Gamma_{59}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 5.2 × 10<sup>-4</sup></b>	95	<sup>1</sup> ACTON	91B OPAL	$E_{\text{cm}}^{ee} = 91.2$ GeV

<sup>1</sup> ACTON 91B looked for isolated photons with  $E > 2\%$  of beam energy ( $> 0.9$  GeV).

### $\Gamma(\mu^+ \mu^- \gamma)/\Gamma_{\text{total}}$

$\Gamma_{60}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 5.6 × 10<sup>-4</sup></b>	95	<sup>1</sup> ACTON	91B OPAL	$E_{\text{cm}}^{ee} = 91.2$ GeV

<sup>1</sup> ACTON 91B looked for isolated photons with  $E > 2\%$  of beam energy ( $> 0.9$  GeV).

### $\Gamma(\tau^+ \tau^- \gamma)/\Gamma_{\text{total}}$

$\Gamma_{61}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 7.3 × 10<sup>-4</sup></b>	95	<sup>1</sup> ACTON	91B OPAL	$E_{\text{cm}}^{ee} = 91.2$ GeV

<sup>1</sup> ACTON 91B looked for isolated photons with  $E > 2\%$  of beam energy ( $> 0.9$  GeV).

### $\Gamma(\ell^+ \ell^- \gamma\gamma)/\Gamma_{\text{total}}$

$\Gamma_{62}/\Gamma$

The value is the sum over  $\ell = e, \mu, \tau$ .

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 6.8 × 10<sup>-6</sup></b>	95	<sup>1</sup> ACTON	93E OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV

<sup>1</sup> For  $m_{\gamma\gamma} = 60 \pm 5$  GeV.

### $\Gamma(q\bar{q}\gamma\gamma)/\Gamma_{\text{total}}$

$\Gamma_{63}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 5.5 × 10<sup>-6</sup></b>	95	<sup>1</sup> ACTON	93E OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV

<sup>1</sup> For  $m_{\gamma\gamma} = 60 \pm 5$  GeV.



$\Gamma(\nu\bar{\nu}\gamma\gamma)/\Gamma_{\text{total}}$   $\Gamma_{64}/\Gamma$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<3.1 \times 10^{-6}$	95	<sup>1</sup> ACTON	93E OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV

<sup>1</sup> For  $m_{\gamma\gamma} = 60 \pm 5$  GeV.

$\Gamma(e^{\pm}\mu^{\mp})/\Gamma_{\text{total}}$   $\Gamma_{65}/\Gamma$

Test of lepton family number conservation. The value is for the sum of the charge states indicated.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<2.62 \times 10^{-7}$	95	AAD	23AQ ATLS	$E_{\text{cm}}^{pp} = 13$ TeV
$<7.5 \times 10^{-7}$	95	AAD	14AU ATLS	$E_{\text{cm}}^{pp} = 8$ TeV
$<2.5 \times 10^{-6}$	95	ABREU	97C DLPH	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
$<1.7 \times 10^{-6}$	95	AKERS	95W OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
$<0.6 \times 10^{-5}$	95	ADRIANI	93I L3	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
$<2.6 \times 10^{-5}$	95	DECAMP	92 ALEP	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV

$\Gamma(e^{\pm}\mu^{\mp})/\Gamma(e^+e^-)$   $\Gamma_{65}/\Gamma_1$

Test of lepton family number conservation. The value is for the sum of the charge states indicated.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<0.07$	90	ALBAJAR	89 UA1	$E_{\text{cm}}^{p\bar{p}} = 546,630$ GeV

$\Gamma(e^{\pm}\tau^{\mp})/\Gamma_{\text{total}}$   $\Gamma_{66}/\Gamma$

Test of lepton family number conservation. The value is for the sum of the charge states indicated.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<5.0 \times 10^{-6}$	95	AAD	21AV ATLS	$E_{\text{cm}}^{pp} = 13$ TeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$<8.1 \times 10^{-6}$	95	AAD	21AO ATLS	$E_{\text{cm}}^{pp} = 13$ TeV
$<5.8 \times 10^{-5}$	95	AABOUD	18CN ATLS	$E_{\text{cm}}^{pp} = 13$ TeV
$<2.2 \times 10^{-5}$	95	ABREU	97C DLPH	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
$<9.8 \times 10^{-6}$	95	AKERS	95W OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
$<1.3 \times 10^{-5}$	95	ADRIANI	93I L3	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
$<1.2 \times 10^{-4}$	95	DECAMP	92 ALEP	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV

$\Gamma(\mu^{\pm}\tau^{\mp})/\Gamma_{\text{total}}$   $\Gamma_{67}/\Gamma$

Test of lepton family number conservation. The value is for the sum of the charge states indicated.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<6.5 \times 10^{-6}$	95	AAD	21AV ATLS	$E_{\text{cm}}^{pp} = 13$ TeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$<9.5 \times 10^{-6}$	95	AAD	21AO ATLS	$E_{\text{cm}}^{pp} = 13$ TeV
$<1.3 \times 10^{-5}$	95	AABOUD	18CN ATLS	$E_{\text{cm}}^{pp} = 8, 13$ TeV
$<1.2 \times 10^{-5}$	95	ABREU	97C DLPH	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
$<1.7 \times 10^{-5}$	95	AKERS	95W OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
$<1.9 \times 10^{-5}$	95	ADRIANI	93I L3	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV
$<1.0 \times 10^{-4}$	95	DECAMP	92 ALEP	$E_{\text{cm}}^{ee} = 88\text{--}94$ GeV

**$\Gamma(pe)/\Gamma_{\text{total}}$**   **$\Gamma_{68}/\Gamma$**   
 Test of baryon number and lepton number conservations. Charge conjugate states are implied.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>&lt;1.8 \times 10^{-6}</math></b>	95	<sup>1</sup> ABBIENDI	99I	OPAL $E_{\text{cm}}^{ee} = 88\text{--}94$ GeV

<sup>1</sup> ABBIENDI 99I give the 95%CL limit on the partial width  $\Gamma(Z^0 \rightarrow pe) < 4.6$  KeV and we have transformed it into a branching ratio.

**$\Gamma(p\mu)/\Gamma_{\text{total}}$**   **$\Gamma_{69}/\Gamma$**   
 Test of baryon number and lepton number conservations. Charge conjugate states are implied.

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>&lt;1.8 \times 10^{-6}</math></b>	95	<sup>1</sup> ABBIENDI	99I	OPAL $E_{\text{cm}}^{ee} = 88\text{--}94$ GeV

<sup>1</sup> ABBIENDI 99I give the 95%CL limit on the partial width  $\Gamma(Z^0 \rightarrow p\mu) < 4.4$  KeV and we have transformed it into a branching ratio.

**AVERAGE PARTICLE MULTIPLICITIES IN HADRONIC Z DECAY**

Summed over particle and antiparticle, when appropriate.

**$\langle N_\gamma \rangle$**

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>20.97 \pm 0.02 \pm 1.15</math></b>	ACKERSTAFF	98A	OPAL $E_{\text{cm}}^{ee} = 91.2$ GeV

**$\langle N_{\pi^\pm} \rangle$**


<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>17.03 \pm 0.16</math> OUR AVERAGE</b>			
$17.007 \pm 0.209$	ABE	04C	SLD $E_{\text{cm}}^{ee} = 91.2$ GeV
$17.26 \pm 0.10 \pm 0.88$	ABREU	98L	DLPH $E_{\text{cm}}^{ee} = 91.2$ GeV
$17.04 \pm 0.31$	BARATE	98V	ALEP $E_{\text{cm}}^{ee} = 91.2$ GeV
$17.05 \pm 0.43$	AKERS	94P	OPAL $E_{\text{cm}}^{ee} = 91.2$ GeV

**$\langle N_{\pi^0} \rangle$**

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>9.76 \pm 0.26</math> OUR AVERAGE</b>			
$9.55 \pm 0.06 \pm 0.75$	ACKERSTAFF	98A	OPAL $E_{\text{cm}}^{ee} = 91.2$ GeV
$9.63 \pm 0.13 \pm 0.63$	BARATE	97J	ALEP $E_{\text{cm}}^{ee} = 91.2$ GeV
$9.90 \pm 0.02 \pm 0.33$	ACCIARRI	96	L3 $E_{\text{cm}}^{ee} = 91.2$ GeV
$9.2 \pm 0.2 \pm 1.0$	ADAM	96	DLPH $E_{\text{cm}}^{ee} = 91.2$ GeV

**$\langle N_\eta \rangle$**

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>1.01 \pm 0.08</math> OUR AVERAGE</b>			Error includes scale factor of 1.3. See the ideogram below.
$1.20 \pm 0.04 \pm 0.11$	HEISTER	02C	ALEP $E_{\text{cm}}^{ee} = 91.2$ GeV
$0.97 \pm 0.03 \pm 0.11$	ACKERSTAFF	98A	OPAL $E_{\text{cm}}^{ee} = 91.2$ GeV
$0.93 \pm 0.01 \pm 0.09$	ACCIARRI	96	L3 $E_{\text{cm}}^{ee} = 91.2$ GeV



$\langle N_{\rho^\pm} \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>2.57 ± 0.15 OUR AVERAGE</b>			
2.59 ± 0.03 ± 0.16	<sup>1</sup> BEDDALL 09		ALEPH archive, $E_{cm}^{ee} = 91.2$ GeV
2.40 ± 0.06 ± 0.43	ACKERSTAFF 98A	OPAL	$E_{cm}^{ee} = 91.2$ GeV

<sup>1</sup> BEDDALL 09 analyse 3.2 million hadronic Z decays as archived by ALEPH collaboration and report a value of  $2.59 \pm 0.03 \pm 0.15 \pm 0.04$ . The first error is statistical, the second systematic, and the third arises from extrapolation to full phase space. We combine the systematic errors in quadrature.

$\langle N_{\rho^0} \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>1.24 ± 0.10 OUR AVERAGE</b>	Error includes scale factor of 1.1.		
1.19 ± 0.10	ABREU	99J	DLPH $E_{cm}^{ee} = 91.2$ GeV
1.45 ± 0.06 ± 0.20	BUSKULIC	96H	ALEP $E_{cm}^{ee} = 91.2$ GeV

$\langle N_\omega \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>1.02 ± 0.06 OUR AVERAGE</b>			
1.00 ± 0.03 ± 0.06	HEISTER	02C	ALEP $E_{cm}^{ee} = 91.2$ GeV
1.04 ± 0.04 ± 0.14	ACKERSTAFF	98A	OPAL $E_{cm}^{ee} = 91.2$ GeV
1.17 ± 0.09 ± 0.15	ACCIARRI	97D	L3 $E_{cm}^{ee} = 91.2$ GeV

$\langle N_{\eta'} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.17 ± 0.05 OUR AVERAGE</b>	Error includes scale factor of 2.4.		
0.14 ± 0.01 ± 0.02	ACKERSTAFF 98A	OPAL	$E_{cm}^{ee} = 91.2 \text{ GeV}$
0.25 ± 0.04	<sup>1</sup> ACCIARRI 97D	L3	$E_{cm}^{ee} = 91.2 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.068 ± 0.018 ± 0.016	<sup>2</sup> BUSKULIC 92D	ALEP	$E_{cm}^{ee} = 91.2 \text{ GeV}$

<sup>1</sup> ACCIARRI 97D obtain this value averaging over the two decay channels  $\eta' \rightarrow \pi^+ \pi^- \eta$  and  $\eta' \rightarrow \rho^0 \gamma$ .

<sup>2</sup> BUSKULIC 92D obtain this value for  $x > 0.1$ .

$\langle N_{f_0(980)} \rangle$


<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.147 ± 0.011 OUR AVERAGE</b>			
0.164 ± 0.021	ABREU 99J	DLPH	$E_{cm}^{ee} = 91.2 \text{ GeV}$
0.141 ± 0.007 ± 0.011	ACKERSTAFF 98Q	OPAL	$E_{cm}^{ee} = 91.2 \text{ GeV}$

$\langle N_{a_0(980)\pm} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.27 ± 0.04 ± 0.10</b>	ACKERSTAFF 98A	OPAL	$E_{cm}^{ee} = 91.2 \text{ GeV}$

$\langle N_{\phi} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.098 ± 0.006 OUR AVERAGE</b>	Error includes scale factor of 2.0. See the ideogram below.		
0.105 ± 0.008	ABE 99E	SLD	$E_{cm}^{ee} = 91.2 \text{ GeV}$
0.091 ± 0.002 ± 0.003	ACKERSTAFF 98Q	OPAL	$E_{cm}^{ee} = 91.2 \text{ GeV}$
0.104 ± 0.003 ± 0.007	ABREU 96U	DLPH	$E_{cm}^{ee} = 91.2 \text{ GeV}$
0.122 ± 0.004 ± 0.008	BUSKULIC 96H	ALEP	$E_{cm}^{ee} = 91.2 \text{ GeV}$



**$\langle N_{f_2(1270)} \rangle$**

VALUE	DOCUMENT ID	TECN	COMMENT
<b><math>0.169 \pm 0.025</math> OUR AVERAGE</b>	Error includes scale factor of 1.4.		
$0.214 \pm 0.038$	ABREU	99J	DLPH $E_{cm}^{ee} = 91.2$ GeV
$0.155 \pm 0.011 \pm 0.018$	ACKERSTAFF	98Q	OPAL $E_{cm}^{ee} = 91.2$ GeV

**$\langle N_{f_1(1285)} \rangle$**

VALUE	DOCUMENT ID	TECN	COMMENT
<b><math>0.165 \pm 0.051</math></b>	<sup>1</sup> ABDALLAH	03H	DLPH $E_{cm}^{ee} = 91.2$ GeV

<sup>1</sup> ABDALLAH 03H assume a  $K\bar{K}\pi$  branching ratio of  $(9.0 \pm 0.4)\%$ .

**$\langle N_{f_1(1420)} \rangle$**

VALUE	DOCUMENT ID	TECN	COMMENT
<b><math>0.056 \pm 0.012</math></b>	<sup>1</sup> ABDALLAH	03H	DLPH $E_{cm}^{ee} = 91.2$ GeV

<sup>1</sup> ABDALLAH 03H assume a  $K\bar{K}\pi$  branching ratio of 100%.

**$\langle N_{f_2'(1525)} \rangle$**


VALUE	DOCUMENT ID	TECN	COMMENT
<b><math>0.012 \pm 0.006</math></b>	ABREU	99J	DLPH $E_{cm}^{ee} = 91.2$ GeV

$\langle N_{K^\pm} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>2.24 ± 0.04 OUR AVERAGE</b>			
2.203 ± 0.071	ABE	04C	SLD $E_{cm}^{ee} = 91.2$ GeV
2.21 ± 0.05 ± 0.05	ABREU	98L	DLPH $E_{cm}^{ee} = 91.2$ GeV
2.26 ± 0.12	BARATE	98V	ALEP $E_{cm}^{ee} = 91.2$ GeV
2.42 ± 0.13	AKERS	94P	OPAL $E_{cm}^{ee} = 91.2$ GeV

$\langle N_{K^0} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>2.039 ± 0.025 OUR AVERAGE</b>	Error includes scale factor of 1.3. See the ideogram below.		
2.093 ± 0.004 ± 0.029	BARATE	00O	ALEP $E_{cm}^{ee} = 91.2$ GeV
2.01 ± 0.08	ABE	99E	SLD $E_{cm}^{ee} = 91.2$ GeV
2.024 ± 0.006 ± 0.042	ACCIARRI	97L	L3 $E_{cm}^{ee} = 91.2$ GeV
1.962 ± 0.022 ± 0.056	ABREU	95L	DLPH $E_{cm}^{ee} = 91.2$ GeV
1.99 ± 0.01 ± 0.04	AKERS	95U	OPAL $E_{cm}^{ee} = 91.2$ GeV



$\langle N_{K^*(892)^\pm} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.72 ± 0.05 OUR AVERAGE</b>			
0.712 ± 0.031 ± 0.059	ABREU	95L	DLPH $E_{cm}^{ee} = 91.2$ GeV
0.72 ± 0.02 ± 0.08	ACTON	93	OPAL $E_{cm}^{ee} = 91.2$ GeV

$\langle N_{K^*(892)^0} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.739±0.022 OUR AVERAGE</b>			
0.707±0.041	ABE	99E	SLD $E_{cm}^{ee} = 91.2$ GeV
0.74 ±0.02 ±0.02	ACKERSTAFF	97S	OPAL $E_{cm}^{ee} = 91.2$ GeV
0.77 ±0.02 ±0.07	ABREU	96U	DLPH $E_{cm}^{ee} = 91.2$ GeV
0.83 ±0.01 ±0.09	BUSKULIC	96H	ALEP $E_{cm}^{ee} = 91.2$ GeV
0.97 ±0.18 ±0.31	ABREU	93	DLPH $E_{cm}^{ee} = 91.2$ GeV

$\langle N_{K_2^*(1430)} \rangle$


<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.073±0.023</b>	ABREU	99J	DLPH $E_{cm}^{ee} = 91.2$ GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.19 ±0.04 ±0.06	<sup>1</sup> AKERS	95X	OPAL $E_{cm}^{ee} = 91.2$ GeV

<sup>1</sup> AKERS 95X obtain this value for  $x < 0.3$ .

$\langle N_{D^\pm} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.187±0.020 OUR AVERAGE</b>	Error includes scale factor of 1.5.		See the ideogram below.
0.170±0.009±0.014	ALEXANDER	96R	OPAL $E_{cm}^{ee} = 91.2$ GeV
0.251±0.026±0.025	BUSKULIC	94J	ALEP $E_{cm}^{ee} = 91.2$ GeV
0.199±0.019±0.024	<sup>1</sup> ABREU	93I	DLPH $E_{cm}^{ee} = 91.2$ GeV

<sup>1</sup> See ABREU 95 (erratum).



$\langle N_{D^0} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.462 ± 0.026 OUR AVERAGE</b>			
0.465 ± 0.017 ± 0.027	ALEXANDER 96R	OPAL	$E_{cm}^{ee} = 91.2$ GeV
0.518 ± 0.052 ± 0.035	BUSKULIC 94J	ALEP	$E_{cm}^{ee} = 91.2$ GeV
0.403 ± 0.038 ± 0.044	<sup>1</sup> ABREU 93I	DLPH	$E_{cm}^{ee} = 91.2$ GeV

<sup>1</sup> See ABREU 95 (erratum).

$\langle N_{D_s^\pm} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.131 ± 0.010 ± 0.018</b>	ALEXANDER 96R	OPAL	$E_{cm}^{ee} = 91.2$ GeV

$\langle N_{D^{*(2010)\pm}} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.183 ± 0.008 OUR AVERAGE</b>			
0.1854 ± 0.0041 ± 0.0091	<sup>1</sup> ACKERSTAFF 98E	OPAL	$E_{cm}^{ee} = 91.2$ GeV
0.187 ± 0.015 ± 0.013	BUSKULIC 94J	ALEP	$E_{cm}^{ee} = 91.2$ GeV
0.171 ± 0.012 ± 0.016	<sup>2</sup> ABREU 93I	DLPH	$E_{cm}^{ee} = 91.2$ GeV

<sup>1</sup> ACKERSTAFF 98E systematic error includes an uncertainty of ±0.0069 due to the branching ratios  $B(D^{*+} \rightarrow D^0 \pi^+) = 0.683 \pm 0.014$  and  $B(D^0 \rightarrow K^- \pi^+) = 0.0383 \pm 0.0012$ .

<sup>2</sup> See ABREU 95 (erratum).

$\langle N_{D_{s1}(2536)^+} \rangle$

<u>VALUE (units 10<sup>-3</sup>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
2.9 <sup>+0.7</sup> <sub>-0.6</sub> ± 0.2	<sup>1</sup> ACKERSTAFF 97W	OPAL	$E_{cm}^{ee} = 91.2$ GeV

<sup>1</sup> ACKERSTAFF 97W obtain this value for  $x > 0.6$  and with the assumption that its decay width is saturated by the  $D^* K$  final states.

$\langle N_{B^*} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.28 ± 0.01 ± 0.03</b>	<sup>1</sup> ABREU 95R	DLPH	$E_{cm}^{ee} = 91.2$ GeV

<sup>1</sup> ABREU 95R quote this value for a flavor-averaged excited state.

$\langle N_{J/\psi(1S)} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.0056 ± 0.0003 ± 0.0004</b>	<sup>1</sup> ALEXANDER 96B	OPAL	$E_{cm}^{ee} = 91.2$ GeV

<sup>1</sup> ALEXANDER 96B identify  $J/\psi(1S)$  from the decays into lepton pairs.

$\langle N_{\psi(2S)} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.0023 ± 0.0004 ± 0.0003</b>	ALEXANDER 96B	OPAL	$E_{cm}^{ee} = 91.2$ GeV



$\langle N_p \rangle$


<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1.046 ± 0.026 OUR AVERAGE</b>			
1.054 ± 0.035	ABE	04C	SLD $E_{cm}^{ee} = 91.2$ GeV
1.08 ± 0.04 ± 0.03	ABREU	98L	DLPH $E_{cm}^{ee} = 91.2$ GeV
1.00 ± 0.07	BARATE	98V	ALEP $E_{cm}^{ee} = 91.2$ GeV
0.92 ± 0.11	AKERS	94P	OPAL $E_{cm}^{ee} = 91.2$ GeV

$\langle N_{\Delta(1232)++} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.087 ± 0.033 OUR AVERAGE</b>	Error includes scale factor of 2.4.		
0.079 ± 0.009 ± 0.011	ABREU	95W	DLPH $E_{cm}^{ee} = 91.2$ GeV
0.22 ± 0.04 ± 0.04	ALEXANDER	95D	OPAL $E_{cm}^{ee} = 91.2$ GeV

$\langle N_\Lambda \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.388 ± 0.009 OUR AVERAGE</b>	Error includes scale factor of 1.7. See the ideogram below.		
0.404 ± 0.002 ± 0.007	BARATE	00O	ALEP $E_{cm}^{ee} = 91.2$ GeV
0.395 ± 0.022	ABE	99E	SLD $E_{cm}^{ee} = 91.2$ GeV
0.364 ± 0.004 ± 0.017	ACCIARRI	97L	L3 $E_{cm}^{ee} = 91.2$ GeV
0.374 ± 0.002 ± 0.010	ALEXANDER	97D	OPAL $E_{cm}^{ee} = 91.2$ GeV
0.357 ± 0.003 ± 0.017	ABREU	93L	DLPH $E_{cm}^{ee} = 91.2$ GeV



$\langle N_{\Lambda(1520)} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.0224 ± 0.0027 OUR AVERAGE</b>			
0.029 ± 0.005 ± 0.005	ABREU 00P	DLPH	$E_{cm}^{ee} = 91.2 \text{ GeV}$
0.0213 ± 0.0021 ± 0.0019	ALEXANDER 97D	OPAL	$E_{cm}^{ee} = 91.2 \text{ GeV}$

$\langle N_{\Sigma^+} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.107 ± 0.010 OUR AVERAGE</b>			
0.114 ± 0.011 ± 0.009	ACCIARRI 00J	L3	$E_{cm}^{ee} = 91.2 \text{ GeV}$
0.099 ± 0.008 ± 0.013	ALEXANDER 97E	OPAL	$E_{cm}^{ee} = 91.2 \text{ GeV}$

$\langle N_{\Sigma^-} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.082 ± 0.007 OUR AVERAGE</b>			
0.081 ± 0.002 ± 0.010	ABREU 00P	DLPH	$E_{cm}^{ee} = 91.2 \text{ GeV}$
0.083 ± 0.006 ± 0.009	ALEXANDER 97E	OPAL	$E_{cm}^{ee} = 91.2 \text{ GeV}$

$\langle N_{\Sigma^{++}\Sigma^-} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.181 ± 0.018 OUR AVERAGE</b>			
0.182 ± 0.010 ± 0.016	<sup>1</sup> ALEXANDER 97E	OPAL	$E_{cm}^{ee} = 91.2 \text{ GeV}$
0.170 ± 0.014 ± 0.061	ABREU 95O	DLPH	$E_{cm}^{ee} = 91.2 \text{ GeV}$

<sup>1</sup>We have combined the values of  $\langle N_{\Sigma^+} \rangle$  and  $\langle N_{\Sigma^-} \rangle$  from ALEXANDER 97E adding the statistical and systematic errors of the two final states separately in quadrature. If isospin symmetry is assumed this value becomes  $0.174 \pm 0.010 \pm 0.015$ .

$\langle N_{\Sigma^0} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.076 ± 0.010 OUR AVERAGE</b>			
0.095 ± 0.015 ± 0.013	ACCIARRI 00J	L3	$E_{cm}^{ee} = 91.2 \text{ GeV}$
0.071 ± 0.012 ± 0.013	ALEXANDER 97E	OPAL	$E_{cm}^{ee} = 91.2 \text{ GeV}$
0.070 ± 0.010 ± 0.010	ADAM 96B	DLPH	$E_{cm}^{ee} = 91.2 \text{ GeV}$

$\langle N_{(\Sigma^+\Sigma^-\Sigma^0)/3} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.084 ± 0.005 ± 0.008</b>	ALEXANDER 97E	OPAL	$E_{cm}^{ee} = 91.2 \text{ GeV}$

$\langle N_{\Sigma(1385)^+} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.0239 ± 0.0009 ± 0.0012</b>	ALEXANDER 97D	OPAL	$E_{cm}^{ee} = 91.2 \text{ GeV}$

$\langle N_{\Sigma(1385)^-} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.0240 ± 0.0010 ± 0.0014</b>	ALEXANDER 97D	OPAL	$E_{cm}^{ee} = 91.2 \text{ GeV}$

$\langle N_{\Sigma(1385)^{++}\Sigma(1385)^{-}} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.046 ± 0.004 OUR AVERAGE</b>	Error includes scale factor of 1.6.		
0.0479 ± 0.0013 ± 0.0026	ALEXANDER	97D OPAL	$E_{cm}^{ee} = 91.2 \text{ GeV}$
0.0382 ± 0.0028 ± 0.0045	ABREU	95O DLPH	$E_{cm}^{ee} = 91.2 \text{ GeV}$

$\langle N_{\Xi^{-}} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.0258 ± 0.0009 OUR AVERAGE</b>			
0.0247 ± 0.0009 ± 0.0025	ABDALLAH	06E DLPH	$E_{cm}^{ee} = 91.2 \text{ GeV}$
0.0259 ± 0.0004 ± 0.0009	ALEXANDER	97D OPAL	$E_{cm}^{ee} = 91.2 \text{ GeV}$

$\langle N_{\Xi(1530)^0} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.0059 ± 0.0011 OUR AVERAGE</b>	Error includes scale factor of 2.3.		
0.0045 ± 0.0005 ± 0.0006	ABDALLAH	05C DLPH	$E_{cm}^{ee} = 91.2 \text{ GeV}$
0.0068 ± 0.0005 ± 0.0004	ALEXANDER	97D OPAL	$E_{cm}^{ee} = 91.2 \text{ GeV}$

$\langle N_{\Omega^{-}} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.00164 ± 0.00028 OUR AVERAGE</b>			
0.0018 ± 0.0003 ± 0.0002	ALEXANDER	97D OPAL	$E_{cm}^{ee} = 91.2 \text{ GeV}$
0.0014 ± 0.0002 ± 0.0004	ADAM	96B DLPH	$E_{cm}^{ee} = 91.2 \text{ GeV}$

$\langle N_{\Lambda_c^+} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.078 ± 0.012 ± 0.012</b>	ALEXANDER	96R OPAL	$E_{cm}^{ee} = 91.2 \text{ GeV}$

$\langle N_{\bar{D}} \rangle$

<u>VALUE (units 10<sup>-6</sup>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------------------------------	--------------------	-------------	----------------


• • • We do not use the following data for averages, fits, limits, etc. • • •

5.9 ± 1.8 ± 0.5	<sup>1</sup> SCHAEL	06A ALEP	$E_{cm}^{ee} = 91.2 \text{ GeV}$
-----------------	---------------------	----------	----------------------------------

<sup>1</sup>SCHAEL 06A obtain this anti-deuteron production rate per hadronic Z decay in the anti-deuteron momentum range from 0.62 to 1.03 GeV/c.

$\langle N_{charged} \rangle$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>20.76 ± 0.16 OUR AVERAGE</b>	Error includes scale factor of 2.1. See the ideogram below.		
20.46 ± 0.01 ± 0.11	ACHARD	03G L3	$E_{cm}^{ee} = 91.2 \text{ GeV}$
21.21 ± 0.01 ± 0.20	ABREU	99 DLPH	$E_{cm}^{ee} = 91.2 \text{ GeV}$
21.05 ± 0.20	AKERS	95Z OPAL	$E_{cm}^{ee} = 91.2 \text{ GeV}$
20.91 ± 0.03 ± 0.22	BUSKULIC	95R ALEP	$E_{cm}^{ee} = 91.2 \text{ GeV}$
21.40 ± 0.43	ACTON	92B OPAL	$E_{cm}^{ee} = 91.2 \text{ GeV}$
20.71 ± 0.04 ± 0.77	ABREU	91H DLPH	$E_{cm}^{ee} = 91.2 \text{ GeV}$
20.7 ± 0.7	ADEVA	91I L3	$E_{cm}^{ee} = 91.2 \text{ GeV}$
20.1 ± 1.0 ± 0.9	ABRAMS	90 MRK2	$E_{cm}^{ee} = 91.1 \text{ GeV}$



## Z HADRONIC POLE CROSS SECTION

OUR EVALUATION is obtained using the fit procedure and correlations as determined by the LEP Electroweak Working Group (see the note “The Z boson” and ref. LEP-SLC 06). Corrections as discussed in VOUSINAS 20 and JANOT 20 are also included. This quantity is defined as

$$\sigma_h^0 = \frac{12\pi}{M_Z^2} \frac{\Gamma(e^+ e^-) \Gamma(\text{hadrons})}{\Gamma_Z^2}$$

It is one of the parameters used in the Z lineshape fit.

VALUE (nb)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>41.4802 ± 0.0325 OUR EVALUATION</b>				
41.4802 ± 0.0325		<sup>1</sup> JANOT 20		
• • • We do not use the following data for averages, fits, limits, etc. • • •				
41.500 ± 0.037		<sup>2</sup> VOUSINAS 20		
41.541 ± 0.037		LEP-SLC 06		$E_{cm}^{ee} = 88-94$ GeV
41.501 ± 0.055	4.10M	<sup>3</sup> ABBIENDI 01A	OPAL	$E_{cm}^{ee} = 88-94$ GeV
41.578 ± 0.069	3.70M	ABREU 00F	DLPH	$E_{cm}^{ee} = 88-94$ GeV
41.535 ± 0.055	3.54M	ACCIARRI 00C	L3	$E_{cm}^{ee} = 88-94$ GeV
41.559 ± 0.058	4.07M	<sup>4</sup> BARATE 00C	ALEP	$E_{cm}^{ee} = 88-94$ GeV
42 ± 4	450	ABRAMS 89B	MRK2	$E_{cm}^{ee} = 89.2-93.0$ GeV

- <sup>1</sup> JANOT 20 applies a correction to LEP-SLC 06 using an updated Bhabha cross section calculation. This result also includes a correction to account for correlated luminosity bias as presented in VOUTSINAS 20.
- <sup>2</sup> VOUTSINAS 20 applies a correction to LEP-SLC 06 to account for correlated luminosity bias.
- <sup>3</sup> ABBIENDI 01A error includes approximately 0.031 due to statistics, 0.033 due to event selection systematics, 0.029 due to uncertainty in luminosity measurement, and 0.011 due to LEP energy uncertainty.
- <sup>4</sup> BARATE 00C error includes approximately 0.030 due to statistics, 0.026 due to experimental systematics, and 0.025 due to uncertainty in luminosity measurement.

## Z VECTOR COUPLINGS

These quantities are the effective vector couplings of the  $Z$  to charged leptons and quarks. Their magnitude is derived from a measurement of the  $Z$  lineshape and the forward-backward lepton asymmetries as a function of energy around the  $Z$  mass. The relative sign among the vector to axial-vector couplings is obtained from a measurement of the  $Z$  asymmetry parameters,  $A_e$ ,  $A_\mu$ , and  $A_\tau$ . By convention the sign of  $g_A^e$  is fixed to be negative (and opposite to that of  $g_V^e$  obtained using  $\nu_e$  scattering measurements). For the light quarks, the sign of the couplings is assigned consistently with this assumption. The LEP/SLD-based fit values quoted below correspond to global nine- or five-parameter fits to lineshape, lepton forward-backward asymmetry, and  $A_e$ ,  $A_\mu$ , and  $A_\tau$  measurements. See the note “The  $Z$  boson” and ref. LEP-SLC 06 for details. Where  $p\bar{p}$  and  $e\bar{p}$  data is quoted, OUR FIT value corresponds to a weighted average of this with the LEP/SLD fit result.

$g_V^e$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>-0.03817 \pm 0.00047</math> OUR FIT</b>				
$-0.058 \pm 0.016 \pm 0.007$	5026	<sup>1</sup> ACOSTA	05M CDF	$E_{cm}^{p\bar{p}} = 1.96$ TeV
$-0.0346 \pm 0.0023$	137.0k	<sup>2</sup> ABBIENDI	010 OPAL	$E_{cm}^{ee} = 88-94$ GeV
$-0.0412 \pm 0.0027$	124.4k	<sup>3</sup> ACCIARRI	00C L3	$E_{cm}^{ee} = 88-94$ GeV
$-0.0400 \pm 0.0037$		BARATE	00C ALEP	$E_{cm}^{ee} = 88-94$ GeV
$-0.0414 \pm 0.0020$		<sup>4</sup> ABE	95J SLD	$E_{cm}^{ee} = 91.31$ GeV

- <sup>1</sup> ACOSTA 05M determine the forward-backward asymmetry of  $e^+e^-$  pairs produced via  $q\bar{q} \rightarrow Z/\gamma^* \rightarrow e^+e^-$  in 15 M( $e^+e^-$ ) effective mass bins ranging from 40 GeV to 600 GeV. These results are used to obtain the vector and axial-vector couplings of the  $Z$  to  $e^+e^-$ , assuming the quark couplings are as predicted by the standard model. Higher order radiative corrections have not been taken into account.
- <sup>2</sup> ABBIENDI 010 use their measurement of the  $\tau$  polarization in addition to the lineshape and forward-backward lepton asymmetries.
- <sup>3</sup> ACCIARRI 00C use their measurement of the  $\tau$  polarization in addition to forward-backward lepton asymmetries.
- <sup>4</sup> ABE 95J obtain this result combining polarized Bhabha results with the  $A_{LR}$  measurement of ABE 94C. The Bhabha results alone give  $-0.0507 \pm 0.0096 \pm 0.0020$ .

### $g_V^\mu$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	-------------	--------------------	-------------	----------------

**-0.0367 ± 0.0023 OUR FIT**

$-0.0388^{+0.0060}_{-0.0064}$	182.8k	<sup>1</sup> ABBIENDI	01O OPAL	$E_{cm}^{ee} = 88-94$ GeV
$-0.0386 \pm 0.0073$	113.4k	<sup>2</sup> ACCIARRI	00C L3	$E_{cm}^{ee} = 88-94$ GeV
$-0.0362 \pm 0.0061$		BARATE	00C ALEP	$E_{cm}^{ee} = 88-94$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$-0.0413 \pm 0.0060$	66143	<sup>3</sup> ABBIENDI	01K OPAL	$E_{cm}^{ee} = 89-93$ GeV
----------------------	-------	-----------------------	----------	---------------------------

<sup>1</sup> ABBIENDI 01O use their measurement of the  $\tau$  polarization in addition to the lineshape and forward-backward lepton asymmetries.

<sup>2</sup> ACCIARRI 00C use their measurement of the  $\tau$  polarization in addition to forward-backward lepton asymmetries.

<sup>3</sup> ABBIENDI 01K obtain this from an angular analysis of the muon pair asymmetry which takes into account effects of initial state radiation on an event by event basis and of initial-final state interference.

### $g_V^\tau$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	-------------	--------------------	-------------	----------------

**-0.0366 ± 0.0010 OUR FIT**

$-0.0365 \pm 0.0023$	151.5k	<sup>1</sup> ABBIENDI	01O OPAL	$E_{cm}^{ee} = 88-94$ GeV
$-0.0384 \pm 0.0026$	103.0k	<sup>2</sup> ACCIARRI	00C L3	$E_{cm}^{ee} = 88-94$ GeV
$-0.0361 \pm 0.0068$		BARATE	00C ALEP	$E_{cm}^{ee} = 88-94$ GeV

<sup>1</sup> ABBIENDI 01O use their measurement of the  $\tau$  polarization in addition to the lineshape and forward-backward lepton asymmetries.

<sup>2</sup> ACCIARRI 00C use their measurement of the  $\tau$  polarization in addition to forward-backward lepton asymmetries.

### $g_V^l$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	-------------	--------------------	-------------	----------------

**-0.03783 ± 0.00041 OUR FIT**

$-0.0358 \pm 0.0014$	471.3k	<sup>1</sup> ABBIENDI	01O OPAL	$E_{cm}^{ee} = 88-94$ GeV
$-0.0397 \pm 0.0020$	379.4k	<sup>2</sup> ABREU	00F DLPH	$E_{cm}^{ee} = 88-94$ GeV
$-0.0397 \pm 0.0017$	340.8k	<sup>3</sup> ACCIARRI	00C L3	$E_{cm}^{ee} = 88-94$ GeV
$-0.0383 \pm 0.0018$	500k	BARATE	00C ALEP	$E_{cm}^{ee} = 88-94$ GeV

<sup>1</sup> ABBIENDI 01O use their measurement of the  $\tau$  polarization in addition to the lineshape and forward-backward lepton asymmetries.

<sup>2</sup> Using forward-backward lepton asymmetries.

<sup>3</sup> ACCIARRI 00C use their measurement of the  $\tau$  polarization in addition to forward-backward lepton asymmetries.

### $g_V^u$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	-------------	--------------------	-------------	----------------

**0.266 ± 0.034 OUR AVERAGE**

$0.270 \pm 0.037$		<sup>1</sup> ANDREEV	18A H1	$e^\pm p$
$0.201 \pm 0.112$	156k	<sup>2</sup> ABAZOV	11D D0	$E_{cm}^{p\bar{p}} = 1.97$ TeV
$0.24^{+0.28}_{-0.11}$		<sup>3</sup> LEP-SLC	06	$E_{cm}^{ee} = 88-94$ GeV
$0.399^{+0.152}_{-0.188} \pm 0.066$	5026	<sup>4</sup> ACOSTA	05M CDF	$E_{cm}^{p\bar{p}} = 1.96$ TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.14	$\begin{matrix} +0.09 \\ -0.09 \end{matrix}$		<sup>5</sup> ABRAMOWICZ16A	ZEUS	
0.144	$\begin{matrix} +0.066 \\ -0.058 \end{matrix}$		<sup>6</sup> ABT	16	
0.27	$\pm 0.13$	1500	<sup>7</sup> AKTAS	06 H1	$e^\pm p \rightarrow \bar{\nu}_e(\nu_e)X,$ $\sqrt{s} \approx 300 \text{ GeV}$

<sup>1</sup> ANDREEV 18A obtain this result in a combined electroweak and QCD analysis using all deep-inelastic  $e^+p$  and  $e^-p$  neutral current and charged current scattering cross sections published by the H1 Collaboration, including data with longitudinally polarized lepton beams.

<sup>2</sup> ABAZOV 11D study  $p\bar{p} \rightarrow Z/\gamma^* e^+e^-$  events using  $5 \text{ fb}^{-1}$  data at  $\sqrt{s} = 1.96 \text{ TeV}$ . The candidate events are selected by requiring two isolated electromagnetic showers with  $E_T > 25 \text{ GeV}$ , at least one electron in the central region and the di-electron mass in the range 50–1000 GeV. From the forward-backward asymmetry, determined as a function of the di-electron mass, they derive the axial and vector couplings of the  $u$ - and  $d$ - quarks and the value of  $\sin^2\theta_{eff}^e = 0.2309 \pm 0.0008(\text{stat}) \pm 0.0006(\text{syst})$ .

<sup>3</sup> LEP-SLC 06 is a combination of the results from LEP and SLC experiments using light quark tagging.  $s$ - and  $d$ -quark couplings are assumed to be identical.

<sup>4</sup> ACOSTA 05M determine the forward-backward asymmetry of  $e^+e^-$  pairs produced via  $q\bar{q} \rightarrow Z/\gamma^* \rightarrow e^+e^-$  in 15  $M(e^+e^-)$  effective mass bins ranging from 40 GeV to 600 GeV. These results are used to obtain the vector and axial-vector couplings of the  $Z$  to the light quarks, assuming the electron couplings are as predicted by the Standard Model. Higher order radiative corrections have not been taken into account.

<sup>5</sup> ABRAMOWICZ 16A determine the  $Z^0$  couplings to  $u$ - and  $d$ -quarks using the ZEUS polarised data from Run II together with the unpolarised data from both ZEUS and H1 Collaborations for Run I and unpolarised H1 data from Run II.

<sup>6</sup> ABT 16 determine the  $Z^0$  couplings to  $u$ - and  $d$ -quarks using the same techniques and data as ABRAMOWICZ 16A but additionally use the published H1 polarised data.

<sup>7</sup> AKTAS 06 fit the neutral current ( $1.5 \leq Q^2 \leq 30,000 \text{ GeV}^2$ ) and charged current ( $1.5 \leq Q^2 \leq 15,000 \text{ GeV}^2$ ) differential cross sections. In the determination of the  $u$ -quark couplings the electron and  $d$ -quark couplings are fixed to their standard model values.

$g_V^d$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>-0.38</b>	<b><math>\begin{matrix} +0.04 \\ -0.05 \end{matrix}</math></b>	<b>OUR AVERAGE</b>		
-0.488	$\pm 0.092$	<sup>1</sup> ANDREEV	18A H1	$e^\pm p$
-0.351	$\pm 0.251$	156k <sup>2</sup> ABAZOV	11D D0	$E_{cm}^{p\bar{p}} = 1.97 \text{ TeV}$
-0.33	$\begin{matrix} +0.05 \\ -0.07 \end{matrix}$	<sup>3</sup> LEP-SLC	06	$E_{cm}^{ee} = 88\text{--}94 \text{ GeV}$
-0.226	$\begin{matrix} +0.635 \\ -0.290 \end{matrix} \pm 0.090$	5026 <sup>4</sup> ACOSTA	05M CDF	$E_{cm}^{p\bar{p}} = 1.96 \text{ TeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

-0.41	$\begin{matrix} +0.25 \\ -0.20 \end{matrix}$		<sup>5</sup> ABRAMOWICZ16A	ZEUS	
-0.503	$\begin{matrix} +0.171 \\ -0.103 \end{matrix}$		<sup>6</sup> ABT	16	
-0.33	$\pm 0.33$	1500	<sup>7</sup> AKTAS	06 H1	$e^\pm p \rightarrow \bar{\nu}_e(\nu_e)X,$ $\sqrt{s} \approx 300 \text{ GeV}$

- <sup>1</sup> ANDREEV 18A obtain this result in a combined electroweak and QCD analysis using all deep-inelastic  $e^+p$  and  $e^-p$  neutral current and charged current scattering cross sections published by the H1 Collaboration, including data with longitudinally polarized lepton beams.
- <sup>2</sup> ABAZOV 11D study  $p\bar{p} \rightarrow Z/\gamma^* e^+e^-$  events using  $5 \text{ fb}^{-1}$  data at  $\sqrt{s} = 1.96 \text{ TeV}$ . The candidate events are selected by requiring two isolated electromagnetic showers with  $E_T > 25 \text{ GeV}$ , at least one electron in the central region and the di-electron mass in the range 50–1000 GeV. From the forward-backward asymmetry, determined as a function of the di-electron mass, they derive the axial and vector couplings of the  $u$ - and  $d$ - quarks and the value of  $\sin^2\theta_{eff}^l = 0.2309 \pm 0.0008(\text{stat}) \pm 0.0006(\text{syst})$ .
- <sup>3</sup> LEP-SLC 06 is a combination of the results from LEP and SLC experiments using light quark tagging.  $s$ - and  $d$ -quark couplings are assumed to be identical.
- <sup>4</sup> ACOSTA 05M determine the forward-backward asymmetry of  $e^+e^-$  pairs produced via  $q\bar{q} \rightarrow Z/\gamma^* \rightarrow e^+e^-$  in 15  $M(e^+e^-)$  effective mass bins ranging from 40 GeV to 600 GeV. These results are used to obtain the vector and axial-vector couplings of the  $Z$  to the light quarks, assuming the electron couplings are as predicted by the Standard Model. Higher order radiative corrections have not been taken into account.
- <sup>5</sup> ABRAMOWICZ 16A determine the  $Z^0$  couplings to  $u$ - and  $d$ -quarks using the ZEUS polarised data from Run II together with the unpolarised data from both ZEUS and H1 Collaborations for Run I and unpolarised H1 data from Run II.
- <sup>6</sup> ABT 16 determine the  $Z^0$  couplings to  $u$ - and  $d$ -quarks using the same techniques and data as ABRAMOWICZ 16A but additionally use the published H1 polarised data.
- <sup>7</sup> AKTAS 06 fit the neutral current ( $1.5 \leq Q^2 \leq 30,000 \text{ GeV}^2$ ) and charged current ( $1.5 \leq Q^2 \leq 15,000 \text{ GeV}^2$ ) differential cross sections. In the determination of the  $d$ -quark couplings the electron and  $u$ -quark couplings are fixed to their standard model values.

## Z AXIAL-VECTOR COUPLINGS

These quantities are the effective axial-vector couplings of the  $Z$  to charged leptons and quarks. Their magnitude is derived from a measurement of the  $Z$  lineshape and the forward-backward lepton asymmetries as a function of energy around the  $Z$  mass. The relative sign among the vector to axial-vector couplings is obtained from a measurement of the  $Z$  asymmetry parameters,  $A_e$ ,  $A_\mu$ , and  $A_\tau$ . By convention the sign of  $g_A^e$  is fixed to be negative (and opposite to that of  $g^{\nu_e}$  obtained using  $\nu_e$  scattering measurements). For the light quarks, the sign of the couplings is assigned consistently with this assumption. The LEP/SLD-based fit values quoted below correspond to global nine- or five-parameter fits to lineshape, lepton forward-backward asymmetry, and  $A_e$ ,  $A_\mu$ , and  $A_\tau$  measurements. See the note “The  $Z$  boson” and ref. LEP-SLC 06 for details. Where  $p\bar{p}$  and  $ep$  data is quoted, OUR FIT value corresponds to a weighted average of this with the LEP/SLD fit result.

$g_A^e$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>−0.50111±0.00035 OUR FIT</b>				
−0.528 ±0.123 ±0.059	5026	<sup>1</sup> ACOSTA	05M CDF	$E_{cm}^{p\bar{p}} = 1.96 \text{ TeV}$
−0.50062±0.00062	137.0k	<sup>2</sup> ABBIENDI	01O OPAL	$E_{cm}^{ee} = 88\text{--}94 \text{ GeV}$
−0.5015 ±0.0007	124.4k	<sup>3</sup> ACCIARRI	00C L3	$E_{cm}^{ee} = 88\text{--}94 \text{ GeV}$
−0.50166±0.00057		BARATE	00C ALEP	$E_{cm}^{ee} = 88\text{--}94 \text{ GeV}$
−0.4977 ±0.0045		<sup>4</sup> ABE	95J SLD	$E_{cm}^{ee} = 91.31 \text{ GeV}$



- <sup>1</sup> ACOSTA 05M determine the forward–backward asymmetry of  $e^+e^-$  pairs produced via  $q\bar{q} \rightarrow Z/\gamma^* \rightarrow e^+e^-$  in 15  $M(e^+e^-)$  effective mass bins ranging from 40 GeV to 600 GeV. These results are used to obtain the vector and axial–vector couplings of the  $Z$  to  $e^+e^-$ , assuming the quark couplings are as predicted by the standard model. Higher order radiative corrections have not been taken into account.
- <sup>2</sup> ABBIENDI 010 use their measurement of the  $\tau$  polarization in addition to the lineshape and forward-backward lepton asymmetries.
- <sup>3</sup> ACCIARRI 00C use their measurement of the  $\tau$  polarization in addition to forward-backward lepton asymmetries.
- <sup>4</sup> ABE 95J obtain this result combining polarized Bhabha results with the  $A_{LR}$  measurement of ABE 94C. The Bhabha results alone give  $-0.4968 \pm 0.0039 \pm 0.0027$ .

### $g_A^\mu$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>-0.50120 \pm 0.00054</math> OUR FIT</b>				
$-0.50117 \pm 0.00099$	182.8k	<sup>1</sup> ABBIENDI	010 OPAL	$E_{cm}^{ee} = 88-94$ GeV
$-0.5009 \pm 0.0014$	113.4k	<sup>2</sup> ACCIARRI	00C L3	$E_{cm}^{ee} = 88-94$ GeV
$-0.50046 \pm 0.00093$		BARATE	00C ALEP	$E_{cm}^{ee} = 88-94$ GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$-0.520 \pm 0.015$	66143	<sup>3</sup> ABBIENDI	01k OPAL	$E_{cm}^{ee} = 89-93$ GeV

- <sup>1</sup> ABBIENDI 010 use their measurement of the  $\tau$  polarization in addition to the lineshape and forward-backward lepton asymmetries.
- <sup>2</sup> ACCIARRI 00C use their measurement of the  $\tau$  polarization in addition to forward-backward lepton asymmetries.
- <sup>3</sup> ABBIENDI 01k obtain this from an angular analysis of the muon pair asymmetry which takes into account effects of initial state radiation on an event by event basis and of initial-final state interference.

### $g_A^\tau$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>-0.50204 \pm 0.00064</math> OUR FIT</b>				
$-0.50165 \pm 0.00124$	151.5k	<sup>1</sup> ABBIENDI	010 OPAL	$E_{cm}^{ee} = 88-94$ GeV
$-0.5023 \pm 0.0017$	103.0k	<sup>2</sup> ACCIARRI	00C L3	$E_{cm}^{ee} = 88-94$ GeV
$-0.50216 \pm 0.00100$		BARATE	00C ALEP	$E_{cm}^{ee} = 88-94$ GeV

- <sup>1</sup> ABBIENDI 010 use their measurement of the  $\tau$  polarization in addition to the lineshape and forward-backward lepton asymmetries.
- <sup>2</sup> ACCIARRI 00C use their measurement of the  $\tau$  polarization in addition to forward-backward lepton asymmetries.

### $g_A^l$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>-0.50123 \pm 0.00026</math> OUR FIT</b>				
$-0.50089 \pm 0.00045$	471.3k	<sup>1</sup> ABBIENDI	010 OPAL	$E_{cm}^{ee} = 88-94$ GeV
$-0.5007 \pm 0.0005$	379.4k	ABREU	00F DLPH	$E_{cm}^{ee} = 88-94$ GeV
$-0.50153 \pm 0.00053$	340.8k	<sup>2</sup> ACCIARRI	00C L3	$E_{cm}^{ee} = 88-94$ GeV
$-0.50150 \pm 0.00046$	500k	BARATE	00C ALEP	$E_{cm}^{ee} = 88-94$ GeV

- <sup>1</sup> ABBIENDI 010 use their measurement of the  $\tau$  polarization in addition to the lineshape and forward-backward lepton asymmetries.
- <sup>2</sup> ACCIARRI 00C use their measurement of the  $\tau$  polarization in addition to forward-backward lepton asymmetries.

### $g_A^u$

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
<b>0.519<sup>+0.028</sup><sub>-0.033</sub> OUR AVERAGE</b>				
0.548 ± 0.036		1 ANDREEV 18A	H1	$e^\pm p$
0.501 ± 0.110	156k	2 ABAZOV 11D	D0	$E_{cm}^{p\bar{p}} = 1.97$ TeV
0.47 <sup>+0.05</sup> <sub>-0.33</sub>		3 LEP-SLC 06		$E_{cm}^{ee} = 88-94$ GeV
0.441 <sup>+0.207</sup> <sub>-0.173</sub> ± 0.067	5026	4 ACOSTA 05M	CDF	$E_{cm}^{p\bar{p}} = 1.96$ TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.50 <sup>+0.12</sup> <sub>-0.05</sub>		5 ABRAMOWICZ16A	ZEUS	
0.532 <sup>+0.107</sup> <sub>-0.063</sub>		6 ABT 16		
0.57 ± 0.08	1500	7 AKTAS 06	H1	$e^\pm p \rightarrow \bar{\nu}_e(\nu_e)X$ , $\sqrt{s} \approx 300$ GeV

<sup>1</sup> ANDREEV 18A obtain this result in a combined electroweak and QCD analysis using all deep-inelastic  $e^+p$  and  $e^-p$  neutral current and charged current scattering cross sections published by the H1 Collaboration, including data with longitudinally polarized lepton beams.

<sup>2</sup> ABAZOV 11D study  $p\bar{p} \rightarrow Z/\gamma^* e^+ e^-$  events using  $5 \text{ fb}^{-1}$  data at  $\sqrt{s} = 1.96$  TeV. The candidate events are selected by requiring two isolated electromagnetic showers with  $E_T > 25$  GeV, at least one electron in the central region and the di-electron mass in the range 50–1000 GeV. From the forward-backward asymmetry, determined as a function of the di-electron mass, they derive the axial and vector couplings of the  $u$ - and  $d$ - quarks and the value of  $\sin^2\theta_{eff}^e = 0.2309 \pm 0.0008(\text{stat}) \pm 0.0006(\text{syst})$ .

<sup>3</sup> LEP-SLC 06 is a combination of the results from LEP and SLC experiments using light quark tagging.  $s$ - and  $d$ -quark couplings are assumed to be identical.

<sup>4</sup> ACOSTA 05M determine the forward-backward asymmetry of  $e^+ e^-$  pairs produced via  $q\bar{q} \rightarrow Z/\gamma^* \rightarrow e^+ e^-$  in 15  $M(e^+ e^-)$  effective mass bins ranging from 40 GeV to 600 GeV. These results are used to obtain the vector and axial-vector couplings of the  $Z$  to the light quarks, assuming the electron couplings are as predicted by the Standard Model. Higher order radiative corrections have not been taken into account.

<sup>5</sup> ABRAMOWICZ 16A determine the  $Z^0$  couplings to  $u$ - and  $d$ -quarks using the ZEUS polarised data from Run II together with the unpolarised data from both ZEUS and H1 Collaborations for Run I and unpolarised H1 data from Run II.

<sup>6</sup> ABT 16 determine the  $Z^0$  couplings to  $u$ - and  $d$ -quarks using the same techniques and data as ABRAMOWICZ 16A but additionally use the published H1 polarised data.

<sup>7</sup> AKTAS 06 fit the neutral current ( $1.5 \leq Q^2 \leq 30,000 \text{ GeV}^2$ ) and charged current ( $1.5 \leq Q^2 \leq 15,000 \text{ GeV}^2$ ) differential cross sections. In the determination of the  $u$ -quark couplings the electron and  $d$ -quark couplings are fixed to their standard model values.

### $g_A^d$

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
<b>-0.527<sup>+0.040</sup><sub>-0.028</sub> OUR AVERAGE</b>				
-0.619 ± 0.108		1 ANDREEV 18A	H1	$e^\pm p$
-0.497 ± 0.165	156k	2 ABAZOV 11D	D0	$E_{cm}^{p\bar{p}} = 1.97$ TeV
-0.52 <sup>+0.05</sup> <sub>-0.03</sub>		3 LEP-SLC 06		$E_{cm}^{ee} = 88-94$ GeV
-0.016 <sup>+0.346</sup> <sub>-0.536</sub> ± 0.091	5026	4 ACOSTA 05M	CDF	$E_{cm}^{p\bar{p}} = 1.96$ TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$-0.56^{+0.41}_{-0.15}$		<sup>5</sup> ABRAMOWICZ16A	ZEUS		
$-0.409^{+0.373}_{-0.213}$		<sup>6</sup> ABT	16		
$-0.80 \pm 0.24$	1500	<sup>7</sup> AKTAS	06	H1	$e^\pm p \rightarrow \bar{\nu}_e(\nu_e)X,$ $\sqrt{s} \approx 300 \text{ GeV}$

<sup>1</sup> ANDREEV 18A obtain this result in a combined electroweak and QCD analysis using all deep-inelastic  $e^+p$  and  $e^-p$  neutral current and charged current scattering cross sections published by the H1 Collaboration, including data with longitudinally polarized lepton beams.

<sup>2</sup> ABAZOV 11D study  $p\bar{p} \rightarrow Z/\gamma^* e^+ e^-$  events using  $5 \text{ fb}^{-1}$  data at  $\sqrt{s} = 1.96 \text{ TeV}$ . The candidate events are selected by requiring two isolated electromagnetic showers with  $E_T > 25 \text{ GeV}$ , at least one electron in the central region and the di-electron mass in the range 50–1000 GeV. From the forward-backward asymmetry, determined as a function of the di-electron mass, they derive the axial and vector couplings of the  $u$ - and  $d$ - quarks and the value of  $\sin^2\theta_{eff}^\ell = 0.2309 \pm 0.0008(\text{stat}) \pm 0.0006(\text{syst})$ .

<sup>3</sup> LEP-SLC 06 is a combination of the results from LEP and SLC experiments using light quark tagging.  $s$ - and  $d$ -quark couplings are assumed to be identical.

<sup>4</sup> ACOSTA 05M determine the forward-backward asymmetry of  $e^+ e^-$  pairs produced via  $q\bar{q} \rightarrow Z/\gamma^* \rightarrow e^+ e^-$  in 15  $M(e^+ e^-)$  effective mass bins ranging from 40 GeV to 600 GeV. These results are used to obtain the vector and axial-vector couplings of the  $Z$  to the light quarks, assuming the electron couplings are as predicted by the Standard Model. Higher order radiative corrections have not been taken into account.

<sup>5</sup> ABRAMOWICZ 16A determine the  $Z^0$  couplings to  $u$ - and  $d$ -quarks using the ZEUS polarised data from Run II together with the unpolarised data from both ZEUS and H1 Collaborations for Run I and unpolarised H1 data from Run II.

<sup>6</sup> ABT 16 determine the  $Z^0$  couplings to  $u$ - and  $d$ -quarks using the same techniques and data as ABRAMOWICZ 16A but additionally use the published H1 polarised data.

<sup>7</sup> AKTAS 06 fit the neutral current ( $1.5 \leq Q^2 \leq 30,000 \text{ GeV}^2$ ) and charged current ( $1.5 \leq Q^2 \leq 15,000 \text{ GeV}^2$ ) differential cross sections. In the determination of the  $d$ -quark couplings the electron and  $u$ -quark couplings are fixed to their standard model values.

## Z COUPLINGS TO NEUTRAL LEPTONS

Averaging over neutrino species, the invisible  $Z$  decay width determines the effective neutrino coupling  $g^{\nu\ell}$ . For  $g^{\nu e}$  and  $g^{\nu\mu}$ ,  $\nu_e e$  and  $\nu_\mu e$  scattering results are combined with  $g_A^e$  and  $g_V^e$  measurements at the  $Z$  mass to obtain  $g^{\nu e}$  and  $g^{\nu\mu}$  following NOVIKOV 93C.

### $g^{\nu\ell}$

VALUE	DOCUMENT ID	TECN	COMMENT
<b><math>0.50076 \pm 0.00076</math></b>	<sup>1</sup> LEP-SLC	06	$E_{cm}^{ee} = 88\text{--}94 \text{ GeV}$

<sup>1</sup> From invisible  $Z$ -decay width.

### $g^{\nu e}$

VALUE	DOCUMENT ID	TECN	COMMENT
<b><math>0.528 \pm 0.085</math></b>	<sup>1</sup> VILAIN	94	CHM2 From $\nu_\mu e$ and $\nu_e e$ scattering

<sup>1</sup> VILAIN 94 derive this value from their value of  $g^{\nu\mu}$  and their ratio  $g^{\nu e}/g^{\nu\mu} = 1.05^{+0.15}_{-0.18}$ .

### $g^{\nu\mu}$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.502±0.017</b>	<sup>1</sup> VILAIN	94	CHM2 From $\nu_\mu e$ scattering

<sup>1</sup> VILAIN 94 derive this value from their measurement of the couplings  $g_A^{e\nu\mu} = -0.503 \pm 0.017$  and  $g_V^{e\nu\mu} = -0.035 \pm 0.017$  obtained from  $\nu_\mu e$  scattering. We have re-evaluated this value using the current PDG values for  $g_A^e$  and  $g_V^e$ .

## Z ASYMMETRY PARAMETERS

For each fermion-antifermion pair coupling to the Z these quantities are defined as

$$A_f = \frac{2g_V^f g_A^f}{(g_V^f)^2 + (g_A^f)^2}$$

where  $g_V^f$  and  $g_A^f$  are the effective vector and axial-vector couplings. For their relation to the various lepton asymmetries see the note "The Z boson" and ref. LEP-SLC 06.

### $A_e$

Using polarized beams, this quantity can also be measured as  $(\sigma_L - \sigma_R) / (\sigma_L + \sigma_R)$ , where  $\sigma_L$  and  $\sigma_R$  are the  $e^+e^-$  production cross sections for Z bosons produced with left-handed and right-handed electrons respectively.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.1515±0.0019 OUR AVERAGE</b>				
0.1454±0.0108±0.0036	144810	<sup>1</sup> ABBIENDI	010	OPAL $E_{cm}^{ee} = 88-94$ GeV
0.1516±0.0021	559000	<sup>2</sup> ABE	01B	SLD $E_{cm}^{ee} = 91.24$ GeV
0.1504±0.0068±0.0008		<sup>3</sup> HEISTER	01	ALEP $E_{cm}^{ee} = 88-94$ GeV
0.1382±0.0116±0.0005	105000	<sup>4</sup> ABREU	00E	DLPH $E_{cm}^{ee} = 88-94$ GeV
0.1678±0.0127±0.0030	137092	<sup>5</sup> ACCIARRI	98H	L3 $E_{cm}^{ee} = 88-94$ GeV
0.162 ±0.041 ±0.014	89838	<sup>6</sup> ABE	97	SLD $E_{cm}^{ee} = 91.27$ GeV
0.202 ±0.038 ±0.008		<sup>7</sup> ABE	95J	SLD $E_{cm}^{ee} = 91.31$ GeV

<sup>1</sup> ABBIENDI 010 fit for  $A_e$  and  $A_\tau$  from measurements of the  $\tau$  polarization at varying  $\tau$  production angles. The correlation between  $A_e$  and  $A_\tau$  is less than 0.03.

<sup>2</sup> ABE 01B use the left-right production and left-right forward-backward decay asymmetries in leptonic Z decays to obtain a value of  $0.1544 \pm 0.0060$ . This is combined with left-right production asymmetry measurement using hadronic Z decays (ABE 00B) to obtain the quoted value.

<sup>3</sup> HEISTER 01 obtain this result fitting the  $\tau$  polarization as a function of the polar production angle of the  $\tau$ .

<sup>4</sup> ABREU 00E obtain this result fitting the  $\tau$  polarization as a function of the polar  $\tau$  production angle. This measurement is a combination of different analyses (exclusive  $\tau$  decay modes, inclusive hadronic 1-prong reconstruction, and a neural network analysis).

<sup>5</sup> Derived from the measurement of forward-backward  $\tau$  polarization asymmetry.

<sup>6</sup> ABE 97 obtain this result from a measurement of the observed left-right charge asymmetry,  $A_Q^{obs} = 0.225 \pm 0.056 \pm 0.019$ , in hadronic Z decays. If they combine this value of  $A_Q^{obs}$  with their earlier measurement of  $A_{LR}^{obs}$  they determine  $A_e$  to be  $0.1574 \pm 0.0197 \pm 0.0067$  independent of the beam polarization.

<sup>7</sup> ABE 95J obtain this result from polarized Bhabha scattering.

### $A_\mu$

This quantity is directly extracted from a measurement of the left-right forward-backward asymmetry in  $\mu^+\mu^-$  production at SLC using a polarized electron beam. This double asymmetry eliminates the dependence on the  $Z$ - $e$ - $e$  coupling parameter  $A_e$ .

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.142±0.015</b>	16844	<sup>1</sup> ABE	01B SLD	$E_{\text{cm}}^{ee} = 91.24$ GeV
••• We do not use the following data for averages, fits, limits, etc. •••				
0.153±0.012	1.7M	<sup>2</sup> AAD	15BT ATLS	$E_{\text{cm}}^{pp} = 7$ TeV

<sup>1</sup> ABE 01B obtain this direct measurement using the left-right production and left-right forward-backward polar angle asymmetries in  $\mu^+\mu^-$  decays of the  $Z$  boson obtained with a polarized electron beam.

<sup>2</sup> AAD 15BT study  $pp \rightarrow Z \rightarrow \ell^+\ell^-$  events where  $\ell$  is an electron or a muon in the dilepton mass region 70–1000 GeV. The background in the  $Z$  peak region is estimated to be < 1% for the muon channel. The muon asymmetry parameter is derived from the measured forward-backward asymmetry assuming the value of the quark asymmetry parameter from the SM. For this reason it is not used in the average.

### $A_\tau$

The LEP Collaborations derive this quantity from the measurement of the  $\tau$  polarization in  $Z \rightarrow \tau^+\tau^-$ . The SLD Collaboration directly extracts this quantity from its measured left-right forward-backward asymmetry in  $Z \rightarrow \tau^+\tau^-$  produced using a polarized  $e^-$  beam. This double asymmetry eliminates the dependence on the  $Z$ - $e$ - $e$  coupling parameter  $A_e$ .

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.143 ±0.004 OUR AVERAGE</b>				
0.1456±0.0076±0.0057	144810	<sup>1</sup> ABBIENDI	010 OPAL	$E_{\text{cm}}^{ee} = 88$ –94 GeV
0.136 ±0.015	16083	<sup>2</sup> ABE	01B SLD	$E_{\text{cm}}^{ee} = 91.24$ GeV
0.1451±0.0052±0.0029		<sup>3</sup> HEISTER	01 ALEP	$E_{\text{cm}}^{ee} = 88$ –94 GeV
0.1359±0.0079±0.0055	105000	<sup>4</sup> ABREU	00E DLPH	$E_{\text{cm}}^{ee} = 88$ –94 GeV
0.1476±0.0088±0.0062	137092	ACCIARRI	98H L3	$E_{\text{cm}}^{ee} = 88$ –94 GeV

<sup>1</sup> ABBIENDI 010 fit for  $A_e$  and  $A_\tau$  from measurements of the  $\tau$  polarization at varying  $\tau$  production angles. The correlation between  $A_e$  and  $A_\tau$  is less than 0.03.

<sup>2</sup> ABE 01B obtain this direct measurement using the left-right production and left-right forward-backward polar angle asymmetries in  $\tau^+\tau^-$  decays of the  $Z$  boson obtained with a polarized electron beam.

<sup>3</sup> HEISTER 01 obtain this result fitting the  $\tau$  polarization as a function of the polar production angle of the  $\tau$ .

<sup>4</sup> ABREU 00E obtain this result fitting the  $\tau$  polarization as a function of the polar  $\tau$  production angle. This measurement is a combination of different analyses (exclusive  $\tau$  decay modes, inclusive hadronic 1-prong reconstruction, and a neural network analysis).

### $A_s$

The SLD Collaboration directly extracts this quantity by a simultaneous fit to four measured  $s$ -quark polar angle distributions corresponding to two states of  $e^-$  polarization (positive and negative) and to the  $K^+K^-$  and  $K^\pm K_S^0$  strange particle tagging modes in the hadronic final states.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.895±0.066±0.062</b>	2870	<sup>1</sup> ABE	00D SLD	$E_{\text{cm}}^{ee} = 91.2$ GeV

<sup>1</sup> ABE 00D tag  $Z \rightarrow s\bar{s}$  events by an absence of  $B$  or  $D$  hadrons and the presence in each hemisphere of a high momentum  $K^\pm$  or  $K_S^0$ .

## $A_c$

This quantity is directly extracted from a measurement of the left-right forward-backward asymmetry in  $c\bar{c}$  production at SLC using polarized electron beam. This double asymmetry eliminates the dependence on the  $Z$ - $e$ - $e$  coupling parameter  $A_e$ . OUR FIT is obtained by a simultaneous fit to several  $c$ - and  $b$ -quark measurements as explained in the note “The  $Z$  boson” and ref. LEP-SLC 06.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.670 ± 0.027 OUR FIT</b>			
0.6712 ± 0.0224 ± 0.0157	<sup>1</sup> ABE	05 SLD	$E_{\text{cm}}^{ee} = 91.24$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.583 ± 0.055 ± 0.055	<sup>2</sup> ABE	02G SLD	$E_{\text{cm}}^{ee} = 91.24$ GeV
0.688 ± 0.041	<sup>3</sup> ABE	01C SLD	$E_{\text{cm}}^{ee} = 91.25$ GeV

<sup>1</sup> ABE 05 use hadronic  $Z$  decays collected during 1996–98 to obtain an enriched sample of  $c\bar{c}$  events tagging on the invariant mass of reconstructed secondary decay vertices. The charge of the underlying  $c$ -quark is obtained with an algorithm that takes into account the net charge of the vertex as well as the charge of tracks emanating from the vertex and identified as kaons. This yields (9970 events)  $A_c = 0.6747 \pm 0.0290 \pm 0.0233$ . Taking into account all correlations with earlier results reported in ABE 02G and ABE 01C, they obtain the quoted overall SLD result.

<sup>2</sup> ABE 02G tag  $b$  and  $c$  quarks through their semileptonic decays into electrons and muons. A maximum likelihood fit is performed to extract simultaneously  $A_b$  and  $A_c$ .

<sup>3</sup> ABE 01C tag  $Z \rightarrow c\bar{c}$  events using two techniques: exclusive reconstruction of  $D^{*+}$ ,  $D^+$  and  $D^0$  mesons and the soft pion tag for  $D^{*+} \rightarrow D^0 \pi^+$ . The large background from  $D$  mesons produced in  $b\bar{b}$  events is separated efficiently from the signal using precision vertex information. When combining the  $A_c$  values from these two samples, care is taken to avoid double counting of events common to the two samples, and common systematic errors are properly taken into account.

## $A_b$

This quantity is directly extracted from a measurement of the left-right forward-backward asymmetry in  $b\bar{b}$  production at SLC using polarized electron beam. This double asymmetry eliminates the dependence on the  $Z$ - $e$ - $e$  coupling parameter  $A_e$ . OUR FIT is obtained by a simultaneous fit to several  $c$ - and  $b$ -quark measurements as explained in the note “The  $Z$  boson” and ref. LEP-SLC 06.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.923 ± 0.020 OUR FIT</b>				
0.9170 ± 0.0147 ± 0.0145		<sup>1</sup> ABE	05 SLD	$E_{\text{cm}}^{ee} = 91.24$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.907 ± 0.020 ± 0.024	48028	<sup>2</sup> ABE	03F SLD	$E_{\text{cm}}^{ee} = 91.24$ GeV
0.919 ± 0.030 ± 0.024		<sup>3</sup> ABE	02G SLD	$E_{\text{cm}}^{ee} = 91.24$ GeV
0.855 ± 0.088 ± 0.102	7473	<sup>4</sup> ABE	99L SLD	$E_{\text{cm}}^{ee} = 91.27$ GeV

<sup>1</sup> ABE 05 use hadronic  $Z$  decays collected during 1996–98 to obtain an enriched sample of  $b\bar{b}$  events tagging on the invariant mass of reconstructed secondary decay vertices. The charge of the underlying  $b$ -quark is obtained with an algorithm that takes into account the net charge of the vertex as well as the charge of tracks emanating from the vertex and identified as kaons. This yields (25917 events)  $A_b = 0.9173 \pm 0.0184 \pm 0.0173$ . Taking into account all correlations with earlier results reported in ABE 03F, ABE 02G and ABE 99L, they obtain the quoted overall SLD result.

<sup>2</sup> ABE 03F obtain an enriched sample of  $b\bar{b}$  events tagging on the invariant mass of a 3-dimensional topologically reconstructed secondary decay. The charge of the underlying  $b$  quark is obtained using a self-calibrating track-charge method. For the 1996–1998 data sample they measure  $A_b = 0.906 \pm 0.022 \pm 0.023$ . The value quoted here is obtained combining the above with the result of ABE 98I (1993–1995 data sample).

- <sup>3</sup> ABE 02G tag  $b$  and  $c$  quarks through their semileptonic decays into electrons and muons. A maximum likelihood fit is performed to extract simultaneously  $A_b$  and  $A_c$ .
- <sup>4</sup> ABE 99L obtain an enriched sample of  $b\bar{b}$  events tagging with an inclusive vertex mass cut. For distinguishing  $b$  and  $\bar{b}$  quarks they use the charge of identified  $K^\pm$ .

## TRANSVERSE SPIN CORRELATIONS IN $Z \rightarrow \tau^+ \tau^-$

The correlations between the transverse spin components of  $\tau^+ \tau^-$  produced in  $Z$  decays may be expressed in terms of the vector and axial-vector couplings:

$$C_{TT} = \frac{|g_A^\tau|^2 - |g_V^\tau|^2}{|g_A^\tau|^2 + |g_V^\tau|^2}$$

$$C_{TN} = -2 \frac{|g_A^\tau| |g_V^\tau|}{|g_A^\tau|^2 + |g_V^\tau|^2} \sin(\Phi_{g_V^\tau} - \Phi_{g_A^\tau})$$

$C_{TT}$  refers to the transverse-transverse (within the collision plane) spin correlation and  $C_{TN}$  refers to the transverse-normal (to the collision plane) spin correlation.

The longitudinal  $\tau$  polarization  $P_\tau (= -A_\tau)$  is given by:

$$P_\tau = -2 \frac{|g_A^\tau| |g_V^\tau|}{|g_A^\tau|^2 + |g_V^\tau|^2} \cos(\Phi_{g_V^\tau} - \Phi_{g_A^\tau})$$

Here  $\Phi$  is the phase and the phase difference  $\Phi_{g_V^\tau} - \Phi_{g_A^\tau}$  can be obtained using both the measurements of  $C_{TN}$  and  $P_\tau$ .

### $C_{TT}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.01 ± 0.12 OUR AVERAGE</b>				
0.87 ± 0.20 <sup>+0.10</sup> <sub>-0.12</sub>	9.1k	ABREU	97G DLPH	$E_{\text{cm}}^{ee} = 91.2$ GeV
1.06 ± 0.13 ± 0.05	120k	BARATE	97D ALEP	$E_{\text{cm}}^{ee} = 91.2$ GeV

### $C_{TN}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.08 ± 0.13 ± 0.04</b>	120k	<sup>1</sup> BARATE	97D ALEP	$E_{\text{cm}}^{ee} = 91.2$ GeV

<sup>1</sup> BARATE 97D combine their value of  $C_{TN}$  with the world average  $P_\tau = -0.140 \pm 0.007$  to obtain  $\tan(\Phi_{g_V^\tau} - \Phi_{g_A^\tau}) = -0.57 \pm 0.97$ .

## FORWARD-BACKWARD $e^+ e^- \rightarrow f\bar{f}$ CHARGE ASYMMETRIES

These asymmetries are experimentally determined by tagging the respective lepton or quark flavor in  $e^+ e^-$  interactions. Details of heavy flavor ( $c$ - or  $b$ -quark) tagging at LEP are described in the note on "The Z boson" and ref. LEP-SLC 06. The Standard Model predictions for LEP data have been (re)computed using the ZFITTER package (version 6.36) with input parameters  $M_Z = 91.187$  GeV,  $M_{\text{top}} = 174.3$  GeV,  $M_{\text{Higgs}} = 150$  GeV,  $\alpha_s = 0.119$ ,  $\alpha^{(5)}(M_Z) = 1/128.877$  and the Fermi constant  $G_F = 1.16637 \times 10^{-5}$  GeV<sup>-2</sup> (see the note on "The Z boson" for references).

For non-LEP data the Standard Model predictions are as given by the authors of the respective publications.

—————  $A_{FB}^{(0,e)}$  CHARGE ASYMMETRY IN  $e^+e^- \rightarrow e^+e^-$  —————

OUR FIT is obtained using the fit procedure and correlations as determined by the LEP Electroweak Working Group (see the note “The Z boson” and ref. LEP-SLC 06). For the Z peak, we report the pole asymmetry defined by  $(3/4)A_e^2$  as determined by the nine-parameter fit to cross-section and lepton forward-backward asymmetry data.

<u>ASYMMETRY (%)</u>	<u>STD. MODEL</u>	<u><math>\sqrt{s}</math> (GeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
<b>1.45 ± 0.25 OUR FIT</b>				
0.89 ± 0.44	1.57	91.2	<sup>1</sup> ABBIENDI	01A OPAL
1.71 ± 0.49	1.57	91.2	ABREU	00F DLPH
1.06 ± 0.58	1.57	91.2	ACCIARRI	00C L3
1.88 ± 0.34	1.57	91.2	<sup>2</sup> BARATE	00C ALEP

<sup>1</sup> ABBIENDI 01A error includes approximately 0.38 due to statistics, 0.16 due to event selection systematics, and 0.18 due to the theoretical uncertainty in *t*-channel prediction.

<sup>2</sup> BARATE 00C error includes approximately 0.31 due to statistics, 0.06 due to experimental systematics, and 0.13 due to the theoretical uncertainty in *t*-channel prediction.

—————  $A_{FB}^{(0,\mu)}$  CHARGE ASYMMETRY IN  $e^+e^- \rightarrow \mu^+\mu^-$  —————

OUR FIT is obtained using the fit procedure and correlations as determined by the LEP Electroweak Working Group (see the note “The Z boson” and ref. LEP-SLC 06). For the Z peak, we report the pole asymmetry defined by  $(3/4)A_e A_\mu$  as determined by the nine-parameter fit to cross-section and lepton forward-backward asymmetry data.

<u>ASYMMETRY (%)</u>	<u>STD. MODEL</u>	<u><math>\sqrt{s}</math> (GeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
<b>1.69 ± 0.13 OUR FIT</b>				
1.59 ± 0.23	1.57	91.2	<sup>1</sup> ABBIENDI	01A OPAL
1.65 ± 0.25	1.57	91.2	ABREU	00F DLPH
1.88 ± 0.33	1.57	91.2	ACCIARRI	00C L3
1.71 ± 0.24	1.57	91.2	<sup>2</sup> BARATE	00C ALEP

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

9 ± 30	−1.3	20	<sup>3</sup> ABREU	95M DLPH
7 ± 26	−8.3	40	<sup>3</sup> ABREU	95M DLPH
−11 ± 33	−24.1	57	<sup>3</sup> ABREU	95M DLPH
−62 ± 17	−44.6	69	<sup>3</sup> ABREU	95M DLPH
−56 ± 10	−63.5	79	<sup>3</sup> ABREU	95M DLPH
−13 ± 5	−34.4	87.5	<sup>3</sup> ABREU	95M DLPH
−29.0 <sup>+</sup> 5.0 <sub>− 4.8</sub> ± 0.5	−32.1	56.9	<sup>4</sup> ABE	90I VNS
− 9.9 ± 1.5 ± 0.5	−9.2	35	HEGNER	90 JADE
0.05 ± 0.22	0.026	91.14	<sup>5</sup> ABRAMS	89D MRK2
−43.4 ± 17.0	−24.9	52.0	<sup>6</sup> BACALA	89 AMY
−11.0 ± 16.5	−29.4	55.0	<sup>6</sup> BACALA	89 AMY
−30.0 ± 12.4	−31.2	56.0	<sup>6</sup> BACALA	89 AMY



-46.2 ± 14.9	-33.0	57.0	<sup>6</sup> BACALA	89	AMY
-29 ± 13	-25.9	53.3	ADACHI	88C	TOPZ
+ 5.3 ± 5.0 ± 0.5	-1.2	14.0	ADEVA	88	MRKJ
-10.4 ± 1.3 ± 0.5	-8.6	34.8	ADEVA	88	MRKJ
-12.3 ± 5.3 ± 0.5	-10.7	38.3	ADEVA	88	MRKJ
-15.6 ± 3.0 ± 0.5	-14.9	43.8	ADEVA	88	MRKJ
- 1.0 ± 6.0	-1.2	13.9	BRAUNSCH...	88D	TASS
- 9.1 ± 2.3 ± 0.5	-8.6	34.5	BRAUNSCH...	88D	TASS
-10.6 + - 2.2 - 2.3 ± 0.5	-8.9	35.0	BRAUNSCH...	88D	TASS
-17.6 + - 4.4 - 4.3 ± 0.5	-15.2	43.6	BRAUNSCH...	88D	TASS
- 4.8 ± 6.5 ± 1.0	-11.5	39	BEHREND	87C	CELL
-18.8 ± 4.5 ± 1.0	-15.5	44	BEHREND	87C	CELL
+ 2.7 ± 4.9	-1.2	13.9	BARTEL	86C	JADE
-11.1 ± 1.8 ± 1.0	-8.6	34.4	BARTEL	86C	JADE
-17.3 ± 4.8 ± 1.0	-13.7	41.5	BARTEL	86C	JADE
-22.8 ± 5.1 ± 1.0	-16.6	44.8	BARTEL	86C	JADE
- 6.3 ± 0.8 ± 0.2	-6.3	29	ASH	85	MAC
- 4.9 ± 1.5 ± 0.5	-5.9	29	DERRICK	85	HRS
- 7.1 ± 1.7	-5.7	29	LEVI	83	MRK2
-16.1 ± 3.2	-9.2	34.2	BRANDELIK	82C	TASS

<sup>1</sup> ABBIENDI 01A error is almost entirely on account of statistics.

<sup>2</sup> BARATE 00C error is almost entirely on account of statistics.

<sup>3</sup> ABREU 95M perform this measurement using radiative muon-pair events associated with high-energy isolated photons.

<sup>4</sup> ABE 90I measurements in the range  $50 \leq \sqrt{s} \leq 60.8$  GeV.

<sup>5</sup> ABRAMS 89D asymmetry includes both  $9 \mu^+ \mu^-$  and  $15 \tau^+ \tau^-$  events.

<sup>6</sup> BACALA 89 systematic error is about 5%.

## ————— $A_{FB}^{(0,\tau)}$ CHARGE ASYMMETRY IN $e^+ e^- \rightarrow \tau^+ \tau^-$ —————

OUR FIT is obtained using the fit procedure and correlations as determined by the LEP Electroweak Working Group (see the note “The Z boson” and ref. LEP-SLC 06). For the Z peak, we report the pole asymmetry defined by  $(3/4)A_e A_\tau$  as determined by the nine-parameter fit to cross-section and lepton forward-backward asymmetry data.

ASYMMETRY (%)	STD. MODEL	$\sqrt{s}$ (GeV)	DOCUMENT ID	TECN
<b>1.88 ± 0.17 OUR FIT</b>				
1.45 ± 0.30	1.57	91.2	<sup>1</sup> ABBIENDI	01A OPAL
2.41 ± 0.37	1.57	91.2	ABREU	00F DLPH
2.60 ± 0.47	1.57	91.2	ACCIARRI	00C L3
1.70 ± 0.28	1.57	91.2	<sup>2</sup> BARATE	00C ALEP
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
-32.8 + - 6.4 - 6.2 ± 1.5	-32.1	56.9	<sup>3</sup> ABE	90I VNS
- 8.1 ± 2.0 ± 0.6	-9.2	35	HEGNER	90 JADE
-18.4 ± 19.2	-24.9	52.0	<sup>4</sup> BACALA	89 AMY
-17.7 ± 26.1	-29.4	55.0	<sup>4</sup> BACALA	89 AMY
-45.9 ± 16.6	-31.2	56.0	<sup>4</sup> BACALA	89 AMY
-49.5 ± 18.0	-33.0	57.0	<sup>4</sup> BACALA	89 AMY

-20 ±14	-25.9	53.3	ADACHI	88C	TOPZ
-10.6 ± 3.1 ±1.5	-8.5	34.7	ADEVA	88	MRKJ
- 8.5 ± 6.6 ±1.5	-15.4	43.8	ADEVA	88	MRKJ
- 6.0 ± 2.5 ±1.0	8.8	34.6	BARTEL	85F	JADE
-11.8 ± 4.6 ±1.0	14.8	43.0	BARTEL	85F	JADE
- 5.5 ± 1.2 ±0.5	-0.063	29.0	FERNANDEZ	85A	MAC
- 4.2 ± 2.0	0.057	29	LEVI	83	MRK2
-10.3 ± 5.2	-9.2	34.2	BEHREND	82	CELL
- 0.4 ± 6.6	-9.1	34.2	BRANDELIK	82C	TASS

<sup>1</sup> ABBIENDI 01A error includes approximately 0.26 due to statistics and 0.14 due to event selection systematics.

<sup>2</sup> BARATE 00C error includes approximately 0.26 due to statistics and 0.11 due to experimental systematics.

<sup>3</sup> ABE 90I measurements in the range  $50 \leq \sqrt{s} \leq 60.8$  GeV.

<sup>4</sup> BACALA 89 systematic error is about 5%.

### ———— $A_{FB}^{(0,\ell)}$ CHARGE ASYMMETRY IN $e^+e^- \rightarrow \ell^+\ell^-$ ————

For the  $Z$  peak, we report the pole asymmetry defined by  $(3/4)A_\ell^2$  as determined by the five-parameter fit to cross-section and lepton forward-backward asymmetry data assuming lepton universality. For details see the note “The  $Z$  boson” and ref. LEP-SLC 06.

<u>ASYMMETRY (%)</u>	<u>STD. MODEL</u>	<u><math>\sqrt{s}</math> (GeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
<b>1.71±0.10 OUR FIT</b>				
1.45±0.17	1.57	91.2	<sup>1</sup> ABBIENDI 01A	OPAL
1.87±0.19	1.57	91.2	ABREU 00F	DLPH
1.92±0.24	1.57	91.2	ACCIARRI 00C	L3
1.73±0.16	1.57	91.2	<sup>2</sup> BARATE 00C	ALEP

<sup>1</sup> ABBIENDI 01A error includes approximately 0.15 due to statistics, 0.06 due to event selection systematics, and 0.03 due to the theoretical uncertainty in  $t$ -channel prediction.

<sup>2</sup> BARATE 00C error includes approximately 0.15 due to statistics, 0.04 due to experimental systematics, and 0.02 due to the theoretical uncertainty in  $t$ -channel prediction.

### ———— $A_{FB}^{(0,u)}$ CHARGE ASYMMETRY IN $e^+e^- \rightarrow u\bar{u}$ ————

<u>ASYMMETRY (%)</u>	<u>STD. MODEL</u>	<u><math>\sqrt{s}</math> (GeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
<b>4.0±6.7±2.8</b>	<b>7.2</b>	<b>91.2</b>	<sup>1</sup> ACKERSTAFF 97T	OPAL

<sup>1</sup> ACKERSTAFF 97T measure the forward-backward asymmetry of various fast hadrons made of light quarks. Then using SU(2) isospin symmetry and flavor independence for down and strange quarks authors solve for the different quark types.

### ———— $A_{FB}^{(0,s)}$ CHARGE ASYMMETRY IN $e^+e^- \rightarrow s\bar{s}$ ————

The  $s$ -quark asymmetry is derived from measurements of the forward-backward asymmetry of fast hadrons containing an  $s$  quark.

<u>ASYMMETRY (%)</u>	<u>STD. MODEL</u>	<u><math>\sqrt{s}</math> (GeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
<b>9.8 ± 1.1 OUR AVERAGE</b>				
10.08 ± 1.13 ± 0.40	10.1	91.2	<sup>1</sup> ABREU 00B	DLPH
6.8 ± 3.5 ± 1.1	10.1	91.2	<sup>2</sup> ACKERSTAFF 97T	OPAL

<sup>1</sup> ABREU 00B tag the presence of an *s* quark requiring a high-momentum-identified charged kaon. The *s*-quark pole asymmetry is extracted from the charged-kaon asymmetry taking the expected *d*- and *u*-quark asymmetries from the Standard Model and using the measured values for the *c*- and *b*-quark asymmetries.

<sup>2</sup> ACKERSTAFF 97T measure the forward-backward asymmetry of various fast hadrons made of light quarks. Then using SU(2) isospin symmetry and flavor independence for down and strange quarks authors solve for the different quark types. The value reported here corresponds then to the forward-backward asymmetry for “down-type” quarks.

————  $A_{FB}^{(0,c)}$  CHARGE ASYMMETRY IN  $e^+ e^- \rightarrow c \bar{c}$  ————

OUR FIT, which is obtained by a simultaneous fit to several *c*- and *b*-quark measurements as explained in the note “The Z boson” and ref. LEP-SLC 06, refers to the **Z pole** asymmetry. The experimental values, on the other hand, correspond to the measurements carried out at the respective energies.

<u>ASYMMETRY (%)</u>	<u>STD. MODEL</u>	<u><math>\sqrt{s}</math> (GeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
<b>7.07 ± 0.35 OUR FIT</b>				
6.31 ± 0.93 ± 0.65	6.35	91.26	<sup>1</sup> ABDALLAH 04F	DLPH
5.68 ± 0.54 ± 0.39	6.3	91.25	<sup>2</sup> ABBIENDI 03P	OPAL
6.45 ± 0.57 ± 0.37	6.10	91.21	<sup>3</sup> HEISTER 02H	ALEP
6.59 ± 0.94 ± 0.35	6.2	91.235	<sup>4</sup> ABREU 99Y	DLPH
6.3 ± 0.9 ± 0.3	6.1	91.22	<sup>5</sup> BARATE 98O	ALEP
6.3 ± 1.2 ± 0.6	6.1	91.22	<sup>6</sup> ALEXANDER 97C	OPAL
8.3 ± 3.8 ± 2.7	6.2	91.24	<sup>7</sup> ADRIANI 92D	L3

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.1 ± 3.5 ± 0.5	−3.5	89.43	<sup>1</sup> ABDALLAH 04F	DLPH
11.0 ± 2.8 ± 0.7	12.3	92.99	<sup>1</sup> ABDALLAH 04F	DLPH
− 6.8 ± 2.5 ± 0.9	−3.0	89.51	<sup>2</sup> ABBIENDI 03P	OPAL
14.6 ± 2.0 ± 0.8	12.2	92.95	<sup>2</sup> ABBIENDI 03P	OPAL
− 12.4 ± 15.9 ± 2.0	−9.6	88.38	<sup>3</sup> HEISTER 02H	ALEP
− 2.3 ± 2.6 ± 0.2	−3.8	89.38	<sup>3</sup> HEISTER 02H	ALEP
− 0.3 ± 8.3 ± 0.6	0.9	90.21	<sup>3</sup> HEISTER 02H	ALEP
10.6 ± 7.7 ± 0.7	9.6	92.05	<sup>3</sup> HEISTER 02H	ALEP
11.9 ± 2.1 ± 0.6	12.2	92.94	<sup>3</sup> HEISTER 02H	ALEP
12.1 ± 11.0 ± 1.0	14.2	93.90	<sup>3</sup> HEISTER 02H	ALEP
− 4.96 ± 3.68 ± 0.53	−3.5	89.434	<sup>4</sup> ABREU 99Y	DLPH
11.80 ± 3.18 ± 0.62	12.3	92.990	<sup>4</sup> ABREU 99Y	DLPH
− 1.0 ± 4.3 ± 1.0	−3.9	89.37	<sup>5</sup> BARATE 98O	ALEP
11.0 ± 3.3 ± 0.8	12.3	92.96	<sup>5</sup> BARATE 98O	ALEP
3.9 ± 5.1 ± 0.9	−3.4	89.45	<sup>6</sup> ALEXANDER 97C	OPAL
15.8 ± 4.1 ± 1.1	12.4	93.00	<sup>6</sup> ALEXANDER 97C	OPAL

-12.9 ± 7.8 ± 5.5	-13.6	35	BEHREND	90D	CELL
7.7 ± 13.4 ± 5.0	-22.1	43	BEHREND	90D	CELL
-12.8 ± 4.4 ± 4.1	-13.6	35	ELSEN	90	JADE
-10.9 ± 12.9 ± 4.6	-23.2	44	ELSEN	90	JADE
-14.9 ± 6.7	-13.3	35	OULD-SAADA	89	JADE

<sup>1</sup> ABDALLAH 04F tag  $b^-$  and  $c^-$  quarks using semileptonic decays combined with charge flow information from the hemisphere opposite to the lepton. Enriched samples of  $c\bar{c}$  and  $b\bar{b}$  events are obtained using lifetime information.

<sup>2</sup> ABBIENDI 03P tag heavy flavors using events with one or two identified leptons. This allows the simultaneous fitting of the  $b$  and  $c$  quark forward-backward asymmetries as well as the average  $B^0-\bar{B}^0$  mixing.

<sup>3</sup> HEISTER 02H measure simultaneously  $b$  and  $c$  quark forward-backward asymmetries using their semileptonic decays to tag the quark charge. The flavor separation is obtained with a discriminating multivariate analysis.

<sup>4</sup> ABREU 99Y tag  $Z \rightarrow b\bar{b}$  and  $Z \rightarrow c\bar{c}$  events by an exclusive reconstruction of several  $D$  meson decay modes ( $D^{*+}$ ,  $D^0$ , and  $D^+$  with their charge-conjugate states).

<sup>5</sup> BARATE 980 tag  $Z \rightarrow c\bar{c}$  events requiring the presence of high-momentum reconstructed  $D^{*+}$ ,  $D^+$ , or  $D^0$  mesons.

<sup>6</sup> ALEXANDER 97C identify the  $b$  and  $c$  events using a  $D/D^*$  tag.

<sup>7</sup> ADRIANI 92D use both electron and muon semileptonic decays.

—————  $A_{FB}^{(0,b)}$  CHARGE ASYMMETRY IN  $e^+e^- \rightarrow b\bar{b}$  —————

OUR FIT, which is obtained by a simultaneous fit to several  $c^-$  and  $b^-$  quark measurements as explained in the note “The  $Z$  boson” and ref. LEP-SLC 06, refers to the **Z pole** asymmetry. The experimental values, on the other hand, correspond to the measurements carried out at the respective energies.

ASYMMETRY (%)	STD. MODEL	$\sqrt{s}$ (GeV)	DOCUMENT ID	TECN
<b>9.92 ± 0.16 OUR FIT</b>				
9.58 ± 0.32 ± 0.14	9.68	91.231	<sup>1</sup> ABDALLAH 05	DLPH
10.04 ± 0.56 ± 0.25	9.69	91.26	<sup>2</sup> ABDALLAH 04F	DLPH
9.72 ± 0.42 ± 0.15	9.67	91.25	<sup>3</sup> ABBIENDI 03P	OPAL
9.77 ± 0.36 ± 0.18	9.69	91.26	<sup>4</sup> ABBIENDI 02I	OPAL
9.52 ± 0.41 ± 0.17	9.59	91.21	<sup>5</sup> HEISTER 02H	ALEP
10.00 ± 0.27 ± 0.11	9.63	91.232	<sup>6</sup> HEISTER 01D	ALEP
7.62 ± 1.94 ± 0.85	9.64	91.235	<sup>7</sup> ABREU 99Y	DLPH
9.60 ± 0.66 ± 0.33	9.69	91.26	<sup>8</sup> ACCIARRI 99D	L3
9.31 ± 1.01 ± 0.55	9.65	91.24	<sup>9</sup> ACCIARRI 98U	L3
9.4 ± 2.7 ± 2.2	9.61	91.22	<sup>10</sup> ALEXANDER 97C	OPAL

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

6.37 ± 1.43 ± 0.17	5.8	89.449	<sup>1</sup> ABDALLAH 05	DLPH
10.41 ± 1.15 ± 0.24	12.1	92.990	<sup>1</sup> ABDALLAH 05	DLPH
6.7 ± 2.2 ± 0.2	5.7	89.43	<sup>2</sup> ABDALLAH 04F	DLPH
11.2 ± 1.8 ± 0.2	12.1	92.99	<sup>2</sup> ABDALLAH 04F	DLPH
4.7 ± 1.8 ± 0.1	5.9	89.51	<sup>3</sup> ABBIENDI 03P	OPAL
10.3 ± 1.5 ± 0.2	12.0	92.95	<sup>3</sup> ABBIENDI 03P	OPAL
5.82 ± 1.53 ± 0.12	5.9	89.50	<sup>4</sup> ABBIENDI 02I	OPAL
12.21 ± 1.23 ± 0.25	12.0	92.91	<sup>4</sup> ABBIENDI 02I	OPAL
-13.1 ± 13.5 ± 1.0	3.2	88.38	<sup>5</sup> HEISTER 02H	ALEP

$5.5 \pm 1.9 \pm 0.1$	5.6	89.38	<sup>5</sup> HEISTER	02H	ALEP
$-0.4 \pm 6.7 \pm 0.8$	7.5	90.21	<sup>5</sup> HEISTER	02H	ALEP
$11.1 \pm 6.4 \pm 0.5$	11.0	92.05	<sup>5</sup> HEISTER	02H	ALEP
$10.4 \pm 1.5 \pm 0.3$	12.0	92.94	<sup>5</sup> HEISTER	02H	ALEP
$13.8 \pm 9.3 \pm 1.1$	12.9	93.90	<sup>5</sup> HEISTER	02H	ALEP
$4.36 \pm 1.19 \pm 0.11$	5.8	89.472	<sup>6</sup> HEISTER	01D	ALEP
$11.72 \pm 0.97 \pm 0.11$	12.0	92.950	<sup>6</sup> HEISTER	01D	ALEP
$5.67 \pm 7.56 \pm 1.17$	5.7	89.434	<sup>7</sup> ABREU	99Y	DLPH
$8.82 \pm 6.33 \pm 1.22$	12.1	92.990	<sup>7</sup> ABREU	99Y	DLPH
$6.11 \pm 2.93 \pm 0.43$	5.9	89.50	<sup>8</sup> ACCIARRI	99D	L3
$13.71 \pm 2.40 \pm 0.44$	12.2	93.10	<sup>8</sup> ACCIARRI	99D	L3
$4.95 \pm 5.23 \pm 0.40$	5.8	89.45	<sup>9</sup> ACCIARRI	98U	L3
$11.37 \pm 3.99 \pm 0.65$	12.1	92.99	<sup>9</sup> ACCIARRI	98U	L3
$-8.6 \pm 10.8 \pm 2.9$	5.8	89.45	<sup>10</sup> ALEXANDER	97C	OPAL
$-2.1 \pm 9.0 \pm 2.6$	12.1	93.00	<sup>10</sup> ALEXANDER	97C	OPAL
$-71 \pm 34 \pm 7$	-58	58.3	SHIMONAKA	91	TOPZ
$-22.2 \pm 7.7 \pm 3.5$	-26.0	35	BEHREND	90D	CELL
$-49.1 \pm 16.0 \pm 5.0$	-39.7	43	BEHREND	90D	CELL
$-28 \pm 11$	-23	35	BRAUNSCH...	90	TASS
$-16.6 \pm 7.7 \pm 4.8$	-24.3	35	ELSEN	90	JADE
$-33.6 \pm 22.2 \pm 5.2$	-39.9	44	ELSEN	90	JADE
$3.4 \pm 7.0 \pm 3.5$	-16.0	29.0	BAND	89	MAC
$-72 \pm 28 \pm 13$	-56	55.2	SAGAWA	89	AMY

<sup>1</sup> ABDALLAH 05 obtain an enriched samples of  $b\bar{b}$  events using lifetime information. The quark (or antiquark) charge is determined with a neural network using the secondary vertex charge, the jet charge and particle identification.

<sup>2</sup> ABDALLAH 04F tag  $b$ - and  $c$ -quarks using semileptonic decays combined with charge flow information from the hemisphere opposite to the lepton. Enriched samples of  $c\bar{c}$  and  $b\bar{b}$  events are obtained using lifetime information.

<sup>3</sup> ABBIENDI 03P tag heavy flavors using events with one or two identified leptons. This allows the simultaneous fitting of the  $b$  and  $c$  quark forward-backward asymmetries as well as the average  $B^0-\bar{B}^0$  mixing.

<sup>4</sup> ABBIENDI 02I tag  $Z^0 \rightarrow b\bar{b}$  decays using a combination of secondary vertex and lepton tags. The sign of the  $b$ -quark charge is determined using an inclusive tag based on jet, vertex, and kaon charges.

<sup>5</sup> HEISTER 02H measure simultaneously  $b$  and  $c$  quark forward-backward asymmetries using their semileptonic decays to tag the quark charge. The flavor separation is obtained with a discriminating multivariate analysis.

<sup>6</sup> HEISTER 01D tag  $Z \rightarrow b\bar{b}$  events using the impact parameters of charged tracks complemented with information from displaced vertices, event shape variables, and lepton identification. The  $b$ -quark direction and charge is determined using the hemisphere charge method along with information from fast kaon tagging and charge estimators of primary and secondary vertices. The change in the quoted value due to variation of  $A_{FB}^C$  and  $R_b$  is given as  $+0.103 (A_{FB}^C - 0.0651) - 0.440 (R_b - 0.21585)$ .

<sup>7</sup> ABREU 99Y tag  $Z \rightarrow b\bar{b}$  and  $Z \rightarrow c\bar{c}$  events by an exclusive reconstruction of several  $D$  meson decay modes ( $D^{*+}$ ,  $D^0$ , and  $D^+$  with their charge-conjugate states).

<sup>8</sup> ACCIARRI 99D tag  $Z \rightarrow b\bar{b}$  events using high  $p$  and  $p_T$  leptons. The analysis determines simultaneously a mixing parameter  $\chi_b = 0.1192 \pm 0.0068 \pm 0.0051$  which is used to correct the observed asymmetry.

<sup>9</sup> ACCIARRI 98U tag  $Z \rightarrow b\bar{b}$  events using lifetime and measure the jet charge using the hemisphere charge.

<sup>10</sup> ALEXANDER 97C identify the  $b$  and  $c$  events using a  $D/D^*$  tag.

### CHARGE ASYMMETRY IN $e^+e^- \rightarrow q\bar{q}$

Summed over five lighter flavors.

Experimental and Standard Model values are somewhat event-selection dependent. Standard Model expectations contain some assumptions on  $B^0-\bar{B}^0$  mixing and on other electroweak parameters.

ASYMMETRY (%)	STD. MODEL	$\sqrt{s}$ (GeV)	DOCUMENT ID	TECN
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
- 0.76 ± 0.12 ± 0.15		91.2	<sup>1</sup> ABREU 92I	DLPH
4.0 ± 0.4 ± 0.63	4.0	91.3	<sup>2</sup> ACTON 92L	OPAL
9.1 ± 1.4 ± 1.6	9.0	57.9	ADACHI 91	TOPZ
- 0.84 ± 0.15 ± 0.04		91	DECAMP 91B	ALEP
8.3 ± 2.9 ± 1.9	8.7	56.6	STUART 90	AMY
11.4 ± 2.2 ± 2.1	8.7	57.6	ABE 89L	VNS
6.0 ± 1.3	5.0	34.8	GREENSHAW 89	JADE
8.2 ± 2.9	8.5	43.6	GREENSHAW 89	JADE

<sup>1</sup> ABREU 92I has 0.14 systematic error due to uncertainty of quark fragmentation.

<sup>2</sup> ACTON 92L use the weight function method on 259k selected  $Z \rightarrow$  hadrons events. The systematic error includes a contribution of 0.2 due to  $B^0-\bar{B}^0$  mixing effect, 0.4 due to Monte Carlo (MC) fragmentation uncertainties and 0.3 due to MC statistics. ACTON 92L derive a value of  $\sin^2\theta_W^{\text{eff}}$  to be  $0.2321 \pm 0.0017 \pm 0.0028$ .

### CHARGE ASYMMETRY IN $p\bar{p} \rightarrow Z \rightarrow e^+e^-$

ASYMMETRY (%)	STD. MODEL	$\sqrt{s}$ (GeV)	DOCUMENT ID	TECN
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
5.2 ± 5.9 ± 0.4		91	ABE 91E	CDF

### ANOMALOUS $Z\gamma$ , $Z\gamma\gamma$ , AND $ZZV$ COUPLINGS

Revised September 2013 by M.W. Gr unewald (U. College Dublin and U. Ghent) and A. Gurtu (Formerly Tata Inst.).

In on-shell  $Z\gamma$  production, deviations from the Standard Model for the  $Z\gamma\gamma^*$  and  $Z\gamma Z^*$  couplings may be described in terms of eight parameters,  $h_i^V$  ( $i = 1, 4; V = \gamma, Z$ ) [1]. The parameters  $h_i^\gamma$  describe the  $Z\gamma\gamma^*$  couplings and the parameters  $h_i^Z$  the  $Z\gamma Z^*$  couplings. In this formalism  $h_1^V$  and  $h_2^V$  lead to  $CP$ -violating and  $h_3^V$  and  $h_4^V$  to  $CP$ -conserving effects. All these anomalous contributions to the cross section increase

rapidly with center-of-mass energy. In order to ensure unitarity, these parameters are usually described by a form-factor representation,  $h_i^V(s) = h_{i_0}^V/(1 + s/\Lambda^2)^n$ , where  $\Lambda$  is the energy scale for the manifestation of a new phenomenon and  $n$  is a sufficiently large power. By convention one uses  $n = 3$  for  $h_{1,3}^V$  and  $n = 4$  for  $h_{2,4}^V$ . Usually limits on  $h_i^V$ 's are put assuming some value of  $\Lambda$ , sometimes  $\infty$ .

In on-shell  $ZZ$  production, deviations from the Standard Model for the  $ZZ\gamma^*$  and  $ZZZ^*$  couplings may be described by means of four anomalous couplings  $f_i^V$  ( $i = 4, 5; V = \gamma, Z$ ) [2]. As above, the parameters  $f_i^\gamma$  describe the  $ZZ\gamma^*$  couplings and the parameters  $f_i^Z$  the  $ZZZ^*$  couplings. The anomalous couplings  $f_5^V$  lead to violation of  $C$  and  $P$  symmetries while  $f_4^V$  introduces  $CP$  violation. Also here, formfactors depending on a scale  $\Lambda$  are used.

All these couplings  $h_i^V$  and  $f_i^V$  are zero at tree level in the Standard Model; they are measured in  $e^+e^-$ ,  $p\bar{p}$  and  $pp$  collisions at LEP, Tevatron and LHC.

## References

1. U. Baur and E.L. Berger, Phys. Rev. **D47**, 4889 (1993).
2. K. Hagiwara *et al.*, Nucl. Phys. **B282**, 253 (1987).

## $h_i^V$

Combining the LEP-2 results taking into account the correlations, the following 95% CL limits are derived [SCHAEL 13A]:

$$\begin{array}{ll}
 -0.12 < h_1^Z < +0.11, & -0.07 < h_2^Z < +0.07, \\
 -0.19 < h_3^Z < +0.06, & -0.04 < h_4^Z < +0.13, \\
 -0.05 < h_1^\gamma < +0.05, & -0.04 < h_2^\gamma < +0.02, \\
 -0.05 < h_3^\gamma < +0.00, & +0.01 < h_4^\gamma < +0.05.
 \end{array}$$

Some of the recent results from the Tevatron and LHC experiments individually surpass the combined LEP-2 results in precision (see below).

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1	AAD	16Q	ATLS	$E_{\text{cm}}^{pp} = 8 \text{ TeV}$
2	KHACHATRYAN	16AE	CMS	$E_{\text{cm}}^{pp} = 8 \text{ TeV}$
3	KHACHATRYAN	15AC	CMS	$E_{\text{cm}}^{pp} = 8 \text{ TeV}$
4	CHATRCHYAN	14AB	CMS	$E_{\text{cm}}^{pp} = 7 \text{ TeV}$
5	AAD	13AN	ATLS	$E_{\text{cm}}^{pp} = 7 \text{ TeV}$
6	CHATRCHYAN	13BI	CMS	$E_{\text{cm}}^{pp} = 7 \text{ TeV}$
7	ABAZOV	12S	D0	$E_{\text{cm}}^{p\bar{p}} = 1.96 \text{ TeV}$
8	AALTONEN	11S	CDF	$E_{\text{cm}}^{p\bar{p}} = 1.96 \text{ TeV}$
9	CHATRCHYAN	11M	CMS	$E_{\text{cm}}^{pp} = 7 \text{ TeV}$
10	ABAZOV	09L	D0	$E_{\text{cm}}^{p\bar{p}} = 1.96 \text{ TeV}$
11	ABAZOV	07M	D0	$E_{\text{cm}}^{p\bar{p}} = 1.96 \text{ TeV}$
12	ABDALLAH	07C	DLPH	$E_{\text{cm}}^{ee} = 183\text{--}208 \text{ GeV}$
13	ACHARD	04H	L3	$E_{\text{cm}}^{ee} = 183\text{--}208 \text{ GeV}$
14	ABBIENDI,G	00C	OPAL	$E_{\text{cm}}^{ee} = 189 \text{ GeV}$
15	ABBOTT	98M	D0	$E_{\text{cm}}^{p\bar{p}} = 1.8 \text{ TeV}$
16	ABREU	98K	DLPH	$E_{\text{cm}}^{ee} = 161, 172 \text{ GeV}$

- <sup>1</sup> AAD 16Q study  $Z\gamma$  production in  $pp$  collisions. In events with no additional jets, 10268 (12738)  $Z$  decays to electron (muon) pairs are selected, with an expected background of  $1291 \pm 340$  ( $1537 \pm 408$ ) events, as well as 1039  $Z$  decays to neutrino pairs with an expected background of  $450 \pm 96$  events. Analyzing the photon transverse momentum distribution above 250 GeV (400 GeV) for lepton (neutrino) events, yields the 95% C.L. limits:  $-7.8 \times 10^{-4} < h_3^Z < 8.6 \times 10^{-4}$ ,  $-3.0 \times 10^{-6} < h_4^Z < 2.9 \times 10^{-6}$ ,  $-9.5 \times 10^{-4} < h_3^\gamma < 9.9 \times 10^{-4}$ ,  $-3.2 \times 10^{-6} < h_4^\gamma < 3.2 \times 10^{-6}$ .
- <sup>2</sup> KHACHATRYAN 16AE determine the  $Z\gamma \rightarrow \nu\bar{\nu}\gamma$  cross section by selecting events with a photon of  $E_T > 145 \text{ GeV}$  and  $\cancel{E}_T > 140 \text{ GeV}$ . 630 candidate events are observed with an expected SM background of  $269 \pm 26$ . The  $E_T$  spectrum of the photon is used to set 95% C.L. limits as follows:  $-1.5 \times 10^{-3} < h_3^Z < 1.6 \times 10^{-3}$ ,  $-3.9 \times 10^{-6} < h_4^Z < 4.5 \times 10^{-6}$ ,  $-1.1 \times 10^{-3} < h_3^\gamma < 0.9 \times 10^{-3}$ ,  $-3.8 \times 10^{-6} < h_4^\gamma < 4.3 \times 10^{-6}$ .
- <sup>3</sup> KHACHATRYAN 15AC study  $Z\gamma$  events in 8 TeV  $pp$  interactions, where the  $Z$  decays into 2 same-flavor, opposite sign leptons ( $e$  or  $\mu$ ) and a photon with  $p_T > 15 \text{ GeV}$ . The  $p_T$  of a lepton is required to be  $> 20 \text{ GeV}/c$ , their effective mass  $> 50 \text{ GeV}$ , and the photon should have a separation  $\Delta R > 0.7$  with each lepton. The observed  $p_T$  distribution of the photons is used to extract the 95% C.L. limits:  $-3.8 \times 10^{-3} < h_3^Z < 3.7 \times 10^{-3}$ ,  $-3.1 \times 10^{-5} < h_4^Z < 3.0 \times 10^{-5}$ ,  $-4.6 \times 10^{-3} < h_3^\gamma < 4.6 \times 10^{-3}$ ,  $-3.6 \times 10^{-5} < h_4^\gamma < 3.5 \times 10^{-5}$ .
- <sup>4</sup> CHATRCHYAN 14AB measure  $Z\gamma$  production cross section for  $p_T^\gamma > 15 \text{ GeV}$  and  $R(\ell\gamma) > 0.7$ , which is the separation between the  $\gamma$  and the final state charged lepton ( $e$  or  $\mu$ ) in the azimuthal angle-pseudorapidity ( $\phi - \eta$ ) plane. The di-lepton mass is required to be  $> 50 \text{ GeV}$ . After background subtraction the number of  $ee\gamma$  and  $\mu\mu\gamma$  events is determined to be  $3160 \pm 120$  and  $5030 \pm 233$  respectively, compatible with expectations from the SM. This leads to a 95% CL limits of  $-1 \times 10^{-2} < h_3^Z < 1 \times 10^{-2}$ ,  $-9 \times 10^{-5} < h_4^Z < 9 \times 10^{-5}$ ,  $-9 \times 10^{-3} < h_3^\gamma < 9 \times 10^{-3}$ ,  $-8 \times 10^{-5} < h_4^\gamma < 8 \times 10^{-5}$ , assuming  $h_1^V$  and  $h_2^V$  have SM values,  $V = \gamma$  or  $Z$ .
- <sup>5</sup> AAD 13AN study  $Z\gamma$  production in  $pp$  collisions. In events with no additional jet, 1417 (2031)  $Z$  decays to electron (muon) pairs are selected, with an expected background of



- 156 ± 54 (244 ± 64) events, as well as 662 Z decays to neutrino pairs with an expected background of 302 ± 42 events. Analysing the photon  $p_T$  spectrum above 100 GeV yields the 95% C.L. limits:  $-0.013 < h_3^Z < 0.014$ ,  $-8.7 \times 10^{-5} < h_4^Z < 8.7 \times 10^{-5}$ ,  $-0.015 < h_3^\gamma < 0.016$ ,  $-9.4 \times 10^{-5} < h_4^\gamma < 9.2 \times 10^{-5}$ . Supersedes AAD 12BX.
- <sup>6</sup> CHATRCHYAN 13BI determine the  $Z\gamma \rightarrow \nu\bar{\nu}\gamma$  cross section by selecting events with a photon of  $E_T > 145$  GeV and a  $\cancel{E}_T > 130$  GeV. 73 candidate events are observed with an expected SM background of  $30.2 \pm 6.5$ . The  $E_T$  spectrum of the photon is used to set 95% C.L. limits as follows:  $|h_3^Z| < 2.7 \times 10^{-3}$ ,  $|h_4^Z| < 1.3 \times 10^{-5}$ ,  $|h_3^\gamma| < 2.9 \times 10^{-3}$ ,  $|h_4^\gamma| < 1.5 \times 10^{-5}$ .
- <sup>7</sup> ABAZOV 12S study  $Z\gamma$  production in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV using  $6.2 \text{ fb}^{-1}$  of data where the Z decays to electron (muon) pairs and the photon has at least 10 GeV of transverse momentum. In data, 304 (308) di-electron (di-muon) events are observed with an expected background of  $255 \pm 16$  ( $285 \pm 24$ ) events. Based on the photon  $p_T$  spectrum, and including also earlier data and the  $Z \rightarrow \nu\bar{\nu}$  decay mode (from ABAZOV 09L), the following 95% C.L. limits are reported:  $|h_{03}^Z| < 0.026$ ,  $|h_{04}^Z| < 0.0013$ ,  $|h_{03}^\gamma| < 0.027$ ,  $|h_{04}^\gamma| < 0.0014$  for a form factor scale of  $\Lambda = 1.5$  TeV.
- <sup>8</sup> AALTONEN 11S study  $Z\gamma$  events in  $p\bar{p}$  interactions at  $\sqrt{s} = 1.96$  TeV with integrated luminosity  $5.1 \text{ fb}^{-1}$  for  $Z \rightarrow e^+e^-/\mu^+\mu^-$  and  $4.9 \text{ fb}^{-1}$  for  $Z \rightarrow \nu\bar{\nu}$ . For the charged lepton case, the two leptons must be of the same flavor with the transverse momentum/energy of one  $> 20$  GeV and the other  $> 10$  GeV. The isolated photon must have  $E_T > 50$  GeV. They observe 91 events with  $87.2 \pm 7.8$  events expected from standard model processes. For the  $\nu\bar{\nu}$  case they require solitary photons with  $E_T > 25$  GeV and missing  $E_T > 25$  GeV and observe 85 events with standard model expectation of  $85.9 \pm 5.6$  events. Taking the form factor  $\Lambda = 1.5$  TeV they derive 95% C.L. limits as  $|h_3^{\gamma,Z}| < 0.022$  and  $|h_4^{\gamma,Z}| < 0.0009$ .
- <sup>9</sup> CHATRCHYAN 11M study  $Z\gamma$  production in  $pp$  collisions at  $\sqrt{s} = 7$  TeV using  $36 \text{ pb}^{-1}$   $pp$  data, where the Z decays to  $e^+e^-$  or  $\mu^+\mu^-$ . The total cross sections are measured for photon transverse energy  $E_T^\gamma > 10$  GeV and spatial separation from charged leptons in the plane of pseudo rapidity and azimuthal angle  $\Delta R(\ell,\gamma) > 0.7$  with the dilepton invariant mass requirement of  $M_{\ell\ell} > 50$  GeV. The number of  $e^+e^-\gamma$  and  $\mu^+\mu^-\gamma$  candidates is 81 and 90 with estimated backgrounds of  $20.5 \pm 2.5$  and  $27.3 \pm 3.2$  events respectively. The 95% CL limits for  $ZZ\gamma$  couplings are  $-0.05 < h_3^Z < 0.06$  and  $-0.0005 < h_4^Z < 0.0005$ , and for  $Z\gamma\gamma$  couplings are  $-0.07 < h_3^\gamma < 0.07$  and  $-0.0005 < h_4^\gamma < 0.0006$ .
- <sup>10</sup> ABAZOV 09L study  $Z\gamma, Z \rightarrow \nu\bar{\nu}$  production in  $p\bar{p}$  collisions at 1.96 TeV C.M. energy. They select 51 events with a photon of transverse energy  $E_T$  larger than 90 GeV, with an expected background of 17 events. Based on the photon  $E_T$  spectrum and including also Z decays to charged leptons (from ABAZOV 07M), the following 95% CL limits are reported:  $|h_{30}^\gamma| < 0.033$ ,  $|h_{40}^\gamma| < 0.0017$ ,  $|h_{30}^Z| < 0.033$ ,  $|h_{40}^Z| < 0.0017$ .
- <sup>11</sup> ABAZOV 07M use 968  $p\bar{p} \rightarrow e^+e^-/\mu^+\mu^-\gamma X$  candidates, at 1.96 TeV center of mass energy, to tag  $p\bar{p} \rightarrow Z\gamma$  events by requiring  $E_T(\gamma) > 7$  GeV, lepton-gamma separation  $\Delta R_{\ell\gamma} > 0.7$ , and di-lepton invariant mass  $> 30$  GeV. The cross section is in agreement with the SM prediction. Using these  $Z\gamma$  events they obtain 95% C.L. limits on each  $h_i^V$ , keeping all others fixed at their SM values. They report:  $-0.083 < h_{30}^Z < 0.082$ ,  $-0.0053 < h_{40}^Z < 0.0054$ ,  $-0.085 < h_{30}^\gamma < 0.084$ ,  $-0.0053 < h_{40}^\gamma < 0.0054$ , for the form factor scale  $\Lambda = 1.2$  TeV.
- <sup>12</sup> Using data collected at  $\sqrt{s} = 183\text{--}208$ , ABDALLAH 07C select 1,877  $e^+e^- \rightarrow Z\gamma$  events with  $Z \rightarrow q\bar{q}$  or  $\nu\bar{\nu}$ , 171  $e^+e^- \rightarrow ZZ$  events with  $Z \rightarrow q\bar{q}$  or lepton pair

(except an explicit  $\tau$  pair), and 74  $e^+e^- \rightarrow Z\gamma^*$  events with a  $q\bar{q}\mu^+\mu^-$  or  $q\bar{q}e^+e^-$  signature, to derive 95% CL limits on  $h_i^V$ . Each limit is derived with other parameters set to zero. They report:  $-0.23 < h_1^Z < 0.23$ ,  $-0.30 < h_3^Z < 0.16$ ,  $-0.14 < h_1^\gamma < 0.14$ ,  $-0.049 < h_3^\gamma < 0.044$ .

<sup>13</sup> ACHARD 04H select 3515  $e^+e^- \rightarrow Z\gamma$  events with  $Z \rightarrow q\bar{q}$  or  $\nu\bar{\nu}$  at  $\sqrt{s} = 189\text{--}209$  GeV to derive 95% CL limits on  $h_i^V$ . For deriving each limit the other parameters are fixed at zero. They report:  $-0.153 < h_1^Z < 0.141$ ,  $-0.087 < h_2^Z < 0.079$ ,  $-0.220 < h_3^Z < 0.112$ ,  $-0.068 < h_4^Z < 0.148$ ,  $-0.057 < h_1^\gamma < 0.057$ ,  $-0.050 < h_2^\gamma < 0.023$ ,  $-0.059 < h_3^\gamma < 0.004$ ,  $-0.004 < h_4^\gamma < 0.042$ .

<sup>14</sup> ABBIENDI,G 00C study  $e^+e^- \rightarrow Z\gamma$  events (with  $Z \rightarrow q\bar{q}$  and  $Z \rightarrow \nu\bar{\nu}$ ) at 189 GeV to obtain the central values (and 95% CL limits) of these couplings:  $h_1^Z = 0.000 \pm 0.100$  ( $-0.190, 0.190$ ),  $h_2^Z = 0.000 \pm 0.068$  ( $-0.128, 0.128$ ),  $h_3^Z = -0.074^{+0.102}_{-0.103}$  ( $-0.269, 0.119$ ),  $h_4^Z = 0.046 \pm 0.068$  ( $-0.084, 0.175$ ),  $h_1^\gamma = 0.000 \pm 0.061$  ( $-0.115, 0.115$ ),  $h_2^\gamma = 0.000 \pm 0.041$  ( $-0.077, 0.077$ ),  $h_3^\gamma = -0.080^{+0.039}_{-0.041}$  ( $-0.164, -0.006$ ),  $h_4^\gamma = 0.064^{+0.033}_{-0.030}$  ( $+0.007, +0.134$ ). The results are derived assuming that only one coupling at a time is different from zero.

<sup>15</sup> ABBOTT 98M study  $p\bar{p} \rightarrow Z\gamma + X$ , with  $Z \rightarrow e^+e^-, \mu^+\mu^-, \bar{\nu}\nu$  at 1.8 TeV, to obtain 95% CL limits at  $\Lambda = 750$  GeV:  $|h_{30}^Z| < 0.36$ ,  $|h_{40}^Z| < 0.05$  (keeping  $h_i^\gamma = 0$ ), and  $|h_{30}^\gamma| < 0.37$ ,  $|h_{40}^\gamma| < 0.05$  (keeping  $h_i^Z = 0$ ). Limits on the  $CP$ -violating couplings are  $|h_{10}^Z| < 0.36$ ,  $|h_{20}^Z| < 0.05$  (keeping  $h_i^\gamma = 0$ ), and  $|h_{10}^\gamma| < 0.37$ ,  $|h_{20}^\gamma| < 0.05$  (keeping  $h_i^Z = 0$ ).

<sup>16</sup> ABREU 98K determine a 95% CL upper limit on  $\sigma(e^+e^- \rightarrow \gamma + \text{invisible particles}) < 2.5$  pb using 161 and 172 GeV data. This is used to set 95% CL limits on  $|h_{30}^\gamma| < 0.8$  and  $|h_{30}^Z| < 1.3$ , derived at a scale  $\Lambda = 1$  TeV and with  $n = 3$  in the form factor representation.

### $f_i^V$

Combining the LEP-2 results taking into account the correlations, the following 95% CL limits are derived [SCHAEL 13A]:

$$\begin{aligned} -0.28 < f_4^Z < +0.32, & & -0.34 < f_5^Z < +0.35, \\ -0.17 < f_4^\gamma < +0.19, & & -0.35 < f_5^\gamma < +0.32. \end{aligned}$$

Some of the recent results from the Tevatron and LHC experiments individually surpass the combined LEP-2 results in precision (see below).

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1	AAD	23CH	$E_{\text{cm}}^{pp} = 13$ TeV
2	SIRUNYAN	21Q CMS	$E_{\text{cm}}^{pp} = 13$ TeV
3	AABOUD	19AY ATLS	$E_{\text{cm}}^{pp} = 13$ TeV
4	AABOUD	18Q ATLS	$E_{\text{cm}}^{pp} = 13$ TeV
5	SIRUNYAN	18BT CMS	$E_{\text{cm}}^{pp} = 13$ TeV
6	KHACHATRY...15B	CMS	$E_{\text{cm}}^{pp} = 8$ TeV

7	KHACHATRYAN...15BC	CMS	$E_{\text{cm}}^{pp} = 7, 8 \text{ TeV}$
8	AAD	13Z ATLS	$E_{\text{cm}}^{pp} = 7 \text{ TeV}$
9	CHATRCHYAN	13B CMS	$E_{\text{cm}}^{pp} = 7 \text{ TeV}$
10	SCHAEEL	09 ALEP	$E_{\text{cm}}^{ee} = 192\text{--}209 \text{ GeV}$
11	ABAZOV	08K D0	$E_{\text{cm}}^{p\bar{p}} = 1.96 \text{ TeV}$
12	ABDALLAH	07C DLPH	$E_{\text{cm}}^{ee} = 183\text{--}208 \text{ GeV}$
13	ABBIENDI	04C OPAL	
14	ACHARD	03D L3	

<sup>1</sup> AAD 23CH measure  $ZZ$  production with the  $Z$  bosons decaying to electrons or muons. Analysing the angular information of the final-state four-lepton system, the following limits are derived at 95% C.L.:  $-0.012 < f_4^Z < 0.012$ ,  $-0.015 < f_4^\gamma < 0.015$ .

<sup>2</sup> SIRUNYAN 21Q measure  $ZZ$  production where both  $Z$  bosons decay in the electron or muon channel. Analysing the four-lepton invariant mass distribution, the following limits are derived at 95% C.L. in units of  $10^{-4}$ :  $-6.6 < f_4^Z < 6.0$ ,  $-5.5 < f_5^Z < 7.5$ ,  $-7.8 < f_4^\gamma < 7.1$ ,  $-6.8 < f_5^\gamma < 7.5$ . This set of parameters is linearly related to a set of EFT parameters, resulting in the following limits at 95% C.L. in units of  $\text{TeV}^{-4}$ :  $-2.3 < c_{\bar{B}W}/\Lambda^4 < 2.5$ ,  $-1.4 < c_{WW}/\Lambda^4 < 1.2$ ,  $-1.4 < c_{BW}/\Lambda^4 < 1.3$ ,  $-1.2 < c_{BB}/\Lambda^4 < 1.2$ .

<sup>3</sup> AABOUD 19AY study  $ZZ$  production in the  $\ell\ell\nu\nu$  decay channel. Events with a pair of isolated high-transverse momentum charged leptons (electron pairs or muon pairs), and with large missing energy, are selected. In the data, 371 (416) di-electron (di-muon) events are found, with a total expected background of  $128 \pm 8$  ( $143 \pm 8$ ) events. Analysing the transverse momentum distribution of the charged dilepton system above 150 GeV, the following 95% C.L. limits are derived in units of  $10^{-3}$ :  $-1.2 < f_4^\gamma < 1.2$ ,  $-1.0 < f_4^Z < 1.0$ ,  $-1.2 < f_5^\gamma < 1.2$ ,  $-1.0 < f_5^Z < 1.0$ .

<sup>4</sup> AABOUD 18Q study  $pp \rightarrow ZZ$  events at  $\sqrt{s} = 13 \text{ TeV}$  with  $Z \rightarrow e^+e^-$  or  $Z \rightarrow \mu^+\mu^-$ . The number of events observed in the  $4e$ ,  $2e2\mu$ , and  $4\mu$  channels is 249, 465, and 303 respectively. Analysing the  $p_T$  spectrum of the leading  $Z$  boson, the following the following 95% C.L. limits are derived in units of  $10^{-4}$ :  $-1.8 < f_4^\gamma < 1.8$ ,  $-1.5 < f_4^Z < 1.5$ ,  $-1.8 < f_5^\gamma < 1.8$ ,  $-1.5 < f_5^Z < 1.5$ .

<sup>5</sup> SIRUNYAN 18BT study  $ppZZ$  events at  $\sqrt{s} = 13 \text{ TeV}$  with  $Z \rightarrow e^+e^-$  or  $Z \rightarrow \mu^+\mu^-$ . The number of events observed in the  $4e$ ,  $2e2\mu$ , and  $4\mu$  channels is 220, 543 and 335 respectively. Analysing the 4-lepton invariant mass spectrum, the following 95% C.L. limits are derived in units of  $10^{-3}$ :  $-1.2 < f_4^\gamma < 1.3$ ,  $-1.2 < f_4^Z < 1.0$ ,  $-1.2 < f_5^\gamma < 1.3$ ,  $-1.0 < f_5^Z < 1.3$ .

<sup>6</sup> KHACHATRYAN 15B study  $ZZ$  production in 8 TeV  $pp$  collisions. In the decay modes  $ZZ \rightarrow 4e$ ,  $4\mu$ ,  $2e2\mu$ , 54, 75, 148 events are observed, with an expected background of  $2.2 \pm 0.9$ ,  $1.2 \pm 0.6$ , and  $2.4 \pm 1.0$  events, respectively. Analysing the 4-lepton invariant mass spectrum in the range from 110 GeV to 1200 GeV, the following 95% C.L. limits are obtained:  $|f_4^Z| < 0.004$ ,  $|f_5^Z| < 0.004$ ,  $|f_4^\gamma| < 0.005$ ,  $|f_5^\gamma| < 0.005$ .

<sup>7</sup> KHACHATRYAN 15BC use the cross section measurement of the final state  $pp \rightarrow ZZ \rightarrow 2\ell 2\nu$ , ( $\ell$  being an electron or a muon) at 7 and 8 TeV to put limits on these triple gauge couplings. Effective mass of the charged lepton pair is required to be in the range 83.5–98.5 GeV and the dilepton  $p_T > 45 \text{ GeV}$ . The reduced missing  $E_T$  is required to be  $> 65 \text{ GeV}$ , which takes into account the fake missing  $E_T$  due to detector effects. The numbers of  $e^+e^-$  and  $\mu^+\mu^-$  events selected are 35 and 40 at 7 TeV and 176 and 271 at 8 TeV respectively. The production cross sections so obtained are in agreement

- with SM predictions. The following 95% C.L. limits are set:  $-0.0028 < f_4^Z < 0.0032$ ,  $-0.0037 < f_4^\gamma < 0.0033$ ,  $-0.0029 < f_5^Z < 0.0031$ ,  $-0.0033 < f_5^\gamma < 0.0037$ . Combining with previous results (KHACHATRYAN 15B and CHATRCHYAN 13B) which include 7 TeV and 8 TeV data on the final states  $p\bar{p} \rightarrow ZZ \rightarrow 2\ell 2\ell'$  where  $\ell$  and  $\ell'$  are an electron or a muon, the best limits are  $-0.0022 < f_4^Z < 0.0026$ ,  $-0.0029 < f_4^\gamma < 0.0026$ ,  $-0.0023 < f_5^Z < 0.0023$ ,  $-0.0026 < f_5^\gamma < 0.0027$ .
- <sup>8</sup> AAD 13Z study  $ZZ$  production in  $p\bar{p}$  collisions at  $\sqrt{s} = 7$  TeV. In the  $ZZ \rightarrow \ell^+ \ell^- \ell'^+ \ell'^-$  final state they observe a total of 66 events with an expected background of  $0.9 \pm 1.3$ . In the  $ZZ \rightarrow \ell^+ \ell^- \nu\nu$  final state they observe a total of 87 events with an expected background of  $46.9 \pm 5.2$ . The limits on anomalous TGCs are determined using the observed and expected numbers of these  $ZZ$  events binned in  $p_T^Z$ . The 95% C.L. are as follows: for form factor scale  $\Lambda = \infty$ ,  $-0.015 < f_4^\gamma < 0.015$ ,  $-0.013 < f_4^Z < 0.013$ ,  $-0.016 < f_5^\gamma < 0.015$ ,  $-0.013 < f_5^Z < 0.013$ ; for form factor scale  $\Lambda = 3$  TeV,  $-0.022 < f_4^\gamma < 0.023$ ,  $-0.019 < f_4^Z < 0.019$ ,  $-0.023 < f_5^\gamma < 0.023$ ,  $-0.020 < f_5^Z < 0.019$ .
- <sup>9</sup> CHATRCHYAN 13B study  $ZZ$  production in  $p\bar{p}$  collisions and select 54  $ZZ$  candidates in the  $Z$  decay channel with electrons or muons with an expected background of  $1.4 \pm 0.5$  events. The resulting 95% C.L. ranges are:  $-0.013 < f_4^\gamma < 0.015$ ,  $-0.011 < f_4^Z < 0.012$ ,  $-0.014 < f_5^\gamma < 0.014$ ,  $-0.012 < f_5^Z < 0.012$ .
- <sup>10</sup> Using data collected in the center of mass energy range 192–209 GeV, SCHAEEL 09 select 318  $e^+e^- \rightarrow ZZ$  events with 319.4 expected from the standard model. Using this data they derive the following 95% CL limits:  $-0.321 < f_4^\gamma < 0.318$ ,  $-0.534 < f_4^Z < 0.534$ ,  $-0.724 < f_5^\gamma < 0.733$ ,  $-1.194 < f_5^Z < 1.190$ .
- <sup>11</sup> ABAZOV 08K search for  $ZZ$  and  $Z\gamma^*$  events with  $1 \text{ fb}^{-1} p\bar{p}$  data at  $\sqrt{s} = 1.96$  TeV in  $(ee)(ee)$ ,  $(\mu\mu)(\mu\mu)$ ,  $(ee)(\mu\mu)$  final states requiring the lepton pair masses to be  $> 30$  GeV. They observe 1 event, which is consistent with an expected signal of  $1.71 \pm 0.15$  events and a background of  $0.13 \pm 0.03$  events. From this they derive the following limits, for a form factor ( $\Lambda$ ) value of 1.2 TeV:  $-0.28 < f_{40}^Z < 0.28$ ,  $-0.31 < f_{50}^Z < 0.29$ ,  $-0.26 < f_{40}^\gamma < 0.26$ ,  $-0.30 < f_{50}^\gamma < 0.28$ .
- <sup>12</sup> Using data collected at  $\sqrt{s} = 183$ –208 GeV, ABDALLAH 07C select 171  $e^+e^- \rightarrow ZZ$  events with  $Z \rightarrow q\bar{q}$  or lepton pair (except an explicit  $\tau$  pair), and 74  $e^+e^- \rightarrow Z\gamma^*$  events with a  $q\bar{q}\mu^+\mu^-$  or  $q\bar{q}e^+e^-$  signature, to derive 95% CL limits on  $f_i^V$ . Each limit is derived with other parameters set to zero. They report:  $-0.40 < f_4^Z < 0.42$ ,  $-0.38 < f_5^Z < 0.62$ ,  $-0.23 < f_4^\gamma < 0.25$ ,  $-0.52 < f_5^\gamma < 0.48$ .
- <sup>13</sup> ABBIENDI 04C study  $ZZ$  production in  $e^+e^-$  collisions in the C.M. energy range 190–209 GeV. They select 340 events with an expected background of 180 events. Including the ABBIENDI 00N data at 183 and 189 GeV (118 events with an expected background of 65 events) they report the following 95% CL limits:  $-0.45 < f_4^Z < 0.58$ ,  $-0.94 < f_5^Z < 0.25$ ,  $-0.32 < f_4^\gamma < 0.33$ , and  $-0.71 < f_5^\gamma < 0.59$ .
- <sup>14</sup> ACHARD 03D study  $Z$ -boson pair production in  $e^+e^-$  collisions in the C.M. energy range 200–209 GeV. They select 549 events with an expected background of 432 events. Including the ACCIARRI 99G and ACCIARRI 99O data (183 and 189 GeV respectively, 286 events with an expected background of 241 events) and the 192–202 GeV ACCIARRI 01I results (656 events, expected background of 512 events), they report the following 95%

CL limits:  $-0.48 \leq f_4^Z \leq 0.46$ ,  $-0.36 \leq f_5^Z \leq 1.03$ ,  $-0.28 \leq f_4^\gamma \leq 0.28$ , and  $-0.40 \leq f_5^\gamma \leq 0.47$ .

---

### ANOMALOUS $W/Z$ QUARTIC COUPLINGS

Revised November 2015 by M.W. Gr unewald (U. College Dublin) and A. Gurtu (Formerly Tata Inst.).

Quartic couplings,  $WWZZ$ ,  $WWZ\gamma$ ,  $WW\gamma\gamma$ , and  $ZZ\gamma\gamma$ , were studied at LEP and Tevatron at energies at which the Standard Model predicts negligible contributions to multiboson production. Thus, to parametrize limits on these couplings, an effective theory approach is adopted which supplements the Standard Model Lagrangian with higher dimensional operators which include quartic couplings. The LEP collaborations chose the lowest dimensional representation of operators (dimension 6) which presumes the  $SU(2)\times U(1)$  gauge symmetry is broken by means other than the conventional Higgs scalar doublet [1–3]. In this representation possible quartic couplings,  $a_0, a_c, a_n$ , are expressed in terms of the following dimension-6 operators [1,2];

$$\begin{aligned} L_6^0 &= -\frac{e^2}{16\Lambda^2} a_0 F^{\mu\nu} F_{\mu\nu} \vec{W}^\alpha \cdot \vec{W}_\alpha \\ L_6^c &= -\frac{e^2}{16\Lambda^2} a_c F^{\mu\alpha} F_{\mu\beta} \vec{W}^\beta \cdot \vec{W}_\alpha \\ L_6^n &= -i\frac{e^2}{16\Lambda^2} a_n \epsilon_{ijk} W_{\mu\alpha}^{(i)} W_\nu^{(j)} W^{(k)\alpha} F^{\mu\nu} \\ \tilde{L}_6^0 &= -\frac{e^2}{16\Lambda^2} \tilde{a}_0 F^{\mu\nu} \tilde{F}_{\mu\nu} \vec{W}^\alpha \cdot \vec{W}_\alpha \\ \tilde{L}_6^n &= -i\frac{e^2}{16\Lambda^2} \tilde{a}_n \epsilon_{ijk} W_{\mu\alpha}^{(i)} W_\nu^{(j)} W^{(k)\alpha} \tilde{F}^{\mu\nu} \end{aligned}$$

where  $F, W$  are photon and  $W$  fields,  $L_6^0$  and  $L_6^c$  conserve  $C, P$  separately ( $\tilde{L}_6^0$  conserves only  $C$ ) and generate anomalous  $W^+W^-\gamma\gamma$  and  $ZZ\gamma\gamma$  couplings,  $L_6^n$  violates  $CP$  ( $\tilde{L}_6^n$  violates both  $C$  and  $P$ ) and generates an anomalous  $W^+W^-Z\gamma$  coupling, and  $\Lambda$  is an energy scale for new physics. For the  $ZZ\gamma\gamma$  coupling the  $CP$ -violating term represented by  $L_6^n$  does not contribute. These couplings are assumed to be real and to vanish at tree level in the Standard Model.

Within the same framework as above, a more recent description of the quartic couplings [3] treats the anomalous parts of the  $WW\gamma\gamma$  and  $ZZ\gamma\gamma$  couplings separately, leading to two sets parametrized as  $a_0^V/\Lambda^2$  and  $a_c^V/\Lambda^2$ , where  $V = W$  or  $Z$ .

With the discovery of a Higgs at the LHC in 2012, it is then useful to go to the next higher dimensional representation (dimension 8 operators) in which the gauge symmetry is broken by the conventional Higgs scalar doublet [3,4]. There are 14 operators which can contribute to the anomalous quartic coupling signal. Some of the operators have analogues in the dimension 6 scheme. The CMS collaboration, [5], have used this parametrization, in which the connections between the two schemes are also summarized:

$$\begin{aligned} \mathcal{L}_{AQGC} = & -\frac{e^2 a_0^W}{8 \Lambda^2} F_{\mu\nu} F^{\mu\nu} W^{+a} W_a^- \\ & -\frac{e^2 a_c^W}{16 \Lambda^2} F_{\mu\nu} F^{\mu a} (W^{+\nu} W_a^- + W^{-\nu} W_a^+) \\ & -e^2 g^2 \frac{\kappa_0^W}{\Lambda^2} F_{\mu\nu} Z^{\mu\nu} W^{+a} W_a^- \\ & -\frac{e^2 g^2 \kappa_c^W}{2 \Lambda^2} F_{\mu\nu} Z^{\mu a} (W^{+\nu} W_a^- + W^{-\nu} W_a^+) \\ & + \frac{f_{T,0}}{\Lambda^4} Tr[\widehat{W}_{\mu\nu} \widehat{W}^{\mu\nu}] \times Tr[\widehat{W}_{\alpha\beta} \widehat{W}^{\alpha\beta}] \end{aligned}$$

The energy scale of possible new physics is  $\Lambda$ , and  $g = e/\sin(\theta_W)$ ,  $e$  being the unit electric charge and  $\theta_W$  the Weinberg angle. The field tensors are described in [3,4].

The two dimension 6 operators  $a_0^W/\Lambda^2$  and  $a_c^W/\Lambda^2$  are associated with the  $WW\gamma\gamma$  vertex. Among dimension 8 operators,  $\kappa_0^W/\Lambda^2$  and  $\kappa_c^W/\Lambda^2$  are associated with the  $WWZ\gamma$  vertex, whereas the parameter  $f_{T,0}/\Lambda^4$  contributes to both vertices. There is a relationship between these two dimension 6 parameters and the dimension 8 parameters  $f_{M,i}/\Lambda^4$  as follows [3]:

$$\frac{a_0^W}{\Lambda^2} = -\frac{4M_W^2}{g^2} \frac{f_{M,0}}{\Lambda^4} - \frac{8M_W^2}{g'^2} \frac{f_{M,2}}{\Lambda^4}$$

$$\frac{a_c^W}{\Lambda^2} = -\frac{4M_W^2}{g^2} \frac{f_{M,1}}{\Lambda^4} - \frac{8M_W^2}{g'^2} \frac{f_{M,3}}{\Lambda^4}$$

where  $g' = e/\cos(\theta_W)$  and  $M_W$  is the invariant mass of the  $W$  boson. This relation provides a translation between limits on dimension 6 operators  $a_{0,c}^W$  and  $f_{M,j}/\Lambda^4$ . It is further required [4] that  $f_{M,0} = 2f_{M,2}$  and  $f_{M,1} = 2f_{M,3}$  which suppresses contributions to the  $WWZ\gamma$  vertex. The complete set of Lagrangian contributions as presented in [4] corresponds to 19 anomalous couplings in total –  $f_{S,i}$ ,  $i = 1, 2$ ,  $f_{M,i}$ ,  $i = 0, \dots, 8$  and  $f_{T,i}$ ,  $i = 0, \dots, 9$  – each scaled by  $1/\Lambda^4$ .

The ATLAS collaboration [6], on the other hand, follows a K-matrix driven approach of Ref. 7 in which the anomalous couplings can be expressed in terms of two parameters  $\alpha_4$  and  $\alpha_5$ , which account for all BSM effects.

It is the early stages in the determination of quartic couplings by the LHC experiments. It is hoped that the two collaborations, ATLAS and CMS, will agree to use at least one common set of parameters to express these limits to enable the reader to make a comparison and allow for a possible LHC combination.

## References

1. G. Belanger and F. Boudjema, Phys. Lett. **B288**, 201 (1992).
2. J.W. Stirling and A. Werthenbach, Eur. Phys. J. **C14**, 103 (2000);  
J.W. Stirling and A. Werthenbach, Phys. Lett. **B466**, 369 (1999);

- A. Denner *et al.*, Eur. Phys. J. **C20**, 201 (2001);  
 G. Montagna *et al.*, Phys. Lett. **B515**, 197 (2001).
3. G. Belanger *et al.*, Eur. Phys. J. **C13**, 283 (2000).
  4. O.J.P. Éboli, M.C. Gonzalez-Garcia, and S.M. Lietti, Phys. Rev. **D69**, 095005 (2004);  
 O.J.P. Éboli, M.C. Gonzalez-Garcia, and J.K. Mizukoshi, Phys. Rev. **D77**, 073005 (2006).
  5. S. Chatrchyan *et al.*, Phys. Rev. **D90**, 032008 (2014);  
 S. Chatrchyan *et al.*, Phys. Rev. Lett. **114**, 051801 (2015).
  6. G. Aad *et al.*, Phys. Rev. Lett. **113**, 141803 (2014).
  7. A. Albateanu, W. Killian, and J. Reuter, JHEP **0811**, 010 (2008).

### $a_0/\Lambda^2, a_c/\Lambda^2$

Combining published and unpublished preliminary LEP results the following 95% CL intervals for the QGCs associated with the  $ZZ\gamma\gamma$  vertex are derived (CERN-PH-EP/2005-051 or hep-ex/0511027):

$$-0.008 < a_0^Z/\Lambda^2 < +0.021$$

$$-0.029 < a_c^Z/\Lambda^2 < +0.039$$

Anomalous  $Z$  quartic couplings have also been measured by the Tevatron and LHC experiments. As discussed in the review on "Anomalous  $W/Z$  quartic couplings," the coupling parameters in the Anomalous QGC Lagrangian may relate to processes involving only the  $W$  or only to the  $Z$  or to both. Thus, results on all other AQGCs are reported together in the  $W$  listings.

VALUE	DOCUMENT ID	TECN
-------	-------------	------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup>	ABBIENDI	04L	OPAL
<sup>2</sup>	HEISTER	04A	ALEP
<sup>3</sup>	ACHARD	02G	L3

<sup>1</sup> ABBIENDI 04L select 20  $e^+e^- \rightarrow \nu\bar{\nu}\gamma\gamma$  acoplanar events in the energy range 180–209 GeV and 176  $e^+e^- \rightarrow q\bar{q}\gamma\gamma$  events in the energy range 130–209 GeV. These samples are used to constrain possible anomalous  $W^+W^-\gamma\gamma$  and  $ZZ\gamma\gamma$  quartic couplings. Further combining with the  $W^+W^-\gamma$  sample of ABBIENDI 04B the following one-parameter 95% CL limits are obtained:  $-0.007 < a_0^Z/\Lambda^2 < 0.023 \text{ GeV}^{-2}$ ,  $-0.029 < a_c^Z/\Lambda^2 < 0.029 \text{ GeV}^{-2}$ ,  $-0.020 < a_0^W/\Lambda^2 < 0.020 \text{ GeV}^{-2}$ ,  $-0.052 < a_c^W/\Lambda^2 < 0.037 \text{ GeV}^{-2}$ .

<sup>2</sup> In the CM energy range 183 to 209 GeV HEISTER 04A select 30  $e^+e^- \rightarrow \nu\bar{\nu}\gamma\gamma$  events with two acoplanar, high energy and high transverse momentum photons. The photon-photon acoplanarity is required to be  $> 5^\circ$ ,  $E_\gamma/\sqrt{s} > 0.025$  (the more energetic photon having energy  $> 0.2\sqrt{s}$ ),  $p_{T\gamma}/E_{\text{beam}} > 0.05$  and  $|\cos\theta_\gamma| < 0.94$ . A likelihood fit to the photon energy and recoil missing mass yields the following one-parameter 95%



CL limits:  $-0.012 < a_0^Z/\Lambda^2 < 0.019 \text{ GeV}^{-2}$ ,  $-0.041 < a_c^Z/\Lambda^2 < 0.044 \text{ GeV}^{-2}$ ,  
 $-0.060 < a_0^W/\Lambda^2 < 0.055 \text{ GeV}^{-2}$ ,  $-0.099 < a_c^W/\Lambda^2 < 0.093 \text{ GeV}^{-2}$ .

<sup>3</sup>ACHARD 02G study  $e^+e^- \rightarrow Z\gamma\gamma \rightarrow q\bar{q}\gamma\gamma$  events using data at center-of-mass energies from 200 to 209 GeV. The photons are required to be isolated, each with energy  $>5 \text{ GeV}$  and  $|\cos\theta| < 0.97$ , and the di-jet invariant mass to be compatible with that of the  $Z$  boson (74–111 GeV). Cuts on  $Z$  velocity ( $\beta < 0.73$ ) and on the energy of the most energetic photon reduce the backgrounds due to non-resonant production of the  $q\bar{q}\gamma\gamma$  state and due to ISR respectively, yielding a total of 40 candidate events of which 8.6 are expected to be due to background. The energy spectra of the least energetic photon are fitted for all ten center-of-mass energy values from 130 GeV to 209 GeV (as obtained adding to the present analysis 130–202 GeV data of ACCIARRI 01E, for a total of 137 events with an expected background of 34.1 events) to obtain the fitted values  $a_0/\Lambda^2 = 0.00_{-0.01}^{+0.02} \text{ GeV}^{-2}$  and  $a_c/\Lambda^2 = 0.03_{-0.02}^{+0.01} \text{ GeV}^{-2}$ , where the other parameter is kept fixed to its Standard Model value (0). A simultaneous fit to both parameters yields the 95% CL limits  $-0.02 \text{ GeV}^{-2} < a_0/\Lambda^2 < 0.03 \text{ GeV}^{-2}$  and  $-0.07 \text{ GeV}^{-2} < a_c/\Lambda^2 < 0.05 \text{ GeV}^{-2}$ .

## Z REFERENCES

AABOUD	23A	JHEP 2312 158 (errat.)	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AAD	23AQ	PR D108 032015	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	23BS	PL B847 138292	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	23CD	EPJ C83 781	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	23CH	JHEP 2312 107	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAIJ	23AM	CP C47 093002	R. Aaij <i>et al.</i>	(LHCb Collab.)
TUMASYAN	23E	PL B842 137563	A. Tumasyan <i>et al.</i>	(CMS Collab.)
AALTONEN	22	SCI 376 170	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AAD	21AO	NATP 17 819	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	21AQ	JHEP 2107 005	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	21AV	PRL 127 271801	G. Aad <i>et al.</i>	(ATLAS Collab.)
SIRUNYAN	21Q	EPJ C81 200	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
JANOT	20	PL B803 135319	P. Janot, S. Jadach	(CERN, CRAC)
VOUSINAS	20	PL B800 135068	G. Voutsinas <i>et al.</i>	(CERN, BOHR)
AABOUD	19AY	JHEP 1910 127	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD	19N	JHEP 1904 048	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
RAINBOLT	19	PR D99 013004	J.L. Rainbolt, M. Schmitt	(NWES)
SIRUNYAN	19AJ	EPJ C79 94	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	19BR	PL B797 134811	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
AABOUD	18AU	JHEP 1807 127	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
Also		JHEP 2312 158 (errat.)	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD	18BL	PL B786 134	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD	18CN	PR D98 092010	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD	18Q	PR D97 032005	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AAIJ	18AR	JHEP 1809 159	R. Aaij <i>et al.</i>	(LHCb Collab.)
ANDREEV	18A	EPJ C78 777	V. Andreev <i>et al.</i>	(H1 Collab.)
SIRUNYAN	18BT	EPJ C78 165	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	18DZ	PRL 121 141801	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
AABOUD	17Q	EPJ C77 367	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AABOUD	16K	PRL 117 111802	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AAD	16L	EPJ C76 210	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	16Q	PR D93 112002	G. Aad <i>et al.</i>	(ATLAS Collab.)
ABRAMOWICZ	16A	PR D93 092002	H. Abramowicz <i>et al.</i>	(ZEUS Collab.)
ABT	16	PR D94 052007	I. Abt <i>et al.</i>	(MPIM, OXF, HAMB, DESY)
KHACHATRYAN...	16AE	PL B760 448	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRYAN...	16CC	PL B763 280	V. Khachatryan <i>et al.</i>	(CMS Collab.)
AAD	15BT	JHEP 1509 049	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	15I	PRL 114 121801	G. Aad <i>et al.</i>	(ATLAS Collab.)
KHACHATRYAN...	15AC	JHEP 1504 164	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRYAN...	15B	PL B740 250	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRYAN...	15BC	EPJ C75 511	V. Khachatryan <i>et al.</i>	(CMS Collab.)
AAD	14AU	PR D90 072010	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	14N	PRL 112 231806	G. Aad <i>et al.</i>	(ATLAS Collab.)
AALTONEN	14E	PRL 112 111803	T. Aaltonen <i>et al.</i>	(CDF Collab.)
CHATRCHYAN	14AB	PR D89 092005	S. Chatrchyan <i>et al.</i>	(CMS Collab.)

AAD	13AN	PR D87 112003	G. Aad <i>et al.</i>	(ATLAS Collab.)
Also		PR D91 119901 (errat.)	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	13Z	JHEP 1303 128	G. Aad <i>et al.</i>	(ATLAS Collab.)
CHATRCHYAN	13B	JHEP 1301 063	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
CHATRCHYAN	13BI	JHEP 1310 164	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
SCHAEEL	13A	PRPL 532 119	S. Schael <i>et al.</i>	
AAD	12BX	PL B717 49	G. Aad <i>et al.</i>	(ATLAS Collab.)
ABAZOV	12S	PR D85 052001	V.M. Abazov <i>et al.</i>	(D0 Collab.)
CHATRCHYAN	12BN	JHEP 1212 034	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
AALTONEN	11S	PRL 107 051802	T. Aaltonen <i>et al.</i>	(CDF Collab.)
ABAZOV	11D	PR D84 012007	V.M. Abazov <i>et al.</i>	(D0 Collab.)
CHATRCHYAN	11M	PL B701 535	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
ABAZOV	09L	PRL 102 201802	V.M. Abazov <i>et al.</i>	(D0 Collab.)
BEDDALL	09	PL B670 300	A. Beddall, A. Beddall, A. Bingul	(UGAZ)
SCHAEEL	09	JHEP 0904 124	S. Schael <i>et al.</i>	(ALEPH Collab.)
ABAZOV	08K	PRL 100 131801	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	07M	PL B653 378	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABDALLAH	07C	EPJ C51 525	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
ABDALLAH	06E	PL B639 179	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
AKTAS	06	PL B632 35	A. Aktas <i>et al.</i>	(H1 Collab.)
LEP-SLC	06	PRPL 427 257	ALEPH, DELPHI, L3, OPAL, SLD and working groups	
SCHAEEL	06A	PL B639 192	S. Schael <i>et al.</i>	(ALEPH Collab.)
ABDALLAH	05	EPJ C40 1	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
ABDALLAH	05C	EPJ C44 299	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
ABE	05	PRL 94 091801	K. Abe <i>et al.</i>	(SLD Collab.)
ABE	05F	PR D71 112004	K. Abe <i>et al.</i>	(SLD Collab.)
ACOSTA	05M	PR D71 052002	D. Acosta <i>et al.</i>	(CDF Collab.)
ABBIENDI	04B	PL B580 17	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBIENDI	04C	EPJ C32 303	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBIENDI	04E	PL B586 167	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBIENDI	04G	EPJ C33 173	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBIENDI	04L	PR D70 032005	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABDALLAH	04F	EPJ C34 109	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
ABE	04C	PR D69 072003	K. Abe <i>et al.</i>	(SLD Collab.)
ACHARD	04C	PL B585 42	P. Achard <i>et al.</i>	(L3 Collab.)
ACHARD	04H	PL B597 119	P. Achard <i>et al.</i>	(L3 Collab.)
HEISTER	04A	PL B602 31	A. Heister <i>et al.</i>	(ALEPH Collab.)
ABBIENDI	03P	PL B577 18	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABDALLAH	03H	PL B569 129	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
ABDALLAH	03K	PL B576 29	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
ABE	03F	PRL 90 141804	K. Abe <i>et al.</i>	(SLD Collab.)
ACHARD	03D	PL B572 133	P. Achard <i>et al.</i>	(L3 Collab.)
ACHARD	03G	PL B577 109	P. Achard <i>et al.</i>	(L3 Collab.)
ABBIENDI	02I	PL B546 29	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABE	02G	PRL 88 151801	K. Abe <i>et al.</i>	(SLD Collab.)
ACHARD	02G	PL B540 43	P. Achard <i>et al.</i>	(L3 Collab.)
HEISTER	02B	PL B526 34	A. Heister <i>et al.</i>	(ALEPH Collab.)
HEISTER	02C	PL B528 19	A. Heister <i>et al.</i>	(ALEPH Collab.)
HEISTER	02H	EPJ C24 177	A. Heister <i>et al.</i>	(ALEPH Collab.)
ABBIENDI	01A	EPJ C19 587	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBIENDI	01G	EPJ C18 447	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBIENDI	01K	PL B516 1	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBIENDI	01N	EPJ C20 445	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBIENDI	01O	EPJ C21 1	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABE	01B	PRL 86 1162	K. Abe <i>et al.</i>	(SLD Collab.)
ABE	01C	PR D63 032005	K. Abe <i>et al.</i>	(SLD Collab.)
ACCIARRI	01E	PL B505 47	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACCIARRI	01I	PL B497 23	M. Acciarri <i>et al.</i>	(L3 Collab.)
HEISTER	01	EPJ C20 401	A. Heister <i>et al.</i>	(ALEPH Collab.)
HEISTER	01D	EPJ C22 201	A. Heister <i>et al.</i>	(ALEPH Collab.)
ABBIENDI	00N	PL B476 256	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBIENDI,G	00C	EPJ C17 553	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABE	00B	PRL 84 5945	K. Abe <i>et al.</i>	(SLD Collab.)
ABE	00D	PRL 85 5059	K. Abe <i>et al.</i>	(SLD Collab.)
ABREU	00	EPJ C12 225	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	00B	EPJ C14 613	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	00E	EPJ C14 585	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	00F	EPJ C16 371	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	00P	PL B475 429	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACCIARRI	00	EPJ C13 47	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACCIARRI	00C	EPJ C16 1	M. Acciarri <i>et al.</i>	(L3 Collab.)

ACCIARRI	00J	PL B479 79	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACCIARRI	00Q	PL B489 93	M. Acciarri <i>et al.</i>	(L3 Collab.)
BARATE	00B	EPJ C16 597	R. Barate <i>et al.</i>	(ALEPH Collab.)
BARATE	00C	EPJ C14 1	R. Barate <i>et al.</i>	(ALEPH Collab.)
BARATE	00O	EPJ C16 613	R. Barate <i>et al.</i>	(ALEPH Collab.)
ABBIENDI	99B	EPJ C8 217	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABBIENDI	99I	PL B447 157	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABE	99E	PR D59 052001	K. Abe <i>et al.</i>	(SLD Collab.)
ABE	99L	PRL 83 1902	K. Abe <i>et al.</i>	(SLD Collab.)
ABREU	99	EPJ C6 19	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	99B	EPJ C10 415	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	99J	PL B449 364	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	99U	PL B462 425	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	99Y	EPJ C10 219	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACCIARRI	99D	PL B448 152	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACCIARRI	99F	PL B453 94	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACCIARRI	99G	PL B450 281	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACCIARRI	99O	PL B465 363	M. Acciarri <i>et al.</i>	(L3 Collab.)
ABBOTT	98M	PR D57 3817	B. Abbott <i>et al.</i>	(D0 Collab.)
ABE	98D	PRL 80 660	K. Abe <i>et al.</i>	(SLD Collab.)
ABE	98I	PRL 81 942	K. Abe <i>et al.</i>	(SLD Collab.)
ABREU	98K	PL B423 194	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	98L	EPJ C5 585	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACCIARRI	98G	PL B431 199	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACCIARRI	98H	PL B429 387	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACCIARRI	98U	PL B439 225	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACKERSTAFF	98A	EPJ C5 411	K. Akerstaff <i>et al.</i>	(OPAL Collab.)
ACKERSTAFF	98E	EPJ C1 439	K. Akerstaff <i>et al.</i>	(OPAL Collab.)
ACKERSTAFF	98O	PL B420 157	K. Akerstaff <i>et al.</i>	(OPAL Collab.)
ACKERSTAFF	98Q	EPJ C4 19	K. Akerstaff <i>et al.</i>	(OPAL Collab.)
BARATE	98O	PL B434 415	R. Barate <i>et al.</i>	(ALEPH Collab.)
BARATE	98T	EPJ C4 557	R. Barate <i>et al.</i>	(ALEPH Collab.)
BARATE	98V	EPJ C5 205	R. Barate <i>et al.</i>	(ALEPH Collab.)
ABE	97	PRL 78 17	K. Abe <i>et al.</i>	(SLD Collab.)
ABREU	97C	ZPHY C73 243	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	97E	PL B398 207	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	97G	PL B404 194	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACCIARRI	97D	PL B393 465	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACCIARRI	97J	PL B407 351	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACCIARRI	97L	PL B407 389	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACCIARRI	97R	PL B413 167	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACKERSTAFF	97M	ZPHY C74 413	K. Akerstaff <i>et al.</i>	(OPAL Collab.)
ACKERSTAFF	97S	PL B412 210	K. Akerstaff <i>et al.</i>	(OPAL Collab.)
ACKERSTAFF	97T	ZPHY C76 387	K. Akerstaff <i>et al.</i>	(OPAL Collab.)
ACKERSTAFF	97W	ZPHY C76 425	K. Akerstaff <i>et al.</i>	(OPAL Collab.)
ALEXANDER	97C	ZPHY C73 379	G. Alexander <i>et al.</i>	(OPAL Collab.)
ALEXANDER	97D	ZPHY C73 569	G. Alexander <i>et al.</i>	(OPAL Collab.)
ALEXANDER	97E	ZPHY C73 587	G. Alexander <i>et al.</i>	(OPAL Collab.)
BARATE	97D	PL B405 191	R. Barate <i>et al.</i>	(ALEPH Collab.)
BARATE	97E	PL B401 150	R. Barate <i>et al.</i>	(ALEPH Collab.)
BARATE	97F	PL B401 163	R. Barate <i>et al.</i>	(ALEPH Collab.)
BARATE	97H	PL B402 213	R. Barate <i>et al.</i>	(ALEPH Collab.)
BARATE	97J	ZPHY C74 451	R. Barate <i>et al.</i>	(ALEPH Collab.)
ABREU	96R	ZPHY C72 31	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	96S	PL B389 405	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	96U	ZPHY C73 61	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACCIARRI	96	PL B371 126	M. Acciarri <i>et al.</i>	(L3 Collab.)
ADAM	96	ZPHY C69 561	W. Adam <i>et al.</i>	(DELPHI Collab.)
ADAM	96B	ZPHY C70 371	W. Adam <i>et al.</i>	(DELPHI Collab.)
ALEXANDER	96B	ZPHY C70 197	G. Alexander <i>et al.</i>	(OPAL Collab.)
ALEXANDER	96F	PL B370 185	G. Alexander <i>et al.</i>	(OPAL Collab.)
ALEXANDER	96N	PL B384 343	G. Alexander <i>et al.</i>	(OPAL Collab.)
ALEXANDER	96R	ZPHY C72 1	G. Alexander <i>et al.</i>	(OPAL Collab.)
BUSKULIC	96D	ZPHY C69 393	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
BUSKULIC	96H	ZPHY C69 379	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
BUSKULIC	96T	PL B384 449	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
BUSKULIC	96Y	PL B388 648	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
ABE	95J	PRL 74 2880	K. Abe <i>et al.</i>	(SLD Collab.)
ABREU	95	ZPHY C65 709 (errat.)	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	95D	ZPHY C66 323	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	95L	ZPHY C65 587	P. Abreu <i>et al.</i>	(DELPHI Collab.)

ABREU	95M	ZPHY C65 603	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	95O	ZPHY C67 543	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	95R	ZPHY C68 353	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	95V	ZPHY C68 541	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	95W	PL B361 207	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	95X	ZPHY C69 1	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACCIARRI	95B	PL B345 589	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACCIARRI	95C	PL B345 609	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACCIARRI	95G	PL B353 136	M. Acciarri <i>et al.</i>	(L3 Collab.)
AKERS	95C	ZPHY C65 47	R. Akers <i>et al.</i>	(OPAL Collab.)
AKERS	95U	ZPHY C67 389	R. Akers <i>et al.</i>	(OPAL Collab.)
AKERS	95W	ZPHY C67 555	R. Akers <i>et al.</i>	(OPAL Collab.)
AKERS	95X	ZPHY C68 1	R. Akers <i>et al.</i>	(OPAL Collab.)
AKERS	95Z	ZPHY C68 203	R. Akers <i>et al.</i>	(OPAL Collab.)
ALEXANDER	95D	PL B358 162	G. Alexander <i>et al.</i>	(OPAL Collab.)
BUSKULIC	95R	ZPHY C69 15	D. Buskalic <i>et al.</i>	(ALEPH Collab.)
MIYABAYASHI	95	PL B347 171	K. Miyabayashi <i>et al.</i>	(TOPAZ Collab.)
ABE	94C	PRL 73 25	K. Abe <i>et al.</i>	(SLD Collab.)
ABREU	94B	PL B327 386	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	94P	PL B341 109	P. Abreu <i>et al.</i>	(DELPHI Collab.)
AKERS	94P	ZPHY C63 181	R. Akers <i>et al.</i>	(OPAL Collab.)
BUSKULIC	94G	ZPHY C62 179	D. Buskalic <i>et al.</i>	(ALEPH Collab.)
BUSKULIC	94J	ZPHY C62 1	D. Buskalic <i>et al.</i>	(ALEPH Collab.)
VILAIN	94	PL B320 203	P. Vilain <i>et al.</i>	(CHARM II Collab.)
ABREU	93	PL B298 236	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	93I	ZPHY C59 533	P. Abreu <i>et al.</i>	(DELPHI Collab.)
Also		ZPHY C65 709 (errat.)	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	93L	PL B318 249	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACTON	93	PL B305 407	P.D. Acton <i>et al.</i>	(OPAL Collab.)
ACTON	93D	ZPHY C58 219	P.D. Acton <i>et al.</i>	(OPAL Collab.)
ACTON	93E	PL B311 391	P.D. Acton <i>et al.</i>	(OPAL Collab.)
ADRIANI	93	PL B301 136	O. Adriani <i>et al.</i>	(L3 Collab.)
ADRIANI	93I	PL B316 427	O. Adriani <i>et al.</i>	(L3 Collab.)
BUSKULIC	93L	PL B313 520	D. Buskalic <i>et al.</i>	(ALEPH Collab.)
NOVIKOV	93C	PL B298 453	V.A. Novikov, L.B. Okun, M.I. Vysotsky	(ITEP)
ABREU	92I	PL B277 371	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	92M	PL B289 199	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACTON	92B	ZPHY C53 539	D.P. Acton <i>et al.</i>	(OPAL Collab.)
ACTON	92L	PL B294 436	P.D. Acton <i>et al.</i>	(OPAL Collab.)
ACTON	92N	PL B295 357	P.D. Acton <i>et al.</i>	(OPAL Collab.)
ADEVA	92	PL B275 209	B. Adeva <i>et al.</i>	(L3 Collab.)
ADRIANI	92D	PL B292 454	O. Adriani <i>et al.</i>	(L3 Collab.)
ALITTI	92B	PL B276 354	J. Alitti <i>et al.</i>	(UA2 Collab.)
BUSKULIC	92D	PL B292 210	D. Buskalic <i>et al.</i>	(ALEPH Collab.)
BUSKULIC	92E	PL B294 145	D. Buskalic <i>et al.</i>	(ALEPH Collab.)
DECAMP	92	PRPL 216 253	D. Decamp <i>et al.</i>	(ALEPH Collab.)
ABE	91E	PRL 67 1502	F. Abe <i>et al.</i>	(CDF Collab.)
ABREU	91H	ZPHY C50 185	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACTON	91B	PL B273 338	D.P. Acton <i>et al.</i>	(OPAL Collab.)
ADACHI	91	PL B255 613	I. Adachi <i>et al.</i>	(TOPAZ Collab.)
ADEVA	91I	PL B259 199	B. Adeva <i>et al.</i>	(L3 Collab.)
AKRAWY	91F	PL B257 531	M.Z. Akrawy <i>et al.</i>	(OPAL Collab.)
DECAMP	91B	PL B259 377	D. Decamp <i>et al.</i>	(ALEPH Collab.)
DECAMP	91J	PL B266 218	D. Decamp <i>et al.</i>	(ALEPH Collab.)
JACOBSEN	91	PRL 67 3347	R.G. Jacobsen <i>et al.</i>	(Mark II Collab.)
SHIMONAKA	91	PL B268 457	A. Shimonaka <i>et al.</i>	(TOPAZ Collab.)
ABE	90I	ZPHY C48 13	K. Abe <i>et al.</i>	(VENUS Collab.)
ABRAMS	90	PRL 64 1334	G.S. Abrams <i>et al.</i>	(Mark II Collab.)
AKRAWY	90J	PL B246 285	M.Z. Akrawy <i>et al.</i>	(OPAL Collab.)
BEHREND	90D	ZPHY C47 333	H.J. Behrend <i>et al.</i>	(CELLO Collab.)
BRAUNSCH...	90	ZPHY C48 433	W. Braunschweig <i>et al.</i>	(TASSO Collab.)
ELSEN	90	ZPHY C46 349	E. Elsen <i>et al.</i>	(JADE Collab.)
HEGNER	90	ZPHY C46 547	S. Hegner <i>et al.</i>	(JADE Collab.)
STUART	90	PRL 64 983	D. Stuart <i>et al.</i>	(AMY Collab.)
ABE	89	PRL 62 613	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	89C	PRL 63 720	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	89L	PL B232 425	K. Abe <i>et al.</i>	(VENUS Collab.)
ABRAMS	89B	PRL 63 2173	G.S. Abrams <i>et al.</i>	(Mark II Collab.)
ABRAMS	89D	PRL 63 2780	G.S. Abrams <i>et al.</i>	(Mark II Collab.)
ALBAJAR	89	ZPHY C44 15	C. Albajar <i>et al.</i>	(UA1 Collab.)
BACALA	89	PL B218 112	A. Bacala <i>et al.</i>	(AMY Collab.)

BAND	89	PL B218 369	H.R. Band <i>et al.</i>	(MAC Collab.)
GREENSHAW	89	ZPHY C42 1	T. Greenshaw <i>et al.</i>	(JADE Collab.)
OULD-SAAD	89	ZPHY C44 567	F. Ould-Saada <i>et al.</i>	(JADE Collab.)
SAGAWA	89	PRL 63 2341	H. Sagawa <i>et al.</i>	(AMY Collab.)
ADACHI	88C	PL B208 319	I. Adachi <i>et al.</i>	(TOPAZ Collab.)
ADEVA	88	PR D38 2665	B. Adeva <i>et al.</i>	(Mark-J Collab.)
BRAUNSCH...	88D	ZPHY C40 163	W. Braunschweig <i>et al.</i>	(TASSO Collab.)
ANSARI	87	PL B186 440	R. Ansari <i>et al.</i>	(UA2 Collab.)
BEHREND	87C	PL B191 209	H.J. Behrend <i>et al.</i>	(CELLO Collab.)
BARTEL	86C	ZPHY C30 371	W. Bartel <i>et al.</i>	(JADE Collab.)
Also		ZPHY C26 507	W. Bartel <i>et al.</i>	(JADE Collab.)
Also		PL 108B 140	W. Bartel <i>et al.</i>	(JADE Collab.)
ASH	85	PRL 55 1831	W.W. Ash <i>et al.</i>	(MAC Collab.)
BARTEL	85F	PL 161B 188	W. Bartel <i>et al.</i>	(JADE Collab.)
DERRICK	85	PR D31 2352	M. Derrick <i>et al.</i>	(HRS Collab.)
FERNANDEZ	85A	PRL 54 1620	E. Fernandez <i>et al.</i>	(MAC Collab.)
LEVI	83	PRL 51 1941	M.E. Levi <i>et al.</i>	(Mark II Collab.)
BEHREND	82	PL 114B 282	H.J. Behrend <i>et al.</i>	(CELLO Collab.)
BRANDELIK	82C	PL 110B 173	R. Brandelik <i>et al.</i>	(TASSO Collab.)

---