

GAUGE AND HIGGS BOSONS

γ (photon)

$$I(J^{PC}) = 0,1(1^{--})$$

Mass $m < 1 \times 10^{-18}$ eV

Charge $q < 1 \times 10^{-46}$ e (mixed charge)

Charge $q < 1 \times 10^{-35}$ e (single charge)

Mean life $\tau =$ Stable

g (gluon)

$$I(J^P) = 0(1^-)$$

Mass $m = 0$ [a]

SU(3) color octet

graviton

$$J = 2$$

Mass $m < 1.76 \times 10^{-23}$ eV

W

$$J = 1$$

Charge = ± 1 e

Mass $m = 80.3692 \pm 0.0133$ GeV [b]

W/Z mass ratio = 0.88136 ± 0.00015

$m_Z - m_W = 10.818 \pm 0.013$ GeV

$m_{W^+} - m_{W^-} = -0.029 \pm 0.028$ GeV

Full width $\Gamma = 2.085 \pm 0.042$ GeV

$\langle N_{\pi^\pm} \rangle = 15.70 \pm 0.35$

$\langle N_{K^\pm} \rangle = 2.20 \pm 0.19$

$\langle N_p \rangle = 0.92 \pm 0.14$

$\langle N_{\text{charged}} \rangle = 19.39 \pm 0.08$

W^- modes are charge conjugates of the modes below.

W^+ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
$\ell^+ \nu$	[c] $(10.86 \pm 0.09) \%$		–
$e^+ \nu$	$(10.71 \pm 0.16) \%$		40185
$\mu^+ \nu$	$(10.63 \pm 0.15) \%$		40185
$\tau^+ \nu$	$(11.38 \pm 0.21) \%$		40165
hadrons	$(67.41 \pm 0.27) \%$		–
$\pi^+ \gamma$	< 7	$\times 10^{-6}$	95% 40184
$D_s^+ \gamma$	< 6	$\times 10^{-4}$	95% 40160
cX	$(33.3 \pm 2.6) \%$		–
$c\bar{s}$	$(31^{+13}_{-11}) \%$		–
invisible	[d] $(1.4 \pm 2.9) \%$		–
$\pi^+ \pi^+ \pi^-$	< 1.01	$\times 10^{-6}$	95% 40184



$$J = 1$$

Charge = 0

Mass $m = 91.1880 \pm 0.0020$ GeV [e]

Full width $\Gamma = 2.4955 \pm 0.0023$ GeV

$\Gamma(\ell^+ \ell^-) = 83.984 \pm 0.086$ MeV [c]

$\Gamma(\text{invisible}) = 499.2 \pm 1.5$ MeV [f]

$\Gamma(\text{hadrons}) = 1744.4 \pm 2.0$ MeV

$\Gamma(\mu^+ \mu^-)/\Gamma(e^+ e^-) = 1.0001 \pm 0.0024$

$\Gamma(\tau^+ \tau^-)/\Gamma(e^+ e^-) = 1.0020 \pm 0.0032$ [g]

Average charged multiplicity

$$\langle N_{\text{charged}} \rangle = 20.76 \pm 0.16 \quad (S = 2.1)$$

Couplings to quarks and leptons

$$g_V^\ell = -0.03783 \pm 0.00041$$

$$g_V^u = 0.266 \pm 0.034$$

$$g_V^d = -0.38^{+0.04}_{-0.05}$$

$$g_A^\ell = -0.50123 \pm 0.00026$$

$$g_A^u = 0.519^{+0.028}_{-0.033}$$

$$g_A^d = -0.527^{+0.040}_{-0.028}$$

$$g^{\nu\ell} = 0.5008 \pm 0.0008$$

$$g^{\nu e} = 0.53 \pm 0.09$$

$$g^{\nu\mu} = 0.502 \pm 0.017$$

Asymmetry parameters [h]

$$\begin{aligned}
 A_e &= 0.1515 \pm 0.0019 \\
 A_\mu &= 0.142 \pm 0.015 \\
 A_\tau &= 0.143 \pm 0.004 \\
 A_s &= 0.90 \pm 0.09 \\
 A_c &= 0.670 \pm 0.027 \\
 A_b &= 0.923 \pm 0.020
 \end{aligned}$$

Charge asymmetry (%) at Z pole

$$\begin{aligned}
 A_{FB}^{(0\ell)} &= 1.71 \pm 0.10 \\
 A_{FB}^{(0u)} &= 4 \pm 7 \\
 A_{FB}^{(0s)} &= 9.8 \pm 1.1 \\
 A_{FB}^{(0c)} &= 7.07 \pm 0.35 \\
 A_{FB}^{(0b)} &= 9.92 \pm 0.16
 \end{aligned}$$

Z DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
$e^+ e^-$	(3.3632±0.0042) %		45594
$\mu^+ \mu^-$	(3.3662±0.0066) %		45594
$\tau^+ \tau^-$	(3.3696±0.0083) %		45559
$\ell^+ \ell^-$	[c] (3.3658±0.0023) %		—
$\ell^+ \ell^- \ell^+ \ell^-$	[j] (4.55 ±0.17) × 10 ⁻⁶		45594
invisible	(20.000 ±0.055) %		—
hadrons	(69.911 ±0.056) %		—
($u\bar{u} + c\bar{c}$)/2	(11.6 ±0.6) %		—
($d\bar{d} + s\bar{s} + b\bar{b}$)/3	(15.6 ±0.4) %		—
$c\bar{c}$	(12.03 ±0.21) %		—
$b\bar{b}$	(15.12 ±0.05) %		—
$b\bar{b}b\bar{b}$	(3.6 ±1.3) × 10 ⁻⁴		—
ggg	< 1.1	% CL=95%	—
$\pi^0 \gamma$	< 2.01	× 10 ⁻⁵ CL=95%	45594
$\eta \gamma$	< 5.1	× 10 ⁻⁵ CL=95%	45592
$\rho^0 \gamma$	< 4.0	× 10 ⁻⁶ CL=95%	45591
$\omega \gamma$	< 3.9	× 10 ⁻⁶ CL=95%	45591
$\eta'(958) \gamma$	< 4.2	× 10 ⁻⁵ CL=95%	45589
$\phi \gamma$	< 7	× 10 ⁻⁷ CL=95%	45588
$\gamma \gamma$	< 1.46	× 10 ⁻⁵ CL=95%	45594
$\pi^0 \pi^0$	< 1.52	× 10 ⁻⁵ CL=95%	45594
$\gamma \gamma \gamma$	< 2.2	× 10 ⁻⁶ CL=95%	45594
$\pi^\pm W^\mp$	[j] < 7	× 10 ⁻⁵ CL=95%	10176
$\rho^\pm W^\mp$	[j] < 8.3	× 10 ⁻⁵ CL=95%	10151

$J/\psi(1S)X$		$(3.51 \begin{smallmatrix} +0.23 \\ -0.25 \end{smallmatrix}) \times 10^{-3}$	$S=1.1$	—
$J/\psi(1S)\gamma$		< 1.2	$\times 10^{-6}$	CL=95% 45541
$\psi(2S)X$		$(1.60 \pm 0.29) \times 10^{-3}$		—
$\psi(2S)\gamma$		< 2.4	$\times 10^{-6}$	CL=95% 45519
$J/\psi(1S)J/\psi(1S)$		< 2.2	$\times 10^{-6}$	CL=95% 45489
$\chi_{c1}(1P)X$		$(2.9 \pm 0.7) \times 10^{-3}$		—
$\chi_{c2}(1P)X$		< 3.2	$\times 10^{-3}$	CL=90% —
$\Upsilon(1S) X + \Upsilon(2S) X$ $+ \Upsilon(3S) X$		$(1.0 \pm 0.5) \times 10^{-4}$		—
$\Upsilon(1S)X$		< 4.4	$\times 10^{-5}$	CL=95% —
$\Upsilon(1S)\gamma$		< 1.1	$\times 10^{-6}$	CL=95% 45103
$\Upsilon(2S)X$		< 1.39	$\times 10^{-4}$	CL=95% —
$\Upsilon(2S)\gamma$		< 1.3	$\times 10^{-6}$	CL=95% 45043
$\Upsilon(3S)X$		< 9.4	$\times 10^{-5}$	CL=95% —
$\Upsilon(3S)\gamma$		< 2.4	$\times 10^{-6}$	CL=95% 45006
$\Upsilon(1, 2, 3S) \Upsilon(1, 2, 3S)$		< 1.5	$\times 10^{-6}$	CL=95% —
$D^0\gamma$		< 2.2	$\times 10^{-3}$	CL=95% 45575
$(D^0/\bar{D}^0) X$		$(20.7 \pm 2.0) \%$		—
$D^\pm X$		$(12.2 \pm 1.7) \%$		—
$D^*(2010)^\pm X$	[j]	$(11.4 \pm 1.3) \%$		—
$D_{s1}(2536)^\pm X$		$(3.6 \pm 0.8) \times 10^{-3}$		—
$D_{sJ}(2573)^\pm X$		$(5.8 \pm 2.2) \times 10^{-3}$		—
$D^{*'}(2629)^\pm X$		searched for		—
$B^+ X$	[k]	$(6.08 \pm 0.13) \%$		—
$B_s^0 X$	[k]	$(1.59 \pm 0.13) \%$		—
$B_c^+ X$		searched for		—
$\Lambda_c^+ X$		$(1.54 \pm 0.33) \%$		—
$\Xi_c^0 X$		seen		—
$\Xi_b X$		seen		—
b -baryon X	[k]	$(1.38 \pm 0.22) \%$		—
anomalous γ + hadrons	[l]	< 3.2	$\times 10^{-3}$	CL=95% —
$e^+ e^- \gamma$	[l]	< 5.2	$\times 10^{-4}$	CL=95% 45594
$\mu^+ \mu^- \gamma$	[l]	< 5.6	$\times 10^{-4}$	CL=95% 45594
$\tau^+ \tau^- \gamma$	[l]	< 7.3	$\times 10^{-4}$	CL=95% 45559
$\ell^+ \ell^- \gamma \gamma$	[n]	< 6.8	$\times 10^{-6}$	CL=95% —
$q\bar{q}\gamma\gamma$	[n]	< 5.5	$\times 10^{-6}$	CL=95% —
$\nu\bar{\nu}\gamma\gamma$	[n]	< 3.1	$\times 10^{-6}$	CL=95% 45594
$e^\pm \mu^\mp$	LF	[j]	< 2.62	$\times 10^{-7}$ CL=95% 45594
$e^\pm \tau^\mp$	LF	[j]	< 5.0	$\times 10^{-6}$ CL=95% 45577
$\mu^\pm \tau^\mp$	LF	[j]	< 6.5	$\times 10^{-6}$ CL=95% 45577
pe	L,B	< 1.8	$\times 10^{-6}$	CL=95% 45589
$p\mu$	L,B	< 1.8	$\times 10^{-6}$	CL=95% 45589



was H^0

$$J = 0$$

$$\text{Mass } m = 125.20 \pm 0.11 \text{ GeV} \quad (S = 1.4)$$

$$\text{Full width } \Gamma = 3.7_{-1.4}^{+1.9} \text{ MeV} \quad (\text{assumes equal on-shell and off-shell effective couplings})$$

H Signal Strengths in Different Channels

$$\text{Combined Final States} = 1.03 \pm 0.04$$

$$W W^* = 1.00 \pm 0.08$$

$$Z Z^* = 1.02 \pm 0.08$$

$$\gamma\gamma = 1.10 \pm 0.06$$

$$c\bar{c} \text{ Final State} < 14, \text{ CL} = 95\%$$

$$b\bar{b} = 0.99 \pm 0.12$$

$$\mu^+\mu^- = 1.21 \pm 0.35$$

$$\tau^+\tau^- = 0.91 \pm 0.09$$

$$Z\gamma = 2.2 \pm 0.7$$

$$\gamma^*\gamma \text{ Final State} = 1.5 \pm 0.5$$

$$\text{Fermion coupling } (\kappa_F) = 0.94 \pm 0.05$$

$$\text{Gauge boson coupling } (\kappa_V) = 1.023 \pm 0.026$$

$$t\bar{t}H \text{ Production} = 1.10 \pm 0.18$$

$$HH \text{ Production Cross Section in } pp \text{ Collisions} < 2.4, \text{ CL} = 95\%$$

$$tH \text{ production} = 6 \pm 4$$

$$H \text{ Production Cross Section in } pp \text{ Collisions at } \sqrt{s} = 13 \text{ TeV} = 56.8 \pm 3.4 \text{ pb}$$

H DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
$W W^*$	(25.7 \pm 2.5) %		—
$Z Z^*$	(2.80 \pm 0.30) %		—
$\gamma\gamma$	(2.50 \pm 0.20) $\times 10^{-3}$		62600
$b\bar{b}$	(53 \pm 8) %		—
$e^+ e^-$	< 3.0 $\times 10^{-4}$	95%	62600
$\mu^+ \mu^-$	(2.6 \pm 1.3) $\times 10^{-4}$		62600
$\tau^+ \tau^-$	(6.0 $^{+0.8}_{-0.7}$) %		62575
$Z\gamma$	(3.4 \pm 1.1) $\times 10^{-3}$		29392
$Z\rho(770)$	< 1.21 %	95%	29384
$Z\phi(1020)$	< 3.6 $\times 10^{-3}$	95%	29378
ZJ/ψ	< 1.9 $\times 10^{-3}$	95%	29267
$Z\psi(2S)$	< 6.6 $\times 10^{-3}$	95%	29214
$J/\psi\gamma$	< 2.0 $\times 10^{-4}$	95%	62561
$J/\psi J/\psi$	< 3.8 $\times 10^{-4}$	95%	62523
$\psi(2S)\gamma$	< 1.05 $\times 10^{-3}$	95%	62546
$\psi(2S)J/\psi$	< 2.1 $\times 10^{-3}$	95%	62507
$\psi(2S)\psi(2S)$	< 3.0 $\times 10^{-3}$	95%	62491
$\Upsilon(1S)\gamma$	< 2.5 $\times 10^{-4}$	95%	62242
$\Upsilon(1S)\Upsilon(1S)$	< 1.7 $\times 10^{-3}$	95%	61881
$\Upsilon(2S)\gamma$	< 4.2 $\times 10^{-4}$	95%	62199
$\Upsilon(3S)\gamma$	< 3.4 $\times 10^{-4}$	95%	62172
$\Upsilon(nS)\Upsilon(mS)$	< 3.5 $\times 10^{-4}$	95%	—
$\rho(770)\gamma$	< 1.04 $\times 10^{-3}$	95%	62597
$\omega(782)\gamma$	< 5.5 $\times 10^{-4}$	95%	62597
$K^*(892)\gamma$	< 2.2 $\times 10^{-4}$	95%	62597
$\phi(1020)\gamma$	< 5 $\times 10^{-4}$	95%	62596
$e\mu$	<i>LF</i> < 4.4 $\times 10^{-5}$	95%	62600
$e\tau$	<i>LF</i> < 2.0 $\times 10^{-3}$	95%	62587
$\mu\tau$	<i>LF</i> < 1.5 $\times 10^{-3}$	95%	62587
invisible	< 10.7 %	95%	—
γ invisible	< 2.9 %	95%	—

Neutral Higgs Bosons, Searches for

Mass limits for heavy neutral Higgs bosons (H_2^0, A^0) in the MSSM

- $m > 1121$ GeV, CL = 95% ($\tan\beta = 10$)
- $m > 1475$ GeV, CL = 95% ($\tan\beta = 20$)
- $m > 1677$ GeV, CL = 95% ($\tan\beta = 30$)
- $m > 1826$ GeV, CL = 95% ($\tan\beta = 40$)
- $m > 1950$ GeV, CL = 95% ($\tan\beta = 50$)

$$m > 2062 \text{ GeV, CL} = 95\% \quad (\tan\beta = 60)$$

Charged Higgs Bosons (H^\pm and $H^{\pm\pm}$), Searches for

Mass limits for $m_{H^\pm} < m(\text{top})$ in the MSSM

$$m > 155 \text{ GeV, CL} = 95\%$$

Mass limits for $m_{H^\pm} > m(\text{top})$ in the MSSM

$$m > 181 \text{ GeV, CL} = 95\% \quad (\tan\beta = 10)$$

$$m > 249 \text{ GeV, CL} = 95\% \quad (\tan\beta = 20)$$

$$m > 390 \text{ GeV, CL} = 95\% \quad (\tan\beta = 30)$$

$$m > 894 \text{ GeV, CL} = 95\% \quad (\tan\beta = 40)$$

$$m > 1017 \text{ GeV, CL} = 95\% \quad (\tan\beta = 50)$$

$$m > 1103 \text{ GeV, CL} = 95\% \quad (\tan\beta = 60)$$

New Heavy Bosons (W' , Z' , leptoquarks, etc.), Searches for

Additional W Bosons

W' with standard couplings

$$\text{Mass } m > 6000 \text{ GeV, CL} = 95\% \quad (pp \text{ direct search})$$

W_R (Right-handed W Boson)

$$\text{Mass } m > 715 \text{ GeV, CL} = 90\% \quad (\text{electroweak fit})$$

Additional Z Bosons

Z'_{SM} with standard couplings

$$\text{Mass } m > 5150 \text{ GeV, CL} = 95\% \quad (pp \text{ direct search})$$

Z_{LR} of $SU(2)_L \times SU(2)_R \times U(1)$ (with $g_L = g_R$)

$$\text{Mass } m > 630 \text{ GeV, CL} = 95\% \quad (p\bar{p} \text{ direct search})$$

$$\text{Mass } m > 1162 \text{ GeV, CL} = 95\% \quad (\text{electroweak fit})$$

Z_χ of $SO(10) \rightarrow SU(5) \times U(1)_\chi$ (with $g_\chi = e/\cos\theta_W$)

$$\text{Mass } m > 4800 \text{ GeV, CL} = 95\% \quad (pp \text{ direct search})$$

Z_ψ of $E_6 \rightarrow SO(10) \times U(1)_\psi$ (with $g_\psi = e/\cos\theta_W$)

$$\text{Mass } m > 4560 \text{ GeV, CL} = 95\% \quad (pp \text{ direct search})$$

Z_η of $E_6 \rightarrow SU(3) \times SU(2) \times U(1) \times U(1)_\eta$ (with $g_\eta = e/\cos\theta_W$)

$$\text{Mass } m > 3.900 \times 10^3 \text{ GeV, CL} = 95\% \quad (pp \text{ direct search})$$

Scalar Leptoquarks

- $m > 1800$ GeV, CL = 95% (1st gen., pair prod., $B(eq)=1$)
- $m > 1755$ GeV, CL = 95% (1st gen., single prod., $B(eq)=1$)
- $m > 1700$ GeV, CL = 95% (2nd gen., pair prod., $B(\mu q)=1$)
- $m > 660$ GeV, CL = 95% (2nd gen., single prod., $B(\mu q)=1$)
- $m > 1460$ GeV, CL = 95% (3rd gen., pair prod., $B(\tau b)=1$)
- $m > 1280$ GeV, CL = 95% (3rd gen., single prod., $B(\tau b)=1$)

(See the Particle Listings for assumptions on leptoquark quantum numbers and branching fractions.)

Diquarks

- Mass $m > 7200$ GeV, CL = 95% (E_6 diquark)

Axigluon

- Mass $m > 6600$ GeV, CL = 95%

Axions (A^0) and Other Very Light Bosons, Searches for

See the review on "Axions and other similar particles."

The best limit for the half-life of neutrinoless double beta decay with Majoron emission is $> 7.2 \times 10^{24}$ years (CL = 90%).

NOTES

- [a] Theoretical value. A mass as large as a few MeV may not be precluded.
- [b] This value does not include the AALTONEN 22 measurement by CDF. See the W mass section in the listings for details.
- [c] ℓ indicates each type of lepton (e , μ , and τ), not sum over them.
- [d] This represents the width for the decay of the W boson into a charged particle with momentum below detectability, $p < 200$ MeV.
- [e] The Z -boson mass listed here corresponds to a Breit-Wigner resonance parameter. It lies approximately 34 MeV above the real part of the position of the pole (in the energy-squared plane) in the Z -boson propagator.
- [f] This partial width takes into account Z decays into $\nu\bar{\nu}$ and any other possible undetected modes.
- [g] This ratio has not been corrected for the τ mass.
- [h] Here $A \equiv 2g_V g_A / (g_V^2 + g_A^2)$.
- [i] Here ℓ indicates e or μ .
- [j] The value is for the sum of the charge states or particle/antiparticle states indicated.
- [k] This value is updated using the product of (i) the $Z \rightarrow b\bar{b}$ fraction from this listing and (ii) the b -hadron fraction in an unbiased sample of weakly decaying b -hadrons produced in Z -decays provided by the Heavy Flavor Averaging Group (HFLAV, <http://www.slac.stanford.edu/xorg/hflav/osc/PDG.2009/#FRACZ>).
- [l] See the Z Particle Listings for the γ energy range used in this measurement.
- [n] For $m_{\gamma\gamma} = (60 \pm 5)$ GeV.