## **QUARKS**

The u-, d-, and s-quark masses are the  $\overline{\rm MS}$  masses at the scale  $\mu$  = 2 GeV. The c- and b-quark masses are the  $\overline{\rm MS}$  masses renormalized at the  $\overline{\rm MS}$  mass, i.e.  $\overline{m} = \overline{m}(\mu = \overline{m})$ . The t-quark mass is extracted from event kinematics (see the review "The Top Quark").

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

 $m_u=2.16\pm0.07$  MeV, CL =90% Charge  $=\frac{2}{3}$  e  $I_z=+\frac{1}{2}$   $m_u/m_d=0.462\pm0.020$ , CL =90%

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

 $m_d=4.70\pm0.07$  MeV, CL =90% Charge  $=-\frac{1}{3}$  e  $I_z=-\frac{1}{2}$   $m_s/m_d=17$ –22  $\overline{m}=(m_u+m_d)/2=3.49\pm0.07$  MeV, CL =90%

$$I(J^P)=0(\tfrac{1}{2}^+)$$

 $m_s = 93.5 \pm 0.8$  MeV, CL = 90% Charge =  $-\frac{1}{3}$  e Strangeness = -1  $m_s$  /  $((m_u+m_d)/2)=27.33^{+0.18}_{-0.14}$ , CL = 90%

$$I(J^P) = 0(\frac{1}{2}^+)$$

 $m_c=1.2730\pm0.0046$  GeV, CL =90% Charge  $=\frac{2}{3}$  e Charm =+1  $m_b-m_c=3.45\pm0.05$  GeV

$$I(J^P)=0(\tfrac{1}{2}^+)$$

 $m_b=4.183\pm 0.007$  GeV, CL =90% Charge  $=-\frac{1}{3}$  e Bottom =-1

$$I(J^P)=0(\tfrac{1}{2}^+)$$

$$\mathsf{Charge} = \tfrac{2}{3} \ e \qquad \qquad \mathsf{Top} = +1$$

Created: 4/29/2024 19:17

Mass (direct measurements)  $m=172.57\pm0.29~{\rm GeV}^{[a,b]}~({\rm S}=1.5)$  Mass (from cross-section measurements)  $m=162.5^{+2.1}_{-1.5}~{\rm GeV}^{[a]}$  Mass (Pole from cross-section measurements)  $m=172.4\pm0.7~{\rm GeV}$   $m_t-m_{\overline{t}}=-0.15\pm0.20~{\rm GeV}~({\rm S}=1.1)$  Full width  $\Gamma=1.42^{+0.19}_{-0.15}~{\rm GeV}~({\rm S}=1.4)$   $\Gamma(W\,b)/\Gamma(W\,q\,(q=b,\,s,\,d))=0.957\pm0.034~({\rm S}=1.5)$ 

#### t-quark EW Couplings

 $F_0 = 0.693 \pm 0.013$   $F_- = 0.315 \pm 0.010$   $F_+ = -0.005 \pm 0.007$  $F_{V+A} < 0.29$ , CL = 95%

t DECAY MODES	Fraction $(\Gamma_i/\Gamma)$	Confidence level	<i>p</i> (MeV/ <i>c</i> )			
$\overline{Wq(q=b, s, d)}$			_			
W b			_			
$e \nu_e b$	(11.10±0.30) % -					
$\mu u_{\mu}$ b	(11.40±0.20) % -					
$ au u_{ au}$ $m{b}$	$(10.7 \pm 0.5)\%$					
q <del>q</del> b	$(66.5 \pm 1.4)$	%	_			
$\gamma q(q=u,c)$	[c] < 4.5	$\times 10^{-5}$ 95%	_			
$\Delta T = 1$ we	ak neutral current ( <i>T</i> .	1) modes				
Zq(q=u,c)	[d] < 1.2	× 10 <sup>-4</sup> 95%	_			
Hu 7	7 < 1.9	$\times10^{-4}$ 95%	_			
Hc 7	7 < 4.3	$\times10^{-4}$ 95%	_			
$\ell^+ \overline{q}  \overline{q}'(q=d,s,b; q'=u,c)$ 7	71 < 1.6	$\times$ 10 <sup>-3</sup> 95%	_			
Lepton Fami	ly number ( <i>LF</i> ) violat	ing modes				

### Lepton Family number (LF) violating modes

	LF	< 8.9	$\times 10^{-7}$	_
$e^{\pm}\mu^{\mp}u$	LF	< 7	× 10 <sup>-8</sup>	_

# b' (4th Generation) Quark, Searches for

Mass m > 190 GeV, CL = 95% ( $p\overline{p}$ , quasi-stable b') Mass m > 1390 GeV, CL = 95% (B( $b' \to Zb$ ) = 1) Mass m > 1350 GeV, CL = 95% (B( $b' \to Wt$ ) = 1) Mass m > 1570 GeV, CL = 95% (B( $b' \to Hb$ ) = 1) Mass m > 46.0 GeV, CL = 95% ( $e^+e^-$ , all decays)

Created: 4/29/2024 19:17

# t' (4<sup>th</sup> Generation) Quark, Searches for

```
m(t'(2/3)) > 1280 GeV, CL = 95% (B(t' \to Zt) = 1) m(t'(2/3)) > 1295 GeV, CL = 95% (B(t' \to Wb) = 1) m(t'(2/3)) > 1310 GeV, CL = 95% (singlet t') m(t'(2/3)) > 1350 GeV, CL = 95% (t' in a weak isospin doublet (t',b')) m(t'(5/3)) > 1.460 \times 10^3 GeV, CL = 95% (t'(5/3) \to tW^+)
```

## Free Quark Searches

All searches since 1977 have had negative results.

#### **NOTES**

- [a] A discussion of the definition of the top quark mass in these measurements can be found in the review "The Top Quark."
- [b] Based on published top mass measurements using data from Tevatron Run-I and Run-II and LHC at  $\sqrt{s}=7$  TeV. Including the most recent unpublished results from Tevatron Run-II, the Tevatron Electroweak Working Group reports a top mass of  $173.2\pm0.9$  GeV. See the note "The Top Quark' in the Quark Particle Listings of this *Review*.
- [c] This limit is for  $\Gamma(t \to \gamma q)/\Gamma(t \to W b)$ .
- [d] This limit is for  $\Gamma(t \to Zq)/\Gamma(t \to Wb)$ .

Created: 4/29/2024 19:17